



THE AGA KHAN UNIVERSITY

eCommons@AKU

---

Section of Otolaryngology, Head & Neck Surgery

Department of Surgery

---

1-1-2021

## Use of endoscope in teaching of otolaryngology residents about site and size of tympanic membrane perforation and its impact on degree of hearing loss in adult patients: A cross sectional study

Muntazir Hussain

Muhammad Wasif


Mohammad Sohail Awan

Shayan Khalid

Zeeshan Sheikh

*See next page for additional authors*

Follow this and additional works at: [https://ecommons.aku.edu/pakistan\\_fhs\\_mc\\_surg\\_otolaryngol\\_head\\_neck](https://ecommons.aku.edu/pakistan_fhs_mc_surg_otolaryngol_head_neck)

 Part of the [Otolaryngology Commons](#), and the [Otorhinolaryngologic Diseases Commons](#)

---

---

**Authors**

Muntazir Hussain, Muhammad Wasif, Mohammad Sohail Awan, Shayan Khalid, Zeeshan Sheikh, and Haissan Iftikhar

---

## Use of endoscope in teaching of otolaryngology residents about site and size of tympanic membrane perforation and its impact on degree of hearing loss in adult patients: A cross sectional study

Muntazir Hussain,<sup>1</sup> Muhammad Wasif,<sup>2</sup> Muhammad Sohail Awan,<sup>3</sup> Shayan Khalid Ghaloo,<sup>4</sup> Zeeshan Sheikh,<sup>5</sup> Haissan Iftikhar<sup>6</sup>

### Abstract

**Objective:** To determine the association between site and size of perforation of the tympanic membrane and the level of conductive hearing.

**Methods:** The cross-sectional study was conducted from November 2015 to October 2016 at Aga Khan University Hospital Karachi and comprised patients with tympanic membrane perforation without any other middle-ear disease. Karl-Storz Rigid Endoscope attached to a camera was used to take pictures of the tympanic membrane. Site of the perforation was determined using a vertical line to divide the membrane into two anterior and posterior halves. Size of the perforation was calculated as a percentage of the total membrane using Image J software. Data analysis was done using Stata 12.

**Results:** Of the 55 patients, 29(53%) were males and 26(47%) were females. The overall mean age was 33+/-15 years. With every 5% increase in the size of perforation, the hearing loss increased by 1 decibel. A difference of 5.5 decibels was noted between anterior and posterior perforations of similar size.

**Conclusion:** Hearing loss increased with increase in the size of perforation.

(JPMA 70: S-14 [Suppl. 1]; 2021)

### Introduction

The external auditory canal and tympanic cavity are separated by tympanic membrane. Vertically, it measures approximately 9-10mm, while, horizontally, it is 8-9mm.<sup>1</sup> Tympanic membrane (TM) acts as a barrier to infections from the external ear to the middle ear, and plays an important role in the transmission of sound from the external to the inner ear. TM perforations can cause conductive hearing loss (CHL) that may range from a negligible loss up to 50 decibels (dB).<sup>2</sup>

There are a number of causes which can lead to TM perforation, including trauma, infections etc. of which infections are the most common cause.<sup>3</sup> Increase in the size of perforation causes greater CHL, and the transformer action of the middle ear is lost in the total absence of TM.<sup>3,4</sup> The magnitude of hearing loss, according to a study, depends significantly on the location of the perforation.<sup>5</sup> However, another study said the location of perforation had no significant effect on hearing.<sup>6</sup>

Knowledge of the anatomical details of TM perforation is

.....  
<sup>1</sup>Cancer Foundation Hospital, Karachi, <sup>2-4,6</sup>Department of Otolaryngology, Head and Neck Surgery, Aga Khan University Hospital, Karachi, <sup>5</sup>Department of Otolaryngology, Head and Neck Surgery, Jinnah Medical and Dental College, Karachi, Pakistan.

**Correspondence:** Muhammad Wasif. Email: wasif\_siddiq313@yahoo.com


of utmost importance for otolaryngology residents, including the size of perforation and, more importantly, the site of perforation. Missing any perforation in the pars flaccida of TM can be critical, as it may harbour underlying cholesteatoma which can lead to serious complications.

The current study was planned to determine the association between the site and the size of TM perforation and the level of CHL. It was also planned to increase the knowledge of residents about the anatomical details of TM perforation by recording and saving it on the monitor attached to a zero-degree rigid endoscope.

### Patients and Methods

The cross-sectional study was conducted from November 2015 to October 2016 at the Department of Otorhinolaryngology and Head and Neck Surgery of Aga Khan University Hospital, Karachi. After exemption from the institutional ethics review committee same hospital, all patients presenting to the outpatient department (OPD) with TM perforation without any other middle-ear disease were enrolled. Those with active middle-ear disease, with history of surgery in the same ear, patients in whom the anterior rim of perforation was not seen, and those with suspected cholesteatoma, ossicular damage, adhesions and tympanosclerosis were excluded.

CHL was measured using pure tone audiogram. Hearing-loss was documented at frequencies of 500Hz, 1000Hz



**Figure:** a) One-third (33%) perforation of tympanic membrane. b) Total area of tympanic membrane.

and 2000Hz. Karl-Storz rigid endoscope attached to a camera and monitor was used to take TM pictures. Site of the perforation was determined using a vertical line to divide the TM into anterior and posterior halves. The size of the perforation was calculated as a percentage of the total TM using Image J software (Figure-1).

Anatomical findings on the still images saved in the monitor by the attending consultant were then explained to the residents without evaluating its impact on residents' understanding.

Data was analysed using Stata 12.0. Univariate analysis was run to determine the associations of hearing-loss with independent variables. Multivariable analysis was carried out using multiple linear regression to assess the factors related to the degree of hearing-loss.  $P < 0.25$  was taken as significant for univariate analysis, while for multivariable analysis the level of significance was set at  $p < 0.05$ .

## Results

Of the 55 patients, 29(53%) were males and 26(47%) were females. The overall mean age was  $33 \pm 15$  years. The left ear was affected in 31(56%) patients, while in 24(44%) patients the right ear was affected. Also, 31(56%) patients had anterior perforation, while 24(44%) patients had posterior perforation. Mean size of TM perforation was  $25.6 \pm 15\%$  (range: 2-72%) of total surface area. Mean CHL loss was  $19.8 \pm 10.1$  dB (range: 0-38dB) (Table-1).

**Table-1:** Quantitative variables.

Variable	Mean $\pm$ SD	Minimum	Maximum
Mean Age(Years)	$33.8 \pm 15.1$	11	75
Size of perforation (%)	$25.6 \pm 15$	2	72
CHL 500Hz(dB)	$25 \pm 12.3$	0	50
CHL 1000Hz(dB)	$20.7 \pm 12.6$	0	45
CHL 2000Hz(dB)	$13.6 \pm 9$	0	30
Average CHL(dB)	$19.8 \pm 10.1$	0	38

SD: Standard deviation; CHL: Conductive hearing loss; dB: Decibels.

**Table-2:** Univariate analysis ( $p < 0.25$ ).

Variable	P value	Result
Age	0.35	Insignificant
Gender	0.19	Significant
Side of perforation	0.69	Insignificant
Size of perforation	0.05	Significant
Site of perforation	0.16	Significant

Univariate analysis showed gender, size of perforation and site of perforation as significant variables (Table-2). On multivariable analysis, only the size and site of perforation were significant (Table-3). For every 5% increase in the size of perforation, hearing-loss increased by 1dB. A difference of 5.5dB was noted between anterior and posterior perforations of similar size with hearing loss being more in the anterior perforations ( $p < 0.05$ ).

**Table-3:** Multiple linear regression analysis ( $p < 0.05$ ).

Factors	P value	Result
Gender and Side of perforation	0.35	Insignificant
Gender and size of perforation	0.14	Insignificant
Size and Site of perforation	0.02	Significant

## Discussion

The middle ear is separated from the external ear by TM, a shield-like structure. Besides transmitting the sound energy from the external to the middle and the internal ear, it also prevents passage of infections from the external to the middle ear. TM can perforate due to various reasons, most commonly by infections and less commonly by trauma and other causes.<sup>3</sup> Once the TM gets perforated, it loses its baffling effect, and the sound is then transferred to both the oval window and the round window simultaneously, resulting in the loss of phase difference and in hearing-loss.<sup>7</sup>

The current study showed that mean CHL was more in lower frequencies compared to the higher frequencies, which is in line with literature.<sup>8,9</sup> An increase in hearing-loss by 1dB was observed for every 5% increase in the size of perforation, and the finding was similar to earlier studies.<sup>10,11</sup>

One important result of the current study is that anterior perforations were associated with higher conductive loss, which was not reported by earlier studies.<sup>12,13</sup> The earlier results can be explained by the fact that a posterior perforation would allow the sound to reach the round window niche situated postero-inferiorly more quickly, resulting in the loss of phase difference compared to anterior perforations. On the contrary, some studies found no difference in hearing-loss between posterior and anterior perforations.<sup>14-16</sup>

The results of the current study can be explained by other less-explored determinants of hearing-loss, such as middle ear volume and malleolar location of TM perforation. One study reported that perforations which involved the malleolar region had higher conductive losses, and another study showed that perforations at the Umbo region were likely to have higher CHL.<sup>17,18</sup> In one study, there was an inverse relationship between the middle-ear volume and the degree of CHL, and a difference up to 35dB could be predicted for perforations of similar size if the middle-ear volume differed substantially.<sup>11</sup>

One of the possibilities behind the variations in findings of the current study compared to earlier ones cited above

could be that there were no major differences in the degree of hearing-loss according to location, and the current results could have been different if the malleolar location and the middle-ear volume parameters were also taken into account. However, the findings do open a door for debate about anterior perforations being associated with higher degree of hearing-loss compared to the posterior location. Further studies, with large sample sizes need to be carried out in which both the middle-ear volume and malleolar location of perforations should also taken into account.

During the course of the current study, it was observed that still images of TM served as a good learning opportunity for the residents and they reported that the anatomical details seen on the monitor with the endoscope were much superior compared to the details they had observed with the otoscope. Further studies focussing on the teaching part of endoscope compared to otoscope are required to see if the endoscope is actually superior. The current study serves as the beginning point of such studies in the future.

## Conclusion

The degree of hearing-loss increased proportionally with increase in the size of TM perforation. Hearing-loss was higher in lower frequencies compared to higher frequencies. Higher degree of hearing-loss was seen in patients with anterior compared to posterior perforations. Also, endoscopes can be considered for routine ear examination and still images can help in better understanding of anatomical details for young residents.

**Disclaimer:** None.

**Conflict of Interests:** None.

**Source of Funding:** None.

## References

1. Isaacson G. Endoscopic anatomy of the pediatric middle ear. *Otolaryngol Head Neck Surg* 2014;150:6-15. doi: 10.1177/0194599813509589.
2. Voss SE, Rosowski JJ, Merchant SN, Peake WT. Non-ossicular signal transmission in human middle ears: Experimental assessment of the "acoustic route" with perforated tympanic membranes. *J Acoust Soc Am* 2007;122:2135-53. doi: 10.1121/1.2769617.
3. Kumar N, Chilke D, Puttewar MP. Clinical Profile of Tubotympanic CSOM and Its Management With Special Reference to Site and Size of Tympanic Membrane Perforation, Eustachian Tube Function and Three Flap Tympanoplasty. *Indian J Otolaryngol Head Neck Surg* 2012;64:5-12. doi: 10.1007/s12070-010-0114-5.
4. Maharjan M, Kafle P, Bista M, Shrestha S, Toran KC. Observation of hearing loss in patients with chronic suppurative otitis media tubotympanic type. *Kathmandu Univ Med J (KUMJ)* 2009;7:397-401. doi: 10.3126/kumj.v7i4.2761.
5. Ahmad SW, Ramani GV. Hearing loss in perforations of the

- tympanic membrane. *J Laryngol Otol* 1979;93:1091-8. doi: 10.1017/s0022215100088162.
6. Sood AS, Pal P, Kumar A. Tympanic membrane perforation: correlation of hearing loss with its site and size. *Int J Otorhinolaryngol Head Neck Surg* 2018;4:397-402.
  7. Ogisi FO, Adobamen P. Type 1 Tympanoplasty in Benin: a 10- year review. *Niger Postgrad Med J* 2004;11:84-7.
  8. Voss SE, Rosowski JJ, Merchant SN, Peake WT. Middle-ear function with tympanic-membrane perforations. I. Measurements and mechanisms. *J Acoust Soc Am* 2001;110:1432-44. doi: 10.1121/1.1394195.
  9. Dawood MR. Frequency Dependence Hearing Loss Evaluation in Perforated Tympanic Membrane. *Int Arch Otorhinolaryngol* 2017;21:336-42. doi: 10.1055/s-0037-1598597.
  10. Vaidya S, Sharma JK, Singh G. Study of outcome of tympanoplasties in relation to size and site of tympanic membrane perforation. *Indian J Otolaryngol Head Neck Surg* 2014;66:341-6. doi: 10.1007/s12070-014-0733-3.
  11. Mehta RP, Rosowski JJ, Voss SE, O'Neil E, Merchant SN. Determinants of hearing loss in perforations of the tympanic membrane. *Otol Neurotol* 2006;27:136-43. doi: 10.1097/01.mao.0000176177.17636.53.
  12. Gaur S, Sinha ON, Bhushan A, Batni G. Observations on Tympanic Membrane Perforations (Safe Type) and Hearing Loss. *Indian J Otolaryngol Head Neck Surg* 2017;69:29-34. doi: 10.1007/s12070-016-1021-1.
  13. Anthony WP, Harrison CW. Tympanic membrane perforation. Effect on audiogram. *Arch Otolaryngol* 1972;95:506-10. doi: 10.1001/archotol.1972.00770080796003.
  14. Ibekwe TS, Nwaorgu OG, Ijaluola TG. Correlating the site of tympanic membrane perforation with Hearing loss. *BMC Ear Nose Throat Disord* 2009;9:1. doi: 10.1186/1472-6815-9-1.
  15. Park H, Hong SN, Kim HS, Han JJ, Chung J, Seo MW, et al. Determinants of conductive hearing loss in tympanic membrane perforation. *Clin Exp Otorhinolaryngol* 2015;8:92-6. doi: 10.3342/ceo.2015.8.2.92.
  16. Saliba I, Abela A, Arcand P. Tympanic membrane perforation: size, site and hearing evaluation. *Int J Pediatr Otorhinolaryngol* 2011;75:527-31. doi: 10.1016/j.ijporl.2011.01.012.
  17. Pannu KK, Chadha S, Kumar D, Preeti. Evaluation of hearing loss in tympanic membrane perforation. *Indian J Otolaryngol Head Neck Surg* 2011;63:208-13. doi: 10.1007/s12070-011-0129-6.
  18. Lerut B, Pfammatter A, Moons J, Linder T. Functional correlations of tympanic membrane perforation size. *Otol Neurotol* 2012;33:379-86. doi: 10.1097/MAO.0b013e318245cea5.
-