



JNCI Cancer Spectrum (2020) 4(2): pkz100


doi: 10.1093/jncics/pkz100

First published online December 11, 2019

Article

## ARTICLE

# BRCA1 Promoter Methylation Status in 1031 Primary Breast Cancers Predicts Favorable Outcomes Following Chemotherapy

Olafur A. Stefansson, PhD<sup>1,2,3</sup>, Holmfrídur Hilmarsdóttir, BSc<sup>2</sup>, Krístrun Olafsdóttir, MSc<sup>1</sup>, Laufey Tryggvadóttir, PhD<sup>1,4</sup>, Asgerdur Sverrisdóttir, MD<sup>5</sup>, Oskar T. Johannsson, MD<sup>5</sup>, Jon G. Jonasson, MD<sup>1,6</sup>, Jorunn E. Eyfjörð, PhD<sup>1,2</sup>, Stefan Sigurdsson , PhD<sup>1,2,\*</sup>

<sup>1</sup>Faculty of Medicine, University of Iceland, Reykjavík, Iceland; <sup>2</sup>Cancer Research Laboratory, Biomedical Center, University of Iceland, Reykjavík, Iceland; <sup>3</sup>Current affiliation deCODE genetics/Amgen Inc., Sturlugata 8, Reykjavík, Iceland; <sup>4</sup>Icelandic Cancer Registry, Icelandic Cancer Society, Reykjavík, Iceland; <sup>5</sup>Department of Oncology, Landspítali University Hospital, Reykjavík, Iceland and <sup>6</sup>Department of Pathology, Landspítali University Hospital, Reykjavík, Iceland

\*Correspondence to: Stefan Sigurdsson, PhD, Faculty of Medicine, Cancer Research Laboratory, Biomedical Center, School of Health Sciences, University of Iceland, Reykjavík, Iceland (e-mail: stefsi@hi.is).

## Abstract

**Background:** Breast Cancer 1 gene (*BRCA1*) is known to be inactivated in breast tumors by promoter methylation. Tumor cells in patients carrying a germline mutation in *BRCA1* are sensitive to cytotoxic drugs that cause DNA double strand breaks. However, very little is known on whether patients with *BRCA1* promoter methylated tumors are similarly sensitive to cytotoxic drugs. In this study, we address this by making use of extensive follow-up data on patients treated with cyclophosphamide, methotrexate, and fluorouracil in Iceland between 1976 and 2007.

**Methods:** We analyzed *BRCA1* promoter methylation by pyrosequencing DNA from tumor samples from 1031 patients with primary breast cancer. Of those, 965 were sporadic cases, 61 were *BRCA2*, and five were *BRCA1* germline mutation carriers. All cases were examined with respect to clinicopathological parameters and breast cancer-specific survival in patients treated with cytotoxic drugs. Information on chemotherapy treatment in noncarriers was available for 26 *BRCA1* methylated tumors and 857 unmethylated tumors.

**Results:** *BRCA1* was promoter methylated in 29 sporadic tumors or in 3.0% of cases (29 of 965), whereas none of the tumors derived from *BRCA* germline mutation carriers were promoter methylated. Important to note, patients with *BRCA1* promoter methylation receiving chemotherapeutic drug treatment show highly improved breast cancer-specific survival compared with unmethylated controls (hazard ratio = 0.10, 95% confidence interval = 0.01 to 0.75, two-sided  $P = .02$ ).

**Conclusions:** *BRCA1* promoter methylation is predictive of improved disease outcome in patients receiving cyclophosphamide, methotrexate, and fluorouracil drug treatment. Our results support the use of markers indicative of “BRCAness” in sporadic breast cancers to identify patients that are likely to benefit from the use of DNA-damaging agents.

Germline mutations in the breast cancer-susceptibility gene, Breast Cancer 1 gene (*BRCA1*), significantly increase the risk of developing breast and ovarian cancer, in addition to other

forms of cancer (1). The *BRCA1* protein product is involved in DNA double strand-break (DSB) repair by homologous recombination, a highly conserved error-free DNA-repair pathway

Received: March 12, 2019; Revised: October 28, 2019; Accepted: December 7, 2019

© The Author(s) 2020. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

that uses an intact sister chromatid in late S or G2 phases of the cell cycle for the repair (2).

Tumors lacking BRCA1 or BRCA2 are homologous recombination deficient (HRD) and are characterized by mutational signatures, including indels, rearrangements, and base substitutions (3,4). Based on these mutational signatures, a model, HRDetect, has been developed to predict the BRCA1/BRCA2 deficiency or “BRCAness” of tumors (3).

Loss of the wild-type allele is seen in most tumors arising in BRCA1 mutation carriers (5). These tumors have high HRDetect scores compared with those that do not show loss of heterozygosity at the BRCA1 loci. Tumors without loss of heterozygosity have similar HRDetect scores as noncarriers.

Germline mutations in BRCA genes and somatic mutations in homologous recombination genes are associated with increased sensitivity to platinum chemotherapy and Poly ADP Ribose Polymerase (PARP) inhibitors in breast cancer (6–10) and ovarian cancer (11–15). These agents induce replication fork stalling, creating DNA substrates that are dependent on homologous recombination for replication restart and are essential for the survival of the cell (16).

BRCA1 is sometimes inactivated in breast tumors by promoter methylation (17,18). BRCA1 methylated tumors are associated with the basal-like or triple-negative subtype that is predominant in germline mutation carriers (19,20). Recently BRCA1 methylated tumors were associated with mutational signatures characteristic of tumors arising in BRCA1 germline mutation carriers (4,21).

It is currently unclear, however, whether BRCA1 promoter methylation translates to clinical benefits from the use of DNA-damaging agents in patients. We therefore carried out a large retrospective study aimed at determining whether BRCA1 methylation is associated with improved outcomes in survival among chemotherapy-treated patients.

## Methods

### Study Group

The study group consisted of 1031 patients (women) diagnosed between 1976 and 2007 previously screened for the local BRCA1 c.5074G>A and BRCA2 c.767-771delCAAAT-germline mutations (22,23). In addition to the two BRCA founder mutations, BRCA2 c.767-771delCAAAT and the much rarer BRCA1 c.5074G>A, the only other BRCA mutation of some frequency, c.9976A>T, is not found to be associated with risk of breast or ovarian cancer but rather risk of small cell lung cancer and squamous cell carcinoma of the skin (24). Out of the 1031 cases, 965 were sporadic, 61 were BRCA2 germline mutation carriers, and five were BRCA1 mutation carriers. DNA samples from these patients were extracted from FF tissue samples (n=417) and adjacent normal breast tissues (n=91) using a standard phenolchloroform (+ proteinase K) method. DNA derived from formalin-fixed and paraffin-embedded tumors (n=615) was extracted by deparaffinization using Octane followed by two rounds of ethanol washes and then overnight incubation in digestion buffer (50 mM Tris pH 8.8, 1 mM EDTA and 0.5% Tween, proteinase K).

This work was carried out according to the permits from the Icelandic Data Protection Commission (2006050307) and Bioethics Committee (VSNb2006050001/03–16).

### DNA-Methylation Analysis

The EZ-96 DNA Methylation-Gold kit (Zymo Research; D5008) was used to carry out bisulfite conversion of DNA samples derived from tumor and normal breast tissues. Primer design was based on the PyroMark Assay Design 2.0 software (Qiagen). Pre-PCR reactions, using the Immolase DNA polymerase (Bioline; Bio-21047), along with pyrosequencing (PyroMark Q24; Qiagen) were carried as previously described (25). Four CpG sites were analyzed at genetic positions: chr17: 43125409(GRCh38.p7), chr17: 43125411(GRCh38.p7), chr17: 43125419(GRCh38.p7), chr17: 43125427(GRCh38.p7).

The primers used for pre-PCR were as follows: Forward primer 5-GTAGGGGTTTAGTTATTTGAGAAATTTT-3; reverse biotinylated primer 5-TATCCCTCCCATCTCTAATTATAC-3. The sequencing primer for the pyrosequencing reaction was as follows: 5-AGTTTAAATTTATTTGTAATTTT-3. Tumor samples were considered to be BRCA1 methylated at median greater than 10% methylation across the four CpGs (median values).

### Clinicopathological Parameters and Treatment

Information on patient age, birth date, and date of diagnosis, tumor grade, staging, and chemotherapy treatment given at time of diagnosis were obtained from the nationwide Icelandic Cancer Registry (26,27). Estrogen-receptor (ER) status was based on tissue-microarray (TMA) analysis (n=664) (20,26) and, where TMA data were not available, we used data derived from routine clinical diagnoses based on dextran-coated charcoal assay used in Iceland in the period 1981 to 1995 (n=98).


### Statistical Analysis

The  $\chi^2$  and Fisher exact contingency table tests were carried out using base functions in R. The Kaplan–Meier method was used for generating univariable survival curves and the log-rank test was used for estimating P values. Hazard ratios (HRs) were estimated by applying the Cox proportional hazards model using R (survival package). Here, BRCA1-methylation status represents the main variable, whereas year of birth, year of diagnosis, ER status, nodal status, and tumor size are introduced as adjustment covariates. Patients were followed from the date of diagnosis until death or date of last follow-up (December 31, 2016). The endpoint was breast cancer-specific survival, defined as the time from diagnosis to death from breast cancer, as registered on death certificates. Patients who died of causes other than breast cancer were censored at the date of death. The cox.zph function in the survival package for R was used to test the proportional hazards assumption for a Cox regression model fit. All tests were two-sided and a P value of less than .05 was considered statistically significant.

## Results

### The Incidence of BRCA1 Methylation in Primary Breast Tumors and Normal Tissue

We analyzed BRCA1 promoter CpG methylation in 1031 primary tumors, along with 91 normal breast tissue samples derived from the same cohort. BRCA1 methylation was identified in 29 tumors, none of which were derived from carries of germline mutations in either BRCA1 (n=5) or BRCA2 (n=61)



**Figure 1.** Box plot of BRCA1-promoter CpG methylation in normal breast tissue and tumors derived from carriers of germline mutations in BRCA1, BRCA2, and noncarriers.

(Figure 1). The frequency of BRCA1 promoter methylation in non-BRCA mutated breast tumors is approximately 3% (29 out of 965; 3.0%).

### BRCA1 Methylation in Relation to Clinicopathological Parameters

Information on ER status was available for 762 out of the 1031 tumors in our cohort; 700 out of the 762 tumors were derived from noncarriers (Table 1, A). We find that 90.5% of tumors with BRCA1 promoter methylation are negative for ER, or 19 out of 21 (see Table 1, A). Similarly, tumors arising in BRCA1 mutation carriers were exclusively ER negative (five out of five) as previously reported (20), and by TMA analysis, we find that the majority of BRCA1 methylated tumors are of the basal-like subtype (eight out of 12; 67%) (see Table 1, B).

BRCA1 promoter-methylated tumors are statistically significantly less differentiated based on histopathological grading ( $\chi^2=11.1$ ,  $P=.0039$ ) (Table 1, C). However, no association was found with respect to clinical staging in terms of tumor size or nodal status (Table 1, D and E).

### The Effects of BRCA1 Methylation with Respect to Responsiveness to Cytotoxic Treatment

To assess the association between BRCA1 methylation and response to cytotoxic chemotherapy, we obtained information on chemotherapy and used time to patient death (breast cancer-specific survival) as a proxy for treatment response. In our cohort, cyclophosphamide, methotrexate, and fluorouracil (CMF) was the most commonly used cytotoxic treatment given at time of diagnosis, and nine of 29 patients with BRCA1 methylated tumors received CMF treatment (Table 2). Information on chemotherapy treatment in noncarriers was available for 26 BRCA1 methylated tumors and 857 unmethylated tumors. In noncarriers, patients with BRCA1 methylated tumors show long-term

survival following cytotoxic treatment (HR = 0.10, 95% CI = 0.014 to 0.751,  $P=.025$ ) (Figure 2, A; Table 3). In contrast, nontreated patients with BRCA1 methylation, compared with nontreated patients without BRCA1 methylated tumors, showed similar time to breast cancer-specific death (Figure 2, B; Table 3). In comparing treated vs nontreated survival curves for non-BRCA mutated cases (Figure 2, A and B), there are notable differences, but these are likely explained by clinical presentation reflecting poor prognosis, and as a result, the treated group is biased toward reduced survival. This, however, does not affect our results, as we find improved survival for patients with BRCA1-methylated tumors after receiving cytotoxic treatment.

In this cohort, nine patients with BRCA1 methylated tumors received chemotherapeutic treatment with CMF (Table 2). However, patients with non-BRCA1 methylated tumors received CMF or other treatments. By restricting our analysis to CMF-treated patients only, we find that the relation between BRCA1 methylation and breast cancer-specific survival holds statistical significance (HR = 0.11, 95% CI = 0.014 to 0.81,  $P=.03$ ).

## Discussion

In this article, we demonstrate that BRCA1 promoter methylation, analyzed in a large cohort of 1031 primary breast tumors, predicts improved breast cancer-specific survival outcomes in patients treated with cytotoxic chemotherapy. No difference is seen in patients that did not receive a cytotoxic treatment, suggesting that BRCA1 promoter methylation is a predictive factor for chemosensitivity but not a prognostic factor. Previous studies have shown that BRCA1 mutation carriers are sensitive to treatment with platinum and PARP inhibitors. However, little is known about the effect of BRCA1 methylation with respect to treatment based on DNA-damaging agents.

The classical CMF combination was the standard treatment for breast cancer patients until it was replaced by anthracyclines and taxanes as adjuvant treatment (28). Although it is less effective than anthracyclines and taxanes, large retrospective studies have shown a clear benefit of CMF, especially in triple-negative breast cancer (29–31). Although both methotrexate and fluorouracil (5-FU) are antimetabolites that block the synthesis of thymidine required for DNA synthesis, cyclophosphamide is an alkylating agent, and its cytotoxic metabolite phosphoramidate mustard leads to inter- and intrastrand cross-links in DNA causing replication fork stalling (32–34). Because of the importance of the DNA DSB repair machinery in DNA-replication restart, phosphoramidate mustard treatment is likely to lead to synergistic lethality in BRCA-deficient cells. This mechanism of action is similar to that currently thought to underlie the effectiveness of PARP1 inhibitors and platinum drugs in killing BRCA deficient cells (16). The mechanism of PARP1 inhibition is however more complex than previously expected. In addition to generating persistent single-strand breaks leading to collapse of the DNA-replication fork, PARP1 becomes trapped on DNA when inhibited, forming a cytotoxic lesion leading to replication fork stalling (35). Recently PARP inhibitors were also shown to affect the recruitment of POLQ to DNA DSBs, inhibiting microhomology-mediated end joining or Alt-EJ pathway (36–38).

Recent data have shown that 40–70% of triple-negative breast cancer is HR deficient (3,4,39) and therefore likely to respond to agents causing DNA-replication fork stalling (40–43). Replication fork stalling possibly explains the efficacy of

**Table 1.** CpG–promoter methylation of BRCA1 gene analyzed with respect to clinicopathological parameters and breast cancer subtypes\*

A) ER status with respect to BRCA1 methylation		
ER status	BRCA1 unmethylated	BRCA1 methylated
ER positive	526 (77.5%)	2 (9.5%)
ER negative	153 (22.5%)	19 (90.4%)
Total (N = 700)	679	21
Fisher test, $P < .0001$		
B) Breast cancers subtypes with respect to BRCA1 methylation		
Subtype	BRCA1 unmethylated	BRCA1 methylated
LumA	78 (40%)	0
LumB	61 (31.3%)	1 (8.3%)
HER2	18 (9.2%)	0
Basal-like	34 (17.4%)	8 (66.7%)
5-negative (5NP)	4 (2.1%)	3 (25%)
Total (N = 207)	195	12
$\chi^2 = 39.0, P < .0001$		
C) Histological grade with respect to BRCA1 methylation		
Histological grade	BRCA1 unmethylated	BRCA1 methylated
1+	104 (23.6%)	0
2+	187 (42.4%)	6 (31.6%)
3+	150 (34%)	13 (68.4%)
Total (N = 460)	441	19
$\chi^2 = 11.1, P = .0039$		
D) Tumor size with respect to BRCA1 methylation		
Tumor size	BRCA1 unmethylated	BRCA1 methylated
T1	462 (54.4%)	8 (30.8%)
T2	319 (37.6%)	16 (61.5%)
T3	54 (6.4%)	2 (7.7%)
T4	14 (1.6%)	0
Total (N = 875)	849	26
$\chi^2 = 6.9, P = .075$		
E) Nodal status with respect to BRCA1 methylation		
Nodal status	BRCA1 unmethylated	BRCA1 methylated
Negative	422 (51.9%)	17 (65.4%)
Positive	391 (48.1%)	8 (30.6%)
Total (N = 839)	813	26
Fisher test, $P = .15$		
F) Age at diagnosis		
Age	BRCA1 unmethylated	BRCA1 methylated
≤55	439 (46.9%)	22 (75.9%)
>55	497 (53.1%)	7 (24.1%)
Total (N = 965)	936	29
Fisher test, $P = .002$		


\*ER = estrogen receptor.

cyclophosphamide in triple-negative tumors (30,31), as these are likely to be HRD.

HRDetect is a recently introduced method that makes use of next-generation sequencing for identifying mutational signatures characteristic of tumors arising in BRCA1 or BRCA2 germline mutation carriers (3). The main idea behind HRDetect is to enable identification of tumors with defective homologous recombination repair. Indeed, Davies et al. found that tumors with BRCA1 methylation were detected as HRD by HRDetect (3). However, it is unclear whether this finding translates to enhanced sensitivity of BRCA1 methylated tumors to treatment

with DNA-damaging agents. Our results suggest that, indeed, this does seem to be the case. It is important to note that HRDetect genomic signatures might still persist after restored homologous recombination function, especially in the metastatic setting. To discriminate between tumors that are truly HRD, functional methods such as RAD51 foci staining have been developed for clinical use to improve selection of patients sensitive to DNA-damaging agents (9,10).

Information on HRDetect scores was available for 24 tumors in our cohort, of which four were BRCA1 methylated as previously described ([3]; Supplementary Figure 1, available online).



**Figure 2.** Breast cancer-specific survival time in patients treated **A)** with and **B)** without cytotoxic chemotherapy analyzed with respect to *BRCA1* methylation status. The *P* values shown reflect log-rank hypothesis testing for differences in survival times with respect to *BRCA1* methylation status.

**Table 2.** Type of chemotherapeutic drugs used during treatment in the cohort of non-*BRCA* mutation carriers listed out with respect to *BRCA1* methylation status\*

	<i>BRCA1</i> unmethylated	<i>BRCA1</i> methylated
CMF	242	9
Vincristine	97	2
Taxanes	9	0
Anthracycline	53	0
Any type	294	9

\*CMF = cyclophosphamide, methotrexate, and fluorouracil.

**Table 3.** The effect of *BRCA1* methylation on time to breast cancer-specific death with and without chemotherapeutic treatment analyzed by multivariate Cox proportional hazards regression\*

Treatment	HR (95% CI)	<i>P</i>
<i>BRCA1</i> methylation in nontreated patients ( <i>n</i> = 391)	1.58 (0.52 to 4.80)	.416
<i>BRCA1</i> methylation in treated patients ( <i>n</i> = 183)	0.10 (0.01 to 0.75)	.025

\*Adjusted for year of diagnosis and birth, estrogen-receptor status, tumor size, and nodal status. CI = confidence interval; HR = hazard ratio.

The four *BRCA1* methylated tumors all show high HRDetect scores, comparable to values seen in *BRCA1* mutation carriers. Two of these four received CMF treatment, and both are still alive today, more than 18 years after their diagnosis.

Recently, in Tutt et al., advanced TNBC tumors were investigated for response to docetaxel and carboplatin, which revealed better response to docetaxel. These tumors, however, were pretreated with drugs that cause DNA lesions that require DNA DSB repair. As suggested by the authors, *BRCA1* methylation detected in the primary tumor may have been lost following this first treatment, thereby explaining the lack of response to carboplatin in the advanced tumors. Our study differs from that of Tutt et al. in that we investigate the response to first treatment in our patient cohort (8).

Of interest, we observe substantially lower frequency of *BRCA1* methylation than previously reported (17,18,44,45). The discrepancy likely reflects either smaller cohorts in previous studies, leading to a situation where the detection of a single methylated tumor greatly influences the frequency, or use of methods known to have a higher false-positive detection rate, compared with pyrosequencing, such as methylation-specific PCR. In our cohort, 3% of primary breast cancer samples are *BRCA1* methylated, which is in agreement with a recent study using data from The Cancer Genome Atlas (21).

Our cohort is based on DNA samples isolated from freshly frozen (FF) tumors (*n* = 417) and formalin-fixed paraffin-embedded (FFPE) tumors (*n* = 615). Previous studies have shown higher fraction of *BRCA1* methylation in FFPE samples (46). Our analysis, based on pyrosequencing, does not show statistically significant differences in frequency of *BRCA1* methylation in noncarriers (*n* = 965) between FFPE (14 out of 587; 2.4%) and FF (15 out of 378; 4.0%).

A limitation of this study lies in the low number of *BRCA1* methylated tumors detected in our cohort. However, we would like to point out that this is the largest cohort studied to date on *BRCA1* methylation. Another limitation lies in the use of time to breast cancer-specific death as a proxy for response to treatment.

In summary, we demonstrate that *BRCA1* methylation is an important predictive factor of chemosensitivity in breast cancer rather than being a prognostic factor. This provides support for the use of methods aimed at defining *BRCA*ness to identify patients that will derive benefits from DNA-damaging



chemotherapy treatment and, possibly, targeted therapy using PARP inhibitors.

## Funding

The authors would like to thank The Icelandic Research Fund ([www.rannis.is](http://www.rannis.is)) (14193–051 and 152077–051) and Gongum saman ([www.gongumsaman.is](http://www.gongumsaman.is)) for funding.

## Notes

The funders had no role in the design of the study; the collection, analysis, and interpretation of the data; the writing of the manuscript; and the decision to submit the manuscript for publication.

The authors have no disclosures.

## References

- Laloo F, Evans DG. Familial breast cancer. *Clin Genet.* 2012;82(2):105–114.
- Jasin M, Rothstein R. Repair of strand breaks by homologous recombination. *Cold Spring Harb Perspect Biol.* 2013;5(11):a012740.
- Davies H, Glodzik D, Morganella S, et al. HRDetect is a predictor of BRCA1 and BRCA2 deficiency based on mutational signatures. *Nat Med.* 2017;23(4):517–525.
- Nik-Zainal S, Davies H, Staaf J, et al. Landscape of somatic mutations in 560 breast cancer whole-genome sequences. *Nature.* 2016;534(7605):47–54.
- Maxwell KN, Wubbenhorst B, Wenz BM, et al. BRCA locus-specific loss of heterozygosity in germline BRCA1 and BRCA2 carriers. *Nat Commun.* 2017;8(1):319.
- Fong PC, Boss DS, Yap TA, et al. Inhibition of poly(ADP-ribose) polymerase in tumors from BRCA mutation carriers. *N Engl J Med.* 2009;361(2):123–134.
- Hahnen E, Lederer B, Hauke J, et al. Germline mutation status, pathological complete response, and disease-free survival in triple-negative breast cancer: secondary analysis of the geparsixto randomized clinical trial. *JAMA Oncol.* 2017;3(10):1378–1385.
- Tutt A, Tovey H, Cheang MCU, et al. Carboplatin in BRCA1/2-mutated and triple-negative breast cancer BRCAness subgroups: the TNT trial. *Nat Med.* 2018;24(5):628–637.
- Cruz C, Castroviejo-Bermejo M, Gutierrez-Enriquez S, et al. RAD51 foci as a functional biomarker of homologous recombination repair and PARP inhibitor resistance in germline BRCA-mutated breast cancer. *Ann Oncol.* 2018;29(5):1203–1210.
- Castroviejo-Bermejo M, Cruz C, Llop-Guevara A, et al. A RAD51 assay feasible in routine tumor samples calls PARP inhibitor response beyond BRCA mutation. *EMBO Mol Med.* 2018;10(12):e9172.
- Bolton KL, Chenevix-Trench G, Goh C, et al. Association between BRCA1 and BRCA2 mutations and survival in women with invasive epithelial ovarian cancer. *JAMA.* 2012;307(4):382–390.
- Mirza MR, Monk BJ, Herrstedt J, et al. Niraparib maintenance therapy in platinum-sensitive, recurrent ovarian cancer. *N Engl J Med.* 2016;375(22):2154–2164.
- Swisher EM, Lin KK, Oza AM, et al. Rucaparib in relapsed, platinum-sensitive high-grade ovarian carcinoma (ARIEL2 Part 1): an international, multicentre, open-label, phase 2 trial. *Lancet Oncol.* 2017;18(1):75–87.
- Vencken PM, Kriege M, Hoogwerf D, et al. Chemosensitivity and outcome of BRCA1- and BRCA2-associated ovarian cancer patients after first-line chemotherapy compared with sporadic ovarian cancer patients. *Ann Oncol.* 2011;22(6):1346–1352.
- Yang D, Khan S, Sun Y, et al. Association of BRCA1 and BRCA2 mutations with survival, chemotherapy sensitivity, and gene mutator phenotype in patients with ovarian cancer. *JAMA.* 2011;306(14):1557–1565.
- Lord CJ, Ashworth A. PARP inhibitors: Synthetic lethality in the clinic. *Science.* 2017;355(6330):1152–1158.
- Birgisdottir V, Stefansson OA, Bodvarsdottir SK, et al. Epigenetic silencing and deletion of the BRCA1 gene in sporadic breast cancer. *Breast Cancer Res.* 2006;8(4):R38.
- Esteller M, Silva JM, Dominguez G, et al. Promoter hypermethylation and BRCA1 inactivation in sporadic breast and ovarian tumors. *J Natl Cancer Inst.* 2000;92(7):564–569.
- Foulkes WD, Stefansson IM, Chappuis PO, et al. Germline BRCA1 mutations and a basal epithelial phenotype in breast cancer. *J Natl Cancer Inst.* 2003;95(19):1482–1485.
- Stefansson OA, Jonasson JG, Olafsdottir K, et al. CpG island hypermethylation of BRCA1 and loss of pRb as co-occurring events in basal/triple-negative breast cancer. *Epigenetics.* 2011;6(5):638–649.
- Polak P, Kim J, Braunstein LZ, et al. A mutational signature reveals alterations underlying deficient homologous recombination repair in breast cancer. *Nat Genet.* 2017;49(10):1476–1486.
- Bergthorsson JT, Jonasdottir A, Johannesdottir G, et al. Identification of a novel splice-site mutation of the BRCA1 gene in two breast cancer families: screening reveals low frequency in Icelandic breast cancer patients. *Hum Mutat.* 1998;(suppl 1):S195–S197.
- Thorlacius S, Olafsdottir G, Tryggvadottir L, et al. A single BRCA2 mutation in male and female breast cancer families from Iceland with varied cancer phenotypes. *Nat Genet.* 1996;13(1):117–119.
- Rafnar T, Sigurjonsdottir GR, Stacey SN, et al. Association of BRCA2 K3326\* with small cell lung cancer and squamous cell cancer of the skin. *J Natl Cancer Inst.* 2018;110(9):967–974.
- Stefansson OA, Hermanowicz S, van der Horst J, et al. CpG promoter methylation of the ALKBH3 alkylation repair gene in breast cancer. *BMC Cancer.* 2017;17(1):469.
- Jonasson JG, Stefansson OA, Johannsson OT, et al. Oestrogen receptor status, treatment and breast cancer prognosis in Icelandic BRCA2 mutation carriers. *Br J Cancer.* 2016;115(7):776–783.
- Sigurdardottir LG, Jonasson JG, Stefansson S, et al. Data quality at the Icelandic Cancer Registry: comparability, validity, timeliness and completeness. *Acta Oncol.* 2012;51(7):880–889.
- Munzone E, Curigliano G, Burstein HJ, et al. CMF revisited in the 21st century. *Ann Oncol.* 2012;23(2):305–311.
- Bonadonna G, Moliterni A, Zambetti M, et al. 30 years' follow up of randomised studies of adjuvant CMF in operable breast cancer: Cohort study. *BMJ.* 2005;330(7485):217.
- Colleoni M, Cole BF, Viale G, et al. Classical cyclophosphamide, methotrexate, and fluorouracil chemotherapy is more effective in triple-negative, node-negative breast cancer: results from two randomized trials of adjuvant chemoendocrine therapy for node-negative breast cancer. *J Clin Oncol.* 2010;28(18):2966–2973.
- Falo C, Moreno A, Benito E, et al. Primary chemotherapy with cyclophosphamide, methotrexate, and 5-fluorouracil in operable breast carcinoma. *Cancer.* 2005;103(4):657–663.
- Anderson D, Bishop JB, Garner RC, et al. Cyclophosphamide: review of its mutagenicity for an assessment of potential germ cell risks. *Mutat Res.* 1995;330(1–2):115–181.
- Ganesan S, Keating AF. Phosphoramidate mustard exposure induces DNA adduct formation and the DNA damage repair response in rat ovarian granulosa cells. *Toxicol Appl Pharmacol.* 2015;282(3):252–258.
- Wilson PM, Danenberg PV, Johnston PG, et al. Standing the test of time: targeting thymidylate biosynthesis in cancer therapy. *Nat Rev Clin Oncol.* 2014;11(5):282–298.
- Murai J, Huang SY, Das BB, et al. Trapping of PARP1 and PARP2 by clinical PARP inhibitors. *Cancer Res.* 2012;72(21):5588–5599.
- Ceccaldi R, Liu JC, Amunugama R, et al. Homologous-recombination-deficient tumours are dependent on Poltheta-mediated repair. *Nature.* 2015;518(7538):258–262.
- D'Andrea AD. Mechanisms of PARP inhibitor sensitivity and resistance. *DNA Repair (Amst).* 2018;71:172–176.
- Mateos-Gomez PA, Gong F, Nair N, et al. Mammalian polymerase theta promotes alternative NHEJ and suppresses recombination. *Nature.* 2015;518(7538):254–257.
- Loibl S, Weber KE, Timms KM, et al. Survival analysis of carboplatin added to an anthracycline/taxane-based neoadjuvant chemotherapy and HRD score as predictor of response-final results from GeparSixto. *Ann Oncol.* 2018;29(12):2341–2347.
- Akashi-Tanaka S, Watanabe C, Takamaru T, et al. BRCAness predicts resistance to taxane-containing regimens in triple negative breast cancer during neoadjuvant chemotherapy. *Clin Breast Cancer.* 2015;15(1):80–85.
- Sharma P. Update on the treatment of early-stage triple-negative breast cancer. *Curr Treat Options Oncol.* 2018;19(5):22.
- Telli ML, Hellyer J, Audeh W, et al. Homologous recombination deficiency (HRD) status predicts response to standard neoadjuvant chemotherapy in patients with triple-negative or BRCA1/2 mutation-associated breast cancer. *Breast Cancer Res Treat.* 2018;168(3):625–630.
- Telli ML, Timms KM, Reid J, et al. Homologous Recombination Deficiency (HRD) score predicts response to platinum-containing neoadjuvant chemotherapy in patients with triple-negative breast cancer. *Clin Cancer Res.* 2016;22(15):3764–3773.
- Shakeri H, Fakhrou A, Nikanfar A, et al. Methylation analysis of BRCA1 and APC in breast cancer and its relationship to clinicopathological features. *Clin Lab.* 2016;62(12):2333–2337.
- Xu X, Gammon MD, Zhang Y, et al. BRCA1 promoter methylation is associated with increased mortality among women with breast cancer. *Breast Cancer Res Treat.* 2009;115(2):397–404.
- Turner NC, Reis-Filho JS, Russell AM, et al. BRCA1 dysfunction in sporadic basal-like breast cancer. *Oncogene.* 2007;26(14):2126–2132.