

Accepted Manuscript

Title: Densities of spotted hyaena (*Crocuta crocuta*) and African golden wolf (*Canis anthus*) increase with increasing anthropogenic influence

Authors: Gidey Yirga, Herwig Leirs, Hans H. De Iongh, Tsehaye Asmelash, Kindeya Gebrehiwot, Matthijs Vos, Hans Bauer



PII: S1616-5047(17)30044-7
DOI: <http://dx.doi.org/doi:10.1016/j.mambio.2017.02.004>
Reference: MAMBIO 40891

To appear in:

Received date: 22-3-2016
Accepted date: 14-2-2017

Please cite this article as: Yirga, Gidey, Leirs, Herwig, De Iongh, Hans H., Asmelash, Tsehaye, Gebrehiwot, Kindeya, Vos, Matthijs, Bauer, Hans, Densities of spotted hyaena (*Crocuta crocuta*) and African golden wolf (*Canis anthus*) increase with increasing anthropogenic influence. *Mammalian Biology* <http://dx.doi.org/10.1016/j.mambio.2017.02.004>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Densities of spotted hyaena (*Crocuta crocuta*) and African golden wolf (*Canis anthus*) increase with increasing anthropogenic influence

Gidey Yirga^{*a,b}, Herwig Leirs^c, Hans H. De Iongh^{c, d}, Tsehaye Asmelash^e, Kindeya Gebrehiwot^f, Matthijs Vos^b and Hans Bauer^g

^aDepartment of Biology, Mekelle University P.O. Box 231, Mekelle, Ethiopia

^bTheoretical and Applied Biodiversity Research, Ruhr Universität Bochum, D-44780 Bochum, Germany

^cEvolutionary Ecology Group, University of Antwerp, Groenenborgerlaan 171, 2020 Antwerp, Belgium

^dInstitute of Environmental Sciences, Leiden University P.O.Box 9518, Leiden, the Netherlands

^eDepartment of Microbiology, Mekelle University P.O. Box 231, Mekelle, Ethiopia

^fDepartment of land resource management and environmental protection, Mekelle University P.O. Box 231, Mekelle, Ethiopia

^gWildlife Conservation Research Unit, Zoology, Reanati-Kaplan Centre, University of Oxford. Tubney House, Tubney OX13 5QL, UK

* Corresponding author: gidey.yirga@yahoo.com

Abstract

We report densities of spotted hyaena (*Crocuta crocuta*) and African golden wolf (*Canis anthus*) in Enderta district in northern Ethiopia with high human and low natural prey densities. We estimated spotted hyaena and African golden wolf abundance and characterized their spatial distribution with three methods: we surveyed four road-transects for 66 nights during dry (n=41) and wet (n=25) seasons, we used 34 calling stations and we mapped all active spotted hyaena dens at the time of the survey. The density of spotted hyaena and African golden wolf increased with proximity to towns where human density was higher. A total of 562 spotted hyaena and 63 African golden wolf responded to calling stations, leading to estimates of 1,145 spotted hyaena and 166 African golden wolf in Enderta district. This method also found a significantly higher spotted hyaena and African golden wolf abundance in high human density areas. Maximum response radius was 2.8 km for spotted hyaena and 2.5 km for African golden wolf, and response probability was 0.83 for spotted hyaena and 0.8 for African golden wolf, respectively. We found 40 active spotted hyaena dens with 1,507 remnants of prey, and the majority of the dens were located close to rivers and villages. Our findings show a positive relationship between spotted hyaena, African golden wolf and human concentrations that might demonstrate a case of exceptional coexistence of humans and carnivores, both at high densities. We suggest further investigations into co-adaptations between humans and predators in the study area.

Key-words: abundance, carnivores, density, human-carnivore coexistence, human-carnivore conflict, prey depleted landscapes

Introduction

In protected areas, carnivore density is strongly related to the available natural prey biomass (Karanth et al. 2004; Khorozyan et al. 2008) while in human-dominated landscapes carnivore density depends on the available anthropogenic resources (Gehrt et al. 2010; Luis Llanez 2012; Yirga et al. 2013). When these anthropogenic food resources are abundant and human persecution is low, large carnivores can reach densities even greater than their densities in the wild (Athreya 2012). Occurrence and persistence of carnivores like spotted hyaena (*Crocuta crocuta*,) and golden jackal (*Canis aureus* and *Canis anthus*) in anthropogenic landscapes is associated with availability of anthropogenic food sources (Woodroffe and Ginsberg 1998; Fuller and Sievert 2001; Basille et al. 2009). Predators including spotted hyaena and golden jackal are able to persist at high human densities and at high levels of landscape transformation (Linnell et al. 2001; Cardillo et al. 2004; Blanco and Cortes 2007; Basille et al. 2009; Agarwal et al. 2010).

Spotted hyaenas show high behavioral and ecological plasticity that help them to live in close proximity to humans (Woodroffe 2000; Sunquist and Sunquist 2001; Boydston et al. 2003). They inhabit very wide historical ranges with relatively stable populations across Africa (Kolowski and Holekamp 2009) and are opportunistic carnivores that feed on locally abundant prey species (Cooper et al. 1999). Spotted hyaenas live in societies that are large (up to 90 hyenas in a clan) and more complex than any other carnivores (Kruuk 1972; Holekamp et al. 2007). The mean clan size across Africa is approximately 21 hyenas (Smith and Holekamp 2010). Sizes of subgroups of spotted hyaenas range from one to tens of animals (Holekamp et al. 2000). These subgroups are important for cooperative activities as territorial defense (Tilson and Hamilton 1984; Frank 1986; Henschel and Skinner 1991; Boydston et al. 2001). Spotted hyaenas often

congregate at a communal den (Boydston et al. 2003). Dens serve important protective functions such as the sheltering of cubs, which usually spend their first 8–12 months of life at dens (Holekamp and Smale 1998). Dens also have important social functions. For instance, cubs form relationships with both peers and adults at dens, and dens serve as meeting points for adults (Holekamp et al. 2000). However, it is poorly known how spotted hyaenas select den sites in human-dominated landscapes.

The golden jackal (*Canis anthus* and *Canis aureus*) is found in many areas of Africa, Europe and Asia (Sillero-Zubiri et al. 2004) and is able to live in human-dominated agricultural landscapes (Šálek et al. 2014). Due to their habitat plasticity and opportunistic feeding habits, golden jackals inhabit a variety of different habitat types (Stoyanov 2012; Giannatos 2004; Jhala and Moehlman 2004). Golden jackals venture into human habitations at night to feed on human organic waste (Jhala 2008). They are common throughout their range and high densities are observed in areas with abundant food and cover (Jhala 2008). Despite its widespread distribution, there is very little quantitative information on golden jackal abundance, habitat use, and ranging patterns in relation to food availability (Sillero-Zubiri et al. 2004; Jhala 2008).

The African golden wolf (*Canis anthus*) and Eurasian golden jackal (*Canis aureus*), were originally thought to be the same species, formerly known as the golden jackal (Koepfli et al. 2015). African and Eurasian golden jackals are genetically distinct lineages and African golden jackals merit recognition as a distinct species (Koepfli et al. 2015). Two recent studies reported that the larger-sized golden jackals from Ethiopia and North and West Africa were more closely related to gray wolves than to other populations of golden jackals (Rueness et al., 2011; Gaubert et al., 2012). The small, golden-like jackal from eastern African is actually a small variety of a new species, distinct from the gray wolf, that has a distribution across North and East Africa, and

this previously unrecognized species is named the African golden wolf (*Canis anthus*) (Koepfli et al. 2015). According to this report taxonomists had mistaken African and Eurasian golden jackals for the same species because of a high degree of similarity in their skull and tooth morphology. According to Koepfli et al. (2015), the golden-like jackal from Enderta district, northern Ethiopia is most probably the African golden wolf (*Canis anthus*) hereafter called golden wolf.

Spotted hyaena and the golden wolf are common carnivores across Enderta district, and can easily be seen almost every night at the outskirts of the city. We sought to establish spotted hyaena, golden wolf and human abundance relationships and map spotted hyaena dens in human-dominated landscapes. We also investigated spotted hyaena and golden wolf spatial distribution patterns.

Study area

The study was conducted within Enderta district of Tigray region, northern Ethiopia (Fig 1). The district is located between 13⁰-14⁰ North and at 39⁰-40⁰ 30' East; it has a population density of 79 persons km⁻² and an average family size of five (Sara 2010). Altitude ranges from 1,400 to 2,700 m.a.s.l. and the landscape is mountainous, with undulating, flat and plain lands, rugged valleys and gorges, and hilly areas (Sara 2010). The district has a total surface area of 1,416 km², of which permanent rainfed agriculture covers 49 %; the remaining area is too arid or too steep for agriculture and most of it is used for grazing. The average land holding size per household is

0.75 hectare and more than 80 % of the population is engaged in subsistence farming (Sara 2010). Due to chronic food insecurity, many of the households are at least partly dependent on food-aid. Rainfall of the area is bimodal: a short rainy season between January and April, and a long rainy season from June to August with an average annual rainfall of 550 mm. The mean maximum temperature ranges between 12 and 27° C. In the northern highlands of Ethiopia, forests have been completely converted into farms and grazing lands over the last few centuries except for patchy remnants of old aged Afromontane forests around most Ethiopian Orthodox Tewahido Churches (Aerts et al. 2007; Alemayehu 2007). These fragmented church forests are widely distributed habitat patches, and spotted hyaenas hide in these church forests during the day. The church forests have traditionally and religiously been protected for centuries.

Methods

Road Transect survey

We monitored four road transects every month from March to August, 2012 to record spotted hyaena and golden wolf sightings. Transects centered around the city of Mekelle (regional capital) and started at the outskirts of town (urban center). The first (23 km) went East to Adikolo; the second (38 km) went South to Adigudom; the third (36 km) went North to Wukro; and the last (19 km) went East from Quiha (Mekele agglomeration) to Aragure. We started the transect survey at dusk and usually ended around 2100-2200h (GMT+3). We noted during the survey all encountered spotted hyaenas and golden wolves as well as moon phase, season, start and finish time, time of the sighting, distance from viewer (estimate by observers) to the animal and the distance from the beginning point of the driving. We surveyed transects during dry (March through May) and wet (June through August) seasons in a vehicle at a speed of 30 km per hour. Two team members standing at the back of the vehicle with powerful torches (Maglite

LED) were scanning right and left sides (covering approximately 600 m width of transect) for spotted hyaena and golden wolves; in case of doubt a team member would run towards these animals carrying a torch to confirm species. The research team had torches, GPS (Garmin E-trex) and night vision Binoculars (Yukon Ranger 5x42 LT). We placed transects along the main roads because other areas were not accessible and were difficult to drive.

Abundance and distribution survey

We surveyed spotted hyaena and golden wolf abundance and distribution using 34 randomly selected calling stations from 25 March to 4 April in 2012. A calling station survey is an effective and inexpensive technique for counting spotted hyaena and lion (Ogutu and Dublin 1998; Mills et al. 2001; Maddox 2003; Bauer 2007) and jackal (Ogutu 1994; Maddox 2003) in African ecosystems. We played continuous gnu- hyaena distress calls and spotted hyaena sounds for an hour on an MP3 player connected to a megaphone (Monacor 45) placed on the roof of a vehicle. We performed two cycles of 20 minutes broadcast followed by ten minutes pause and rotated the speaker 90° after 5 minutes broadcast to provide 360° coverage. For optimal visibility, we placed calling stations in open areas. Four observers counted responding spotted hyaena and golden wolf using torches immediately after the last broadcast. We placed calling stations at least 7 km apart to ensure that the same individual was not called in at multiple sites.

Estimation of predator abundance and density

The total population size of spotted hyaena and golden wolf was estimated using the model of Mills et al. (2001) and Ogutu et al. (2005). The expected number of spotted hyaena and golden wolf within the response range (u) was estimated by

$$u_s = y_s / S_s \quad (1)$$

Where y is the mean number of carnivores for species s responding per station and S is the response probability of each species from the calibration experiments (Mills et al. 2001; Ogutu et al. 2005). The total number of spotted hyaena (NT) and golden wolf in Enderta district was estimated using the relation

$$NT = (AH/AS) u \quad (2)$$

Where AH =the area of the entire district and AS =the area sampled around a call-in station (Mills et al. 2001; Ogutu et al. 2005).

Calibration experiments

We estimated call-up maximum response range and probability for spotted hyaena and golden wolf following Mills et al. (2001) and Ogutu et al. (2005) through calibration experiments. We performed independent call-ups to estimate the distance from which spotted hyaena and golden wolf were attracted to broadcasts in the following habitats: area enclosure (partially protected areas of a few hectares dotted across the landscape), open agricultural area, human settlement and garbage dumping area. We conducted calibration experiments for spotted hyaenas and golden wolf to estimate maximum response range and response probability (% of spotted hyaenas and golden wolves responding within range) after the abundance and distribution survey. Calibration experiments were conducted in non-survey areas (places where broadcasts had not been done during the abundance and distribution survey).

To minimize the likelihood of habituation (Ogutu and Dublin 1998; Mills et al 2001; Maddox 2003), we looked for spotted hyaenas and golden wolves in non-survey areas. One vehicle stayed with the animals to record their reaction while the other drove away and conducted the broadcast as done during the call-in surveys. We played the tape at 4.0, 3.5, 3, 2.8, 2.6, 2.5, 2.3, and 2 km,

with the megaphone facing in the direction of the target species. These tests were conducted by six observers in mobile communication.

Spotted hyaena den mapping

We located spotted hyaena dens that were active at the time of the survey across the district with the help of villagers, extension workers and wildlife experts; we probably mapped all dens in the study area since it is unlikely that any den would go undetected or unreported by people, even away from villages. We recorded habitat type, vegetation cover, grazing intensity, distance to the nearest watercourse/river, village and urban center, den condition (active or abandoned; determined by the presence of fresh scats, fresh food remnants and fresh traces), human disturbance, protection circumstance as well as livestock damage in and around the spotted hyaena dens. We examined fragments of bone, tooth, hoof, horn and skin found within the dens and outside within a 10 m radius and identified them in situ whenever possible (implying surface collection of all visible remains). We recorded GPS coordinates of the exact locations of all spotted hyaena dens.

We used the JMP 5 software package (SAS Institute, Cary, NC) for statistical analyses. Nonparametric Wilcoxon tests were used to compare spotted hyaena and golden wolf encounters along road transects during the wet and dry seasons. We correlated spotted hyaena and golden wolf encounters and distance traveled along road transects from high human density areas.

Results

Relationship between carnivore and human abundance

We encountered 871 spotted hyaenas (441 dry season, 430 wet season) and 307 golden wolves (215 dry season, 92 wet season) in 66 survey nights (41 dry season, 25 wet season) along four road transects. The number of observations was significantly higher in the dry season for golden

wolf ($p < 0.001$) compared to the wet season (Table 1). The numbers of observations for spotted hyaena in the dry and wet seasons were similar ($p > 0.05$) (Table 1). The mean distances from the viewer to spotted hyaena and golden wolf were approximately 88 and 45 m (dry season) and 69 and 38 m (wet season), respectively. Densities of spotted hyaena and golden wolf increased with increasing anthropogenic influence (presence of organic waste and livestock) in all roads transect. The number of spotted hyaena and golden wolf encountered was higher with proximity to towns where human density was higher (Fig 2, 3). Wukro, Agula, Adigudom, and Meremieti were towns with very high human density in the section of transect (Fig 2).

Response range and response probability

We conducted 18 calibration experiments for spotted hyaenas and one calibration experiment for golden wolves and found a total of 102 spotted hyaenas and 15 golden wolves. We used one independent calibration experiment to estimate golden wolf response range and probability where 12 out of 15 golden wolves responded at a radius of 2.5 km. No response from golden wolves was observed beyond that. It was quite difficult to get more independent experiments for golden wolves due to the mobility of the animal. Maximum response radius was 2.8 km for spotted hyaenas (Table 2). Thus, each calling station covered an area of 24.62 km² for spotted hyaena and 19.63 km² for golden wolf. Response probabilities were 0.83 ± 0.17 (mean \pm SD) for spotted hyaena ($n=15$) and 0.8 for golden wolf ($n=1$), respectively (Table 2). The majority (67 %) of the calibration experiments were conducted in open agricultural areas. We assumed that both the response range and the response probability were the same in all habitats irrespective of the landscape of the district.

Abundance and density of predators

A total of 562 spotted hyaena and 63 golden wolf responded to the calling stations; the majority of spotted hyaena (55 %, n=309) were found in human settlements and waste dumping areas while the majority of golden jackals (67 % n=42) were found in human dominated open agricultural areas (Table 3). Abundance of spotted hyaenas and golden wolves increased with increasing anthropogenic influence (presence of human organic waste and livestock). The mean number of spotted hyaenas and golden wolves responding per calling station were 19 (range 0-150) and 3 (0-7), respectively. Spotted hyaena and golden wolf were seen at 30 (88 %) and 24 (70.6 %) of the calling stations during the survey, respectively. The highest number of spotted hyaenas (150) was in a garbage dumping site; this was the only calling station where it was difficult to accurately count them and 150 is a conservative estimate.

Approximately 837 km² (59 %) and 667 (47 %) km² were covered by 34 calling stations for estimating spotted hyaena and golden wolf, respectively. Therefore, assuming a uniform response range and probability across the district, the overall spotted hyaena density was 0.8 spotted hyaena/km² for a total of 1,145 spotted hyaena (17 ± 26) (mean ± SD). Golden wolf density over the entire district was estimated as 0.12 golden wolf /km² or a total population estimate of 166 golden wolf (1.6 ± 1.8) (mean ± SD). Spotted hyaena density was 74 times higher in the garbage dumping site than in natural areas. Similarly, golden wolf density was highest in the garbage dumping site and lowest in the natural bushy area (Table 3).

Spotted hyena den mapping

A total of 40 active spotted hyaena dens at the time of the survey were assessed and most (85 %) were located approximately 1 to 2 km from a nearby village (Table 4). They were primarily located in close proximity to high human density areas (1.49 km ± 0.94) and the nearest water source (1.15 km ± 1.48) (mean ± SD). Each den had more than one entrance and almost all the

cave and burrow den types had sections. Vegetation cover for most (72.5 %) of the dens was relatively very high and most dens were located (62.5 %) less than 1 km from the nearest stream (Table 4).

A total of 1507 (96 % unidentified) pieces of prey remains were found, and the rate of accumulation varied based upon the type of den (Fig 4, 5). The prey remains accumulated in caves and church forest were highest (Fig 5). Very few dens were found to contain large concentrations of bone: we found 24 cattle skulls, 10 donkey skulls, 534 bone fragments and 50 tooth fragments at a church forest located approximately 3 km from the waste dumping place of Mekelle city. Prey remnant counts ranged from 0 to 584. The church forest dens (n=6) had the highest bone fragment accumulation, with an average of 95.7 bone fragments. The cave dens (n=4) had an average of 93.8 bone fragments each, the burrow dens (n=2) had an average of 90 bone fragments each, and the area enclosure dens (n=6) had an average of 34.4 bone fragments each. The bush dens (n=4) had the lowest bone fragment accumulation, with an average of 7.3 bone fragments each. Most (91.7 %) of the fragments identified were livestock remains that included 27 skulls, five horns, three hooves of cattle and ten skulls of donkeys. Four teeth and two jaws of camels, and four hooves of sheep, goat and donkey were identified.

Discussion

Relationship between carnivores and human inhabitants

Our results of road transect monitoring, calling station survey and den mapping demonstrate a positive relationship between human and spotted hyaena density in prey depleted human-dominated landscapes. Spotted hyaena dens were located in close proximity to high human density areas. This relationship is likely driven by food resources such as livestock and waste. In contrast to Woodroffe (2000) who found a negative relationship between carnivore survival and

human population density, we found a positive relationship between spotted hyaena, golden wolf and human concentrations in areas accompanied by high levels of human activity. The spatial dispersion of human settlements and their density across the landscapes might influence the density and occurrence of spotted hyaena and golden wolf. This suggests a unique preference of spotted hyaena and golden wolf to very high human density areas over areas with lower human density. There are two possible explanations: first, the local communities have a positive attitude towards spotted hyaena because spotted hyaenas are important in sanitizing the environment by foraging on household waste and removing potentially infected organic waste from human settlements (Abay et al. 2011; Yirga et al. 2015a,b) and as a consequence spotted hyaenas are more abundant in areas where human density is relatively high. Second, spotted hyaena survival in the area is almost entirely dependent on anthropogenic food around human concentrations (Yirga et al. 2012) because the natural prey base of the area is highly depleted. This unique coexistence of spotted hyaena and people is not found elsewhere in Africa and has not been reported for other large predators. Our findings support the conclusion that spotted hyena in Tigray region depend exclusively on anthropogenic resources and prefer anthropogenic over natural landscapes (Abay et al. 2011; Yirga et al. 2014). Our findings are in stark contrast to various studies indicating that carnivores including spotted hyaena avoid areas of elevated human activity mainly due to persecution and retaliation (Treves and Karanth 2003; Colyn et al. 2004; Reed and Merenlender 2008; Baker et al. 2008; Van Meter et al. 2009; Packer et al. 2009; Croes et al. 2011; Burton et al. 2011).

Response range, response probability and abundance

The sample size for estimating response probability and range for golden wolf is quite low (one calibration experiment) and is due to the mobility of the animals. Golden wolves were highly

mobile in the landscape and this creates a problem for a vehicle to stay with the animals to record their reaction during the calibration experiment. However, we feel that the estimated response range and response probability of golden wolves is appropriate because the number of golden wolves we encountered during the calibration experiment was high enough to estimate response range and probability.

Response probability and range for spotted hyaena and golden wolf in human-dominated landscapes has not previously been reported. There are no estimates of response probability for predators in human-dominated areas and no comparisons of responses in protected and unprotected populations (Ogutu et al. 2005). Few attempts have been made to calibrate calling stations despite wide utilization to establish abundance, density and distribution of carnivores (Kruuk 1972; Mills 1985; Sillero-Zubiri and Gottelli 1992; Creel and Creel 1996; Mills and Gorman 1997; Ogutu and Dublin 1998).

The estimated maximum response range of 2.8 km for spotted hyaena in the human-dominated landscape was somewhat shorter than either the 3.2 km obtained by Mills et al. (2001) for Kruger spotted hyaenas, 3.7 km estimated by Creel and Creel (1996) for spotted hyaenas in Selous Game Reserve or 4 km estimated by Ogutu et al. (2005) for spotted hyaenas in the Mara ecosystem of Kenya, but similar to the 3 km estimated by Maddox (2003) for Kruger spotted hyaenas. A number of factors can affect response range such as topography, vegetation cover, battery used, volume of mp3 and megaphone, and human density.

The estimated response rate of 0.83 for spotted hyaena in the Enderta district was somewhat similar to the 0.88 estimated by Maddox (2003) in the Serengeti and 1.0 estimated by Creel and Creel (1996) in Selous Game Reserve in Tanzania. The probability of responding to calling stations might be inversely related to response range (Ogutu et al. 2005). The response

probability of spotted hyaena in the Mara ecosystem of Kenya was estimated to be 0.583 by Ogotu et al. (2005), similar to the estimate of 0.61 by Mills et al. (2001). Response probabilities were similar in protected and in unprotected areas with human land uses (Ogotu et al. 2005). However, Boydston et al. (2003) noted that predator behavior can be affected by the presence of people.

Spotted hyaena and golden wolf abundance and density varied considerably among different locations of calling stations in the study area. Human settlements and garbage dumping areas supported a relatively high density of spotted hyaena among others perhaps due to availability of scavengeable food. The high number of spotted hyaenas at the garbage dumping areas suggests that spotted hyaenas may occupy different group territories around different den sites but then commute across other territories to a source of super-rich food. Spotted hyaenas are frequent visitors to garbage dumps and other sources of scavengeable food (Horwitz and Smith 1988; Leakey et al. 1999).

Spotted hyaena den selection

In the current study bush dens had the lowest bone fragment accumulation suggesting that spotted hyaena bone accumulation around bush dens is much more limited than around cave and church forest den locations. In this study 1507 pieces of prey remains predominantly occurred in caves and church forests. This is perhaps due to the permanence of the areas; the longer the time span of the den the more bones will accumulate (Pokines and Peterhans 2007). Surface bone concentration around a spotted hyaena cave den located in Amboseli National Park, Kenya, reached up to 75 bone pieces per m²; from 2,017 specimens, 56 individuals were identified (Hill 1983, 1989). Surface bone concentration around burrow dens is less compared to around caves.

Caves shelter adult spotted hyaenas and encourage bone accumulation while burrow dens are often more difficult to enter and are used to deposit juveniles (Pokines and Peterhans 2007).

Spotted hyaenas in our study area transport carcass portions into church forests. We found 24 skulls of cattle and ten skulls of donkey in a church forest that was approximately 3 km from the garbage dumping area of Mekelle. It is more likely that these skulls of livestock are collected from the garbage dumping area where spotted hyaenas are known to scavenge every night. This indicates that spotted hyaenas transport bones into dens, perhaps for their offspring. Juvenile spotted hyaenas are vulnerable and do not leave the den until 9–12 months old (Lansing et al. 2009). Other studies have shown significant accumulation of bones at spotted hyaenas dens (Bunn 1983; Hill 1989; Boydston et al. 2006). Human remains in a spotted hyaenas cave den located approximately 3.2 km from a hospital cemetery were found in Kajiado, Kenya (Sutcliffe 1969, 1970). This shows that spotted hyenas can also accumulate human remains from a nearby cemetery; there were no reports of spotted hyaenas killing people.

In this study a few dens were found to contain large concentrations of bone, and most of these dens were located within 1 to 3 km from the garbage dumping area of Mekelle. This suggests that the bones accumulated were collected from waste dumps, not obtained from predation. Dens were predominantly located along water resources and villages. Other studies also found that spotted hyaena dens were located close to permanent water resources (Mills 1990; Boydston et al. 2003; Boydston et al. 2006). The location of water sources is suggested to influence movements of spotted hyaenas in arid areas (Tilson and Henschel 1986; Cooper 1989). In the dry season, spotted hyaenas in the Serengeti National Park, Tanzania selected dens located in the direction of large prey herds (Kruuk 1972). These confirm that spotted hyaenas are restricted by water and food distribution. In our study spotted hyaena dens were located with significant

proximity to human settlements where spotted hyaenas have access to human waste and livestock.

Conclusion

Our assessment of road transects, calling station surveys and den mapping have shown spotted hyaena-human relationship and distribution patterns along human-dominated areas. Our results demonstrate the value of road transect monitoring which provides reliable information to establish relationships between spotted hyaena, golden wolf and human concentrations in anthropogenic landscapes. Road transect surveys are relatively expensive and time consuming. Calling stations have the potential to offer much more reliable information on spotted hyaena and golden wolf abundance and density comparatively at low costs and short time, and would also be important to assess distribution patterns. Den mapping provides insights about den selection and location. It is through these more intensive studies that we establish the relationship between spotted hyaena, golden wolf and human inhabitants in anthropogenic landscapes. We conclude that there is a positive relationship between spotted hyaena, golden wolf and human abundance and density in prey depleted anthropogenic landscapes in northern Ethiopia. Our findings show that spotted hyaena persist at very large densities with great affinity for high human density locations, a case of exceptional coexistence of humans and carnivores, both at large densities.

Acknowledgments

We are grateful to the Institutional Collaboration between Mekelle University and the Norwegian University of Life Sciences for financial assistance. We thank MU-IUC project and VLIR UOS for logistical support. We thank Mr. Solomon Tesfay, Mr. Solomon Amare and Ms. Fana Welde-Anenya for their respective contributions.

Ethical statement

- The manuscript has not been submitted to more than one journal for simultaneous consideration.
- The manuscript has not been published previously (partly or in full).
- A single study is not split up into several parts to increase the quantity of submissions and submitted to various journals or to one journal over time (e.g. “salami-publishing”).
- No data have been fabricated or manipulated
- No data, text, or theories by others are presented as if they were the author’s own (“plagiarism”).
- Proper acknowledgements to other works are given.

References

- Abay GY, Bauer H, Gebrihiwot K, Deckers J (2011) Peri-urban spotted hyena (*Crocuta crocuta*) in northern Ethiopia: diet, economic impact and abundance. *European Journal of wildlife Research*, 57: 759-765.
- Aerts R, Negussie A, Maes W, November E, Hermy M, Muys B (2007) Restoration of dry Afromontane forest using pioneer shrubs as nurse plants for *Olea europaea* ssp. *Euspidata*. *Restoration Ecology*, 15: 129-138.
- Agarwala M, Kumar S, Treves A, Naughton-Treves L (2010) Paying for wolves in Solapur, India and Wisconsin, USA: comparing compensation rules and practice to understand the goals and politics of wolf conservation. *Biological Conservation*, 143: 2945–2955.

- Alemayehu WE (2007) *Ethiopian Church Forests: opportunities and challenges for restoration*.
PhD thesis, Wageningen University, Wageningen, The Netherlands.
- Athreya V (2012) Conflict resolution and leopard conservation in a human-dominated landscape.
PhD thesis, Manipal University, India.
- Baker PJ, Boitani L, Harris S, Saunders G, White PCL (2008) Terrestrial carnivores and human food production: impact and management. *Mammal Review*, 38: 123–166.
- Basille M, Herfindal I, Santin-Janin H, Linnell JDC, Odden J, Andersen R, Arild Høgda K, Gaillard J (2009) What shapes Eurasian lynx distribution in human dominated landscapes: selecting prey or avoiding people? *Ecography*, 32: 683–691.
- Bauer H (2007) Status of large carnivores in Bouba Ndjida National Park, Cameroon. *African Journal of Ecology*, 45: 448-450.
- Blanco JC, Cortés Y (2007) Dispersal patterns, social structure and mortality of wolves living in agricultural habitats in Spain. *Journal of Zoology*, 273: 114–124.
- Boydston EE, Kapheim KM, Holekamp KE (2006) Patterns of den occupation by the spotted hyena (*Crocuta crocuta*). *African Journal of Ecology*, 44: 77-86.
- Boydston EE, Kapheim KM, Watts HE, Szychman M, Holekamp KE (2003) Altered behavior in spotted hyenas associated with increased human activity. *Animal Conservation*, 6: 1–10.
- Boydston EE, Morelli TL, Holekamp KE (2001) Sex differences in territorial behavior exhibited by the spotted hyena (*Crocuta crocuta*) *Ethology*, 107:369-385.
- Bunn HT (1983) Comparative analysis of modern bone assemblages from a San hunter-gatherer camp in the Kalahari Desert, Botswana, and from a spotted hyena den near Nairobi,

- Kenya. *Animals and Archaeology: 1. Hunters and Their Prey*, Series 163. BAR International (eds J. Clutton-Brock & C. Grigson), pp. 143-148. Oxford, United Kingdom.
- Burton AC, Sam MK, Kpelle DG, Balangtaa C, Buedi EB, Brashares JS (2011) Evaluating persistence and its predictors in a West African carnivore community. *Biological Conservation*, 144: 2344–2353.
- Cardillo M, Purvis A, Sechrest W, Gittleman JL, Bielby J, Mace GM (2004) Human population density and extinction risk in the world's carnivores. *PLoS Biology*, 2: 909–0914.
- Colyn M, Dufour S, Condé PC, Van Rompaey H (2004) The importance of small carnivores in forest bushmeat hunting in the Classified Forest of Diécké, Guinea. *Small Carnivore Conservation*, 31: 15–18.
- Cooper SM (1989) Clan sizes of spotted hyaenas in the Savuti Region of the Chobe National Park, Botswana. *Botswana Notes and Records*, 21:121-131.
- Cooper SM, Holecamp KE, Smale L (1999) A seasonal feast: long-term analysis of feeding behavior in the spotted hyena (*Crocuta crocuta*). *African Journal of Ecology*, 37: 149-160.
- Creel S, Creel NM (1996) Limitation of African wild dogs by competition with larger carnivores. *Conservation Biology*, 10: 526-538.
- Croes BM, Funston PJ, Rasmussen G, Buij R, Saleh A, Tumenta PN, De Iongh HH (2011) The impact of trophy hunting on lions (*Panthera leo*) and other large carnivores in the Bénoué Complex, northern Cameroon. *Biological Conservation*, 144: 3064–3072.

- Frank, L.G. (1986). Social organization of the spotted hyena (*Crocuta crocuta*). II. Dominance and reproduction. *Animal Behavior*, 34: 1510-1527.
- Fuller TK, Sievert PR (2001) Carnivore demography and the consequences of changes in prey availability. *Carnivore conservation* (eds J.L. Gittleman, S.M. Funk, D. Macdonald & R.K. Wayne), pp. 163–178. Cambridge University Press, Cambridge, United Kingdom.
- Gaubert P, Bloch C, Benyacoub S, Abdelhamid A, Pagani P, Djagoun CAMS, Couloux A, Dufour S (2012) Reviving the African wolf *Canis lupus lupaster* in North and West Africa: a mitochondrial lineage ranging more than 6,000 km wide. *PLoS ONE* 7: e42740.
- Gehrt SD, Riley SPD (2010) Coyotes (*Canis latrans*). *Urban carnivores: ecology, conflict, and conservation* (eds S.D. Gehrt, S.P.D. Riley & B.L. Cypher), pp. 79–95. The Johns Hopkins University Press, Baltimore, Maryland, USA.
- Giannatos G (2004) Conservation action plan for the golden jackal *Canis aureus* L. in Greece. WWF Greece, Athens.
- Henschel JR, Skinner JD (1991) Territorial behavior by a clan of spotted hyenas *Crocuta crocuta*. *Ethology*, 88: 223-235.
- Hill A (1983) Hyenas and early hominids. *Animals and Archaeology: 1. Hunters and Their Prey*, Series 163. BAR International (eds J.C. Clutton-Brock & C. Grigson), pp. 87-92. Oxford, United Kingdom.
- Hill A (1989) Bone modification by modern spotted hyenas. *Bone Modification*. (eds R. Bonnicksen & M.H. Sorg), pp. 169-178. Center for the Study of the First Americans, Orono, Maine.

- Holekamp KE, Boydston EE, Smale L (2000) Group travel in social carnivores. *On the Move: How and Why Animals Travel in Groups* (eds S. Boinski & P.A. Garber), pp. 587–627. University of Chicago Press, Chicago, Illinois.
- Holekamp KE, Sakai ST, Lundrigan BL (2007) Social intelligence in the spotted hyena (*Crocuta crocuta*). *Philosophical Transactions of the Royal Society, London B*, 362: 523–538.
- Holekamp KE, Smale L (1998) Behavioral development in the spotted hyena. *Bioscience*, 48: 997–1005.
- Horwitz L, Smith P (1988) The effects of striped hyena activity on human remains. *Journal of Archeological Science*, 15: 471-481.
- Jhala Y, Moehlman PD (2008) *Canis aureus*. The IUCN Red List of Threatened Species. Version 2015.2. <www.iucnredlist.org>. Downloaded on 11 July 2015.
- Jhala YV, Moehlman PD (2004) Golden jackal *Canis aureus*. In: Sillero-Zubiri C, Hoffmann M, Macdonald D (eds) *Canids: Foxes, Wolves, Jackals and Dogs Status Survey and Conservation Action Plan*. IUCN/SSC Canid Specialist Group Gland, Switzerland, pp 156–161.
- Karanth KU, Nichols JD, Kumar NS, Link WA, Hines, JE (2004) Tigers and their prey: predicting carnivore densities from prey abundance. *Proceedings of the National Academy of Sciences of the USA*, 14: 4854-4858.
- Khorozyan IG, Malkhasyan AG, Abramov AV (2008) Presence-absence surveys of prey and their use in predicting leopard (*Panthera pardus*) densities: a case study from Armenia. *Integrative Zoology*, 3: 322-332.

- Koepfli KP, Pollinger J, Godinho R, Robinson J, Lea A, Hendricks S, Schweizer RM, Thalmann O, Silva P, Fan Z, et al. (2015) Genome-wide evidence reveals that African and Eurasian golden jackals are distinct species. *Current Biology*, 25: 2158–2165.
- Kolowski JM, Holekamp KE (2009) Ecological and anthropogenic influences on space use by spotted hyenas. *Journal of Zoology*, 277: 23-36.
- Kruuk H (1972) *The Spotted Hyena: A Study of Predation and Social Behavior*. The University of Chicago Press: Chicago, USA.
- Lansing SW, Cooper SM, Boydston EE, Holekamp KE (2009) Taphonomic and zooarchaeological implications of spotted hyena (*Crocuta crocuta*) bone accumulations in Kenya: A modern behavioral ecological approach. *Paleobiology*, 35: 289–309.
- Leakey LN, Milledge SAH, Leakey SM, Edung J, Haynes P, Kiptoo DK, McGeorge A (1999) Diet of striped hyena in northern Kenya. *African Journal of Ecology*, 37: 314-326.
- Linnell JDC, Swenson J, Andersen R (2001) Predators and people: conservation of large carnivores is possible at high human densities if management policy is favorable. *Animal Conservation*, 4: 345–349.
- Luis Llanaez L, López-Bao JV, Sazatornil V (2012) Insights into wolf presence in human dominated landscapes: the relative role of food availability, humans and landscape attributes. *Diversity and Distributions*, 18: 459–469.
- Maddox TM (2003) *The ecology of cheetahs and other carnivores in pastoralist-dominated buffer zone*. Ph.D Thesis. University College London and Institute of Zoology, Zoological Society of London, London, United Kingdom.

- Mills MG, Juritz JM, Zuccini W (2001) Estimating the size of spotted hyena (*Crocuta crocuta*) populations through playback recordings allowing for non-response. *Animal Conservation*, 4: 335-343.
- Mills MGL (1985) Hyena survey of Kruger National Park, August-October, 1984 IUCN-SSC Hyena Specialist Group, *Newsletter*, 2: 15–25.
- Mills MGL (1990) Kalahari hyenas: *the comparative behavioral ecology of two species*. Unwin Hyman Pres, London, United Kingdom.
- Mills MGL, Gorman ML (1997) Factors affecting the density and distribution of wild dogs in the Kruger National Park. *Conservation Biology*, 11: 1397-1406.
- Ogutu JO (1994) Test of a call-in technique for estimating lion (*Panthera leo*, Linnaeus 1758) population size in the Masai Mara National reserve, Kenya, Department of Wildlife Management, Moi University, p. 69.
- Ogutu JO, Bhola N, Reid R (2005) The effects of pastoralism and protection on the density and distribution of carnivores and their prey in the Mara ecosystem of Kenya. *Journal of Zoology (London)*, 265: 281-293.
- Ogutu JO, Dublin HT (1998) The response of lions and spotted hyenas to sound playbacks as a technique for estimating population size. *African Journal of Ecology*, 36: 83–95.
- Packer C, Kosmala M, Cooley HS, Brink H, Pintea L, Garshelis D, Purchase G, Strauss M, Swanson A, Balme G, Hunter L, Nowell K (2009) Sport Hunting, Predator Control and Conservation of Large Carnivores. *PLoS ONE*, 4: e5941.

- Pokines JT, Kerbis Peterhans JC (2007) Spotted hyena (*Crocuta crocuta*) den use and taphonomy in the Masai Mara National Reserve, Kenya. *Journal of Archaeological Sciences*, 34: 1914-1931.
- Reed SE, Merenlender AM (2008) Quiet, non-consumptive recreation reduces protected area effectiveness. *Conservation Letters*, 1: 146–154.
- Rueness EK, Asmyhr MG, Sillero-Zubiri C, Macdonald DW, Bekele A, Atickem A, Stenseth, NC (2011) The cryptic African wolf: *Canis aureus lupaster* is not a golden jackal and is not endemic to Egypt. *PLoS ONE* 6: e16385.
- Šálek M, Červinka J, Banea OC, Krofel M, Čirović D, Selanec I, Penezić A, Grill S, Riegert J (2014) Population densities and habitat use of the golden jackal (*Canis aureus*) in farmlands across the Balkan Peninsula. *European Journal of wildlife Research*, 60:193–200.
- Sara AA (2010) *Mitigating drought: Policy Impact Evaluation A case of Tigray region, Ethiopia*, Msc. Thesis, University of Twente, the Netherlands.
- Sillero-Zubiri C, Gottelli D (1992) Population ecology of spotted hyenas in an equatorial mountain forest. *African Journal of Ecology*, 30: 292–300.
- Sillero-Zubiri C, Hoffmann M, Macdonald DW (2004). *Canids: Foxes, Wolves, Jackals and Dogs: Status Survey and Conservation Action Plan*, 2nd ed. IUCN Canid Specialist Group, Gland, Switzerland and Cambridge, UK.
- Smith JE, Holekamp KE (2010) *Spotted Hyenas* Michigan State University, East Lansing, MI, USA a 2010 Elsevier Ltd. All rights reserved.
- Stoyanov S (2012) Golden jackal (*Canis aureus*) in Bulgaria. Current status, distribution, demography and diet. International symposium on hunting, “Modern aspects of

- sustainable management of game population” Zemun-Belgrade, Serbia, 22–24 June, 2012. pp 48–56
- Sunquist ME, Sunquist F (2001) Changing landscapes: consequences for carnivores. In *Carnivore conservation*: 399 – range and area reached by animal vocalizations. *Journal of experimental Biology*, 200: 421-431.
- Sutcliffe A (1970) Spotted hyena: crusher, gnawer, digester and collector of bones. *Nature*, 227: 1110-1113.
- Sutcliffe AJ (1969) Adaptations of spotted hyenas to living in the British Isles. *Mammal Society Bulletin*, 31: 1-4.
- Tilson RT, Hamilton WJ (1984) Social dominance and feeding patterns of spotted hyenas. *Animal Behavior*, 32: 715-724.
- Tilson RT, Henschel JR (1986) Spatial arrangement of spotted hyena groups in a desert environment, Namibia. *African Journal of Ecology*, 24: 173-180.
- Treves A, Karanth KU (2003) Human-carnivore conflict and perspectives on carnivore management worldwide. *Conservation Biology*, 17: 1491–1499.
- Van Meter PE, French JA, Dloniak SM, Watts HE, Holekamp KE (2009) Fecal glucocorticoids reflect socio-ecological and anthropogenic stressors in the lives of wild spotted hyena. *Hormones and Behavior*, 55: 329–337.
- Woodroffe R (2000) Predators and people: using human densities to interpret declines of large carnivores. *Animal Conservation*, 3: 165–173.
- Woodroffe R, Ginsberg JR (1998) Edge effects and the extinction of populations inside protected areas. *Science*, 280: 2126–2128.

- Yirga G, De Iongh HH, Leirs H, Gebrehiwot K, Deckers J, Bauer H (2012) Adaptability of large carnivores to changing anthropogenic food sources: diet change of spotted hyena (*Crocuta crocuta*) during Christian fasting period in northern Ethiopia. *Journal of Animal Ecology*, 81: 1052–1055.
- Yirga G, Ersino W, De Iongh H.H, Leirs H, Gebrehiwot K, Deckers J, Bauer H (2013) Spotted hyena (*Crocuta crocuta*) coexisting at high density with people in Wukro district northern Ethiopia. *Mammalian Biology*, 78: 193– 197.
- Yirga G, Imam E, De Iongh H.H, Leirs H, Kiros S, G/ Yohannes T, Teferi M, Bauer H (2014) Local spotted hyena abundance and community tolerance of depredation in human-dominated landscapes in Northern Ethiopia. *Mammalian Biology*, 79: 325–330.
- Yirga G, Leirs H, De Iongh H.H, Asmelash T, Gebrehiwot K, Deckers J, Bauer H (2015a) Spotted hyena (*Crocuta crocuta*) concentrate around urban waste dumps across Tigray, northern Ethiopia. *Wildlife Research*, 42: 563-569.
- Yirga, G., De Iongh H.H., Leirs H, Gebrihiwot, K., Deckers, J., Bauer, H. (2015b) Food base of the spotted hyena (*Crocuta crocuta*) in Ethiopia, *Wildlife Research*, 42: 19–24.




Figure 1 Map of Ethiopia showing the location of Tigray and map of Enderta district showing the locations of calling stations and spotted hyaena dens represented by circles and stars, respectively. The dot highlighted in map of Enderta district represents Mekelle City.






Figure 2 Road transects showing relationship between spotted hyaena and human abundance in northern Ethiopia. Names under the x-axis are towns found in that section of transect, no name means that section was purely rural.




Figure 3 Spatial distribution of golden wolf and human along four road-transects in 2012 in northern Ethiopia. Names under the x-axis are towns found in that section of the transect, no name means that section was purely rural.




Figure 4 Total prey remnants found at 22 spotted hyaena dens in 2012 in Enderta district, northern Ethiopia




Figure 5 Total and average prey remnants at different spotted hyaena den types (n=22) in 2012 in Enderta district, northern Ethiopia

Table 1 Spotted hyaena and golden wolf counts during wet and dry season in four road-transects in 2012 in Enderta district, northern Ethiopia

| Season | Transect | Length in km | Hyaena | Golden wolf | Frequency | Mean viewing distance (in m) |
|--------|------------------|--------------|------------|-------------|-----------|------------------------------|
| Dry | Mekelle-Adikolo | 23 | 157 | 25 | 11 | 90 |
| | Mekelle-Adigudom | 38 | 86 | 62 | 11 | 96 |
| | Mekelle-Wukro | 36 | 140 | 96 | 10 | 97 |
| | Quiha-Aragure | 19 | 58 | 32 | 9 | 70 |
| | Total | | 441 | 215 | 41 | |
| Wet | Mekelle-Adikolo | 23 | 299 | 25 | 10 | 59 |
| | Mekelle-Adigudom | 38 | 39 | 25 | 5 | 72 |
| | Mekelle-Wukro | 36 | 43 | 29 | 4 | 59 |
| | Quiha-Aragure | 19 | 49 | 13 | 6 | 87 |
| | Total | | 430 | 92 | 25 | |

* The Frequency column refers to the number of survey nights

Table 2 Number of spotted hyaenas that responded during calibration experiments at various distances from the calling stations in Enderta district, northern Ethiopia

| Distance (km) | Number played to | Number responding | Response probability |
|----------------------|-------------------------|--------------------------|-----------------------------|
| 2 | 6 | 3 | 0.5 |
| 2 | 11 | 10 | 0.9 |
| 2 | 1 | 1 | 1 |
| 2 | 7 | 6 | 0.9 |
| 2.3 | 6 | 5 | 0.8 |
| 2.5 | 5 | 5 | 1 |
| 2.5 | 15 | 14 | 0.9 |
| 2.5 | 3 | 3 | 1 |
| 2.5 | 5 | 4 | 0.8 |
| 2.5 | 3 | 3 | 1 |
| 2.6 | 4 | 3 | 0.7 |
| 2.8 | 6 | 5 | 0.8 |
| 2.8 | 3 | 2 | 0.7 |
| 2.8 | 2 | 2 | 1 |
| 2.8 | 6 | 3 | 0.5 |
| 3 | 6 | 0 | 0 |
| 3.5 | 8 | 0 | 0 |
| 4 | 5 | 0 | 0 |

Table 3 The total number of spotted hyaena and golden wolves (n) and estimates of the expected number of individuals (μ) in each location, estimated spotted hyaena and golden wolf population size (P_s) and population density (D) (number/km²) for each of the five locations of calling stations in Enderta district, northern Ethiopia.

| Location | Calling stations | Spotted hyaena | | | | | Golden wolf | | | | |
|------------------------|------------------|-------------------------|------------|-------------|------------|------------|-------------------------|-----------|------------|-----------|-------------|
| | | Area (km ²) | n | μ | P_s | D | Area (km ²) | n | μ | P_s | D |
| Open agricultural area | 20 | 492.4 | 198 | 11.9 | 239 | 0.49 | 392.6 | 42 | 2.6 | 52 | 0.13 |
| Human settlement | 9 | 221.6 | 159 | 21.3 | 192 | 0.87 | 176.7 | 8 | 1.1 | 10 | 0.06 |
| Area enclosure | 2 | 49.2 | 49 | 29.5 | 59 | 1.2 | 39.3 | 7 | 4.4 | 9 | 0.23 |
| Bushy natural area | 2 | 49.2 | 6 | 3.6 | 7 | 0.1 | 39.3 | 1 | 0.6 | 1 | 0.03 |
| Garbage dumping site | 1 | 24.6 | 150 | 181 | 181 | 7.4 | 19.6 | 5 | 5 | 6 | 0.31 |
| Total | 34 | 837.1 | 562 | 19.9 | 677 | 0.8 | 667.5 | 63 | 2.3 | 79 | 0.12 |

*n is the total number of hyaenas and golden wolves that responded in each location of call-ups. μ is the expected number of hyaenas and golden wolves within the response range. Response range and probability for hyaenas and golden wolves were 2.8 and 2.5 km and 0.83 and 0.8, respectively. $u_s = y_s/S_s$ and $P_s = n/\text{response probability}$. *NT = (AH/AS)u where AH =the area of the entire district (1416 km²) and AS =the area sampled around a call-in station (24.62 km² for hyaenas and 19.63 km² for golden wolf) and u (19.9 for hyaena and 2.3 for golden wolf) and this gives the final population numbers of spotted hyaena (1,145) and golden wolf (166).

Table 4 Characteristics of spotted hyaena dens (n=40) located in human-dominated landscapes in Enderta district, northern Ethiopia

| Items | Category | Count | Percentage |
|-------------------------------------|-----------------|---------------|-------------------|
| Distance from the nearest town | <=5km | 22 | 55 |
| | 6-10km | 10 | 25 |
| | >=11km | 8 | 20 |
| Distance from the nearest hyena den | <=3km | 28 | 70 |
| | >=4km | 12 | 30 |
| Distance from the nearest village | 1-2km | 34 | 85 |
| | 3-4km | 6 | 15 |
| Distance from the nearest stream | <=1km | 25 | 62.5 |
| | 1.5-2km | 10 | 25 |
| | >=3km | 5 | 12.5 |
| Grazing intensity | None | 16 | 40 |
| | Low | 6 | 15 |
| | Medium | 0 | 0 |
| | High | 4 | 10 |
| | Very high | 14 | 35 |
| Human disturbance | None | 12 | 30 |
| | Low | 6 | 15 |
| | Medium | 2 | 5 |
| | High | 4 | 10 |
| | Very high | 16 | 40 |
| Vegetation cover | Low | 6 | 15 |
| | Medium | 2 | 5 |
| | High | 3 | 7.5 |
| | Very high | 29 | 72.5 |
| | Habitat | Church forest | 11 |
| Enclosure | | 13 | 32.5 |
| Caves | | 6 | 15 |
| Bush | | 7 | 17.5 |
| Burrow den | | 3 | 7.5 |
| Protection | | Safe | 35 |
| | Unsafe | 5 | 12.5 |
| Livestock attack in and around | Yes | 36 | 90 |
| | No | 4 | 10 |

*Distance was measured by driving from the reference point. Human disturbance and protection of livestock were determined with local people living nearby the dens.