

RESEARCH

Open Access



New approaches to achieve high level enzyme production in *Streptomyces lividans*

Laura Sevillano¹, Erik Vijgenboom², Gilles P. van Wezel², Margarita Díaz^{1*} and Ramón I. Santamaría^{1*}

Abstract

Background: Actinomycetes are saprophytic soil bacteria, and a rich source of industrial enzymes. While some of these enzymes can be produced using well-characterized production platforms such as *Escherichia coli* or *Bacillus subtilis*, *Streptomyces lividans* may be the preferred host for proper folding and efficient secretion of active enzymes. A combination of promoters, signal peptides and hosts were tested in order to obtain the best protein expression in this actinomycete. The xylanase, Xys1, from *S. halstedii*, the α -amylase, Amy, from *S. griseus* and the small laccase, SLAC, from *S. coelicolor* were used as reporters.

Results: The promoters *xysAp* from *S. halstedii* JM8 and *pstSp* from *S. lividans* were the most efficient among those tested. An improvement of 17 % was obtained in xylanase activity when the signal peptide of the α -amylase protein (Amy) of *S. griseus* IMRU3570 was used to direct its secretion. Enhanced expression of SsgA, a protein that plays a role in processes that require cell-wall remodelling, resulted in a improvement of 40 and 70 % of xylanase and amylase production, respectively. Deletion of genes *SLI7232* and *SLI4452* encoding putative repressors of *xysAp* provided improvement of production up to 70 % in the *SLI7232* deletion strain. However, full derepression of this promoter activity was not obtained under the conditions assayed.

Conclusions: *Streptomyces lividans* is a frequently used platform for industrial enzyme production and a rational strain-development approach delivered significant improvement of protein production by this host.

Keywords: *Streptomyces*, Protein production, Xylanase, Amylase, Laccase, Vector optimization

Background

The production of proteins of industrial interest at low cost is one of the main goals of biotechnology. *Escherichia coli* has usually been a preferred host for protein production, for reasons of high expression levels and short fermentation time. However, frequently insoluble inclusion bodies are formed, which frustrates downstream processing [1–3]. In such cases, alternative production platforms are required, and microorganisms of the genus *Streptomyces* have been successfully applied as a good alternative expression platform for heterologous protein production [4–10]. Streptomycetes are aerobic,

filamentous Gram-positive soil bacteria that secrete a wide range of extracellular enzymes, which they require to be able to feast on natural polymers such as starch, cellulose, chitin, mannan or xylan. Their use as a production hosts could overcome some of the problems encountered with other systems: streptomycetes do not readily form inclusion bodies, they have a relatively low level of extracellular protease activity, are well suited to expressing GC-rich genes, and have a high secretion capacity. These properties are very useful for directing the expressed protein in a soluble configuration to the culture supernatant, a major advantage in terms of downstream processing.

Nevertheless, the mycelial lifestyle of actinomycetes results in heteromorphous and viscous cultures, which are unfavourable for industrial fermentation, due to mass-related mechanical stress, heat transfer problems and oxidative stress [11]. Enhanced fragmentation of the mycelia has a major impact on growth and product

*Correspondence: mardi@usal.es; santa@usal.es

¹ Instituto de Biología Funcional y Genómica/Departamento de Microbiología y Genética, Consejo Superior de Investigaciones Científicas (CSIC)/Universidad de Salamanca, C/Zacarías González nº 2, 37007 Salamanca, Spain

Full list of author information is available at the end of the article

formation by these organisms and is therefore expected to have an impact on biotechnological applications requiring *Streptomyces* as the production host [12, 13]. The morphology of liquid-grown mycelia is dictated by external factors (media and fermentation conditions), but also by genetic factors. An important example in this respect is the effect of SsgA, which controls all processes that require remodelling of the cell wall, such as germination, tip growth, branching and septum formation [14, 15]. Overexpression of SsgA not only leads to fragmentation of the mycelia, but also to enhanced secretion, with almost three-fold increase of tyrosinase production by *Streptomyces lividans* during batch fermentation [16].

In this work, we show that selection of strong promoters, modification of signal peptides and use of different *S. lividans* hosts result in improved production of putative industrial proteins such as the xylanase Xys1 from *S. halstedii* JM8, the α -amylase Amy from *S. griseus* IMRU3570 and the small laccase SLAC from *S. coelicolor*.


In addition, two transcriptional repressors, XlnR and BlxR, have been identified in *S. lividans* that take part in the control of the *xysA* promoter activity. This provides new handles for the application and further development of the *xysA* promoter as a tool in enzyme production.

Results

Promoter choice has a drastic effect on protein expression

In order to find suitable promoters to express proteins in *Streptomyces* we tested six *Streptomyces* promoters for their ability to enable strong gene expression. These promoters are active during different phases of the developmental cycle or regulated by different carbon sources. Two of these were strong promoters used frequently in other studies to express proteins in *Streptomyces*: *vsip* from *S. venezuelae* [17] and *ermE**p from *Saccharopolyspora erythraea* [18], the activity of which was compared to that of promoters of four highly expressed genes that are studied extensively in our laboratory. These promoters are the *xysAp* promoter belonging to the *xysA* gene from *S. halstedii* JM8 that encodes the xylanase Xys1 (U41627) [19], the *pstS* promoter (*pstSp*) from *S. lividans* (AJ698727), which drives transcription of the high-affinity phosphate-binding protein (PstS) [20], and the promoter *xylAp* of *xylA* (SCO1169, encoding xylose isomerase) and *glpQp* of *glpQ* (SCO1968, encoding a glycerophosphoryl diester phosphodiesterase) from *S. coelicolor*.

The different promoter sequences were inserted upstream of the xylanase gene *xysA*, which was cloned in a multicopy derivative of the bifunctional *Streptomyces* plasmid pN702GEM3 [21] (see “Methods”). A diagram of the plasmid and of the different constructs is presented in Fig. 1a, b. The constructs pNX24 (*xysAp*) [22], pNUF5



(*pstSp*) [20], pNX26 (*xylAp*), and pNXHid (*glpQp*), pNErmX (*ermE**p), pNvsi (*vsip*) as well as the promoterless control plasmid pNX30 were introduced into *S. lividans* 1326 by protoplast transformation. Transformants were cultured at 28 °C in liquid YES medium with 1 % xylose or YE with 5 %fructose, as indicated in the Methods section and xylanase production was analyzed by SDS-PAGE of culture supernatants after 3, 5 and 7 days. Extracellular processing results in two bands for xylanase, Xys1L and Xys1S, with a mass of 43.5 and 33.7 kDa

respectively [19]. The Xys1S species has lost the C-terminal CBM2 domain but both xylanase species have similar specific activity with oat spelt xylan as substrate [19, 21].

The natural xylanase promoter, *xysAp*, and the *pstS* promoter, *pstSp*, resulted in the highest xylanase production (Fig. 2a lanes 2 and 3). The xylanase production

under the control of *ermE**p was poor (Fig. 2a lane 6) and, although production under the control of *vsip* was high, this was lower than that obtained with *xysAp* and *pstSp* (Fig. 2a lane 7 versus lanes 2 and 3). Maximum xylanase levels were reached after 5 days, the production did not increase by longer incubation (data not shown).

The production of xylanase under control of *xysAp* and *pstSp* was strongly dependent on the culture medium. For *xysAp* (pNX24) the highest protein production was obtained in YES medium supplemented with 1 % xylose (Fig. 2b), while for *pstSp* (pNUF5) the highest production was obtained in YE medium in the presence of 5 % of fructose [20] (Fig. 2c).

To demonstrate that *xysAp* and *pstSp* are generally applicable for enzyme production, they were used to drive the expression of α -amylase (Amy) from *S. griseus* obtaining the plasmids pNXAmy and pNUFAmy respectively (Fig. 2d). High production levels of this enzyme were also obtained demonstrating the potential of these promoters to express other proteins of interest (Fig. 2e).

Modification of the secretion signal peptide improves protein export

Changes in the signal peptide might be advantageous for secretion efficiency and therefore for increasing the production of heterologous proteins [23, 24]. In order to compare the efficiency of the α -amylase (Amy) signal peptide with that of xylanase (Xys1), we replaced the codons for the Xys1 signal peptide in pNX24 (first 45 codons) by those for the corresponding Amy signal peptide (28 codons) [25], resulting in pNXA1 (Table 1, Fig. 3a). A xylanase derivative with the complete Amy signal peptide coding sequence including the first three codons of the mature amylase was also constructed in order to maintain the environment of the amylase signal peptide-cleavage site (pNXA2, Fig. 3a).

Cultures of *S. lividans* harbouring either pNX24 or its derivatives pNXA1 or pNXA2 were grown at 28 °C for 5 days in liquid YES medium supplemented with 1 % xylose; the amount of xylanase in the supernatant was analyzed by SDS-PAGE, xylanase bands intensities were determined with the ImageJ software and the activity quantified using the DNS assay ("Methods") (Fig. 3b, c and Additional file 1: Tables S1, S2). Replacing the original xylanase signal peptide with that of amylase resulted in a similar degree of xylanase secretion (pNXA1, Fig. 3c). However, an increase of 17 % in xylanase activity was obtained when the amylase signal peptide with the three aminoacids extension was used (pNXA2) (Fig. 3c and Additional file 1: Table S2).

Engineered hosts for improved protein yield

The host used as platform for enzyme production is a major determinant for the yield that can be obtained.





Table 1 Plasmids used in this work


Plasmid	Characteristics	Reference
pN702GEM3	<i>E. coli</i> - <i>Streptomyces</i> shuttle vector; contains the pJ702 origin of replication (for <i>Streptomyces</i>), the pUC origin of replication (for <i>E. coli</i>) and a polylinker derived from pGEM3Zf(+) (Promega). It carries the bifunctional Neo/Kan resistance marker from Tn5 (for both <i>E. coli</i> and <i>Streptomyces</i>)	[21]
pNX30	pN702GEM3 derivative. <i>xysA</i> xylanase ORF without any promoter and flanked by transcriptional terminators. Used as negative control	[35]
pNX24	pN702GEM3 derivative. <i>xysA</i> promoter from <i>S. halstedii</i> controls xylanase expression	[22]
pNUF5	pN702GEM3 derivative. <i>pstS</i> promoter from <i>S. lividans</i> controls xylanase expression	[20]
pNErmX	pN702GEM3 derivative. Erythromycin resistance promoter from <i>S. erythraeus</i> controls xylanase expression	This work
pNvsi	pN702GEM3 derivative. <i>vsj</i> promoter from <i>S. venezuelae</i> controls xylanase expression	This work
pNX26	pN702GEM3 derivative. Xylose isomerase promoter from <i>S. coelicolor</i> controls xylanase expression	This work
pNXHid	pN702GEM3 derivative. Promoter of a putative hydrolase (<i>glpQ</i>) from <i>S. coelicolor</i> controls xylanase expression	This work
pNXA1	pNX24 derivative. <i>xysA</i> promoter from <i>S. halstedii</i> controls xylanase expression with amylase signal peptide	This work
pNXA2	pNX24 derivative. <i>xysA</i> promoter from <i>S. halstedii</i> . Controls xylanase expression with the amylase signal peptide with three additional amino acids	This work
pNXAmy	pN702GEM3 derivative. <i>xysA</i> promoter from <i>S. halstedii</i> controls amylase expression	This work
pNUFAmy	pN702GEM3 derivative. <i>pstS</i> promoter from <i>S. lividans</i> controls amylase expression	This work
pEPOS101	pHJL401 derivative harbouring <i>xysAp</i> driving the expression of SLAC (<i>SCO6712</i>)	This work



We explored two ways of improving the host: (1) the effect of host morphology on xylanase production and (2) the effect of putative repressors on *xysA* expression. To change the morphology we selected a *S. lividans* 1326 strain overexpressing SsgA (GSAL1) [16]. The *ssgA* gene is a morphogene that pleiotropically affects growth and cell division [26]; it activates sporulation-specific cell division by ensuring the correct localization of SsgB, which in turn recruits FtsZ to future septum sites [27]. The enhanced expression of *ssgA* leads to strongly enhanced septation in vegetative hyphae, which results in fragmented growth and much wider hyphae, a phenotype that favours protein production as well as secretion [16].

In order to establish the effect of SsgA overexpression on xylanase production, pNX24 was introduced to *S. lividans* 1326 and its derivative overexpressing SsgA (GSAL1). The transformants were grown for 5 days in YES medium with 1 % xylose and the amount of xylanase in the culture supernatants was analysed by SDS-PAGE (Fig. 4a) and the xylanase activity was quantified by the DNS protocol (Fig. 4b). Both, the total amount of xylanase (Xys1L + Xys1S) bands, as determined with ImageJ, and the enzyme activity increased in the SsgA-overexpressing host GSAL1. The band intensities in the SDS-PAGE increased by 70 % while the enzyme activity increased by 40 % (Fig. 4a, b and Additional file 1: Tables S3, S4).

Streptomyces lividans 1326 and GSAL1 were also transformed with pNXAmy harbouring the α -amylase gene



under control of *xysAp*, and the culture supernatants were analysed by SDS-PAGE (Fig. 4c) and the amylase activity of the samples was quantified by the DNS protocol (Fig. 4d). The GSAL1 transformant showed an increase of 45 % in the band intensities while the amylase activity increased by 70 % (Fig. 4c, d and Additional file 1: Tables S5 and S6). Both strains were also transformed with the empty plasmid (pN702GEM3) as a control showing no detectable xylanase or amylase activity in the quantification assays (data not shown).

The second approach was prompted by the observation that maximum enzyme production, controlled by *xysAp*, was obtained after 4 days of growth in liquid media with low accumulation after 24 h (data not shown). This suggests that a repressor(s) bound to *xysAp* might be controlling its expression during the first days of growth. Carbon catabolite repression of *xysAp* was demonstrated previously and the role of different regions of the promoter studied [28]. Among them, a motif 5'-CGAACTTTCG-3', also present in promoter regions of a number of *S. lividans* genes, all of them related to xylan metabolism (data not shown), seems to be important in the glucose control of this promoter [28]. This palindromic sequence is the binding site for BxlR,


a xylobiose responsive repressor in *S. thermoviolaceus* [29]. The homologue of BxlR in *S. lividans* is encoded by SLI7232. In addition to BxlR, another putative repressor, XlnR (SLI4215) has been implicated in xylanase expression in *S. lividans* TK24 [30].

In order to test whether these two repressors may affect the temporal accumulation pattern of an enzyme which gene is controlled by *xysAp*, the gene for the small lactase, SLAC [31] was used as reporter under the control of the *xysAp* (plasmid pEPOS101). SLAC was used as reporter to eliminate potential interference of the induction of host xylanases and xylose metabolism related enzymes in *bxlR* and *xlnR* deletion strains.

Both repressor mutant strains (obtained as indicated in “Methods”) showed an increased SLAC expression compared to the wild type parent, *S. lividans* 1326. The highest increase was observed in the $\Delta bxlR$ strain with 70 % more SLAC after 72 h of growth and 30 % after 96 h. However, the profile of enzyme activity over time was similar in all three strains and complete derepression was not obtained during the first 2 days of incubation suggesting additional levels of control (Fig. 5).

Discussion

The high-level production of heterologous proteins is a key objective in biotechnology. *S. lividans* is a preferred heterologous host for industrial production of proteins from prokaryotic and eukaryotic organisms [5, 10, 32]. It has several major advantages over the more traditional platforms such as *E. coli* and *B. subtilis* as well as compared to other *Streptomyces* species. These advantages include: (i) low endogenous extracellular proteolytic activity in comparison with other *Streptomyces* species



or *Bacillus* [33, 34]; (ii) the fact that *S. lividans* does not form inclusion bodies, as is often the case in *E. coli*; and (iii) its ability to produce bioactive proteins [5]. Furthermore, the *Streptomyces* protein secretion machinery very efficiently exports proteins of interest into the culture supernatant, which facilitates subsequent protein purification. However, *S. lividans* also has some significant drawbacks, which relate to the mycelial growth associated with slow growth and viscous cultures and the lack of well-established expression systems. Thus, the implementation of *Streptomyces* as a production platform requires the optimization of promoter/vector/host systems.

In this work, we expand the toolbox of available promoters in combination with secretion signals to improve protein production in *S. lividans*. Several strong, constitutive promoters have been previously described for *Streptomyces*, but the number of suitable inducible promoters is scarce [3]. We compared six promoters and obtained good expression of the reporter proteins Xys1 from *S. halstedii* and Amy from *S. griseus* using the promoters *xysAp*, which is induced by different carbon sources such as xylose, xylan and fructose and repressed by glucose [28], and *pstSp*, induced by low phosphate conditions and by different carbon sources such as fructose, xylose or galactose [20, 35, 36]. These promoters resulted in more efficient and reproducible protein production than the widely applied *Streptomyces* promoters *ermE**p and *vsip* (Fig. 2a lanes 2 and 3 versus lanes 6 and 7). Therefore, the high-copy number *E. coli* and *Streptomyces* shuttle plasmids developed in this work; equipped with strongly inducible promoters (*xysAp* or *pstSp*), represent a good option for producing high levels of proteins of interest in *Streptomyces* as has been demonstrated for the xylanase, (Xys1) from *S. halstedii*, the α -amylase (Amy) from *S. griseus* and laccase (SLAC) from *S. coelicolor*.

The elimination of the putative *xysAp* repressors XlnR or BxlR from the host strains increased the protein production of the reporter SLAC in these mutants (up to 70 % after 72 h in the Δ *bxlR* strain) indicating the suitability of these strains. Moreover, in the Δ *bxlR* strain, the activity is clearly induced earlier with an almost two-fold higher SLAC production at the onset of stationary growth at 48 h. The fact that repression is not completely relieved in these single mutants during logarithmic growth (24 h) suggests that multiple (negative) control mechanisms act on *xysAp* in *S. lividans*.

We then proceeded to test the effect of altering the signal peptide, the adaptation of which could be a valuable additional tool in the production process to obtain more efficient secretion. As reviewed by Lammertyn and

Anné [23], several modifications have been found that improve *Streptomyces* signal peptides, and some of these were applied successfully in the vectors developed in this study. We observed an improvement of 17 % in xylanase secretion by changing the xylanase signal peptide for an amylase signal peptide with three additional amino acids from the mature amylase to maintain the sequence around the signal peptidase cleavage site (Fig. 3c). However, modification of the net charge of the amylase signal peptide as it has been described in amylase secretion [25] did not further improve xylanase production (data not shown). Thus, changes in the secretion signal that work for one protein do not necessarily also work for others because this effect is highly sequence dependent and may be also affected by the level of expression reached. Additionally, changing the N-terminal part of the gene may also have repercussions at the level of translation.

As discussed above, protein production is not only about providing the optimal expression system, but also depends on the morphology of the production host, which is a major factor in determining production level as well as fermentation time. Problems associated with the fermentation of filamentous organisms include slow growth rates, high viscosity, mixing problems due to the formation of large mycelial clumps and complex downstream processing [16]. Enhanced expression of the SsgA protein in *S. lividans* reduces such problems significantly as it favours fragmented growth due to enhanced cell division [15]. Interestingly, SsgA is also linked to the expression of all of the components of the secretion machinery. For example, enhanced expression of SsgA results in a strong increase in the secretion of tyrosinase (which is a substrate for the Tat system) in much shorter fermentation time, and expression of genes encoding components of the Tat and Sec secretion machinery is strongly upregulated in *ssgA* null mutants [14]. The latter is most likely the result of a feedback response to compensate for the absence of the important morphoprotein SsgA [14]. As a result of the improved growth and secretion capacity, the enhanced expression of SsgA has been shown to be potentially very advantageous for protein production [16]. Indeed, xylanase and amylase production increased by 40 and 70 %, respectively, in *S. lividans* GSAL1, which overexpresses SsgA when compared to the wild type strain (Fig. 4b, d).

Conclusions

The use of *Streptomyces* as a platform for high-level proteins production requires proper optimization for each protein of interest. We constructed a set of bifunctional plasmids (*E. coli-Streptomyces*) with strong and inducible promoters that are convenient for protein

production in *Streptomyces*. Combination with hosts with either enhanced expression of SsgA or lacking specific transcriptional repressors further improved enzyme production.

Methods

Strains and culture conditions

Escherichia coli DH5 α [37] was used for the cloning and isolation of plasmids. It was grown in Luria–Bertani liquid broth or on LB agar. DNA manipulations in *E. coli* were done following standard procedures [38].

Streptomyces lividans 1326 and its derivative strains, GSAL1 [16], $\Delta xlnR$, and $\Delta bxlR$ were used as hosts. The genome sequence of *S. lividans* 1326 has been published [39] and is available at StrepDB (<http://strepdb.streptomyces.org.uk>). GSAL1 is a transformant of *S. lividans* harbouring the integrative plasmid pGWS4-SD, which results in the effective overproduction of the SsgA protein [15]. The $\Delta xlnR$ and $\Delta bxlR$ mutants were prepared and isolated according to protocols described previously [40]. In the $\Delta xlnR$ mutant, nucleotides -146 to $+756$ relative to the start codon of *SLI4452* were replaced by a 62 nt scar of the *loxP* recombination site including two XbaI sites. In the $\Delta bxlR$ mutant, nts $+1$ to $+1127$ relative to the start of *SLI7232* were replaced. Strains were grown on R2YE agar plates for transformation, on MSA agar plates for sporulation [41] and in YE (1 % (w/v) Yeast extract, 5 mM MgCl₂ pH 7.2) or YES (1 % (w/v) Yeast extract, 10.3 % (w/v) sucrose, 5 mM MgCl₂ pH 7.2) liquid media supplemented with 1 % (w/v) of xylose or 5 % (w/v) of fructose for protein expression [20]. Neomycin, 15 μ g/mL, or thiostrepton, 2.5 μ g/mL, were added to the culture depending of the plasmid used. Liquid-grown cultures were carried out in baffled flasks at 28 °C and 200 rpm. DNA manipulations in *Streptomyces* were carried out essentially as described by Kieser et al. [41].

Plasmid constructions

All plasmids used in the present work are listed in Table 1. PCR amplification of each promoter was accomplished with oligonucleotides including EcoRI and NdeI restriction sites (Table 2). PCR products were purified by agarose gel electrophoresis, digested with EcoRI and NdeI, and cloned in plasmid pNX30 [5] digested with the same restriction enzymes.

pNXA1 and pNXA2 were constructed by PCR amplification of the signal peptide of the α -amylase (EMBL X57568) from *S. griseus* IMRU 3570 with oligonucleotides including NdeI and MreI restriction sites (MreI cuts just after the sequence that encodes the signal peptide of the xylanase and the correct frame was conserved in the

construction) (Table 2). PCR products were purified by agarose gel electrophoresis, digested with NdeI and MreI and cloned in plasmid pNX24 [22] digested with the same restriction enzymes. The new constructs have the α -amylase signal peptide fused in-frame with the *xysA* gene.

The plasmids pNXAmy and pNUFAmy were constructed by replacing xylanase gene by the α -amylase ORF in pNX24 and pNUF5 respectively. The α -amylase gene (*X57568.1*) was amplified from *S. griseus* genome with oligonucleotides MRG11 and MRG12 (Table 2).

The inserts of all plasmid constructions were sequenced on both strands using a Perkin–Elmer ABI Prism 377 DNA sequencer. *In silico* plasmids were done with the Gene Construction Kit (GCK, Textco).

Protein analysis

Protein profiles were analysed by denaturing polyacrylamide gel electrophoresis (SDS-PAGE) in a MiniProtean II system (Bio-Rad). Proteins were detected by 0.5 % Coomassie brilliant blue R staining and Low-Range SDS-PAGE Standards (Bio-Rad) were used as markers. Xylanase and amylase band intensities were analysed with ImageJ software [44].

Xylanase and amylase activities assays

Xylanase and amylase activities were measured by the dinitrosalicylic acid (DNS) method, using xylose or maltose as standards respectively [42]. One unit of xylanase or amylase activities was defined as the amount of enzyme required releasing 1 μ mol of reducing sugars, (expressed as xylose equivalents or maltose equivalents), in 1 min. All data shown are means of at least three different experiments.

SLAC in-gel assay

Secreted SLAC activity was determined in the supernatant of liquid cultures of the wild type *S. lividans* 1326, $\Delta xlnR$ and $\Delta bxlR$ strains harbouring the SLAC reporter plasmid pEPOS101. Cells were grown in YES medium supplemented with 1 % xylose, 20 μ M Cu(II) and 2.5 μ g/mL thiostrepton and samples (1.5 mL) were taken after 24, 48, 72 and 96 h of incubation at 28 °C. In-gel SLAC activity stain was done essentially according to Endo et al. [43]. Samples were mixed 1:1 with SDS-PAGE loading buffer without β -mercaptoethanol and without boiling and loaded directly on the gels. Following electrophoresis and detection of SLAC with N,N-Dimethyl-p-phenylenediamine sulfate salt (DMPPDA, Sigma D-4790) and naphthol, digital images of the gels were taken and analysed with ImageJ [44]. Band intensities were normalized

Table 2 Oligonucleotides used in this work

Name	Sequence 5'–3'	Use
LS-001	TTTTTTGAATTCTGTGCGGCTGCCCTCCGCGC	Forward oligonucleotide for cloning the <i>glpQp</i> . The sequence recognized by EcoRI is in italics
LS-002	TTTTTTCATATGCGTACTCTCGCGTCGAACG	Reverse oligonucleotide for cloning the <i>glpQp</i> . The sequence recognized by NdeI is in italics
LS-Amy	TTTTTTCATATGCCCCGAGACTCGCCACC	Forward oligonucleotide to introduce the amylase signal peptide. The sequence recognized by NdeI is in italics
LS-003	TTTTTTCGCCCCGCGCAGCGCGGGTGTG	Reverse oligonucleotide to introduce the amylase signal peptide. The sequence recognized by MreI is in italics
LS-004	TTTTTTCGCCCCGCGCGGGCGCAGCGCGC	Reverse oligonucleotide to introduce the amylase signal peptide with three additional amino acids. The sequence recognized by MreI is in italics
LS-023	TTTTTTGAATTCTGGTACCAGCCCGACCCGAGC	Forward oligonucleotide for cloning the <i>ermE</i> * <i>p</i> . The sequence recognized by EcoRI is in italics
LS-024	TTTTTTCATATGACCAACCGGCACGATTGTGCC	Reverse oligonucleotide for cloning the <i>ermE</i> * <i>p</i> . The sequence recognized by NdeI is in italics
LS-026	TTTTTTGAATTCGGGGATGACCACCGCGGGAG	Forward oligonucleotide for cloning the <i>vsip</i> . The sequence recognized by EcoRI is in italics
LS-027	TTTTTTCATATCGGTGAACCTCCTTCGATCGATG	Reverse oligonucleotide for cloning the <i>vsip</i> . The sequence recognized by EcoRV is in italics
RS-003	TTTTTTGAATTCGGGCTTCTCCCTCTCCGCGGG	Forward oligonucleotide for cloning the <i>xyIp</i> . The sequence recognized by EcoRI is in italics
RS-004	TTTTTTCATATGCGCGGCTCCTACTCGCTGC	Reverse oligonucleotide for cloning the <i>xyIp</i> . The sequence recognized by NdeI is in italics
LS- <i>celF</i>	TTTTTTGAATTCGGCCGCGCTCCCGTCTGGC	Forward oligonucleotide for cloning the <i>celAp</i> . The sequence recognized by EcoRI is in italics
LS- <i>celR</i>	TTTTTTCATATGAGTACCTCGATTCAGAGGA	Reverse oligonucleotide for cloning the <i>celAp</i> . The sequence recognized by NdeI is in italics
MRG11	TTTTTTCATATGCCCCGAGACTCCGCACC	Forward oligonucleotide for cloning amylase ORF. The sequence recognized by NdeI is in italics
MRG12	TTTTTTCCTGAGGCCGCGCCAGGTGTCGTTGAG	Reverse oligonucleotide for cloning amylase ORF. The sequence recognized by XhoI is in italics
4215UpF	GCGGAATTCAGTGCTCAAGGACGCCGGC	Forward oligonucleotide for the cloning upstream flank of XlnR. The sequence recognized by EcoRI is in italics
4215UpR	GCGGGATCCCATCCCGTGGCCTCCTCTCC	Reverse oligonucleotide for the cloning upstream flank of XlnR. The sequence recognized by BamHI is in italics
4215DwF	GCGTCTAGAGTAACTCGAGCGGTCTCGCCC	Forward oligonucleotide for the cloning downstream flank of XlnR. The sequence recognized by XbaI is in italics
4215DwR	CGCAAGCTTCCGCATCTGCTGGAGCCGG	Reverse oligonucleotide for the cloning downstream flank of XlnR. The sequence recognized by HindIII is in italics
7232UpF	GCGGAATTCGTGAGGTGTGTGTCATGAGCC	Forward oligonucleotide for the cloning of upstream flank of BxIR. The sequence recognized by EcoRI is in italics
7232UpR	CGCTCTAGACCGGGCGCCACCTCAAC	Reverse oligonucleotide for the cloning upstream flank of BxIR. The sequence recognized by XbaI is in italics
7232DwF	GCGTCTAGAGGCTCCTACCGCCGGGC	Forward oligonucleotide for the cloning downstream flank of BxIR. The sequence recognized by XbaI is in italics
7232DwR	CGCAAGCTTGTGCCGCGCCGGAGCCG	Reverse oligonucleotide for the cloning downstream flank of BxIR. The sequence recognized by HindIII is in italics

for the biomass (dry weight) and expressed in arbitrary units per dry weight.

Additional file

Additional file 1. Supplementary tables (S1–S6) containing enzyme quantification by activity or by ImageJ.

Authors' contributions

LS performed most of the experimental work; GPVW and EV carried out the studies on the SsgA overproducer strain and the repressor mutant strains; MD and RS selected the different promoters to be studied, directed the work and

wrote the manuscript with contributions from all authors. All authors read and approved the final manuscript.

Author details

¹ Instituto de Biología Funcional y Genómica/Departamento de Microbiología y Genética, Consejo Superior de Investigaciones Científicas (CSIC)/Universidad de Salamanca, C/Zacarías González nº 2, 37007 Salamanca, Spain. ² Molecular Biotechnology, IBL, Sylvius Laboratory, Leiden University, Leiden, The Netherlands.

Acknowledgements

This work has been supported by ERA-IB grants EUJ2008-03631 and PCIN-2014-067 of the Ministerio de Economía y Competitividad, to RS. EV and GPVW acknowledge the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO)/Advanced Chemical Technologies for Sustainability (ACTS) for grant

053.80.703 in the ERA-IB framework (EIB.08.013 EPOS). The authors also thank the CSIC for the co-financing of this publication in Open Access through its Unit of Information Resources for Research (URICI). Thanks are due to Dr. JA Gil for the *S. griseus* DNA, to Dr. J. Anne for his suggestions on signal peptide constructions, to MJ Jiménez Rufo for her excellent technical work and to N. Skinner for supervising the English version of the manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 26 October 2015 Accepted: 19 January 2016

Published online: 04 February 2016

References

- de Marco A. Recombinant polypeptide production in *E. coli*: towards a rational approach to improve the yields of functional proteins. *Microb Cell Fact*. 2013;12:101.
- Hochkoeppler A. Expanding the landscape of recombinant protein production in *Escherichia coli*. *Biotechnol Lett*. 2013;35:1971–81.
- Vrancken K, Anne J. Secretory production of recombinant proteins by *Streptomyces*. *Future Microbiol*. 2009;4:181–8.
- Díaz M, Adham SA, Ramón D, Gil JA, Santamaría RI. *Streptomyces lividans* and *Brevibacterium lactofermentum* as heterologous hosts for the production of X₂₂ xylanase from *Aspergillus nidulans*. *Appl Microbiol Biotechnol*. 2004;65:401–6.
- Díaz M, Ferreras E, Moreno R, Yepes A, Berenguer J, Santamaría R. High-level overproduction of *Thermus* enzymes in *Streptomyces lividans*. *Appl Microbiol Biotechnol*. 2008;79:1001–8.
- Zhu Y, Wang L, Du Y, Wang S, Yu T, Hong B. Heterologous expression of human interleukin-6 in *Streptomyces lividans* TK24 using novel secretory expression vectors. *Biotechnol Lett*. 2011;33:253–61.
- Ayadi DZ, Chouayekh H, Mhiri S, Zerria K, Fathallah DM, Bejar S. Expression by *Streptomyces lividans* of the rat alpha integrin CD11b A-domain as a secreted and soluble recombinant protein. *J Biomed Biotechnol*. 2007;2007:54327.
- Kim MR, Choeng YH, Chi WJ, Kang DK, Hong SK. Heterologous production of streptokinase as a secretory form in *Streptomyces lividans* and nonsecretory form in *Escherichia coli*. *J Microbiol Biotechnol*. 2010;20:132–7.
- Noda S, Ito Y, Shimizu N, Tanaka T, Ogino C, Kondo A. Over-production of various secretory-form proteins in *Streptomyces lividans*. *Protein Expr Purif*. 2010;73:198–202.
- Anné J, Vrancken K, Van Mellaert L, Van Impe J, Bernaerts K. Protein secretion biotechnology in Gram-positive bacteria with special emphasis on *Streptomyces lividans*. *Biochim Biophys Acta*. 2014;1843:1750–61.
- Nielsen J. Modelling the morphology of filamentous microorganisms. *Trends Biotechnol*. 1996;14:438–43.
- van Wezel GP, McKenzie NL, Nodwell JR. Chapter 5. Applying the genetics of secondary metabolism in model actinomycetes to the discovery of new antibiotics. *Methods Enzymol*. 2009;458:117–41.
- van Dissel D, Claessen D, Roth M, van Wezel GP. A novel locus for mycelial aggregation forms a gateway to improved *Streptomyces* cell factories. *Microb Cell Fact*. 2015;14:44.
- Noens EE, Mersinias V, Willemsse J, Traag BA, Laing E, Chater KF, Smith CP, Koerten HK, van Wezel GP. Loss of the controlled localization of growth stage-specific cell-wall synthesis pleiotropically affects developmental gene expression in an *ssgA* mutant of *Streptomyces coelicolor*. *Mol Microbiol*. 2007;64:1244–59.
- van Wezel GP, van der Meulen J, Kawamoto S, Luiten RG, Koerten HK, Kraal B. *ssgA* is essential for sporulation of *Streptomyces coelicolor* A3(2) and affects hyphal development by stimulating septum formation. *J Bacteriol*. 2000;182:5653–62.
- van Wezel GP, Krabben P, Traag BA, Keijser BJ, Kerste R, Vijgenboom E, Heijnen JJ, Kraal B. Unlocking *Streptomyces* spp. for use as sustainable industrial production platforms by morphological engineering. *Appl Environ Microbiol*. 2006;72:5283–8.
- Van Mellaert L, Lammertyn E, Schacht S, Proost P, Van Damme J, Wroblowski B, Anne J, Scarcez T, Sablon E, Raeymaeckers J, Van Broekhoven A. Molecular characterization of a novel subtilisin inhibitor protein produced by *Streptomyces venezuelae* CBS762.70. *DNA Seq*. 1998;9:19–30.
- Bibb MJ, Janssen GR, Ward JM. Cloning and analysis of the promoter region of the erythromycin resistance gene (*ermE*) of *Streptomyces erythraeus*. *Gene*. 1985;38:215–26.
- Ruiz-Arribas A, Sánchez P, Calvete JJ, Raida M, Fernández-Abalos JM, Santamaría RI. Analysis of *xysA*, a gene from *Streptomyces halstedii* JM8 that encodes a 45-kilodalton modular xylanase, Xys1. *Appl Environ Microbiol*. 1997;63:2983–8.
- Díaz M, Esteban A, Fernández-Ábalos JM, Santamaría RI. The high-affinity phosphate-binding protein PstS is accumulated under high fructose concentrations and mutation of the corresponding gene affects differentiation in *Streptomyces lividans*. *Microbiology*. 2005;151:2583–92.
- Fernández-Abalos JM, Reviejo V, Díaz M, Rodríguez S, Leal F, Santamaría RI. Posttranslational processing of the xylanase Xys1L from *Streptomyces halstedii* JM8 is carried out by secreted serine proteases. *Microbiology*. 2003;149:1623–32.
- Adham SA, Honrubia P, Díaz M, Fernández-Ábalos JM, Santamaría RI, Gil JA. Expression of the genes coding for the xylanase Xys1 and the cellulase Cel1 from the straw-decomposing *Streptomyces halstedii* JM8 cloned into the amino-acid producer *Brevibacterium lactofermentum* ATCC13869. *Arch Microbiol*. 2001;177:91–7.
- Lammertyn E, Desmyter S, Schacht S, Van Mellaert L, Anne J. Influence of charge variation in the *Streptomyces venezuelae* alpha-amylase signal peptide on heterologous protein production by *Streptomyces lividans*. *Appl Microbiol Biotechnol*. 1998;49:424–30.
- Guan C, Cui W, He X, Hu X, Xu J, Du G, Chen J, Zhou Z. Construction and development of a novel expression system of *Streptomyces*. *Protein Expr Purif*. 2015;113:17–22.
- Vigal T, Gil JA, Daza A, García-González MD, Villadas P, Martín JF. Effects of replacement of promoters and modification of the leader peptide region of the *amy* gene of *Streptomyces griseus* on synthesis and secretion of alpha-amylase by *Streptomyces lividans*. *Mol Gen Genet*. 1991;231:88–96.
- Jakimowicz D, van Wezel GP. Cell division and DNA segregation in *Streptomyces*: how to build a septum in the middle of nowhere? *Mol Microbiol*. 2012;85:393–404.
- Willemsse J, Borst JW, de Waal E, Bisseling T, van Wezel GP. Positive control of cell division: FtsZ is recruited by SsgB during sporulation of *Streptomyces*. *Genes Dev*. 2011;25:89–99.
- Rodríguez S, Santamaría RI, Fernández-Ábalos JM, Díaz M. Identification of the sequences involved in the glucose-repressed transcription of the *Streptomyces halstedii* JM8 *xysA* promoter. *Gene*. 2005;351:1–9.
- Tsujibo H, Kosaka M, Ikenishi S, Sato T, Miyamoto K, Inamori Y. Molecular characterization of a high-affinity xylobiose transporter of *Streptomyces thermoviolaceus* OPC-520 and its transcriptional regulation. *J Bacteriol*. 2004;186:1029–37.
- Rigali S, Derouaux A, Giannotta F, Dusart J. Subdivision of the helix-turn-helix GntR family of bacterial regulators in the FadR, HutC, MocR, and YtrA subfamilies. *J Biol Chem*. 2002;277:12507–15.
- Machczynski MC, Vijgenboom E, Samyn B, Canters GW. Characterization of SLAC: a small laccase from *Streptomyces coelicolor* with unprecedented activity. *Protein Sci*. 2004;13:2388–97.
- Nakashima N, Mitani Y, Tamura T. Actinomycetes as host cells for production of recombinant proteins. *Microb Cell Fact*. 2005;4:7.
- Binnie C, Cossar JD, Stewart DI. Heterologous biopharmaceutical protein expression in *Streptomyces*. *Trends Biotechnol*. 1997;15:315–20.
- Nijland R, Kuipers OP. Optimization of protein secretion by *Bacillus subtilis*. *Recent Pat Biotechnol*. 2008;2:79–87.
- Esteban A, Díaz M, Yepes A, Santamaría RI. Expression of the *pstS* gene of *Streptomyces lividans* is regulated by the carbon source and is partially independent of the PhoP regulator. *BMC Microbiol*. 2008;8:201.
- Sola-Landa A, Moura RS, Martín JF. The two-component PhoR-PhoP system controls both primary metabolism and secondary metabolite biosynthesis in *Streptomyces lividans*. *Proc Natl Acad Sci U S A*. 2003;100:6133–8.
- Hanahan D. Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol*. 1983;166:557–80.
- Sambrook J, Fritsch E, Maniatis T. Molecular cloning: a laboratory manual. 2nd ed. Cold Spring Harbor: Cold Spring Harbor Laboratory; 1989.

39. Cruz-Morales P, Vijgenboom E, Iruegas-Bocardo F, Girard G, Yanez-Guerra LA, Ramos-Aboites HE, Pernodet JL, Anne J, van Wezel GP, Barona-Gomez F. The genome sequence of *Streptomyces lividans* 66 reveals a novel tRNA-dependent peptide biosynthetic system within a metal-related genomic island. *Genome Biol Evol*. 2013;5:1165–75.
40. Blundell KL, Wilson MT, Svistunenko DA, Vijgenboom E, Worrall JA. Morphological development and cytochrome c oxidase activity in *Streptomyces lividans* are dependent on the action of a copper bound Sco protein. *Open biology*. 2013;3:120163.
41. Kieser T, Hopwood DA, Bibb JM, Chater KF, Buttner MJ. *Practical Streptomyces genetics*. Norwich: John Innes Foundation; 2000.
42. Biely P, Mislovicova D, Toman R. Soluble chromogenic substrates for the assay of endo-1,4- β -xylanases and endo-1,4- β -glucanases. *Anal Biochem*. 1985;144:142–6.
43. Endo K, Hosono K, Beppu T, Ueda K. A novel extracytoplasmic phenol oxidase of *Streptomyces*: its possible involvement in the onset of morphogenesis. *Microbiology*. 2002;148:1767–76.
44. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods*. 2012;9:671–5.
45. García-González MD, Martín JF, Vigal T, Liras P. Characterization, expression in *Streptomyces lividans*, and processing of the amylase of *Streptomyces griseus* IMRU 3570: two different amylases are derived from the same gene by an intracellular processing mechanism. *J Bacteriol*. 1991;173:2451–8.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

