

THE [N II] 205 μm EMISSION IN LOCAL LUMINOUS INFRARED GALAXIES*

YINGHE ZHAO (赵应和)^{1,2,3}, NANYAO LU¹, C. KEVIN XU¹, YU GAO (高煜)^{2,3}, STEVEN D. LORD⁴, VASSILIS CHARMANDARIS^{5,6},
TANIO DIAZ-SANTOS^{7,8}, AARON EVANS^{9,10}, JUSTIN HOWELL¹, ANDREEA O. PETRIC¹¹, PAUL P. VAN DER WERF¹², AND
DAVID B. SANDERS¹³

¹ Infrared Processing and Analysis Center, California Institute of Technology 100-22, Pasadena, CA 91125, USA; zhaoyinghe@gmail.com

² Purple Mountain Observatory, Chinese Academy of Sciences, Nanjing 210008, China

³ Key Laboratory of Radio Astronomy, Chinese Academy of Sciences, Nanjing 210008, China

⁴ The SETI Institute, 189 Bernardo Avenue, Suite 100, Mountain View, CA 94043, USA

⁵ Department of Physics and ITCP, University of Crete, GR-71003 Heraklion, Greece

⁶ IAASARS, National Observatory of Athens, GR-15236, Penteli, Greece

⁷ Spitzer Science Center, California Institute of Technology, MS 220-6, Pasadena, CA 91125, USA

⁸ Nucleo de Astronomia de la Facultad de Ingenieria, Universidad Diego Portales, Av. Ejercito Libertador 441, Santiago, Chile

⁹ Department of Astronomy, University of Virginia, 530 McCormick Road, Charlottesville, VA 22904, USA

¹⁰ National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, VA 22903, USA

¹¹ Gemini Observatory, Northern Operations Center, 670 N. A'ohoku Place, Hilo, HI 96720, USA

¹² Leiden Observatory, Leiden University, P.O. Box 9513, 2300 RA Leiden, The Netherlands

¹³ University of Hawaii, Institute for Astronomy, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

Received 2015 April 6; accepted 2016 January 6; published 2016 February 29

ABSTRACT

In this paper, we present the measurements of the [N II] 205 μm line for a flux-limited sample of 122 (ultra-) luminous infrared galaxies [(U)LIRGs] and 20 additional normal galaxies, obtained with the *Herschel Space Observatory* (*Herschel*). We explore the far-infrared (FIR) color dependence of the [N II] 205 μm ($L_{[\text{N II}]205 \mu\text{m}}$) to the total infrared (L_{IR}) luminosity ratio, and find that $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}$ only depends modestly on the 70–160 μm flux density ratio (f_{70}/f_{160}) when $f_{70}/f_{160} \lesssim 0.6$, whereas such dependence becomes much steeper for $f_{70}/f_{160} > 0.6$. We also investigate the relation between $L_{[\text{N II}]205 \mu\text{m}}$ and star formation rate (SFR), and show that $L_{[\text{N II}]205 \mu\text{m}}$ has a nearly linear correlation with SFR, albeit the intercept of such a relation varies somewhat with f_{60}/f_{100} , consistent with our previous conclusion that [N II] 205 μm emission can serve as an SFR indicator with an accuracy of ~ 0.4 dex, or ~ 0.2 dex if f_{60}/f_{100} is known independently. Furthermore, together with the *Infrared Space Observatory* measurements of [N II], we use a total of ~ 200 galaxies to derive the local [N II] 205 μm luminosity function (LF) by tying it to the known IR LF with a bivariate method. As a practical application, we also compute the local SFR volume density ($\dot{\rho}_{\text{SFR}}$) using the newly derived SFR calibrator and LF. The resulting $\log \dot{\rho}_{\text{SFR}} = -1.96 \pm 0.11 M_{\odot} \text{ yr}^{-1} \text{ Mpc}^{-3}$ agrees well with previous studies. Finally, we determine the electron densities (n_e) of the ionized medium for a subsample of 12 (U)LIRGs with both [N II] 205 μm and [N II] 122 μm data, and find that n_e is in the range of ~ 1 –100 cm^{-3} , with a median value of 22 cm^{-3} .

Key words: galaxies: evolution – galaxies: ISM – galaxies: luminosity function, mass function – galaxies: starburst – infrared: ISM

Supporting material: machine-readable table

1. INTRODUCTION

Emission from the forbidden atomic fine-structure transitions in the far-infrared (FIR), such as the [C II] 158 μm , [N II] 122 and 205 μm , [O I] 63 and 145 μm , and [O III] 52 μm and 88 μm lines, is important for cooling the interstellar medium (ISM) and for providing critical diagnostic tools for the study of the star-forming ISM (e.g., Stacey et al. 1991; Lord et al. 1995; Malhotra et al. 2001; Farrah et al. 2013; De Looze et al. 2014; Fischer et al. 2014; Sargsyan et al. 2014). Among these lines, the [C II] 158 μm emission is probably the most important and well studied since it is the brightest single line in most galaxies and accounts for 0.1%–1% of the total FIR luminosity (e.g., Stacey et al. 1991; Díaz-Santos et al. 2013).

The [N II] 205 μm line is of particular interest for the following reasons. First, this $^3\text{P}_1 \rightarrow ^3\text{P}_0$ transition (205.197 μm ; hereafter [N II] 205 μm) of singly ionized nitrogen is expected to be an excellent indicator of the star

formation rate (SFR) based on the following facts. (1) The ionization potential of nitrogen is only 14.53 eV. Thus the [N II] 205 μm emission arises only from H II regions and essentially traces all warm ionized ISMs. It can be utilized to estimate the ionizing photon rate (e.g., Bennett et al. 1994). (2) The [N II] 205 μm transition can be easily collisionally excited because of its low critical density (44 cm^{-3} ; Oberst et al. 2006) and excitation energy (~ 70 K). (3) This emission is usually optically thin and suffers much less dust extinction than optical and near-infrared lines. Indeed, Zhao et al. (2013) have shown that the [N II] 205 μm line can serve as an SFR indicator, which is especially useful for studying high-redshift galaxies for which the redshifted [N II] 205 μm line is readily obtainable with modern submillimeter telescopes such as the Atacama Large Millimeter/submillimeter Array (ALMA).

Second, this line can provide complementary information on the origin of the [C II] 158 μm emission (e.g., Oberst et al. 2006; Walter et al. 2009; Parkin et al. 2013, 2014; Decarli et al. 2014; Hughes et al. 2015). The [C II] line can arise from both neutral and ionized gases since it takes only 11.3 eV to form C^+ , while the ionization potential of hydrogen is 13.60 eV.

* Based on *Herschel* observations. *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.

Therefore, it is important to know what fraction of the observed [C II] line emission is from the ionized gas for the study of star-forming regions such as photodissociation region (PDR) modeling. Fortunately, the critical densities for the [N II] 205 μm and [C II] 158 μm lines in ionized regions are nearly identical (44 and 46 cm^{-3} at $T = 8000$ K, respectively; Oberst et al. 2006), and both require similar ionization potentials to further form N^{++} (29.6 eV) and C^{++} (24.4 eV). As a result, the [C II]/[N II] 205 μm line ratio from ionized gas is a function of only the assumed N^+/C^+ abundances within the H II region, and therefore the observed [C II]/[N II] 205 μm line ratio yields the fraction of the [C II] emission that arises from the ionized gas (Oberst et al. 2006, 2011).

Third, the ratio of the [N II] 122–205 μm lines (hereafter $R_{122/205}$) is an excellent density probe of low-density ionized gas due to their different critical densities (n_{crit}) required for collisional excitations and being at the same ionization level. At an electron temperature of 8000 K, n_{crit} are ~ 293 and 44 cm^{-3} for the [N II] 122 and 205 μm lines, respectively (Oberst et al. 2006). Therefore, $R_{122/205}$ is sensitive to gas densities of $10 \lesssim n_e \lesssim 300 \text{ cm}^{-3}$ (Oberst et al. 2006, 2011; also see Section 3.3).

However, the [N II] 205 μm line is generally inaccessible to ground-based facilities for local galaxies, and, for extragalactic objects, only a handful were observed using satellite and airborne platforms (Petuchowski et al. 1994; Lord et al. 1995) prior to the advent of the *Herschel Space Observatory* (hereafter, *Herschel*; Pilbratt et al. 2010). These studies have shown that the [N II] 205 μm line is fairly bright, and the luminosity of the [N II] 205 μm line ($L_{[\text{N II}]205 \mu\text{m}}$) may be up to $\sim 10^{-3.5}$ times the total infrared luminosity (L_{IR} ; 8–1000 μm ; also see Zhao et al. 2013). With such a high luminosity, this line offers an excellent method for studying SFR and ionized gas properties in galaxies at high redshifts. The advantage of the [N II] 205 μm line over other FIR lines, such as [C II] 158 μm , [N II] 122 μm , and [O III] 88 μm , is that it starts to fall into atmospheric sub/millimeter windows that have higher transmission at lower- z , due to its longer wavelength. The detectability of the [N II] 205 μm line and its potential for important astrophysical applications at high- z have already been demonstrated by a few experimental ALMA observing campaigns of a galaxy at $z = 4.76$ (Nagao et al. 2012), and by the IRAM 30 m telescope and Plateau de Bure Interferometer detection of distant, strongly lensed galaxies (Combes et al. 2012, $z \sim 5.2$; Decarli et al. 2012, 2014, $z \sim 3.9$ and 4.7).

In Zhao et al. (2013), we reported our first results on the [N II] 205 μm line emission for an initial set of 70 (ultra-)luminous infrared galaxies [(U)LIRGs; $L_{\text{IR}} \geq 10^{11(12)} L_{\odot}$]¹⁴, observed with the Fourier-transform spectrometer (FTS) of the Spectral and Photometric Imaging Receiver (SPIRE; Griffin et al. 2010) on board *Herschel*. In that Letter we focused on the possibility of using the [N II] 205 μm emission as an SFR indicator. Here we expand our analysis to our full *Herschel* sample of 122 LIRGs, which is a flux-limited subsample with $f_{\text{IR}}(8\text{--}1000 \mu\text{m}) > 6.5 \times 10^{-13} \text{ W m}^{-2}$ from the Great Observatories All-Sky LIRGs Survey (GOALS; Armus et al. 2009). In addition, we include 20 additional nearby large galaxies for

which SPIRE/FTS mapping observations covering the entire galaxy disk are available from the *Herschel* Science Archive (HSA).

In this paper, besides further investigating the relation between $L_{[\text{N II}]205 \mu\text{m}}$ and SFR, we also derive the luminosity function (LF) of the [N II] 205 μm line and SFR density in the local universe. It is important to constrain the LF of the [N II] emission locally since now it becomes possible to build a large sample for studying the [N II] LF at high redshift using modern facilities such as ALMA. The local [N II] LF can serve as a benchmark necessary for observational and theoretical (e.g., Orsi et al. 2014) studies on its evolution. Given the unprecedented sensitivity of *Herschel* at $\sim 200 \mu\text{m}$, and the large number of galaxies in the local universe already observed, for the first time we can derive the local [N II] LF (see Section 3.2), using a bivariate method and by utilizing the local IR LF, which has been studied extensively in the literature with *IRAS* observations (e.g., Soifer et al. 1986; Sanders et al. 2003).

In addition, we estimate the electron densities for a subsample of our (U)LIRGs by comparing the observed $R_{122/205}$ with theoretical predications. As shown in Rubin et al. (1994; also see Oberst et al. 2006), $R_{122/205}$ varies from 3 for $n_e \sim 100 \text{ cm}^{-3}$ to 10 for $n_e \gtrsim 10^3 \text{ cm}^{-3}$. Although Petuchowski et al. (1994) and Lord et al. (1995) observed both the [N II] 205 μm and [N II] 122 μm lines in M82, the use of $R_{122/205}$ for estimating n_e was mostly limited to our own Galaxy (e.g., Wright et al. 1991; Bennett et al. 1994; Oberst et al. 2006, 2011) prior to the advent of *Herschel*. Furthermore, so far there are only a handful of normal galaxies (e.g., M51 and Cen A; Parkin et al. 2013, 2014; NGC 891; Hughes et al. 2015) for which n_e of the low-density gas has been derived using the ratio of these two lines. For (U)LIRGs, it is still unclear what the typical n_e for the low-density ionized gas is.

The remainder of this paper is organized as follows. We give a brief introduction of the sample, observations, and data reduction in Section 2, present the results and discussion in Section 3, and briefly summarize the main conclusions in the last section. Throughout the paper, we adopt a Hubble constant of $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.28$, and $\Omega_\Lambda = 0.72$, which are based on the five-year *WMAP* results (Hinshaw et al. 2009), and are the same as those used by the GOALS project (Armus et al. 2009).

2. SAMPLE, OBSERVATIONS, AND DATA REDUCTION

2.1. (Ultra-)Luminous Infrared Galaxies

The primary sample studied in this paper is from the *Herschel* open time project *Herschel Spectroscopic Survey of Warm Molecular Gas in Local Luminous Infrared Galaxies* (OT1_nlu_1; PI: N. Lu). This project aims primarily at studying the dense and warm molecular gas properties of 125 LIRGs (e.g., Lu et al. 2014, 2015), which comprise a flux-limited subset of the GOALS sample (Armus et al. 2009). N. Lu et al. (2016, in preparation) will give the program details and complete set of spectra for individual galaxies. The [N II] observations were available for 123 targets, one of which is a multiple-source system, and our targeted object turned out to be a Galactic source according to its SPIRE/FTS spectrum, and was consequently excluded from our analysis. Here we present the [N II] 205 μm data for the 122 galaxies (hereafter GOALS-FTS sample), including 111 LIRGs and 11 ULIRGs. Of these

¹⁴ L_{IR} is calculated by using the *IRAS* four-band fluxes and the equation given in Sanders & Mirabel (1996), i.e., $L_{\text{IR}}(8\text{--}1000 \mu\text{m}) = 4\pi D_L^2 f_{\text{IR}}$, where D_L is the luminosity distance, and $f_{\text{IR}} = 1.8 \times 10^{-14} (13.48f_{12} + 5.16f_{25} + 2.58f_{60} + f_{100}) (\text{W m}^{-2})$.

Table 1
Fluxes of the [N II] 205 μm Emission

Galaxy Name	R.A. (hh:mm:ss)	Decl. (dd:mm:ss)	Obs ID	$f_{[\text{N II}] 205 \mu\text{m}}^a$ ($10^{-17} \text{ W m}^{-2}$)	f_{corr} ($R_{122/205}$)	SFR ($M_{\odot} \text{ yr}^{-1}$)	f_{60} (Jy)	f_{100} (Jy)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
GOALS-FTS Sample								
NGC0023	00:09:53.4	+25:55:26.2	1342247622	10.79 ± 0.75	1.58	23.0	9.03	15.66
NGC0034	00:11:06.5	-12:06:24.9	1342199253	2.55 ± 0.49	2.33	55.0	17.05	16.86
MCG-02-01-051	00:18:50.9	-10:22:37.6	1342247617	3.80 ± 0.54	1.00	50.3	7.48	9.66
Mapping Galaxy Sample								
NGC1266	03:16:00.7	-02:25:38	1342239353	2.85 ± 0.42	...	5.6	13.13	16.89
NGC1377	03:36:39.1	-20:54:08	1342239352	0.58 ± 0.10	7.43	5.95
NGC1482	03:54:38.9	-20:30:10	1342248233	36.1 ± 2.4	...	13.1	33.96	46.73
ISO Sample								
NGC0520	01:24:34.90	+03:47:30.0	77702295	39.2 ± 16.3	1.2	15.4	31.10	47.12
NGC0986	02:33:34.10	-39:02:41.0	74300187	43.3 ± 18.1	1.2	12.6	25.14	51.31
NGC1222	03:08:56.80	-02:57:18.0	82400836	8.7 ± 4.0	1.5	8.3	13.07	15.38

Note. Columns: (1) galaxy name; (2) and (3) R.A. and decl. (J2000); for the GOALS-FTS sample, the coordinate gives the position where the *Herschel* SPIRE/FTS observation was pointed; (4) observation ID (number) for the *Herschel* (ISO) observation; (5) [N II] 205 μm flux: measured from the SPIRE/FTS spectra for the GOALS-FTS and Mapping Galaxy samples; obtained from [N II] 122 μm emission for the ISO sample; (6) for the GOALS-FTS sample, correction factor (f_{corr}) applied to Column (5) to obtain the total [N II] 205 μm flux (see Section 2.3 for details); for the ISO sample, the [N II] 122 μm -to-[N II] 205 μm conversion factor ($R_{122/205}$); (7) star formation rate (Section 3.1); (8) and (9) *IRAS* fluxes at 60 and 100 μm respectively.

^a For the ISO sample, the listed error has been taken into account for the uncertainty of $R_{122/205}$.

(This table is available in its entirety in machine-readable form.)

sources, 48 galaxies are point-like sources with respect to the $\sim 17''$ *Herschel* SPIRE/FTS beam at $\sim 210 \mu\text{m}$, and 74 are extended sources, which were determined according to their flux fractions of the FIR continuum emissions at both 70 and 160 μm , observed within the SPIRE/FTS beam (see Section 2.3 for details).

The observations were conducted with the SPIRE/FTS in its point source spectroscopy mode and high spectral resolution configuration, yielding a spectral resolution of 0.04 cm^{-1} (or 1.2 GHz) over the spectral coverage of 194–672 μm . The data were reduced using the default version of the standard SPIRE reduction and calibration pipeline for point source mode included in the *Herschel* Interactive Processing Environment (HIPE; Ott 2010) version 11.0.

In most cases, the [N II] 205 μm line is the brightest line in the SPIRE/FTS wavelength range (N. Lu et al. 2016, in preparation), and it has high signal-to-noise ratios (S/N). As shown in Zhao et al. (2013), the line fluxes were obtained by fitting the observed profile using the instrumental *Sinc* function convolved with a free-width Gaussian profile. This is because the line widths of most (U)LIRGs are $\gtrsim 200 \text{ km s}^{-1}$ (e.g., see Arribas et al. (2014) for ionized gas and Rosenberg et al. (2015) for molecular gas). Given the instrumental resolution ($\sim 300 \text{ km s}^{-1}$ at 210 μm), the observed line might be marginally resolved and could not be modeled by a pure *Sinc* function. Therefore, we adopted the *Sinc*-convolved-Gaussian (SCG) profiles for the integrated [N II] line flux measurements except for a few galaxies where a pure *Sinc* profile was a better choice due to the intrinsically narrow line width and/or relatively low S/N in the line. During the fitting process, the width of the *Sinc* function was fixed to be the SPIRE/FTS resolution (1.2 GHz), while the width of the Gaussian function was allowed to vary. The resulting full width at half maximum (FWHM) of the [N II] 205 μm line, which was obtained from the Gaussian part of the

SCG profile, is $\sim 100\text{--}600 \text{ km s}^{-1}$, with a median value of $\sim 300 \text{ km s}^{-1}$. Based on the 1σ statistical uncertainties, the lines in most ($>80\%$) sources are detected at better than 7σ , with the median at $\sim 14\sigma$. The measured line fluxes are given in Table 1.

2.2. Local Normal Galaxies

Almost all of our sample galaxies are (U)LIRGs, and hence have a rather limited dynamic range of several physical parameters such as luminosity, FIR color, etc. To increase the sample size and dynamic range of our study, we also include in our analysis 20 nearby normal galaxies having *Herschel* SPIRE/FTS mapping observations that cover the *entire* galaxy. As in Zhao et al. (2013), we further include another 53 unresolved galaxies (23 detections and 30 upper limits) observed by the *Infrared Space Observatory* (ISO; Kessler et al. 1996, 2003) Long Wavelength Spectrometer (LWS), for which the [N II] 122 μm fluxes were measured by Brauher et al. (2008; hereafter, the ISO sample).

2.2.1. *Herschel* SPIRE/FTS Mapping Observations

These observations were carried out by various *Herschel* projects, e.g., the *Very Nearby Galaxies Survey* (KPGT_cwilso01_1; PI: C. D. Wilson; e.g., Spinoglio et al. 2012b; Parkin et al. 2013; Schirm et al. 2014; Hughes et al. 2015) and the *Beyond the Peak: Resolved Far-Infrared Spectral Mapping of Nearby Galaxies with SPIRE/FTS* (OT1_jsmith01_1; PI: J. D. Smith). The level 0.5 raw data were obtained through HSA, and reduced with the default pipeline for mapping observations provided in HIPE 11. We then added up all of the pixels of the level 2 product which have valid data, and measured the integrated [N II] 205 μm fluxes using the method described in Section 2.1 above. During




Figure 1. [N II] 122 μm to [N II] 205 μm emission ratio ($R_{122/205}$) plotted against the *IRAS* FIR color. The upward and downward arrows represent lower and upper limits, respectively.

this process, we have converted the units of the mapping data from MJy/sr to Jy/pixel using the area of an individual pixel.

2.2.2. *ISO* LWS Observations

For the *ISO* sample, we derived their [N II] 205 μm fluxes ($f_{[\text{N II}]205 \mu\text{m}}$) from the observed [N II] 122 μm fluxes ($f_{[\text{N II}]122 \mu\text{m}}$) using the following empirical method. The $R_{122/205}$ used in our conversion was estimated on the basis of the actual observations of these two lines for a sample consisting of 7 normal galaxies and 26 (U)LIRGs. Besides our GOALS-FTS sources, about half of these (U)LIRGs are from the Farrah et al. (2013) sample, for which the [N II] 122 μm were observed with the *Herschel* Photodetector Array Camera and Spectrometer (PACS; Poglitsch et al. 2010), and $f_{[\text{N II}]122 \mu\text{m}}$ were adopted from Farrah et al. (2013); whereas $f_{[\text{N II}]205 \mu\text{m}}$ were measured from the SPIRE/FTS data observed in the program “OT1_d-farrah_1” (PI: D. Farrah). For our GOALS-FTS objects, the aperture-corrected (see Section 2.3) $f_{[\text{N II}]205 \mu\text{m}}$ were used.

In Figure 1, we plotted $R_{122/205}$ against the *IRAS* FIR color, f_{60}/f_{100} . It seems that $R_{122/205}$ shows some dependence on f_{60}/f_{100} . Kewley et al. (2000) found that electron density tends to correlate with f_{25}/f_{60} . Therefore, the weak dependence of $R_{122/205}$ on f_{60}/f_{100} appears to be understandable. To further check whether there is a correlation between f_{60}/f_{100} and $R_{122/205}$, we computed the Kendall’s τ correlation coefficient using the *cenken* function in the *NADA* package within the public domain **R** statistical software environment.¹⁵ For the whole data set presented in Figure 1, we have $\tau = 0.18$, with a p -value of 0.13, and thus we do not reject the null hypothesis that these two parameters are uncorrelated at the 0.05

significance level. However, we have $\tau = 0.35$ with a p -value of 0.03 if we limit the sample to $f_{60}/f_{100} < 0.9$. Therefore, there exists a weak correlation between f_{60}/f_{100} and $R_{122/205}$ within this color range. Since almost all of the *ISO* galaxies fall within this color range, we adopt FIR color-dependent $R_{122/205}$. Nevertheless, we caution that such an analysis is possibly limited by the small size of the sample. However, this (un)correlation will not affect our main conclusions since the two $R_{122/205}$ values adopted in the following only differ by ~ 0.3 , which is negligible compared to the overall uncertainties.

For sources with $f_{60}/f_{100} < 0.7$, we adopt $R_{122/205} = 1.2 \pm 0.5$, and for the other warmer galaxies (with $f_{60}/f_{100} < 1.0$), $R_{122/205} = 1.5 \pm 0.7$. These adopted $R_{122/205}$ values are the median of the corresponding detections, and are lower than the single value of 2.6 adopted in Zhao et al. (2013). The latter was based on the theoretical prediction of an electron density of $n_e = 80 \text{ cm}^{-3}$, i.e., the median value of H II regions in late-type galaxies (Ho et al. 1997). However, the adopted n_e in Zhao et al. (2013) was measured from H II regions in the centers of nearby galaxies, and thus might be an overestimate of the mean n_e for the entire galaxy. As a result, the overall $f_{[\text{N II}]205 \mu\text{m}}$ obtained from $f_{[\text{N II}]122 \mu\text{m}}$ in Zhao et al. (2013) was somewhat underestimated.

For these nearby galaxies, the redshift-independent distance was adopted if a direct primary measurement distance could be found in the NED¹⁶ database, otherwise it was derived with the same method as used for our (U)LIRG sample (e.g., Armus et al. 2009), i.e., by correcting the heliocentric velocity for the 3-attractor flow model of Mould et al. (2000).

2.3. Aperture Corrections

Around 205 μm , the SPIRE/FTS beam can be well represented by a symmetrical Gaussian profile with an FWHM of 17'' (Makiwa et al. 2013; Swinyard et al. 2014). However, this beam cannot fully cover the entire [N II] emission region in most of our targets assuming that their FIR sizes indicate the extent of the [N II] emission. To define a source as “extended” compared to the SPIRE/FTS beam, we calculated the fractional 70 and 160 μm fluxes within a Gaussian beam with an FWHM of 17'' (see below). An extended source will have both fractions less than 90%. Based on this definition, 74 galaxies are classified as extended. Therefore, to achieve our ultimate goals of deriving the LF of the [N II] 205 μm emission, as well as of further exploring the applicability of $L_{[\text{N II}]205 \mu\text{m}}$ as an SFR indicator, we need to apply an aperture correction to the observed [N II] 205 μm fluxes for most sources.

Zhao et al. (2013) found that $L_{[\text{N II}]205 \mu\text{m}}$ correlates almost linearly with L_{IR} . Hence, the aperture correction can be done by utilizing PACS photometry images and estimating the L_{IR} measured within the region outside of the SPIRE/FTS beam. However, as already shown in Zhao et al. (2013), the $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}$ ratio also depends somewhat on the FIR color. To account for this dependence and to minimize the uncertainty in the final, total $L_{[\text{N II}]205 \mu\text{m}}$, we used the FIR color-dependent $L_{[\text{N II}]205 \mu\text{m}}-L_{\text{IR}}$ relation (see below) to correct the $L_{[\text{N II}]205 \mu\text{m}}$, which was measured directly from the SPIRE/FTS spectra.

To measure the L_{IR} and FIR color within the 17'' SPIRE/FTS beam near 205 μm for our sample of (U)LIRGs, we applied the following steps. First, in order to have the same resolution as the SPIRE/FTS, we convolved the 70 and

¹⁵ <http://www.R-project.org/>

¹⁶ <http://ned.ipac.caltech.edu>




Figure 2. Distributions of (a) the FIR colors measured inside (dotted line) and outside (solid line) of the SPIRE/FTS beam and (b) the ratios of aperture-corrected $L_{[\text{N II}]205 \mu\text{m}}$ to $L_{[\text{N II}]205 \mu\text{m,beam}}$ (solid) and the total $L_{\text{IR,PACS}}$ to $L_{\text{IR,PACS,beam}}$ (dotted), for the subsample of our extended (U)LIRGs.

160 μm images (e.g., J. Chu et al. 2016, in preparation), which were obtained with PACS, with kernels computed through the algorithm described in Aniano et al. (2011). These convolution kernels were generated by comparing the PACS PSFs at 70 and 160 μm with a Gaussian profile of FWHM of $17''$ (as a representative of the SPIRE/FTS beam around 210 μm). Then, we converted the units of the convolved images from $\text{Jy}/\text{arcsec}^2$ to Jy/beam by multiplying the area of the Gaussian profile. The fluxes within the SPIRE/FTS beam at 70 and 160 μm (hereafter, $f_{70,\text{beam}}$ and $f_{160,\text{beam}}$, respectively) were measured from the convolved PACS images at the SPIRE/FTS pointing position. The total fluxes ($f_{70,\text{tot}}$ and $f_{160,\text{tot}}$), were also measured from the convolved images by doing aperture photometry. Therefore, the fluxes outside of the SPIRE/FTS beam are $f_{70,\text{out}} = f_{70,\text{tot}} - f_{70,\text{beam}}$ and $f_{160,\text{out}} = f_{160,\text{tot}} - f_{160,\text{beam}}$, for the 70 and 160 μm respectively. Note that in this subsection the *IR luminosity* ($L_{\text{IR,PACS}}$) is calculated using the f_{70} and f_{160} fluxes and the formula ($L_{\text{IR,PACS}} = 1.010\nu L(70 \mu\text{m}) + 1.218\nu L(160 \mu\text{m})$) presented in Galametz et al. (2013) since the PACS data have much higher angular resolution than the *IRAS* data, which is necessary for our purpose. However, the L_{IR} used for the remainder of our analysis is derived from the *IRAS* four-band fluxes and the well known equation given in Sanders & Mirabel (1996).

Since the galaxies in our sample are (U)LIRGs, and our SPIRE observations usually were targeted at the center of each object, the measured FIR color within the beam is very warm, whereas the part missed by the SPIRE beam, which needs to be corrected for, is generally much colder. This is illustrated in Figure 2(a), in which we plotted the distributions of the FIR color inside ($(f_{70}/f_{160})_{\text{beam}}$; dotted histogram) and outside ($(f_{70}/f_{160})_{\text{out}}$; solid histogram) of the SPIRE/FTS beam. We can see that $(f_{70}/f_{160})_{\text{beam}}$ and $(f_{70}/f_{160})_{\text{out}}$ peak at ~ 1.41 and ~ 0.45 respectively. Therefore, it is necessary to include more fiducial data with cooler FIR colors to better establish the $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}$ -FIR color relationship.

For this purpose, we include in our analysis a dozen nearby, spatially resolved galaxies, which have SPIRE/FTS mapping

observations in the HSA and are mainly from the same projects listed in Section 2.2.1. These SPIRE/FTS observations (3 of them are in the sample of the 20 galaxies mentioned in Section 2.2) were reduced with the same method as described in Section 2.2.1. The PACS imaging data of these nearby, very extended galaxies were reduced using the Scanamorphos technique (Roussel 2013) provided in HIPE 12.1, and then were convolved from their native resolutions to the $17''$ resolution of SPIRE at $\sim 210 \mu\text{m}$ using the same method as for our GOALS-FTS sample. The convolved images were rebinned to maps with pixel sizes corresponding to the SPIRE/FTS mapping observations. In order to increase the S/N for the SPIRE/FTS mapping observations, we stacked spectra from regions with similar f_{70}/f_{160} colors. Also, to reduce the uncertainties in the stacked spectrum and IR flux for each color bin, only pixels with S/N > 3 both at 70 and 160 μm were used. The [N II] 205 μm fluxes of the stacked spectrum from these galaxies were measured using the same method as for the GOALS-FTS sample described in Section 2.1 above.

The final $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR,PACS}}$ -FIR color relation is shown in Figure 3, where $L_{[\text{N II}]205 \mu\text{m}}$, $L_{\text{IR,PACS}}$, and FIR color were measured within the SPIRE/FTS beam for all of our GOALS-FTS (U)LIRGs, and for other galaxies these were measured within the stacked spaxels. This relation is rather flat for $\log(f_{70}/f_{160}) \lesssim -0.2$ (equivalent to $f_{60}/f_{100} \lesssim 0.46$; after Dale et al. 2001), but becomes much steeper when the FIR color is warmer, and has the largest scatter at the warmest end. To investigate this relation, we used the Kaplan–Meier estimate (Kaplan & Meier 1958) for censored data.¹⁷ The resulting $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR,PACS}}-f_{70}/f_{160}$ relation is shown by the solid line in Figure 3, with a scatter of 0.22 dex (compared to the observed $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR,PACS}}$). We used this relation to calculate the $L_{[\text{N II}]205 \mu\text{m}}$ outside of the SPIRE/FTS beam for our GOALS-FTS sample, and the uncertainty of 0.22 dex was propagated to the final uncertainty values for $L_{[\text{N II}]205 \mu\text{m}}$ after taking a quadratic sum of all errors.

¹⁷ Implemented in the *locfit.censor* function in the *locfit* package in *R*.




Figure 3. Correlation between the [N II] 205 μm to IR luminosity (see the text for the derivation of the IR luminosity used here) ratio and FIR color. For (U) LIRGs, $L_{[\text{N II}]205 \mu\text{m}}$, $L_{\text{IR,PACS}}$, and FIR color were measured within the SPIRE beam, whereas for other labelled individual galaxies, we used the SPIRE mapping observations and stacked spectra for similar FIR colors to measure $L_{[\text{N II}]205 \mu\text{m}}$ and $L_{\text{IR,PACS}}$ within the region of the mapped pixel (see the text for more details). The dashed (red) line shows the best polynomial fit to the results shown by the solid (black) line, which were computed using the *locfit.censor* function.

We also fitted the relation estimated by *locfit.censor* (i.e., the solid line in Figure 3) with a third-order polynomial function because an analytical form could be more convenient for future studies. The best-fit gives

$$\log(L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}) = -3.83 - 1.26x - 1.86x^2 - 0.90x^3, \quad (1)$$

with $x = \log(f_{70}/f_{160})$, and a scatter of 0.01 dex, and it is plotted as a dashed line in Figure 3. Please note this relation is only valid for the color range we have investigated, i.e., $-0.9 \leq \log(f_{70}/f_{160}) \leq 0.4$. An uncertainty of 0.23 dex should be adopted if this best-fit relation is used to compute $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}$ from f_{70}/f_{160} . As seen in Figure 3, there is a strong relation between $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}$ and FIR color for $\log(f_{70}/f_{160})_{\text{out}} > -0.2$. Given that about one-third of our extended sources have $\log(f_{70}/f_{160})_{\text{out}} > -0.2$ (see Figure 2(a)) it is necessary to use a color-dependent aperture correction for the [N II] 205 μm emission.

The aperture-corrected, total $L_{[\text{N II}]205 \mu\text{m}}$ for the extended GOALS-FTS sources were, $L_{[\text{N II}]205 \mu\text{m}} = L_{[\text{N II}]205 \mu\text{m,beam}} + L_{[\text{N II}]205 \mu\text{m,out}}$, where $L_{[\text{N II}]205 \mu\text{m,beam}}$ and $L_{[\text{N II}]205 \mu\text{m,out}}$ represent the [N II] 205 μm luminosities measured inside and outside of the SPIRE/FTS beam, respectively. $L_{[\text{N II}]205 \mu\text{m,beam}}$ was measured directly from the SPIRE/FTS spectrum, while $L_{[\text{N II}]205 \mu\text{m,out}}$ was obtained using $L_{\text{IR,PACS,out}}$ and $(f_{70}/f_{160})_{\text{out}}$. As shown in Figure 2(a), most of our galaxies have $\log(f_{70}/f_{160})_{\text{out}} < -0.2$, so the aperture correction for these sources should not be very sensitive to the FIR color as indicated by Figure 3 (and Equation (1)). The solid histogram

in Figure 2(b) shows the distribution of the $L_{[\text{N II}]205 \mu\text{m}}/L_{[\text{N II}]205 \mu\text{m,beam}}$ ratios ($\equiv f_{\text{corr}}$) for the extended sources. For about 70% of all the cases, f_{corr} is less than 2, i.e., the SPIRE/FTS beam captured more than half of the total [N II] 205 μm emission from a galaxy.

To further examine whether the color-dependent aperture correction is essential, we also plotted the $L_{\text{IR,PACS}}/L_{\text{IR,PACS,beam}}$ distribution (dotted line) in Figure 2(b). The median values of f_{corr} and $L_{\text{IR,PACS}}/L_{\text{IR,PACS,beam}}$ are 1.66 and 1.44, respectively, which indicates that the overall $L_{[\text{N II}]205 \mu\text{m}}$ would be underestimated by about 12% if we used a constant $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}$ ratio to do the aperture correction. This is insignificant compared to the scatter of the $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}$ -FIR color relation. However, the underestimation will reach 30% for sources with $\log(f_{70}/f_{160})_{\text{out}} > -0.2$. Therefore, it is still worth using a color-dependent $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}$ relation to do the aperture correction since the underestimation is systematic.

3. RESULTS AND DISCUSSION

3.1. The [N II] 205 μm Emission as an SFR Indicator

3.1.1. $L_{[\text{N II}]}$ -SFR Correlation

To estimate the SFR of our sources, we used the algorithm of Dale et al. (2007), e.g., $\text{SFR}(M_{\odot} \text{yr}^{-1}) = 4.5 \times 10^{-37} L_{\text{IR}}(W) + 7.1 \times 10^{-37} \nu L_{\nu}(1500 \text{ \AA})(W)$, which takes into account dust obscuration by combining the *IRAS* IR and *GALEX* FUV fluxes. Here L_{IR} was calculated with the *IRAS* four-band data. Without taking into account the dependence of SFR on the assumed initial mass function, the uncertainty in SFR from this composite calibrator is dominated by the uncertainty in the coefficient of the first term on the right side of the equation. Here we adopted an uncertainty of 40% (e.g., Kennicutt & Evans 2012), and it was propagated to the final SFR after taking a quadratic sum of all errors. For the GOALS-FTS sample, the FUV data were adopted from Howell et al. (2010), while for the normal galaxy and *ISO* samples, the FUV data were compiled from the literature (mainly from, e.g., Dale et al. 2007; Gil de Paz et al. 2007) and the *GALEX* data release GR7.¹⁸ Since our SFR estimate relies on the availability of UV observations, we restricted our sample to 121 galaxies with available UV photometric data (hereafter SFR sample).

Before further analysis, however, it is instructive to check whether our data set is capable of exploring a relation between $L_{[\text{N II}]205 \mu\text{m}}$ and SFR. This is due to the fact that (1) for the GOALS-FTS sample, the aperture correction is essentially the conversion of L_{IR} into $L_{[\text{N II}]205 \mu\text{m}}$, and (2) the SFRs for the GOALS-FTS galaxies are dominated by L_{IR} . Therefore, $L_{[\text{N II}]205 \mu\text{m}}$ and SFR may artificially correlate with each other even if they do not have an intrinsic relationship. As shown in Figures 4(a) and (b), sources with $f_{\text{corr}} \leq 1.5$ and $f_{\text{corr}} > 1.5$ for our GOALS-FTS sample reside in a similar phase space of $L_{[\text{N II}]205 \mu\text{m}}$ and L_{IR} . In addition, very extended sources with $f_{\text{corr}} > 2.0$ only occupy a small fraction ($\sim 17\%$) of the SFR sample. Therefore, we conclude that our data set will not artificially make a correlation between $L_{[\text{N II}]205 \mu\text{m}}$ and SFR.

In Figure 5, we plot SFR against $L_{[\text{N II}]205 \mu\text{m}}$ for both (U) LIRGs and normal galaxies. Squares represent the (U)LIRGs in the GOALS-FTS sample. Circles show normal galaxies observed by *Herschel*, whereas diamonds are the *ISO* sources from Brauer et al. (2008). The solid symbols in Figure 5

¹⁸ <http://galex.stsci.edu/GR6/#5>




Figure 4. Distributions of (a) and (b): $L_{[\text{N II}]205 \mu\text{m}}$ and L_{IR} for the GOALS-FTS sample, respectively. (c): FIR color for the SFR sample. The solid and dotted histograms in panels (a) and (b) show the results for the $f_{\text{corr}} \leq 1.5$ and $f_{\text{corr}} > 1.5$ subsamples respectively.




Figure 5. Correlation between the $[\text{N II}]205 \mu\text{m}$ luminosity and SFR. The squares and circles are galaxies with *Herschel* observations, while the diamonds are galaxies from Brauher et al. (2008), whose $L_{[\text{N II}]205 \mu\text{m}}$ were derived with the $[\text{N II}]122 \mu\text{m}$ emission. The solid symbol indicates that the AGN contributes more than 35% to the total bolometric luminosity, and are excluded from the fit. The points are color-coded according to their f_{60}/f_{100} . For each FIR color bin, the best-fit relation (slope of 1) is shown by the dashed line, with the black line showing the relation for the entire sample.

indicate that the fractional contribution from a possible active galactic nucleus (AGN) to the bolometric luminosity, f_{AGN} , is greater than 0.35. Here f_{AGN} was derived from a set of the mid-

IR diagnostics based on $[\text{Ne V}]/[\text{Ne II}]$, $[\text{O IV}]/[\text{Ne II}]$, continuum slope, polycyclic aromatic hydrocarbon equivalent width, and the diagram of Laurent et al. (2000), following the prescriptions in Armus et al. (2007; see also Veilleux et al. 2009; Petric et al. 2011; Stierwalt et al. 2013). These galaxies are excluded from our fitting procedures for the $L_{[\text{N II}]205 \mu\text{m}}$ -SFR relation(s).

From Figure 5, we can see that the scatter in the $L_{[\text{N II}]205 \mu\text{m}}$ -SFR relation becomes larger with the increase of SFR, consistent with Zhao et al. (2013). In this figure, we demonstrate that the increase in scatter is traced to the individual galaxy colors (f_{60}/f_{100}). To isolate the color dependence (and thus reduce the scatter) of the $L_{[\text{N II}]205 \mu\text{m}}$ -SFR relation, we divide our sample galaxies into three subsamples according to their f_{60}/f_{100} , i.e., a “cold” one with $0.2 \leq f_{60}/f_{100} < 0.6$ (i.e., a blackbody temperature of $30 \lesssim T \lesssim 50$ K), a “warm” one with $0.6 \leq f_{60}/f_{100} < 0.9$ ($50 \lesssim T \lesssim 60$ K), and a “hot” one with $0.9 \leq f_{60}/f_{100} < 1.4$ ($60 \lesssim T \lesssim 90$ K). These color bins were chosen according to the FIR color distribution (three peaks in Figure 4(c)) of our sample galaxies. Additional considerations for the separation of cold and warm/hot samples are that (1) starburst galaxies usually have $f_{60}/f_{100} > 0.55$ (Buat & Burgarella 1998) and (2) the turnover of the $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}-f_{60}/f_{100}$ relation happens at $f_{60}/f_{100} \sim 0.5$.

To investigate the relationship between $L_{[\text{N II}]205 \mu\text{m}}$ and SFR, we fitted each subsample using a least-squares, geometrical mean functional relationship (Isobe et al. 1990) with a linear form, i.e.,

$$\log \text{SFR}(M_{\odot} \text{ yr}^{-1}) = a + b \log L_{[\text{N II}]}(L_{\odot}). \quad (2)$$

to all galaxies, except those with $f_{\text{AGN}} > 0.35$. Using the same method, we also fitted the whole galaxy sample. To take into account the uncertainties, both in $L_{[\text{N II}]205 \mu\text{m}}$ and SFR, we used two independent approaches that allow us to evaluate their reliabilities, and to estimate the final coefficients (a , b) and associated errors in Equation (2). The first method (M1) was




Figure 6. Distributions of the fitted intercepts and slopes, which were obtained through the Monte Carlo simulations (see the text for details), for each (sub-)sample. In each panel, the numbers in parentheses give the best-fit parameters for a Gaussian function.

carried out by using a Monte Carlo simulation described as follows. First, we generated a simulated sample by assuming a Gaussian error using the measured data points and uncertainties. Second, we fitted this sample using Equation (2) and the geometrical mean method. Third, we repeated the previous two steps 10,000 times. The distributions of the fitted results from this process are shown in Figure 6. The second method (M2) is

that the observed data points were fitted by using a weighted least-squares, geometrical mean regression. The weighting is defined after Williams et al. (2010), namely, $1/\sigma^2 \equiv 1/(b^2\sigma_{L_{[NII]}}^2 + \sigma_{\text{SFR}}^2)$, where $\sigma_{L_{[NII]}}$ and σ_{SFR}^2 are the errors in $L_{[NII]205\ \mu\text{m}}$ and SFR, respectively.

Table 2 lists the number of objects in each sample, the fitting coefficients, 1σ errors, and scatters from both methods, for the

Table 2
Summary of the Fitted Coefficients for Equation (2)

Sample	N	Intercept (a)	Slope (b)	Scatter (dex)	χ^2_{red}	Method	ρ	Sig	Including <i>ISO</i> gal?
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
Cold	62	-5.24(0.24)	0.89(0.03)	0.18	...	M1	0.88	$\gg 5\sigma$	Yes
	62	-5.17(0.22)	0.88(0.03)	0.18	1.13	M2	0.88	$\gg 5\sigma$	Yes
	46	-5.26(0.38)	0.90(0.05)	0.19	...	M1	0.59	4.76 σ	No
	46	-5.12(0.24)	0.88(0.04)	0.18	1.29	M2	0.59	4.76 σ	No
Warm	39	-5.47(0.49)	0.97(0.07)	0.25	...	M1	0.83	$\gg 5\sigma$	Yes
	39	-5.53(0.53)	0.98(0.07)	0.25	1.76	M2	0.83	$\gg 5\sigma$	Yes
	35	-5.65(0.64)	1.00(0.09)	0.25	...	M1	0.78	$\gg 5\sigma$	No
	35	-5.68(0.64)	1.00(0.09)	0.25	1.83	M2	0.78	$\gg 5\sigma$	No
Hot	14	-5.00(0.80)	0.97(0.11)	0.20	...	M1	0.61	2.51 σ	...
	14	-5.04(0.85)	0.98(0.12)	0.20	1.23	M2	0.61	2.51 σ	...
All	115	-6.29(0.22)	1.07(0.03)	0.37	...	M1	0.72	$\gg 5\sigma$	Yes
	115	-6.61(0.23)	1.12(0.03)	0.37	3.23	M2	0.72	$\gg 5\sigma$	Yes
	95	-6.29(0.33)	1.07(0.03)	0.38	...	M1	0.50	5.49 σ	No
	95	-6.56(0.26)	1.11(0.03)	0.38	3.63	M2	0.50	5.49 σ	No
Fixed Slope									
Cold	62	-5.99	1.0	0.22	1.31	M2	0.88	$\gg 5\sigma$	Yes
	46	-5.99	1.0	0.22	1.52	M2	0.59	4.76 σ	No
Warm	39	-5.64	1.0	0.25	1.72	M2	0.83	$\gg 5\sigma$	Yes
	35	-5.64	1.0	0.25	1.77	M2	0.78	$\gg 5\sigma$	No
Hot	14	-5.20	1.0	0.20	1.13	M2	0.61	2.5 σ	...
All	115	-5.78	1.0	0.35	3.33	M2	0.72	$\gg 5\sigma$	Yes
	95	-5.77	1.0	0.36	3.71	M2	0.50	5.49 σ	No

Note. Column (1): samples defined according to their f_{60}/f_{100} , i.e., $0.2 \leq f_{60}/f_{100} < 0.6$, $0.6 \leq f_{60}/f_{100} < 0.9$, and $0.9 \leq f_{60}/f_{100} < 1.4$ for “Cold,” “Warm,” and “Hot,” respectively. “All” indicates that the sample includes all of the galaxies. Column (2): number of sources in each sample. Columns (3) and (4): coefficients (and the 1σ uncertainties) between $L_{[\text{N II}]205 \mu\text{m}}$ and SFR, as defined in Equation (2). Column (5): rms scatter of the relation. Column (6): reduced chi-square. Column (7): the method used to fit the relation. Columns (8) and (9): the Spearman’s rank correlation coefficient (ρ) and the level of significance (Sig; computed from the p -value using a Student’s t distribution, which approaches the normal distribution as the sample size increases), respectively. Column (10): whether the *ISO* galaxies were included in the sample. None of the *ISO* galaxies have $0.9 \leq f_{60}/f_{100} < 1.4$. The bold rows show the results that we used to calculate the 3σ range of the slope and discussed in more detail in the main text.

$L_{[\text{N II}]205 \mu\text{m}}$ -SFR relation. From the table, we can see that M1 and M2 give consistent results. Hence, we only discuss the results from M2 hereafter. In Table 2, we also show the Spearman’s rank correlation coefficient (ρ , assessing how well an arbitrary monotonic function could describe the relationship between two variables), and the level of significance (Sig), which was computed from the p -value using a Student’s t distribution. These ρ and Sig indicate that there exists a very strong correlation between $L_{[\text{N II}]205 \mu\text{m}}$ and SFR.

On average, SFR scales with $L_{[\text{N II}]205 \mu\text{m}}^b$ with b between 0.62 and 1.34 at 3σ significance. The slopes in the current work are consistent with the result (0.95 ± 0.05) found in Zhao et al. (2013) within 1σ - 2σ uncertainty ranges. The nearly linear relation between $L_{[\text{N II}]205 \mu\text{m}}$ and SFR indicates that the power source of the $[\text{N II}]205 \mu\text{m}$ emission may be related to the details of the star formation processes that take place in each galaxy. Given such a strong correlation, and to reduce any systematic uncertainties caused by the sample itself (such as sample size, dynamic range, etc.), we also fitted the $L_{[\text{N II}]205 \mu\text{m}}$ -SFR relation with a fixed slope of 1. These results are also given in Table 2, and plotted in Figure 5 as a dashed line for each sample. The reduced χ^2 from both fixed and varying slopes, as listed in Table 2, agree with each other within $< 15\%$, and thus the fitted results with a fixed slope of 1 are recommended to be used for computing SFRs.

Are the fitted relations sensitive to $R_{122/205}$ for our (sub) samples? To further check this, we fitted the $L_{[\text{N II}]205 \mu\text{m}}$ -SFR relations for the (sub)samples by excluding the *ISO* galaxies and using the method M2. We found that the resultant slopes and intercepts only have tiny changes, as shown in Table 2. Therefore, we conclude that our results are not affected substantially by including the *ISO* galaxies.

3.1.2. The Scatter in the $L_{[\text{N II}]205 \mu\text{m}}$ -SFR Relation

Table 2 shows that the scatter of each subsample is a factor of ~ 1.5 smaller compared to that of the full sample. This suggests that the color dependence of the $[\text{N II}]205 \mu\text{m}$ emission contributes significantly to the total scatter of the $L_{[\text{N II}]205 \mu\text{m}}$ -SFR relation. This is further confirmed by the following two checks: (1) a principal component analysis indicates that the FIR color accounts for 41% of the total variance of the entire sample and (2) we simply normalized the $L_{[\text{N II}]205 \mu\text{m}}$ by the galaxy FIR color, i.e., $\log L_{[\text{N II}]205 \mu\text{m}, \text{norm}} = \log L_{[\text{N II}]205 \mu\text{m}} + \log(f_{60}/f_{100})$, and then fitted the $\log L_{[\text{N II}]205 \mu\text{m}, \text{norm}}$ -SFR relations with method M2 (varying slope) and calculated the scatters, which are 0.15, 0.23, 0.20, and 0.26 dex for the “Cold,” “Warm,” “Hot,” and “All” samples, respectively. For the former three samples, these values are almost the same as those for the $L_{[\text{N II}]205 \mu\text{m}}$ -SFR




Figure 7. (a) $L_{[\text{N II}]205 \mu\text{m}}/\text{SFR}$ (in units of $L_{\odot}/(M_{\odot} \text{ yr}^{-1})$) plotted against f_{60}/f_{100} ; (b) $[\text{O III}] 88 \mu\text{m}/[\text{N II}] 122 \mu\text{m}$ vs. f_{60}/f_{100} ; and (c), $L_{[\text{N II}]205 \mu\text{m}}/\text{SFR}$ vs. $[\text{O III}] 88 \mu\text{m}/[\text{N II}] 122 \mu\text{m}$. The symbols are the same as in Figure 5. The points in panels (b) and (c), which have both $[\text{O III}] 88 \mu\text{m}$ and $[\text{N II}] 122 \mu\text{m}$ data, are outlined in panel (a) with colored symbols. In panel (a), the flat line is the mean value of $\log(L_{[\text{N II}]205 \mu\text{m}}/\text{SFR})$ for the sources with $\log f_{60}/f_{100} \leq -0.29$, and the other line gives the result of a least-squares, geometrical mean fit to the subsample with $\log f_{60}/f_{100} \geq -0.29$. The line in panel (b) is $\log(L_{[\text{O III}]88 \mu\text{m}}/L_{[\text{N II}]122 \mu\text{m}}) = 1.2 + 3.1 \log f_{60}/f_{100}$. The values of the x - and y -axis in panel (c) are obtained using Equations (4) and (5) respectively.

relation, while for the “All” sample, it is reduced by a factor of 1.3. If we “corrected” the $L_{[\text{N II}]205 \mu\text{m}}$ using the dotted lines in Figure 7(a), i.e.,

$$y = \begin{cases} 6.07 & \text{if } x \leq -0.29 \\ 5.20 - 3.00x & \text{otherwise} \end{cases} \quad (3)$$

where $y = \log(L_{[\text{N II}]205 \mu\text{m}}/\text{SFR})$, in units of $L_{\odot}/(M_{\odot} \text{ yr}^{-1})$, and $x = \log(f_{60}/f_{100})$, their scatters would become 0.15, 0.21, 0.22, and 0.18 dex respectively, and are comparable to the measurement uncertainty of 0.19 dex (median value of the entire sample).

As discussed in detail in Zhao et al. (2013), the scatter and/or color dependence in the $L_{[\text{N II}]205 \mu\text{m}}-\text{SFR}$ (or $L_{[\text{N II}]}$

$205 \mu\text{m}-L_{\text{IR}}$) relation is mainly due to the variation of ionization conditions in different galaxies. This is because the FIR color is tightly correlated with the ionization parameter, U (Abel et al. 2009; Fischer et al. 2014; Cormier et al. 2015). Adopting the $[\text{O III}] 88 \mu\text{m}$ -to- $[\text{N II}] 205 \mu\text{m}$ flux ratio as an indicator of the hardness of the radiation field, Zhao et al. (2013) also suggested that the hardness variation can largely account for the scatter. However, the $[\text{O III}] 88 \mu\text{m}/[\text{N II}] 205 \mu\text{m}$ ratio is sensitive to electron density (Rubin 1985) since the levels that emit these two lines have their critical densities differing by a factor of >10 . Therefore, here we used the $[\text{O III}] 88 \mu\text{m}/[\text{N II}] 122 \mu\text{m}$ ratio, which is insensitive to density, as a hardness indicator (Ferkinhoff et al. 2011) to further check the hardness effect.

However, we note that the $[\text{O III}] 88 \mu\text{m}/[\text{N II}] 122 \mu\text{m}$ ratio is only a good hardness indicator for a fixed U . It is correlated with U at a given hardness (Cormier et al. 2015). Therefore, for the sample having both $[\text{O III}] 88 \mu\text{m}$ and $[\text{N II}] 122 \mu\text{m}$ data (hereafter “OIII sample”), we correct $\log(L_{[\text{N II}]205 \mu\text{m}}/\text{SFR})$ (then the scatter changed from 0.36 dex to 0.28 dex), and the $[\text{O III}] 88 \mu\text{m}/[\text{N II}] 122 \mu\text{m}$ ratios using the following equations, respectively.

$$\log(L_{[\text{N II}]205 \mu\text{m}}/\text{SFR})_{\text{corr}} = \begin{cases} y & \text{if } x \leq -0.29 \\ y - (5.20 - 3.00x) + 6.07 & \text{otherwise} \end{cases} \quad (4)$$

and

$$\log(L_{[\text{O III}]88 \mu\text{m}}/L_{[\text{N II}]122 \mu\text{m}})_{\text{corr}} = y_1 - (1.2 + 3.1x) \quad (5)$$

where $y = \log(L_{[\text{N II}]205 \mu\text{m}}/\text{SFR})$, $y_1 = \log(L_{[\text{O III}]88 \mu\text{m}}/L_{[\text{N II}]122 \mu\text{m}})$, and $x = \log(f_{60}/f_{100})$. Equation (5) is the result of a least-square fit to the $[\text{O III}] 88 \mu\text{m}/[\text{N II}] 122 \mu\text{m} - f_{60}/f_{100}$ relation (see Figure 7(b)). In this way, we may eliminate/reduce the effect of U on both parameters. As shown in 7(c), there exists a weak correlation between $\log(L_{[\text{N II}]205 \mu\text{m}}/\text{SFR})$ and $\log(L_{[\text{O III}]88 \mu\text{m}}/L_{[\text{N II}]122 \mu\text{m}})$, with the Spearman’s rank coefficient $\rho = -0.5$ at a 2.1σ level of significance. The scatter of the OIII sample is reduced from 0.28 dex to 0.20 dex after further correcting for this hardness dependence. Therefore, the variations of ionization parameter and hardness can be mainly responsible for the scatter in the $[\text{N II}] 205 \mu\text{m}-\text{SFR}$ relation. Nevertheless, the OIII sample is small and biased toward higher $\log(L_{[\text{N II}]205 \mu\text{m}}/\text{SFR})$ at a given FIR color (see Figure 7(a)). A larger and more unbiased sample is needed in future studies.

3.1.3. Comparisons with Other FIR Fine-structure Line-based SFR Indicators

Other FIR atomic and ionic fine-structure lines have been studied for their application as SFR tracers (e.g., Farrah et al. 2013; De Looze et al. 2014). In comparison, our $[\text{N II}] 205 \mu\text{m}$ line-based SFR tracer fares relatively well in terms of the overall uncertainty and the systematic dependence on the FIR color. Farrah et al. (2013) found that among the six lines ($[\text{O III}] 52 \mu\text{m}$, $[\text{N III}] 57 \mu\text{m}$, $[\text{O I}] 63 \mu\text{m}$, $[\text{N II}] 122 \mu\text{m}$, $[\text{O I}] 145 \mu\text{m}$, and $[\text{C II}] 158 \mu\text{m}$) studied, the $[\text{O I}]$ and $[\text{N II}] 122 \mu\text{m}$ are the most reliable tracers of SFR for a sample of ULIRGs, and the derived SFRs for a given object are consistent to within a factor of three (0.48 dex). De Looze et al. (2014) found that the scatters are 0.46, 0.46, and 0.66 dex, for

the [C II], [O I] 63 μm and [O III] 88 μm lines respectively. Whereas the [N II] 205 μm indicator has an accuracy of <0.4 dex, which can be further improved to ~ 0.2 dex for a source with a known f_{60}/f_{100} . Therefore, the [N II] 205 μm line seems to have the smallest uncertainty, though Farrah et al. (2013) did not report the exact values of the scatter in their SFR indicators.

Many studies (i.e., Malhotra et al. 2001; Brauher et al. 2008; Graciá-Carpio et al. 2011; De Looze et al. 2014; Cormier et al. 2015) have demonstrated that most of the FIR fine-structure lines show some dependence on the dust temperature or FIR color. From ISM modeling, the [O I] 63 and 145 μm lines may be the most FIR color-insensitive lines at a given density, and the line-to-IR continuum luminosity ratios only vary by about 0.3–0.5 dex for $0.3 \leq f_{60}/f_{100} \lesssim 2.2$ (Abel et al. 2009; Cormier et al. 2015). However, De Looze et al. (2014) showed that the ratio between [O I] 63 μm -based and FUV+IR-based SFRs has a span of ~ 1.3 dex for $f_{100}/f_{160} \sim 0.5$ –2.1 (equal to $f_{60}/f_{100} \sim 0.1$ –1 for a graybody spectral energy distribution (SED) assuming a dust emissivity index $\beta = 1.5$), and the span is even larger for [C II]-based SFRs. For the [N II] 205 μm line, the $L_{[\text{N II}]205 \mu\text{m}}/\text{SFR}$ ratios vary about 1.7 dex (see Figure 7(a)) for $0.2 \leq f_{60}/f_{100} \leq 1.4$. Therefore, it seems that the [N II] 205 μm emission is not more sensitive to the FIR color than other lines, and we can conclude that the [N II] 205 μm emission is one of the most reliable SFR tracers among these fine-structure lines, in terms of its relatively small uncertainty.

3.1.4. Applicability of the [N II] 205 μm Emission as an SFR Indicator at High Redshift

The quantitative correlation between the [N II] 205 μm line and SFR for the local LIRGs implies that the [N II] 205 μm can be a particularly useful SFR tracer at high redshifts because of the following reasons. First, the [N II] 205 μm emission has an important advantage over the usual SFR calibrator, L_{IR} , for high-redshift galaxies: it might be relatively easier to be detected, since (a) it is resource intensive to construct an FIR SED to measure L_{IR} at high redshift (especially when $\lambda_{\text{rest}} \leq 100 \mu\text{m}$, it needs very good weather to observe a source at $z = 6$ even with ALMA, and gets harder for a lower redshift), but L_{IR} derived with a single-band flux is very uncertain (see the review of Carilli & Walter 2013), and (b) for sources having very little dust, the IR continuum is very weak, but the FIR fine-structure lines might still be strong (e.g., Decarli et al. 2014; Capak et al. 2015; Ferkinhoff et al. 2015).

Second, as already mentioned in Section 1, the [N II] 205 μm line has an important advantage over other shorter-wavelength FIR lines: it starts to fall into atmospheric sub/millimeter windows that have higher transmission at lower- z . Furthermore, as shown in Lu et al. (2015), the combination of the [C II] 158 μm [or CO(7-6)] and [N II] 205 μm emissions can be used to estimate the f_{60}/f_{100} (see their Equations (4)–(6)), and hence we are able to obtain a more accurate SFR for those high-redshift galaxies if both lines are measured.

However, for the application to high- z galaxies, an issue that should be explored further is whether our SFR indicator depends on metallicity because some sources at higher redshifts might have significantly lower metallicities. Indeed, low-metallicity systems ($\sim 0.25Z_{\odot}$) at high- z have already been observed (Capak et al. 2015). In our sample, there is one galaxy, ESO 350-IG 038 (Haro 11), that has a relatively low

gaseous metallicity value of $\sim 0.48 Z_{\odot}$ (Guseva et al. 2012)¹⁹, and $\log(\text{SFR}/L_{[\text{N II}]}) = -5.09 (M_{\odot} \text{ yr}^{-1} L_{\odot}^{-1})$. We also found that another dwarf galaxy, He 2–10 ($Z = 0.51Z_{\odot}$; Guseva et al. 2011), has *Herschel* FTS observations (Obs ID: 1342245083; PI: V. Leboutteiller). The derived $\log(\text{SFR}/L_{[\text{N II}]})$ for He 2–10 is $-5.28 (M_{\odot} \text{ yr}^{-1} L_{\odot}^{-1})$, where the SFR was calculated using the FIR and UV fluxes adopted from Leitherer et al. (2011). These two $\text{SFR}/L_{[\text{N II}]205 \mu\text{m}}$ ratios are both within 1σ of the fitted relation at their colors $f_{60}/f_{100} > 0.9$. Hence, it seems that the [N II] 205 μm emission is not very sensitive to metallicity. It might also be simply because the UV radiation field has a stronger influence on the [N II] emission than metallicity does, given the fact that a lower metallicity will result in a stronger UV radiation field (e.g., Binney & Merrifield 1998).

In addition, Cormier et al. (2015) found that the average $L_{[\text{N II}]122 \mu\text{m}}/L_{\text{IR}}$ ratios for their dwarf galaxy sample with $7.0 < 12 + (\text{O}/\text{H}) < 8.5$ and for the metal-rich sample in Brauher et al. (2008) only differ by a factor of ~ 2 , which is very small compared to the effects on the [O III] 88 μm line. Hence, the [N II] 205 μm line is not expected to strongly depend on metallicity down to $Z = 1/15Z_{\odot}$ if we assume that [N II] 205 $\mu\text{m}/[\text{N II}] 122 \mu\text{m}$ is not a strong function of metallicity, which is reasonable. Therefore, the [N II] 205 μm emission is one of the FIR tracers least affected by metallicity effects, and can serve as a useful SFR indicator for high-redshift galaxies.

3.2. Local LF and SFR Density

3.2.1. Local LF of the [N II] 205 μm Emission

As shown in Section 3.1, the [N II] emission can be used as an SFR calibrator, which is particularly useful for high-redshift galaxies observable with submillimeter spectroscopic facilities such as ALMA. Therefore, besides its own evolution, studying the [N II] LFs at different redshifts can probe the evolution of SFR density. In the following, we derive the local [N II] LF using the data observed by *Herschel* and *ISO*. The final sample (hereafter LF sample) used to compute the local [N II] LF contains 195 sources, in which 163 galaxies have a [N II] detection, and the remaining 32 objects only have an upper limit on the [N II] flux. For the *ISO* galaxies, $f_{[\text{N II}]205 \mu\text{m}}$ was converted from $f_{[\text{N II}]122 \mu\text{m}}$ using the method described in Section 2.2.2.

Although our GOALS-FTS sample is IR flux-limited, it is not a [N II] flux-limited sample. Also, we have added more normal galaxies into the study to improve the extent of our GOALS-FTS sample. Therefore, considering the breadth of our sample, we are able to use the so-called bivariate method (see, e.g., Xu et al. 1998) to construct the local [N II] LF, in which we transferred the local IR LF of the *IRAS* Revised Bright Galaxy Sample (RBGS) studied in Sanders et al. (2003) to a local [N II] LF utilizing the $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}$ ratio versus the L_{IR} relation. However, as demonstrated in Figure 3, the $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}$ ratio is dependent on the FIR color. Therefore, we further checked whether ours can be a representative sample of the *IRAS* RBGS. To this end, we plot the fractional distribution of L_{IR} , and the cumulative fraction of f_{60}/f_{100} in three L_{IR} bins, in the left and right panels of Figure 8, respectively. Due to the fact that our GOALS-FTS sample is complete and unbiased for

¹⁹ In the current work, we adopt $12 + (\text{O}/\text{H})_{\odot} = 8.69$ (Asplund et al. 2009).




Figure 8. Distributions of L_{IR} (left panel) and f_{60}/f_{100} (right panel) for the IRAS RBG sample (solid), which was used to construct the local infrared LF in Sanders et al. (2003), and our sample (dashed), which is used to derive the local LF of the [N II] 205 μm emission. In the right panel, the labels give the L_{IR} bins, and the number of sources in each bin (in the format of $N = \text{our sample}, \text{RBG sample}$). Only sources with $\log L_{\text{IR}} < 11 L_{\odot}$ are shown in the plots.

galaxies of $L_{\text{IR}} > 10^{11} L_{\odot}$, we plot only the sources with $\log L_{\text{IR}} < 11 L_{\odot}$ in Figure 8.

In Figure 8, we can see that, in general, our sources have similar FIR color distributions to the IRAS RBGS galaxies, albeit the discrepancy is relatively larger in the low-luminosity bins, which could be attributed to the small number statistics involved. Using the *ad. test* task in the *kSamples* package within *R*, we performed an Anderson–Darling test (Anderson & Darling 1952) on these two samples, and got a p -value of 0.17. Therefore, we do not reject the null hypothesis that these two samples arise from a common distribution. Although the discrepancy between the cumulative fractions of the f_{60}/f_{100} distribution for the least luminous L_{IR} bin with $f_{60}/f_{100} < \sim 0.6$ appears large, the $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}$ ratio is much less sensitive to the FIR color for these sources (see Figure 3). Nevertheless, one should keep this caveat in mind when using the local [N II] LF derived in the current work.

The algorithm and the formulation used in this work are the same as those presented in Section 3 of Xu et al. (1998)²⁰, which were based on the Kaplan–Meier estimator (Kaplan & Meier 1958) and therefore can take into account the information contained in the upper limits. Here we adopted the local infrared bolometric (i.e., $L_{\text{IR}}(8\text{--}1000 \mu\text{m})$) LF derived by Sanders et al. (2003). In Table 3, we list the derived local [N II] LF with bin width $\delta \log(L_{[\text{N II}]}) = 0.5 L_{\odot}$, and the associated uncertainties. In Figure 9, we plot this new local [N II] LF, as well as the best fitted Schechter (1976) function,

²⁰ The correct versions of Equations (14) and (15) of Xu et al. (1998) should have read $\text{Covar}(F_{i-1}, F_i) = F_i \text{Var}(F_{i-1})/F_{i-1}$, and $\text{Var}[\log(L_{15 \mu\text{m}}) = L_i] = \sum_j \{ \text{Var}(P_{ij}) \rho [\log(L_{25 \mu\text{m}}) = L_j]^2 + P_{ij}^2 \text{Var}[\rho \log(L_{25 \mu\text{m}}) = L_j] \} (\Delta_j/\delta_i)^2$, respectively.

Table 3
[N II] Local Luminosity Function

$\log(L_{[\text{N II}]} / L_{\odot})$	$\log[\phi / (\text{Mpc}^{-3} \text{dex}^{-1})]$	1σ Error
5.0	-2.17 (-1.96)	-2.50 (-2.46)
5.5	-2.60 (-2.87)	-2.85 (-3.16)
6.0	-2.73 (-2.19)	-3.10 (-2.91)
6.5	-2.99 (-2.85)	-3.47 (-3.20)
7.0	-3.39 (-3.50)	-3.88 (-3.78)
7.5	-4.22 (-4.29)	-4.76 (-4.59)
8.0	-5.66 (-5.66)	-6.11 (-6.10)
Fitted Parameters ^a		
$\log[\phi^* / (\text{Mpc}^{-3} \text{dex}^{-1})]$	$\log(L^* / L_{\odot})$	α
-3.83 ± 0.09	7.36 ± 0.04	-1.55 ± 0.06

Note. Numbers in parentheses give the results excluding *ISO* galaxies whose $L_{[\text{N II}]205 \mu\text{m}}$ are converted from $L_{[\text{N II}]122 \mu\text{m}}$.

^a For the Schechter (1976) function.

i.e.,

$$\phi(L)dL = \phi^* \left(\frac{L}{L^*}\right)^{\alpha} \exp\left(-\frac{L}{L^*}\right) d\left(\frac{L}{L^*}\right) \quad (6)$$

where L is the galaxy luminosity and $\phi(L)dL$ is the number density of galaxies in the luminosity range $L + dL$. The parameters, L^* , ϕ^* , and α , determined from the fit, describe the characteristic luminosity, the normalization factor at L^* , and the slope of the LF at faint luminosities, respectively. The fitted parameters are given in Table 3.

As described earlier, we included the *ISO* sources in our analysis to make our sample larger and more representative of the low- L_{IR} bins. To check how our results will change, we




Figure 9. [N II] 205 μm local differential luminosity function (squares; Table 3). The black solid line shows the smooth fit with the Schechter (1976) function to the data. In addition, we plot the predicted LFs of [N II] 205 μm at different redshifts, derived with the method in Section 3.2.3 (solid colorful lines) and the method presented in Spinoglio et al. (2012a; dotted lines) from the Gruppioni et al. (2013) IR LFs, as well as the modeling results from Orsi et al. (2014; dashed lines). See the text for details.

also derived the [N II] 205 μm LF by excluding the *ISO* galaxies (see Table 3), and found that the differences are pretty large (~ 0.2 – 0.5 dex) for $\log(L_{[\text{N II}]} / L_{\odot}) < 6.5$. This is because only 13 galaxies with $L_{\text{IR}} < 10^{10.5} L_{\odot}$ have [N II] 205 μm data. Furthermore, L_{IR} of these sources do not cover the full range of the RBG sample. However, the inclusion of the *ISO* galaxies in our analysis is not unreasonable since the uncertainty of ~ 0.2 dex in the resultant $L_{[\text{N II}]205 \mu\text{m}}$ (converted from $L_{[\text{N II}]122 \mu\text{m}}$) is much less than the luminosity bin we used to derived the LF. Nevertheless, we caution the reader to keep this caveat in mind when interpreting/using our results.

3.2.2. Local Star Formation Rate Density

It appears that the Schechter function provides a good fit to our data, with a reduced Chi-square of $\chi^2 = 0.31$ (see also Figure 9). We can further use this LF to compute the volume-averaged SFR density (SFRD) for the local universe, by integrating the fitted Schechter function and multiplying by the SFR calibration provided in Section 3.1, i.e.,

$$\begin{aligned} \log \dot{\rho}_{\text{SFR}} (M_{\odot} \text{ yr}^{-1} \text{ Mpc}^{-3}) &= \log \left(\int_0^{\infty} \phi(L) L dL \right) - 5.78 \\ &= \log \phi^* L^* \Gamma(2 + \alpha) - 5.78 \\ &= -1.96, \end{aligned} \quad (7)$$

where the adopted $L_{[\text{N II}]205 \mu\text{m}}$ –SFR relation is based on the total sample with a fixed slope (see the last line in Table 2). The quoted uncertainty in $\dot{\rho}_{\text{SFR}}$ is about 0.11 dex when only the fitted errors of the local LF are taken into account. However, in the worst case scenario, i.e., the full amount of uncertainty in the SFR calibrator propagates to the final error, the resulting uncertainty in $\dot{\rho}_{\text{SFR}}$ could be as large as 0.37 dex.

The local SFRD obtained here is in good agreement with those of Gallego et al. (1995) and Yun et al. (2001), which give $\log \dot{\rho}_{\text{SFR}}$ of -2.02 ± 0.2 and $-1.86 \pm 0.14 M_{\odot} \text{ yr}^{-1} \text{ Mpc}^{-3}$, respectively. These values were derived by assuming the same cosmological model as in this paper, and using the Kennicutt & Evans (2012) SFR calibrators of H α and IR respectively. Our [N II] 205 μm -based $\dot{\rho}_{\text{SFR}}$ is also consistent with other UV- and IR-based SFRD (e.g., Takeuchi et al. 2003; Robotham & Driver 2011), as well as the best-fitting results given by Madau & Dickinson (2014).

3.2.3. Predictions and Comparisons of the [N II] 205 μm LFs at Different Redshifts

Our results provide a local benchmark for studying the evolution of the [N II] 205 μm emission and $\dot{\rho}_{\text{SFR}}$. Here we further predict the [N II] 205 μm LFs at higher redshifts using the method described below. We will also compare our results with those from simulations given by Orsi et al. (2014), and those derived with the method presented in Spinoglio et al. (2012a).

While nearby (U)LIRGs show a [N II] deficit (relative to IR continuum emission) as compared to local normal galaxies (e.g., Graciá-Carpio et al. 2011), observations at high- z indicate that not all of the high-redshift (U)LIRGs show this similar deficit (Ferkinhoff et al. 2011, 2015; Combes et al. 2012; Decarli et al. 2012; Nagao et al. 2012). This is expected if these (U)LIRGs at high redshifts have lower FIR colors than the local galaxies of comparable luminosities. Indeed, Casey et al. (2014 and references therein) show that, for a fixed L_{IR} , the average dust temperature of a galaxy decreases as redshift increases. This implies that the “turning” point in Figure 3 might occur at a higher characteristic L_{IR} if the trend shown in that figure remains independent of redshift. Inspired by these observations, here we have derived the [N II] 205 μm LFs at different

redshifts from the IR LFs given by Gruppioni et al. (2013), by making the following three assumptions:

1. The $L_{[\text{N II}]205\ \mu\text{m}}/L_{\text{IR}}$ ratio has the same T_{dust} -dependence (see Equation (1)) at any redshift.
2. The $L_{\text{IR}}-\lambda_{\text{peak,dust}}$ relation has the same power law (i.e., $\lambda_{\text{peak,dust}} = KL_{\text{IR}}^{-0.06}$, where $\lambda_{\text{peak,dust}}$ is the SED peak wavelength and is inversely proportional to T_{dust}) at any redshift, but with the scaling factor K depending on redshift (Casey et al. 2014).
3. In order to determine the scaling factor K in (2), we used a fixed T_{dust} at $L_{\text{IR}} = L^*$, where L^* is the characteristic luminosity in the LF defined by Gruppioni et al. (2013) and is an increasing function of redshift. Here we related T_{dust} to $\lambda_{\text{peak,dust}}$ using a graybody with $\beta = 1.5$ and $\tau = 1$ at $200\ \mu\text{m}$ (e.g., Casey 2012). Generally, the measured T_{dust} at L^* for $z = 0$ and higher redshifts (Casey et al. 2014) are consistent with our assumption here.

The LFs predicted by our model calculation at various redshifts are shown in Figure 9 with solid colored lines. Our measured $[\text{N II}]205\ \mu\text{m}$ LF ($z = 0$; solid black line) from Table 3 generally agrees (within 1σ – 3σ) with the one at $0 < z < 0.3$ predicted by our method. The difference may be caused by different IR LFs (Sanders et al. LF versus Gruppioni et al. 2013 LF), and/or methods employed. For comparison, we also plotted the results (dotted lines) derived with the Spinoglio et al. (2012a) method, i.e., using an empirical $L_{\text{IR}}-L_{[\text{N II}]122\ \mu\text{m}}$ relation calibrated locally (here we have converted the $L_{[\text{N II}]122\ \mu\text{m}}$ to $L_{[\text{N II}]205\ \mu\text{m}}$ by assuming $R_{122/205} = 1$ ($n_e \sim 10\ \text{cm}^{-3}$); it would not change the LF shapes when using a different $R_{122/205}$). We can see that, both our measured and predicted LFs decline faster when $L_{[\text{N II}]205\ \mu\text{m}} > L^*$ than that based on the Spinoglio et al. (2012a) method. This difference is expected since our $L_{[\text{N II}]205\ \mu\text{m}}/L_{\text{IR}}$ ratios for warmer (and more luminous) galaxies are smaller than Spinoglio et al. (2012a).

In Figure 9, the dashed lines show the model predictions for $z = 0$ – 5 from Orsi et al. (2014) with $n_e = 10\ \text{cm}^{-3}$, which were obtained by using a semi-analytical model of galaxy formation and photoionization code. Except for one or two points, their model predicted LF at $z = 0$ is consistent with our results within 1σ – 3σ . However, the Orsi et al. (2014) LF at $z = 0$ is systematically larger than ours. The difference might be reduced by adopting a larger n_e and/or varying n_e . From the figure, we can see that, for higher redshifts, although the Orsi et al. (2014) results in the low- $L_{[\text{N II}]205\ \mu\text{m}}$ bins are much different from our predicted LFs, they show better agreement in the high-luminosity bins. The most probable reason for this disagreement is that too few points were used to construct the IR LFs at higher redshifts (Gruppioni et al. 2013), which could result in large uncertainties in the IR ($[\text{N II}]205\ \mu\text{m}$) LFs in the low-luminosity bins.

3.3. Electron Density of the Ionized Gas in (U)LIRGs

In Figure 10, the solid curve is the theoretically expected $R_{122/205}$ as a function of electron density. This was derived by assuming collisional excitation by electrons from the ground-state levels of N^+ , and using the collision strengths calculated with the fitted results (Draine 2011b) of Hudson & Bell (2005) with $T = 8000\ \text{K}$. For galaxies in our GOALS-FTS sample, 16 have $[\text{N II}]122\ \mu\text{m}$ data (12 detections and 4 upper limits), as listed in Table 2. Among these 16 galaxies, Mrk 231 has PACS observations and the flux has been measured by Fischer et al.




Figure 10. Theoretical relation between the electron density (n_e) and $[\text{N II}]122\ \mu\text{m}$ to $[\text{N II}]205\ \mu\text{m}$ ratio ($R_{122/205}$). Our measurements are shown by circles, whereas upper limits are indicated by arrows. Two data points with the smallest $R_{122/205}$ are shifted to $n_e = 1\ \text{cm}^{-3}$ since their $R_{122/205}$ are smaller than (or equal to) the theoretical limit.

(2010), whereas the other sources were observed by *ISO* and the fluxes are adopted from Brauher et al. (2008). To derive n_e , we match our observed $R_{122/205}$ (circles in Figure 10) to the theoretically expected curve. The derived n_e ranges from < 1 to tens of cm^{-3} (also see Table 4), and has a median value of $22\ \text{cm}^{-3}$ (corresponding $R_{122/205} = 1.27$) for the 12 galaxies having $[\text{N II}]122\ \mu\text{m}$ detections. This median $R_{122/205}$ for our sample is a bit larger than those of the Milky Way (0.9; Wright et al. 1991; Bennett et al. 1994) and the nearby spiral galaxy M51 (0.8–0.9; Parkin et al. 2013) as well as Cen A (0.8; Parkin et al. 2014), which might suggest that the density of the ionized medium traced by the $[\text{N II}]$ emission is higher in these (U)LIRGs. However, more data are needed to reach a solid conclusion since the current sample is too small and not representative of the (U)LIRG population.

Our results indicate that the $[\text{N II}]205\ \mu\text{m}$ emission is dominated by the low-density ionized gas even in (U)LIRGs. This might be due to the fact that, for dusty H II regions, the ionization parameter is so high in the high-density regime (U is almost proportional to density; see, e.g., Draine 2011a), such as (Ultra-)compact H II regions, that (1) the UV photons are preferentially absorbed by dust (Voit 1992; Bottorff et al. 1998; Draine 2011a), and (2) Most N^+ become N^{++} ; producing much less $[\text{N II}]$ emissions (e.g., Abel et al. 2009; Fischer et al. 2014). Hence, most of the $[\text{N II}]$ emissions should come from the low-density surrounding halos of these dense H II regions in (U)LIRGs. Indeed, high angular resolution ALMA observations of (U)LIRGs have revealed that compact regions of high gas density ($\sim 10^4\ \text{cm}^{-3}$) are surrounded by more extended low-density regions (Xu et al. 2014, 2015). This scenario is also consistent with our finding that sources with warmer f_{60}/f_{100} colors generally have lower $L_{[\text{N II}]205\ \mu\text{m}}$ -to-SFR ratios.

4. SUMMARY

We present *Herschel* SPIRE/FTS observations of the $[\text{N II}]205\ \mu\text{m}$ emission for an IR flux-limited sample of 122 (U)LIRGs. Combining them with SPIRE/FTS mapping observations of nearby normal galaxies, along with *GALEX* UV observations, PACS and *IRAS* photometry, and *ISO* $[\text{N II}]122\ \mu\text{m}$ spectra, we have studied the dependence of the

Table 4
Observed and Derived Parameters for Galaxies Observed at Both [N II] Lines

Galaxy	[N II] 122 μm^a	[N II] 205 μm	$R_{122/205}$	n_e^b (cm^{-3})
IRAS17208-0014	<30	5.4 ± 0.8	<5.52	<334.2
IRASF10565+2448	<16	3.7 ± 0.7	<4.38	<203.5
Mrk 231	4.1 ± 0.5	2.9 ± 0.5	1.40 ± 0.29	26.7 ± 1.1
Mrk 273	<14	3.7 ± 0.4	<3.83	<158.0
Mrk 331	23 ± 6	10.0 ± 0.3	2.30 ± 0.61	65.8 ± 3.1
NGC 0695	15 ± 3	15.2 ± 2.7^c	0.99 ± 0.26	11.7 ± 0.5
NGC 1068	363 ± 30	304.6 ± 124.0^c	1.19 ± 0.49	19.0 ± 1.3
NGC 1614	29 ± 6	15.6 ± 3.5^c	1.86 ± 0.56	45.6 ± 1.6
NGC 2388	36 ± 7	20.5 ± 3.5^c	1.75 ± 0.45	41.1 ± 2.6
NGC 3256	123 ± 25	90.9 ± 23.3^c	1.35 ± 0.44	25.0 ± 0.2
NGC 4194	<19	9.9 ± 0.9	<1.92	<48.4
NGC 6240	23.0 ± 2.7	15.3 ± 1.5	1.51 ± 0.23	31.0 ± 1.5
NGC 6286a	17 ± 5	33.6 ± 5.3^c	0.51 ± 0.17	<1 ^d
NGC 7552	76 ± 10	88.8 ± 27.1^c	0.86 ± 0.28	7.1 ± 0.1
NGC 7771a	47 ± 10	41.8 ± 9.7^c	1.12 ± 0.36	16.6 ± 0.3
UGC 02238	13 ± 4	20.5 ± 2.2^c	0.56 ± 0.21	<1 ^d

Notes. Fluxes are in units of $10^{-17} \text{ W m}^{-2}$.

^a All data, except Mrk 231, are adopted from Brauher et al. (2008). The data for Mrk 231 are adopted from Fischer et al. (2010).

^b The quoted error is obtained from the Monte Carlo simulation by assuming a Gaussian noise distribution.

^c Aperture corrections have been applied.

^d The observed ratios are smaller than (or approach) the theoretical lower limit.

$L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}$ ratio on the FIR color and $L_{[\text{N II}]205 \mu\text{m}}$ -SFR correlation. We also derived the [N II] 205 μm LF and SFR density for the local universe, and also investigated the electron densities for a subsample of (U)LIRGs. Our main results are summarized in the following.

1. The $L_{[\text{N II}]205 \mu\text{m}}/L_{\text{IR}}$ ratio depends on the FIR colors, while the dependence is much more sensitive when $\log(f_{70}/f_{160}) > -0.2$ (equal to $f_{60}/f_{100} > 0.46$) than $\log(f_{70}/f_{160}) \lesssim -0.2$.
2. $L_{[\text{N II}]205 \mu\text{m}}$ shows a strong, almost linear correlation with SFR, though the intercept of such a correlation varies slightly according to the $IRAS f_{60}/f_{100}$. This agrees with the conclusion in Zhao et al. (2013) that [N II] 205 μm emission can be a useful SFR indicator. The overall uncertainty is about 0.4 dex, whereas the accuracy could be improved to ~ 0.2 dex if we know the f_{60}/f_{100} independently, and use the color-dependent $L_{[\text{N II}]205 \mu\text{m}}$ -SFR relation to derive the SFR (see Table 2).
3. Using the bivariate method and combining the ISO [N II] 122 μm data, which were converted to [N II] 205 μm emission based on the empirical conversion factors, we have derived the local [N II] 205 μm LF, which can be well fitted using a Schechter (1976) function with $\phi^* = 10^{-3.83 \pm 0.09} \text{ Mpc}^{-3}$, $\alpha = -1.55 \pm 0.06$ and $L^* = 10^{7.36 \pm 0.04} L_{\odot}$.
4. As a practical application of the [N II] 205 μm LF, we have also computed the SFR volume density for the local universe using the newly derived SFR calibrator, which gives $\log \dot{\rho}_{\text{SFR}} = 1.96 \pm 0.11 \text{ M}_{\odot} \text{ yr}^{-1} \text{ Mpc}^{-3}$, in good agreement with the results of previous studies.
5. For a subsample of 12 LIRGs, which have both [N II] 205 μm and [N II] 122 μm lines observed, we have determined the electron density of the low-density ionized medium in these galaxies by matching the observed [N II] 122 $\mu\text{m}/[\text{N II}] 205 \mu\text{m}$ ratios ($R_{122/205}$) with theoretical grids, and find that n_e is from $< \sim 1 \text{ cm}^{-3}$ to $\sim 66 \text{ cm}^{-3}$, with a median value of 22 cm^{-3} .

We thank the anonymous referee for a careful reading and constructive comments. We thank Dr Álvaro Orsi for providing his simulation results of the [N II] 205 μm LFs. Support for this work was provided in part by NASA through an award issued by JPL/Caltech. Y.Z. and Y.G. are partially supported by the National Natural Science Foundation of China under grant Nos. 11173059, 11390373, and 11420101002, and the CAS pilot-b project #XDB09000000. V.C. would like to acknowledge partial support from the EU FP7 Grant PIRSES-GA-2012-316788. T.D.-S. acknowledges support from ALMA-CONICYT project 31130005 and FONDECYT project 1151239. This research has made use of the NASA/IPAC Extragalactic Database (NED), which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Abel, N. P., Dudley, C., Fischer, J., Satyapal, S., & van Hoof, P. A. M. 2009, *ApJ*, **701**, 1147
- Anderson, T. W., & Darling, D. A. 1952, *Annals of Mathematical Statistics*, **23**, 193
- Aniano, G., Draine, B. T., Gordon, K. D., & Sandstrom, K. 2011, *PASP*, **123**, 1218
- Armus, L., Charmandaris, V., Bernard-Salas, J., et al. 2007, *ApJ*, **656**, 148
- Armus, L., Mazzarella, J. M., Evans, A. S., et al. 2009, *PASP*, **121**, 559
- Arribas, S., Colina, L., Bellocchi, E., Maiolino, R., & Villar-Martín, M. 2014, *A&A*, **568**, A14
- Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, *ARA&A*, **47**, 481
- Bennett, C. L., Fixsen, D. J., Hinshaw, G., et al. 1994, *ApJ*, **434**, 587
- Binney, J., & Merrifield, M. 1998, *Galactic Astronomy* (Princeton, NJ: Princeton Univ. Press)
- Bottoff, M., Lamothe, J., Momjian, E., et al. 1998, *PASP*, **110**, 1040
- Brauher, J. R., Dale, D. A., & Helou, G. 2008, *ApJS*, **178**, 280
- Buat, V., & Burgarella, D. 1998, *A&A*, **334**, 772
- Capak, P., Carilli, C., Jones, G., et al. 2015, *Natur*, **522**, 455
- Carilli, C. L., & Walter, F. 2013, *ARA&A*, **51**, 105
- Casey, C. M. 2012, *MNRAS*, **425**, 3094
- Casey, C. M., Narayanan, D., & Cooray, A. 2014, *PhR*, **541**, 45
- Combes, F., Rex, M., Rawle, T. D., et al. 2012, *A&A*, **538**, L4
- Cormier, D., Madden, S. C., Leboutteiller, S. C., et al. 2015, *A&A*, **578**, A53

- Dale, D., Gil de Paz, A., Gordon, K. D., et al. 2007, *ApJ*, **655**, 863
- Dale, D., Helou, G., Contursi, A., Silbermann, N., & Kolhatkar, S. 2001, *ApJ*, **549**, 215
- Decarli, R., Walter, F., Carilli, C., et al. 2014, *ApJL*, **782**, L17
- Decarli, R., Walter, F., Neri, R., et al. 2012, *ApJ*, **752**, 2
- De Looze, I., Cormier, D., Leboutteiller, V., et al. 2014, *A&A*, **568**, 62
- Díaz-Santos, T., Armus, L., Charmandaris, V., et al. 2013, *ApJ*, **774**, 68
- Draine, B. T. 2011a, *ApJ*, **732**, 100
- Draine, B. T. 2011b, *Physics of the Interstellar and Intergalactic Medium* (Princeton, NJ: Princeton Univ. Press)
- Farrah, D., Leboutteiller, V., Spoon, H. W. W., et al. 2013, *ApJ*, **776**, 38
- Ferkinhoff, C., Brisbin, D., Nikola, T., et al. 2011, *ApJL*, **740**, L29
- Ferkinhoff, C., Brisbin, D., Nikola, T., et al. 2015, *ApJ*, **806**, 260
- Fischer, J., Sturm, E., González-Alfonso, E., et al. 2010, *A&A*, **518**, L41
- Fischer, J., Abel, N. P., González-Alfonso, E., et al. 2014, *ApJ*, **795**, 117
- Galametz, M., Kennicutt, R. C., Calzetti, D., et al. 2013, *MNRAS*, **431**, 1956
- Gallego, J., Zamorano, J., Aragón-Salamanca, A., & Rego, M. 1995, *ApJL*, **455**, L1
- Gil de Paz, A., Boissier, S., Madore, B. F., et al. 2007, *ApJS*, **173**, 185
- Graciá-Carpio, J., Sturm, E., Hailey-Dunsheath, S., et al. 2011, *ApJL*, **728**, L7
- Griffin, M. J., Abergel, A., Abreu, A., et al. 2010, *A&A*, **518**, L3
- Gruppioni, C., Pozzi, F., Rodighiero, G., et al. 2013, *MNRAS*, **432**, 23
- Guseva, N. G., Izotov, Y. I., Fricke, K. J., & Henkel, C. 2012, *A&A*, **541**, A115
- Guseva, N. G., Izotov, Y. I., Stasinska, G., et al. 2011, *A&A*, **529**, A149
- Hinshaw, G., Weiland, J. L., Hill, R. S., et al. 2009, *ApJS*, **180**, 225
- Ho, L. C., Filippenko, A. V., & Sargent, W. L. W. 1997, *ApJ*, **487**, 579
- Howell, J. H., Armus, L., Mazzarella, J. M., et al. 2010, *ApJ*, **715**, 572
- Hudson, C. E., & Bell, K. L. 2005, *A&A*, **430**, 725
- Hughes, T. M., Foyle, K., Schirm, M. R. P., et al. 2015, *A&A*, **575**, A17
- Isobe, T., Feigelson, E. D., Akritas, M. G., & Babu, G. J. 1990, *ApJ*, **364**, 104
- Kaplan, E. L., & Meier, P. 1958, *J. Am. Stat. Assoc.*, **53**, 457
- Kennicutt, R. C., & Evans, N. J. 2012, *ARA&A*, **50**, 531
- Kessler, M. F., Steinz, J. A., Anderegg, M. E., et al. 1996, *A&A*, **315**, L27
- Kessler, M. F., Mueller, T. G., Leech, K., et al. 2003, *The ISO Handbook*, Vol. I: Mission & Satellite Overview, Ver. 2.0 (Noordwijk: ESA)
- Kewley, L. J., Heisler, C. A., Dopita, M. A., et al. 2000, *ApJ*, **530**, 704
- Laurent, O., Mirabel, I. F., Charmandaris, V., Gallais, P., et al. 2000, *A&A*, **359**, 887
- Leitherer, C., Tremonti, C. A., Heckman, T. M., & Calzetti, D. 2011, *AJ*, **141**, 37
- Lord, S. D., Hollenbach, D. J., Colgan, S. W. J., et al. 1995, in *ASP 73, Airborne Astronomy Symposium on the Galactic Ecosystem: From Gas to Stars to Dust* (San Francisco, CA: ASP), 151
- Lu, N., Zhao, Y., Xu, C. K., et al. 2014, *ApJL*, **787**, L23
- Lu, N., Zhao, Y., Xu, C. K., et al. 2015, *ApJL*, **802**, L11
- Madau, P., & Dickinson, M. 2014, *ARA&A*, **52**, 415
- Makiwa, G., Naylor, D. A., Ferlet, M., et al. 2013, *ApOpt*, **52**, 3864
- Malhotra, S., Kaufman, M. J., Hollenbach, D., et al. 2001, *ApJ*, **561**, 766
- Mould, J. R., Huchra, J. P., Freedman, W. L., et al. 2000, *ApJ*, **529**, 786
- Nagao, T., Maiolino, R., De Breuck, C., et al. 2012, *A&A*, **542**, L34
- Oberst, T. E., Parshley, S. C., Nikola, T., et al. 2011, *ApJ*, **739**, 100
- Oberst, T. E., Parshley, S. C., Stacey, G. J., et al. 2006, *ApJL*, **652**, L125
- Ott, S. 2010, in *ASP Conf. Ser. 434, in Astronomical Data Analysis Software and Systems XIX*, ed. Y. Mizumoto, K.-I. Morita, & M. Ohishi (San Francisco, CA: ASP), 139
- Orsi, Á., Padilla, N., Groves, B., et al. 2014, *MNRAS*, **443**, 799
- Parkin, T. J., Wilson, C. D., Schirm, M. R. P., et al. 2013, *ApJ*, **776**, 65
- Parkin, T. J., Wilson, C. D., Schirm, M. R. P., et al. 2014, *ApJ*, **787**, 16
- Petric, A. O., Armus, L., Howell, J., et al. 2011, *ApJ*, **730**, 28
- Petuchowski, S. J., Bennett, C. L., Haas, M. R., et al. 1994, *ApJL*, **427**, L17
- Pilbratt, G. L., Riedinger, J. R., Passvogel, T., et al. 2010, *A&A*, **518**, L1
- Poglitich, A., Waelkens, C., Geis, N., et al. 2010, *A&A*, **518**, L2
- Robotham, A. S. G., & Driver, S. P. 2011, *MNRAS*, **413**, 2570
- Rosenberg, M. J. F., van der Werf, P. P., Aalto, S., et al. 2015, *ApJ*, **801**, 72
- Roussel, H. 2013, *PASP*, **125**, 1126
- Rubin, R. H. 1985, *ApJS*, **57**, 349
- Rubin, R. H., Simpson, J. P., Lord, S., et al. 1994, *ApJ*, **420**, 772
- Sanders, D. B., Mazzarella, J. M., Kim, D.-C., Surace, J. A., & Soifer, B. T. 2003, *AJ*, **126**, 1607
- Sanders, D. B., & Mirabel, I. F. 1996, *ARA&A*, **34**, 749
- Sargsyan, L., Samsyan, A., Leboutteiller, V., et al. 2014, *ApJ*, **790**, 15
- Schechter, P. 1976, *ApJ*, **203**, 297
- Schirm, M. R. P., Wilson, C. D., Parkin, T. J., et al. 2014, *ApJ*, **781**, 101
- Soifer, B. T., Sanders, D. B., Neugebauer, G., et al. 1986, *ApJL*, **303**, L41
- Spinoglio, L., Dasyra, K. M., Franceschini, A., et al. 2012a, *ApJ*, **745**, 171
- Spinoglio, L., Pereira-Santaella, M., Busquet, G., et al. 2012b, *ApJ*, **758**, 108
- Stacey, G. J., Geis, N., Genzel, R., et al. 1991, *ApJ*, **373**, 42
- Stierwalt, S., Armus, L., Surace, J. A., et al. 2013, *ApJS*, **206**, 1
- Swinyard, B. M., Polehampton, E. T., Hopwood, R., et al. 2014, *MNRAS*, **440**, 3658
- Takeuchi, T. T., Yoshikawa, K., & Ishii, T. T. 2003, *ApJL*, **587**, L89
- Veilleux, S., Rupke, D. S. N., Kim, D.-C., et al. 2009, *ApJS*, **182**, 628
- Voit, G. M. 1992, *ApJ*, **399**, 495
- Walter, F., Weiß, A., Riechers, D. A., et al. 2009, *ApJL*, **691**, L1
- Williams, M. J., Bureau, M., & Cappellari, M. 2010, *MNRAS*, **409**, 1330
- Wright, E. L., Mather, J. C., Bennett, C. L., et al. 1991, *ApJ*, **381**, 200
- Xu, C., Hacking, P. B., Fang, F., et al. 1998, *ApJ*, **508**, 576
- Xu, C. K., Cao, C., Lu, N., et al. 2014, *ApJ*, **787**, 48
- Xu, C. K., Cao, C., Lu, N., et al. 2015, *ApJ*, **799**, 11
- Yun, M. S., Reddy, N. A., & Condon, J. J. 2001, *ApJ*, **554**, 803
- Zhao, Y., Lu, N., Xu, C. K., et al. 2013, *ApJL*, **765**, L13