

THE EXTENDED He II $\lambda 4686$ -EMITTING REGION IN IZw 18 UNVEILED: CLUES FOR PECULIAR IONIZING SOURCES

C. KEHRIG¹, J. M. VÍLCHEZ¹, E. PÉREZ-MONTERO¹, J. IGLESIAS-PÁRAMO^{1,2},
J. BRINCHMANN³, D. KUNTH⁴, F. DURRET⁴, AND F. M. BAYO¹

¹Instituto de Astrofísica de Andalucía (CSIC), Glorieta de la Astronomía s/n, E-18008 Granada, Spain

²Estación Experimental de Zonas Áridas (CSIC), Ctra. de Sacramento s/n, La Caada, Almería, Spain

³Leiden Observatory, Leiden University, P.O. Box 9513, NL-2300 RA Leiden, The Netherlands

⁴Institut d'Astrophysique de Paris, UMR 7095, CNRS and UPMC, 98 bis Bd Arago, F-75014 Paris, France

Received 2014 December 1; accepted 2015 January 20; published 2015 March 12

ABSTRACT

New integral field spectroscopy has been obtained for IZw 18, the nearby lowest-metallicity galaxy considered to be our best local analog of systems forming at high redshift (z). Here we report the spatially resolved spectral map of the nebular He II $\lambda 4686$ emission in IZw 18, from which we derived for the first time its total He II-ionizing flux. Nebular He II emission implies the existence of a hard radiation field. He II-emitters are observed to be more frequent among high- z galaxies than for local objects. Therefore, investigating the He II-ionizing source(s) in IZw 18 may reveal the ionization processes at high z . He II emission in star-forming galaxies has been suggested to be mainly associated with Wolf-Rayet stars (WRs), but WRs cannot satisfactorily explain the He II-ionization at all times, particularly at the lowest metallicities. Shocks from supernova remnants, or X-ray binaries, have been proposed as additional potential sources of He II-ionizing photons. Our data indicate that conventional He II-ionizing sources (WRs, shocks, X-ray binaries) are not sufficient to explain the observed nebular He II $\lambda 4686$ emission in IZw 18. We find that the He II-ionizing radiation expected from models for either low-metallicity super-massive O stars or rotating metal-free stars could account for the He II-ionization budget measured, while only the latter models could explain the highest values of He II $\lambda 4686/H\beta$ observed. The presence of such peculiar stars in IZw 18 is suggestive and further investigation in this regard is needed. This letter highlights that some of the clues of the early universe can be found here in our cosmic backyard.

Key words: galaxies: dwarf – galaxies: individual (IZw 18) – galaxies: ISM – galaxies: stellar content – ISM: lines and bands

1. INTRODUCTION

He II recombination emission indicates the presence of very hard ionizing radiation with photon energies ≥ 54 eV. Star-forming galaxies with lower metallicities tend to have larger nebular He II $\lambda 4686$ line intensities compared to those with higher metallicities (e.g., Guseva et al. 2000; Schaerer 2003). While nebular He II emission has been observed in some local low metallicity (Z) starbursts (e.g., Schaerer et al. 1999; Guseva et al. 2000; Kehrig et al. 2004; Thuan & Izotov 2005), He II-emitters are apparently more frequent among high-redshift (z) galaxies than for local objects. Recent work has found that $\geq 3\%$ of the global galaxy population at $z \sim 3$ shows narrow He II lines (Cassata et al. 2013), while this number is much lower at $z \sim 0$ (Kehrig et al. 2011). The He II lines have been suggested as a good tracer of Population III stars (PopIII-stars; the first very hot metal-free stars) in high- z galaxies (e.g., Schaerer 2003, 2008). These stars, which should produce a large amount of hard ionizing radiation, are believed to have contributed significantly to the universe's reionization, a challenging subject in contemporary cosmology (e.g., Bromm 2013). Before interpreting the emission-line spectra of distant star-forming galaxies, it is crucial to understand the formation of high-ionization lines in the nearby universe. The ideal place to perform this study is in extremely metal-poor nearby galaxies with nebular He II emission, which are the natural local counterparts of distant He II-emitters.

In this regard, we have been carrying out a program to investigate nearby low- Z starburst systems using the integral field spectroscopy technique (e.g., Kehrig et al. 2008, 2013;

Pérez-Montero et al. 2011, 2013). As a part of this program, we have recently obtained new deep integral field spectroscopic (IFS) data of IZw 18. This is a nearby ($D = 18.2$ Mpc; Aloisi et al. 2007)⁵ HII galaxy, well known for its extremely low $Z \sim 1/32$ solar (e.g., Vílchez & Iglesias-Páramo 1998), which places IZw 18 among the three most metal-poor galaxies known in the local universe (e.g., Thuan et al. 2004). Its observational characteristics make IZw 18 an excellent local analog of primeval systems (see e.g., Leboutteiller et al. 2013 and references therein).

The presence of the nebular He II $\lambda 4686$ line in the spectrum of IZw 18 has been reported before, although the precise location and extension of this He II $\lambda 4686$ emission is not known (e.g., Garnett et al. 1991; Izotov et al. 1997; Legrand et al. 1997; Vílchez & Iglesias-Páramo 1998). Our unique IFS data unveil for the first time the entire He II $\lambda 4686$ -emitting region and its structure in IZw 18 (see Section 3.2).

Despite various attempts to explain the origin of the nebular He II emission in HII galaxies/regions, it still remains difficult to understand in many cases; several potential mechanisms (e.g., hot Wolf-Rayet (WR) stars, shocks from supernova remnants, X-ray sources) have been proposed to account (in part or fully) for the He II ionization in these objects (e.g., Garnett et al. 1991; Schaerer 1996; Dopita & Sutherland 1996; Cerviño, Mas-Hesse & Kunth 2002; Thuan & Izotov 2005; Kehrig et al. 2011; Shirazi & Brinchmann 2012). Although hot WRs have previously been suggested as the source of

⁵ A distance of 18.2 Mpc is assumed in this work.




Figure 1. Color-composite image of IZw 18 (blue = $H\alpha$ from Palomar, green = far-UV/GALEX, red = SDSS r'). The box represents the FOV ($16'' \times 16''$) of the PMAS spectrograph over the galaxy main body and the extended $H\alpha$ halo. The PMAS FOV is centered on the coordinates R.A. (J2000.0) = $09^{\text{h}}:34^{\text{m}}:02.^{\text{s}}.2$ and decl. (J2000.0) = $+55^{\circ}:14':25''$.

He II -ionizing photons in IZw 18 (e.g., Izotov et al. 1997; de Mello et al. 1998), the main mechanism powering the nebular He II emission in this galaxy is still an open issue.

In this Letter, using new IFS data, we derive for the first time the total $\text{He II}\lambda 4686$ -ionizing flux in IZw 18 and provide new clues to constrain the sources of high ionization.

2. INTEGRAL FIELD SPECTROSCOPIC DATA

We carried out new IFS observations of IZw 18 using the Potsdam Multi-Aperture Spectrophotometer (PMAS; Roth et al. 2005) on the 3.5 m telescope at the Calar Alto Observatory (Almeria, Spain). The data were taken in 2012 December with a typical seeing of $1''$. Each spaxel has a spatial sampling of $1'' \times 1''$ on the sky resulting in a field of view (FOV) of $16'' \times 16''$ ($\sim 1.4 \text{ kpc} \times 1.4 \text{ kpc}$ on IZw 18; see Figure 1). One pointing of IZw 18, encompassing its main body which hosts the two brightest stellar clusters (referred to as the NW and SE knots), was taken during a 2.5 hr integration split into six exposures of 1500 s each. We used the V500 grating, which covers from ~ 3640 to 7200 \AA and provides a linear dispersion of $\sim 2 \text{ \AA}/\text{pixel}$ and an FWHM effective spectral resolution of $\sim 3.6 \text{ \AA}$. Calibration images (exposures of standard star, arc, and continuum lamps) were also obtained. The data reduction was performed following the procedure described in Kehrig et al. (2013).

3. RESULTS

3.1. Emission-line Flux Maps

The emission-line fluxes were measured using the IRAF⁶ task SPLOT. The flux of each emission line was derived by integrating between two points given by the position of a local continuum placed by eye. For each line, this procedure was repeated several times by varying the local continuum position. The final flux of each line and its associated uncertainty were assumed to be the average and standard deviation of the independent, repeated measurements (e.g., Kehrig et al. 2006).

⁶ IRAF is distributed by the National Optical Astronomical Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

We used our own IDL scripts to create the emission-line maps presented in Figure 2. The spaxels where we measure $\text{He II}\lambda 4686$ are indicated with pluses.

3.2. The Spatially Resolved $\text{He II}\lambda 4686$ -emitting Region

An extended $\text{He II}\lambda 4686$ -emitting region with a diameter of $\approx 5''$ ($\approx 440 \text{ pc}$) is revealed from our IFS data (see Figure 2). The narrow line profile for the $\text{He II}\lambda 4686$ emission and its spatial extent are evidence of its nebular nature. The spectra of certain Of stars may exhibit somewhat narrow $\text{He II}\lambda 4686$ lines (e.g., Massey et al. 2004), so He II emission observed in starburst galaxies could be thought to arise in the atmospheres of such stars (e.g., Bergeron 1977). However, if these Of stars were present in appreciable numbers in IZw 18, then broad $H\alpha$ emission would be expected (Massey et al. 2004). Thus, the non-detection of this feature in our spectra supports the nebular origin for the $\text{He II}\lambda 4686$ line in IZw 18.

By adding the emission from the spaxels showing $\text{He II}\lambda 4686$ (see Figure 2), we created the one-dimensional spectrum for the $\text{He II}\lambda 4686$ region (see Figure 3). Using this spectrum, we obtained a very small logarithmic extinction coefficient $c(H\beta) = 0.08 \pm 0.02$ from the observed ratio $H\alpha/H\beta = 2.92 \pm 0.04$, assuming an intrinsic case B recombination $H\alpha/H\beta = 2.75$ (Osterbrock & Ferland 2006, OF06). Using PYNEB (Luridiana et al. 2015), we derived the electron temperature, $T_e = (2.18 \pm 0.05) \times 10^4 \text{ K}$, and density, $n_e \leq 100 \text{ cm}^{-3}$, from the $[\text{O III}]\lambda 4363/[\text{O III}]\lambda \lambda 4959, 5007$ and $[\text{S II}]\lambda 6717/[\text{S II}]\lambda 6731$ ratios, respectively. The integrated flux of the $\text{He II}\lambda 4686$ line, corrected for reddening, is $(2.84 \pm 0.18) \times 10^{-15} \text{ erg s}^{-1} \text{ cm}^{-2}$ which translates to an $\text{He II}\lambda 4686$ luminosity of $L_{\text{HeII}\lambda 4686} = (1.12 \pm 0.07) \times 10^{38} \text{ erg s}^{-1}$. The corresponding He II ionizing photon flux, $Q(\text{He II})_{\text{obs}} = (1.33 \pm 0.08) \times 10^{50} \text{ photon s}^{-1}$, was derived from the measured $L_{\text{HeII}\lambda 4686}$ using the relation $Q(\text{He II}) = L_{\text{HeII}\lambda 4686}/[j(\lambda 4686)/\alpha_B(\text{He II})]$ (assuming case B recombination, and $T_e = 2 \times 10^4 \text{ K}$; OF06). We checked that different CLOUDY models (Ferland et al. 2013) computed for the typical n_e and T_e in the $\text{He II}\lambda 4686$ region, and considering several effective temperatures and geometries with no dust, provide a ratio $L(\text{He II}\lambda 4686)/Q(\text{He II})$ which agrees with the assumed ratio (OF06) within 10%. The total $Q(\text{He II})_{\text{obs}}$, a quantity not reported before for IZw 18, will allow us to constrain possible ionizing sources of He II in IZw 18.

4. DISCUSSION

4.1. Ionizing Sources of He II

One widely favored mechanism for He II ionization in H II galaxies involves hot WRs (e.g., Schaerer 1996). Nevertheless, it has been demonstrated that nebular $\text{He II}\lambda 4686$ does not appear to be always associated with WRs, as is the case of the He II nebulae LMC N44C, LMC N159F, and M33 BCLMP651, among others (Kehrig et al. 2011 and references therein). This indicates that WRs cannot explain He II ionization in all cases, particularly at low Z (e.g., Guseva et al. 2000; Kehrig et al. 2008, 2013; Schaerer 2003; Shirazi & Brinchmann 2012). Besides, it has been shown that for a sample of $\text{He II}\lambda 4686$ -emitting star-forming galaxies, current models of massive stars predict $\text{He II}\lambda 4686$ emission and $\text{He II}\lambda 4686/H\beta$ ratios only for $Z > 0.20 Z_{\odot}$ instantaneous bursts (Shirazi & Brinchmann 2012). In the particular case of IZw 18, previous work claimed that the ratio $\text{He II}\lambda 4686/H\beta$ could be




Figure 2. Emission-line flux maps of IZw 18. Maps are displayed in logarithmic scale and the fluxes are in units of $\text{erg s}^{-1} \text{cm}^{-2}$; the area of each spaxel is 1 arcsec^2 on the sky. *Top row:* $\text{H}\alpha$ and $\text{He II } \lambda 4686$ maps. *Middle row:* for display purposes, the maps of $\text{H}\alpha$ and $\text{He II } \lambda 4686$ are presented as color-filled contour plots and were smoothed using bilinear interpolation. Isocontours of the $\text{H}\alpha$ emission line flux are shown overplotted for reference. *Bottom row:* $[\text{O I}] \lambda 6300$ and $[\text{S II}] \lambda \lambda 6717 + 6731$ maps. The spaxels where we detect nebular $\text{He II } \lambda 4686$ are marked with pluses on the maps of $\text{H}\alpha$ (*top row*), and $[\text{O I}] \lambda 6300$ and $[\text{S II}] \lambda \lambda 6717 + 6731$ (*bottom row*). The spaxels with no measurements available are left blank.




Figure 3. Portion ($\sim 4425\text{--}5100 \text{ \AA}$) of the integrated spectrum of the He II $\lambda 4686$ -emitting region of IZw 18.

reproduced using highly density-bounded photoionization models while underpredicting the electron temperature measurement (Stasińska & Schaerer 1999); these models have been challenged by Péquignot (2008).

Faint broad emission signatures, attributable to WRs, are observed in the spectrum of IZw 18 despite its low Z (e.g., Izotov et al. 1997; Legrand et al. 1997; Brown et al. 2002). A comprehensive study of WRs in IZw 18, using UV STIS spectroscopy, revealed signatures of carbon-type WRs (WC) in two clusters: one in the NW star-forming region and a second one on the outskirts of this region (Brown et al. 2002). Here we deal with the NW one, since it is in the He II $\lambda 4686$ -emitting region (see Figure 2). The C IV $\lambda 1550$ flux measured from the NW WR cluster (Brown et al. 2002) provides a C IV $\lambda 1550$ luminosity of $L_{1550} = 4.67 \times 10^{37} \text{ erg s}^{-1}$. Taking the L_{1550} luminosity of the metal-poor, early-type WC (WCE) model by Crowther and Hadfield (2006, CH06), which mimics a single WCE star in IZw 18, this implies ~ 9 IZw 18-like WCE stars present in the NW cluster. From these nine WCE stars, a total flux of $Q(\text{He II}) = 2.8 \times 10^{48} \text{ photon s}^{-1}$ is expected (assuming $Q(\text{He II}) = 10^{47.5} \text{ photon s}^{-1}$ for one IZw 18-like WCE; CH06), i.e., about 48 times lower than the $Q(\text{He II})_{\text{obs}} = (1.33 \pm 0.08) \times 10^{50} \text{ photon s}^{-1}$ derived from our data (see Section 3.2).

Based on the He II-ionizing flux expected from these IZw 18-like WRs, a very large WR population is required to explain the He II-ionization budget measured; for instance, taking the $Q(\text{He II}) = 10^{47.5} \text{ photon s}^{-1}$ for one IZw 18-like WCE (CH06), the number of these WCEs needed to explain our derived $Q(\text{He II})_{\text{obs}} = (1.33 \pm 0.08) \times 10^{50} \text{ photon s}^{-1}$ would be >400 . In principle, the presence of hundreds of WRs in IZw 18 should not be discarded on the basis of empirical arguments for reduced WR line luminosity at low Z (CH06). However, assuming a Salpeter initial mass function (IMF; Salpeter 1955; $M_{\text{up}} = 150 M_{\odot}$) and the initial mass needed for a star to certainly become a WC (Meynet & Maeder 2005), a cluster with >8 times the total stellar mass of the NW region ($M_{*,\text{NW}} = 2.9 \times 10^5 M_{\odot}$ from Stasińska & Schaerer 1999 scaled to 18.2 Mpc distance) is required to provide >400 WC in IZw 18. Also, such a high number of WRs is not supported by state-of-the-art stellar evolutionary models for single (rotating and non-rotating) massive stars in metal-poor environments (Leitherer et al. 2014). Furthermore, given the decrease in the ratio of WR/O stars with decreasing metallicity, shown by observations and theoretical models (Maeder & Meynet 2012 and references therein), such a large number of WRs appears clearly unreasonable considering the extremely low Z and

the O star content of IZw 18 (CH06). All this suggests that WRs are not solely responsible for the He II $\lambda 4686$ emission in IZw 18.

The binary channel in massive star evolution is suggested to increase the WR population (e.g., Eldridge et al. 2008), but the WR population in Local Group galaxies does not show an increased binary rate at lower Z (e.g., Foellmi et al. 2003; Neugent & Massey 2014); thus, the binary channel does not seem to favor the formation of WRs at lower- Z , in contrast to what we need. Nevertheless, we should bear in mind the possible uncertainties still unsolved in the models (Maeder & Meynet 2012). Further investigation awaits the calculation of evolutionary models for binary stars at very low Z .

As mentioned before, nebular He II $\lambda 4686$ emission observed in the spectra of H II galaxies and extragalactic H II regions has also often been attributed to shocks and X-ray sources (e.g., Pakull & Mirioni 2002; Garnett et al. 1991; Thuan & Izotov 2005). In the following, we discuss these two candidate sources for He II ionization in IZw 18.

The X-ray emission from IZw 18 is dominated by a single X-ray binary apparently located in the field of the NW knot (Thuan et al. 2004). We have computed a CLOUDY photoionization model (Ferland et al. 2013) using as input a power-law spectral energy distribution (SED) with the same X-ray luminosity, column density, and slope that have been reported for IZw 18 (Thuan et al. 2004). This CLOUDY model provides an He II $\lambda 4686$ luminosity of $L_{\text{HeII}\lambda 4686} = 10^{35.7} \text{ erg s}^{-1}$, which is ~ 100 times lower than the $L_{\text{HeII}\lambda 4686}$ measured. This result rules out the X-ray binary as the main source of He II ionizing photons in IZw 18. We note here that the emission from X-ray ionized nebulae has been successfully reproduced by CLOUDY models before (e.g., Pakull & Mirioni 2002).

Guided by the existence of He II $\lambda 4686$ emission associated with supernova remnants (e.g., Kehrig et al. 2011) we explored the conjecture that the He II $\lambda 4686$ region represents such a shock-ionized nebula. The [O I] $\lambda 6300$ line, often strong in remnants, has frequently been used as a sensitive shock-emission test (e.g., Skillman 1985). We checked that, in fact, most of the [O I] $\lambda 6300$ emission in IZw 18 is concentrated on the SE knot (see the [O I] $\lambda 6300$ map in Figure 2) with only 12% of the He II $\lambda 4686$ -emitting spectra showing [O I] $\lambda 6300$ flux above the 3σ detection limit. Additionally, we find no evidence for [S II] enhancement (a usual sign of shock excitation; e.g., Dopita & Sutherland 1996) associated with the He II $\lambda 4686$ region (see the [S II] map in Figure 2). Therefore, the He II $\lambda 4686$ -emitting zone in IZw 18 is unlikely to be produced by shocks.

4.2. Peculiar Very Hot Stars in IZw 18?

Our new observations have allowed us to empirically demonstrate why conventional He II-ionizing sources (e.g., WRs, shocks, X-ray binaries) cannot account for the total He II-ionization budget in IZw 18. What could the nebular He II $\lambda 4686$ emission in IZw 18 originate from?

We have also explored the possibility of very massive, metal-poor O stars to account for our observations of IZw 18. Using current wind models of very massive O stars at low Z , we can derive their He II-ionizing fluxes (Kudritzki 2002). According to the hottest models ($T_e = 60,000\text{K}$), between 10 and 20 super-massive stars with $300 M_{\odot}$ (with $Q(\text{He II}) \approx (0.70\text{--}1.4) \times 10^{49} \text{ photon s}^{-1}$ each) would be sufficient to

explain the derived $Q(\text{He II})_{\text{obs}}$ budget. Very massive stars of up to $300 M_{\odot}$ were claimed to exist in the LMC R136 cluster (Crowther et al. 2010); however, the existence of such super-massive $300 M_{\odot}$ stars remains heavily debated (Vink 2014). Additionally, assuming a Salpeter IMF, 10–20 stars with $290 \leq M_{*}/M_{\odot} \leq 310$ would imply a cluster mass of $\sim 10\text{--}20 \times M_{*,\text{NW}}$. We should bear in mind that an extrapolation of the IMF predicting $300 M_{\odot}$ stars remains unchecked up to now. If we instead consider the $150 M_{\odot}$ star hottest models (with $Q(\text{He II}) \leq 1.9 \times 10^{47}$ photons $^{-1}$ each, for $Z \leq 1/32 Z_{\odot}$; K02), the number of these stars required to explain the $Q(\text{He II})_{\text{obs}}$ would be >650 . For a Salpeter IMF, 650 stars with $145 \leq M_{*}/M_{\odot} \leq 155$ would require a cluster mass $\sim 200 \times M_{*,\text{NW}}$. Besides the $Q(\text{He II})_{\text{obs}}$ budget, in the He II $\lambda 4686$ region we have measured He II $\lambda 4686/H\beta$ ratios as high as 0.08. These values appear too big to be explained even by the models for the hottest, most metal-poor super-massive $300 M_{\odot}$ stars (K02) under ionization-bounded conditions. Further constraints to the observations should await the calculation of new evolutionary tracks and SEDs for single O stars at the metallicity of IZw 18 including rotation.

Searches for very metal-poor starbursts and PopIII-hosting galaxies have been carried out in the distant universe using He II lines (e.g., Schaerer 2008; Cassata et al. 2013). This search is based on the high effective temperature for PopIII-stars which will emit a large number of photons with energy above 54eV, and also on the expected increase of the He II recombination lines with decreasing Z (e.g., Guseva et al. 2000; Schaerer 2003, 2008). Predictions for burst models of different metallicities show how their corresponding $Q(\text{He II})$ can increase by up to $\sim 10^3$ when going from $Z = 10^{-5}$ to 0 (Schaerer 2003). These models cannot explain the $Q(\text{He II})_{\text{obs}}$ when $Z \geq 10^{-5}$ (for a Salpeter IMF, $M_{\text{up}} = 100 M_{\odot}$; see table 3 in Schaerer 2003) even assuming that the total $M_{*,\text{NW}}$ would come from He II-ionizing stars. So another more speculative possibility to explain the derived $Q(\text{He II})_{\text{obs}}$ could be based on nearly metal-free ionizing stars. These stars should ionize He II via their strong UV radiation expected at nearly zero metallicity (e.g., Tumlinson & Shull 2000; Schaerer 2003).

As an approximation of nearly metal-free single stars, we have compared our observations with the He II-ionizing radiation expected from state-of-the-art models for rotating $Z = 0$ stars (Yoon et al. 2012). According to these models, we found that a handful of such stars could explain our derived $Q(\text{He II})_{\text{obs}}$ (e.g., $\sim 8\text{--}10$ stars with mass $M_{\text{ini}} = 150 M_{\odot}$ or $\sim 13\text{--}15$ stars with $M_{\text{ini}} = 100 M_{\odot}$; with $Q(\text{He II}) \approx 1.4 \times 10^{49}$ (0.9×10^{49}) photon s^{-1} for each star with $M_{\text{ini}} = 150 M_{\odot}$ ($100 M_{\odot}$)). Additionally, we note that the ionizing spectra produced by these star models are harder than the ones expected from the hottest models of super-massive $300 M_{\odot}$ stars (K02), so they would also explain the highest He II $\lambda 4686/H\beta$ values observed, providing that ionization-bounded conditions are met. While gas in IZw 18 is very metal-deficient but not primordial, Leboutteiller et al. (2013) have pointed out that the H I envelope of IZw 18 near the NW knot contains essentially metal-free gas pockets. These gas pockets could provide the raw material for making such nearly metal-free stars. Clearly, in this hypothetical scenario, these extremely metal-poor stars cannot belong to the NW cluster, which hosts more chemically evolved stars.

5. SUMMARY AND CONCLUDING REMARKS

This letter reports on new optical IFS observations of the nearby dwarf galaxy IZw 18. This is an extremely metal-poor system, which is our best local laboratory for probing the conditions dominating in distant low- Z starbursts. Our IFS data reveal for the first time the total spatial extent (≈ 440 pc diameter) of the He II $\lambda 4686$ -emitting region and corresponding total He II-ionizing photon flux in IZw 18. The metal-poor sensitivity of the He II line is a primary motivation to develop diagnostics for unevolved starbursts, and strong nebular He II emission is expected to be one of the best signatures of massive PopIII-stars (e.g., Schaerer 2003, 2008). He II emission has been observed to be more frequent at higher- z than locally (Kehrig et al. 2011; Cassata et al. 2013). Thus the analysis of the origin of the He II $\lambda 4686$ nebular line in relatively close ionized regions, which can be studied in more detail, can yield insight into the ionizing sources in the distant universe.

Our observations combined with stellar model predictions point out that conventional excitation sources (e.g., WRs, shocks, X-ray binaries) cannot convincingly explain the He II-ionizing energy budget derived for IZw 18. Other mechanisms are probably also at work. If the He II-ionization in IZw 18 is due to stellar sources, these might be peculiar very hot stars (perhaps uncommon in local starbursts but somewhat more frequent in distant galaxies): according to theoretical stellar models, either super-massive O stars or nearly metal-free ionizing stars could in principle account for the total $Q(\text{He II})_{\text{obs}}$ of IZw 18. However, the super-massive O stars scenario would imply a cluster mass much higher than the mass of the NW knot derived from observations. On the other hand, though metal-free gas pockets were previously reported in IZw 18 (Leboutteiller et al. 2013), we highlight that the existence of nearly metal-free ionizing stars is not yet confirmed observationally. The work presented here can help in the preparation of prospective searches for primeval objects, one of the main drivers for next-generation telescopes (e.g., Bromm 2013).

This work has been partially funded by research projects AYA2010-21887-C04-01 from the Spanish PNA YA, and PEX2011-FQM7058 from Junta de Andalucia. F.D. and D.K. gratefully acknowledge support from the Centre National d'Etudes Spatiales. We express our appreciation to Leslie Sage for his help and suggestions. We also thank Manfred Pakull for his useful comments on this letter. Thanks are due to Jose Luis Ortiz for his careful reading of the manuscript.

REFERENCES

- Aloisi, A., Clementini, G., Tosi, M., et al. 2007, *ApJL*, 667, L151
 Bergeron, J. 1977, *ApJ*, 211, 62
 Bromm, V. 2013, *RPPH*, 76, 112901
 Brown, T. M., Heap, S. R., Hubeny, I., et al. 2002, *ApJL*, 579, L75
 Cassata, P., Le Fèvre, O., Charlot, S., et al. 2013, *A&A*, 556, A68
 Cerviño, M., Mas-Hesse, J. M., & Kunth, D. 2002, *A&A*, 392, 19
 Crowther, P. A., & Hadfield, L. J. 2006, *A&A*, 449, 711
 Crowther, P. A., Schnurr, O., Hirschi, R., et al. 2010, *MNRAS*, 408, 731
 de Mello, D. F., Schaerer, D., Heldmann, J., & Leitherer, C. 1998, *ApJ*, 507, 199
 Dopita, M. A., & Sutherland, R. S. 1996, *ApJS*, 102, 161
 Eldridge, J. J., Izzard, R. G., & Tout, C. A. 2008, *MNRAS*, 384, 1109
 Ferland, G. J., Porter, R. L., van Hoof, P. A. M., et al. 2013, *Rev. Mex. Astron. Astrofis.*, 49, 137
 Foellmi, C., Moffat, A. F. J., & Guerrero, M. A. 2003, *MNRAS*, 338, 360
 Garnett, D. R., Kennicutt Jr., R. C., You-Hua, C., Skillman, E. D., et al. 1991, *ApJ*, 373, 458

- Guseva, N. G., Izotov, Y. I., & Thuan, T. X. 2000, [ApJ](#), **531**, 776
- Izotov, Y. I., Foltz, C. B., Green, R. F., Guseva, N. G., & Thuan, T. X. 1997, [ApJL](#), **487**, L37
- Kehrig, C., Telles, E., & Cuisinier, F. 2004, [AJ](#), **128**, 1141
- Kehrig, C., Vílchez, J. M., Telles, E., Cuisinier, F., & Pérez-Montero, E. 2006, [A&A](#), **457**, 477
- Kehrig, C., Vílchez, J. M., Sánchez, S. F., et al. 2008, [A&A](#), **477**, 813
- Kehrig, C., Oey, M. S., Crowther, P. A., et al. 2011, [A&A](#), **526**, A128
- Kehrig, C., Pérez-Montero, E., Vílchez, J. M., et al. 2013, [MNRAS](#), **432**, 2731
- Kudritzki, R. P. 2002, [ApJ](#), **577**, 389
- Lebouteiller, V., Heap, S., Hubeny, I., & Kunth, D. 2013, [A&A](#), **553**, A16
- Legrand, F., Kunth, D., Roy, J.-R., Mas-Hesse, J. M., & Walsh, J. R. 1997, [A&A](#), **326**, L17
- Leitherer, C., Ekström, S., Meynet, G., et al. 2014, [ApJS](#), **212**, 14
- Luridiana, V., Morisset, C., & Shaw, R. A. 2015, [A&A](#), **573**, A42
- Maeder, A., & Meynet, G. 2012, [RvMP](#), **84**, 25
- Massey, P., et al. 2004, [ApJ](#), **608**, 1001
- Meynet, G., & Maeder, A. 2005, [A&A](#), **429**, 581
- Neugent, K. F., & Massey, P. 2014, [ApJ](#), **789**, 10
- Osterbrock, D. E., & Ferland, G. J. 2006, in *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei* (CA: University Science Books)
- Pakull, M. W., & Mirioni, L. 2002, Proc. Symp. “New Visions of the X-ray universe in the XMM-Newton and *Chandra* Era” (Noordwijk: ESTEC)
- Péquignot, D. 2008, [A&A](#), **478**, 371
- Pérez-Montero, E., et al. 2011, [A&A](#), **532**, A141
- Pérez-Montero, E., Kehrig, C., Brinchmann, J., et al. 2013, [AdAst](#), **837**, 392
- Roth, M. M., Kelz, A., Fechner, T., et al. 2005, [PASP](#), **117**, 620
- Salpeter, E. E. 1955, [ApJ](#), **121**, 161
- Schaerer, D. 1996, [ApJ](#), **467**, L17
- Schaerer, D., Contini, T., & Pindao, M. 1999, [A&AS](#), **136**, 35
- Schaerer, D. 2003, [A&A](#), **397**, 527
- Schaerer, D. 2008, [IAUSS](#), **255**, 66
- Shirazi, M., & Brinchmann, J. 2012, [MNRAS](#), **421**, 1043
- Skillman, E. D. 1985, [ApJ](#), **290**, 449
- Stasińska, G., & Schaerer, D. 1999, [A&A](#), **351**, 72
- Thuan, T. X., Bauer, F. E., Papaderos, P., & Izotov, Y. I. 2004, [ApJ](#), **606**, 213
- Thuan, T. X., & Izotov, Y. I. 2005, [ApJS](#), **161**, 240
- Tumlinson, J., & Shull, J. M. 2000, [ApJL](#), **528**, L65
- Vílchez, J. M., & Iglesias-Paramo, J. 1998, [ApJ](#), **508**, 248
- Vink, J. S. 2014, arXiv:1406.4836
- Yoon, S.-C., Dierks, A., & Langer, N. 2012, [A&A](#), **542**, A113