

BJR



■ RESEARCH

Hip and knee replacement patients prefer pen-and-paper questionnaires

IMPLICATIONS FOR FUTURE PATIENT-REPORTED OUTCOME MEASURE STUDIES

**J. C. Keurentjes,
M. Fiocco,
C. So-Osman,
R. Onstenk,
A. W. M. M.
Koopman-Van
Gemert,
R. G. Pöll,
R. G. H. H. Nelissen**

From Leiden
University Medical
Center, Leiden, the
Netherlands

■ J. C. Keurentjes, MD, Research
■ R. G. H. H. Nelissen, MD, PhD,
Professor, Orthopaedic Surgeon
Leiden University Medical Center,
Department of Orthopaedic
Surgery, Albinusdreef 2, PO Box
9600, Postzone J-11-S, 2300 RC
Leiden, the Netherlands.

■ M. Fiocco, PhD, Assistant
Professor in Medical Statistics
Leiden University Medical Center,
Department of Medical Statistics
and Bioinformatics, Albinusdreef
2, PO Box 9600, Postzone J-11-S,
2300 RC Leiden, the Netherlands.

■ C. So-Osman, MD, PhD,
Clinical Consultant in Transfusion
Medicine
Sanguin Blood Bank,
Plesmanlaan 1A, 2333 BZ Leiden,
the Netherlands.

■ R. Onstenk, MD, Orthopaedic
Surgeon
Groene Hart Ziekenhuis,
Department of Orthopaedic
Surgery, Bleulandweg 10, 2803
HH Gouda, the Netherlands.

■ A. W. M. M. Koopman-Van
Gemert, MD, PhD,
Anaesthesiologist
Albert Schweitzer Ziekenhuis,
Department of Anaesthesiology,
Albert Schweitzerplaats 25, 3318
AT Dordrecht, the Netherlands.

■ R. G. Pöll, MD, PhD, Professor,
Orthopaedic Surgeon
Slotervaart Ziekenhuis,
Louwesweg 6, 1066 EC
Amsterdam, the Netherlands.

Correspondence should be sent
to Dr J. C. Keurentjes; e-mail:
J.C.Keurentjes@lumc.nl

10.1302/2046-3758.211.2000219
\$2.00

Bone Joint Res 2013;2:238–44.
Received 22 July 2013; Accepted
after revision 18 October 2013

Objectives

Electronic forms of data collection have gained interest in recent years. In orthopaedics, little is known about patient preference regarding pen-and-paper or electronic questionnaires. We aimed to determine whether patients undergoing total hip (THR) or total knee replacement (TKR) prefer pen-and-paper or electronic questionnaires and to identify variables that predict preference for electronic questionnaires.

Methods

We asked patients who participated in a multi-centre cohort study investigating improvement in health-related quality of life (HRQoL) after THR and TKR using pen-and-paper questionnaires, which mode of questionnaire they preferred. Patient age, gender, highest completed level of schooling, body mass index (BMI), comorbidities, indication for joint replacement and pre-operative HRQoL were compared between the groups preferring different modes of questionnaire. We then performed logistic regression analyses to investigate which variables independently predicted preference of electronic questionnaires.

Results

A total of 565 THR patients and 387 TKR patients completed the preference question. Of the THR patients, 81.8% (95% confidence interval (CI) 78.4 to 84.7) preferred pen-and-paper questionnaires to electronic questionnaires, as did 86.8% (95% CI 83.1 to 89.8) of TKR patients. Younger age, male gender, higher completed level of schooling and higher BMI independently predicted preference of electronic questionnaires in THR patients. Younger age and higher completed level of schooling independently predicted preference of electronic questionnaires in TKR patients.

Conclusions

The majority of THR and TKR patients prefer pen-and-paper questionnaires. Patients who preferred electronic questionnaires differed from patients who preferred pen-and-paper questionnaires. Restricting the mode of patient-reported outcome measures to electronic questionnaires might introduce selection bias.

Cite this article: *Bone Joint Res* 2013;2:238–44.

Keywords: Health-related quality of life, Total hip replacement, Total knee replacement, Patient-reported outcome measure, PROM, Questionnaire mode

Article focus

- Based on the recent literature, we hypothesised that patients would prefer electronic questionnaires to pen-and-paper questionnaires after total hip and total knee replacement
- We hypothesised that elderly patients would prefer pen-and-paper questionnaires

Key messages

- The vast majority of patients prefer pen-and-paper questionnaires

- Patients who prefer electronic questionnaires are generally younger and have completed a higher level of schooling

Strengths and limitations

- Strengths: a large sample size and a limited extent of optimism in our statistical models
- Limitations: This study was performed using pen-and-paper questionnaires, which might lead to an overestimation of the actual preference of the pen-and-paper questionnaires

Introduction

Traditionally, the assessment of outcome in orthopaedics has focussed on technical aspects. In total hip (THR) or knee replacement (TKR), the cumulative incidence of revision surgery is often used to compare the outcome of different implants or surgical techniques.¹ The underlying assumption of the traditional orthopaedic approach is that the technical aspects are the most important determinants of clinical success. However, a technically well-performed joint replacement does not guarantee clinical success, as no information is provided on functional status and pain. Additionally, the indication for revision surgery varies widely between orthopaedic surgeons.² Patient-reported outcome measures (PROMs), defined as questionnaires that are completed by patients, provide complementary information as they give an impression of a patient's experience of the surgical procedure and their concerns with regard to health status, health-related quality of life (HRQoL) and the results of the treatment received.³

PROMs can be measured using traditional pen-and-paper questionnaires or various electronic counterparts including touch screens,⁴ personal digital assistants,^{5,6} tablets or mobile phones.⁷ Expected advantages of electronic questionnaires include more complete data capturing, immediate availability of results and lower costs of administrating and entering data.^{6,8}

On the other hand, electronic questionnaires may induce selection bias. A meta-analysis performed in 2008 showed that mail surveys had higher response rates than those based online.⁹ A recent randomised controlled trial, in which 2400 patients were randomised to receive either a pen-and-paper questionnaire or an internet-based questionnaire at four years after THR, revealed an enormous difference in response rate: 92% for the pen-and-paper group versus 49% for the internet-based group.¹⁰ Selection bias can occur if the association between exposure and outcome differs between participants and all eligible patients.⁸

To our knowledge, no study has investigated patient preference for electric questionnaires after THR and TKR. The majority of members of a senior citizens club prefers electronic to pen-and-paper questionnaires.¹¹ Given the similar age of THR/TKR patients, we would expect a preference for electronic questionnaires. We aimed to estimate the proportion of patients who prefer pen-and-paper questionnaires to electronic questionnaires and to estimate predictors of electronic questionnaire preference.

Materials and Methods

The current study is part of a multi-centre cohort study of HRQoL after THR/TKR (NTR2190), performed from August 2010 to August 2011.¹²⁻¹⁵ Institutional review board approval was obtained from all participating centres and all patients gave written informed consent (CCMO-Nr:NL29018.058.09;MEC-Nr:P09.189). The data used in this report constitutes a subset of patients who

underwent primary THR or TKR and who completed pre-operative HRQoL questionnaires along with a question regarding their preference for a mode of questionnaire at a mean of three years (1.5 to 6) after surgery.

We performed this study in order to investigate the preference for a mode of questionnaire for future studies in HRQoL after THR or TKR in a Dutch population. A prerequisite for such future studies is that patients can participate without outpatient department visits, thereby facilitating participation and forestalling the occurrence of selection bias. We selected a web-based questionnaire as the most feasible electronic option. At follow-up, we asked all THR and TKR patients which mode of questionnaire they preferred: pen-and-paper questionnaires or web-based electronic questionnaires, each completed at home.

In order to judge whether patients who preferred pen-and-paper questionnaires differed from patients who preferred electronic questionnaires, we compared age, gender, highest completed level of schooling, body mass index (BMI) categories (< 25 kg/m², 25 to 30 kg/m², 30 to 35 kg/m², > 35 kg/m²), comorbidity, indication for joint replacement (osteoarthritis vs other indications) and pre-operative HRQoL between both groups.

We have aggregated the levels of schooling into an approximation of the social classes, on the assumption that level of schooling indexes the type of qualifications obtained, which in turn indicates the type of occupations available to the subject and hence their own adult social class.¹³ Thus: 'university, higher vocational education and preparatory higher vocational and scientific education' have been aggregated as indicating the professional and managerial social classes; 'middle vocational education and preparatory middle vocational education' have been aggregated as indicating the skilled non-manual and manual social classes; and 'lower vocational education, elementary schooling and no formal education' have been aggregated as indicating the semi- and unskilled manual social classes.

Comorbidity was measured using a patient-reported Charnley classification (A, patients in which the index operated hip or knee are affected only; B, patients in which the other hip or knee is affected as well; and C, patients with a hip or knee replacement and other affected joints and/or a medical condition which affects the patients' ability to ambulate).^{16,17}

HRQoL was measured two weeks before TKR/THR, using the Dutch version of the Short-Form 36 (SF-36).^{18,19} This questionnaire comprises 36 items covering eight domains (physical function, role physical, bodily pain, general health, vitality, social function, role emotional and mental health), for each of which a subscale score is calculated (100 indicating no symptoms and 0 indicating extreme symptoms). Additionally, these scales are incorporated into two summary measures: a physical component summary (PCS) and a mental component summary (MCS). Missing items were imputed whenever possible

Table I. Patient characteristics (THR, total hip replacement; TKR, total knee replacement)

Characteristic	THR (n = 565)	TKR (n = 387)
Mean (SD) age at THR/TKR (yrs)	65.9 (10.6)	68.9 (9.7)
Male (n, %)	198 (35.0)	126 (32.6)
Osteoarthritis (n, %)	486 (86.0)	346 (89.4)
Mean (SD) pre-operative Short-Form 36		
Physical component summary	38.9 (9.61)	40.6 (9.53)
Mental component summary	51.8 (10.8)	51.5 (10.2)
Mean follow-up (yrs) (SD; range)	3.20 (1.13; 1.5 to 6.0)	3.14 (1.12; 1.3 to 6.0)
Body mass index at follow-up (n, %)		
< 25 kg/m ²	194 (34.3)	69 (17.8)
25 to 30 kg/m ²	242 (42.9)	171 (44.1)
30 to 35 kg/m ²	97 (17.1)	92 (23.8)
> 35 kg/m ²	32 (5.7)	55 (14.2)
Charnley comorbidity classification (n, %)		
A	123 (23.3)	54 (14.6)
B	75 (14.2)	39 (10.5)
C	331 (62.6)	278 (74.9)
Highest completed level of education (n, %)		
University, higher vocational education and preparatory higher vocational & scientific education	115 (22.6)	52 (15.4)
Middle vocational education and preparatory middle vocational education	186 (36.6)	120 (35.5)
Lower vocational education, elementary schooling and no formal education	207 (40.7)	166 (49.1)

Table II. Proportion of patients who prefer pen-and-paper questionnaires to electronic questionnaires (THR, total hip replacement; TKR, total knee replacement; CI, confidence interval)

Preferred questionnaire	THR (n = 565)		TKR (n = 387)	
	Patients (n)	Proportion (95% CI)	Patients (n)	Proportion (95% CI)
Pen-and-paper	462	81.8 (78.4 to 84.7)	336	86.8 (83.1 to 89.8)
Electronic	103	18.2	51	13.2

according to Ware et al.¹⁹ We compared pre-operative PCS and MCS between both preference groups.

Statistical analysis. We performed all analyses separately for THR and TKR patients, as clinically important differences vary considerably between these patient groups.²⁰ We performed descriptive analyses of baseline patient characteristics. In order to predict which factors increased the probability of preference for electronic questionnaires, we performed multivariate mixed model logistic regression analyses. We considered the following potential predictors: age, gender, highest completed level of schooling category, BMI category, Charnley classification of comorbidity, indication for joint replacement and pre-operative PCS and MCS scores. In the mixed model regression analyses, patient preference was the dependent variable, all potential predictors were included as fixed effects and centre was included as a random effect. The explained variation was estimated using Nagelkerke's generalised r^2 and the discriminative ability was estimated using the area under the receiver operating characteristic (ROC) curve (AUC).²¹ The extent of optimism in the r^2 and AUC estimates was estimated using bootstrap

resampling (n = 1000 bootstrap samples).²²⁻²⁴ All analyses were performed using R v2.15.2 (R Development Core Team, Vienna, Austria).²⁵

Results

Patient characteristics are shown in Table I. A total of 565 THR patients and 387 TKR patients completed the preference question. Pen-and-paper questionnaires were preferred by 462 THR patients (81.8% (95% confidence interval (CI) 78.4 to 84.7) and by 336 TKR patients (86.8% (95% CI 83.1 to 89.8)) (Table II).

Patient characteristics per preference group are shown in Table III and Table IV for THR and TKR patients, respectively. THR patients who preferred electronic questionnaires tended to be younger, more often male, more often obese, less comorbid, more often highly educated and had worse pre-operative physical health. Age, gender and highest completed level of education remained associated with mode of questionnaire preference while adjusting for age and gender (Table III). TKR patients who preferred electronic questionnaires were younger, more often male, less often morbidly obese,

Table III. Comparison of total hip replacement (THR) patients preferring pen-and-paper *versus* electronic questionnaires (OR, odds ratio; CI, confidence interval)

Characteristic	Pen-and-paper	Electronic	Age- and gender-adjusted OR (95% CI)
Patients (n)	462	103	
Mean (SD) age at THR (yrs)	67.5 (9.5)	58.5 (12.2)	0.93 (0.91 to 0.95)
Male (n, %)	140 (30.3)	58 (56.3)	0.35 (0.22 to 0.56)
Osteoarthritis (n, %)	406 (88.5)	77 (75.5)	0.75 (0.40 to 1.43)
Mean (SD) pre-operative Short-Form 36			
Physical component summary	39.5 (8.9)	35.8 (11.9)	1.00 (0.97 to 1.03)
Mental component summary	51.2 (10.6)	54.8 (10.9)	1.01 (0.99 to 1.04)
Mean (SD) follow-up (yrs)	3.17 (1.13)	3.31 (1.11)	-
Body mass index at follow-up (n, %)			
< 25 kg/m ²	157 (35.4)	29 (29.3)	Reference
25 to 30 kg/m ²	187 (42.1)	46 (46.5)	1.32 (0.75 to 2.32)
30 to 35 kg/m ²	78 (17.6)	15 (15.2)	0.88 (0.42 to 1.83)
> 35 kg/m ²	22 (5.0)	9 (9.1)	2.18 (0.84 to 5.69)
Charnley comorbidity classification (n, %)			
A	100 (23.2)	23 (23.5)	Reference
B	57 (13.2)	18 (18.4)	1.28 (0.59 to 2.79)
C	274 (63.6)	57 (58.2)	1.10 (0.61 to 1.98)
Highest completed level of education (n, %)			
University, higher vocational education and preparatory higher vocational & scientific education	78 (19.1)	37 (37.0)	Reference
Middle vocational education and preparatory middle vocational education	140 (34.3)	46 (46.0)	0.82 (0.47 to 1.45)
Lower vocational education, elementary schooling and no formal education	190 (46.6)	17 (17.0)	0.24 (0.12 to 0.47)

Table IV. Comparison of total knee replacement (TKR) patients preferring pen-and-paper *versus* electronic questionnaires (OR, odds ratio; CI, confidence interval)

Characteristic	Pen-and-paper	Electronic	Age- and gender-adjusted OR (95% CI)
Patients (n)	336	51	
Mean (SD) age at TKR (yrs)	70.3 (8.9)	59.9 (9.7)	0.90 (0.86 to 0.93)
Male (n, %)	103 (30.7)	23 (45.1)	0.61 (0.31 to 1.18)
Osteoarthritis (n, %)	302 (90.4)	40 (81.6)	1.29 (0.48 to 3.45)
Mean (SD) pre-operative Short-Form 36			
Physical component summary	41.1 (8.8)	36.7 (12.9)	1.02 (0.98 to 1.06)
Mental component summary	51.2 (10.4)	53.5 (8.9)	1.02 (0.99 to 1.06)
Mean (SD) follow-up (yrs)	3.10 (1.09)	3.44 (1.32)	-
Body mass index at follow-up (n, %)			
< 25 kg/m ²	55 (17.5)	10 (19.6)	Reference
25 to 30 kg/m ²	136 (43.3)	25 (49.0)	0.86 (0.34 to 2.20)
30 to 35 kg/m ²	75 (23.9)	12 (23.5)	0.65 (0.23 to 1.83)
> 35 kg/m ²	48 (15.3)	4 (7.8)	0.41 (0.11 to 1.57)
Charnley comorbidity classification (n, %)			
A	48 (14.9)	6 (12.2)	Reference
B	36 (11.2)	3 (6.1)	0.83 (0.18 to 3.83)
C	238 (73.9)	40 (81.6)	1.35 (0.52 to 3.52)
Highest completed level of education (n, %)			
University, higher vocational education and preparatory higher vocational & scientific education	34 (11.7)	18 (37.5)	Reference
Middle vocational education and preparatory middle vocational education	96 (33.1)	24 (50.0)	0.55 (0.24 to 1.26)
Lower vocational education, elementary schooling and no formal education	160 (55.2)	6 (12.5)	0.08 (0.03 to 0.25)

less often Charnley class B and more often Charnley class C, more often highly educated and had worse pre-operative physical health. Age and highest completed

level of education remained associated with mode of questionnaire preference while adjusting for age and gender (Table IV).

Table V. Multivariate prediction for preference of electronic questionnaires in total hip and knee replacement patients. Odds ratios > 1 indicate higher odds of preferring an electronic questionnaire, per increasing predictor unit (CI, confidence interval; AUC, area under curve)

Characteristic ^a	Odds ratio (95% CI)	p-value
TOTAL HIP REPLACEMENT		
r² = 0.31; AUC = 0.81		
Age at operation	0.93 (0.90 to 0.96)	< 0.001
Male vs female gender	0.31 (0.17 to 0.56)	< 0.001
Body mass index		
25 to 30 kg/m ² vs < 25 kg/m ²	2.06 (1.03 to 4.11)	0.04
30 to 35 kg/m ² vs < 25 kg/m ²	1.17 (0.48 to 2.81)	0.73
> 35 kg/m ² vs < 25 kg/m ²	5.49 (1.74 to 17.3)	0.004
Other indications vs osteoarthritis	0.59 (0.28 to 1.26)	0.17
Charnley comorbidity classification		
A vs B	0.99 (0.40 to 2.42)	0.98
A vs C	0.87 (0.43 to 1.78)	0.70
Education		
U+HVE+PHVSE vs MVE+PMVE	0.89 (0.45 to 1.77)	0.74
U+HVE+PHVSE vs LVE+ES+NFE	0.27 (0.12 to 0.59)	< 0.001
Short-Form 36 score		
Physical component summary	1.00 (0.97 to 1.03)	0.80
Mental component summary	1.00 (0.98 to 1.03)	0.87
TOTAL KNEE REPLACEMENT		
r² = 0.41; AUC = 0.88		
Age at operation	0.89 (0.84 to 0.94)	< 0.001
Male vs female gender	0.53 (0.21 to 1.34)	0.18
Body mass index		
25 to 30 kg/m ² vs < 25 kg/m ²	1.05 (0.26 to 4.28)	0.94
30 to 35 kg/m ² vs < 25 kg/m ²	1.27 (0.30 to 5.38)	0.75
> 35 kg/m ² vs < 25 kg/m ²	1.59 (0.28 to 8.91)	0.60
Other indications vs osteoarthritis	2.05 (0.53 to 7.89)	0.30
Charnley comorbidity classification		
A vs B	1.40 (0.23 to 8.58)	0.72
A vs C	2.07 (0.58 to 7.31)	0.26
Education		
U+HVE+PHVSE vs MVE+PMVE	0.33 (0.13 to 0.85)	0.02
U+HVE+PHVSE vs LVE+ES+NFE	0.04 (0.01 to 0.15)	< 0.001
Short-Form 36 score		
Physical component summary	1.01 (0.96 to 1.06)	0.63
Mental component summary	0.99 (0.95 to 1.04)	0.75

* U+HVE+PHVSE, university, higher vocational education and preparatory higher vocational and scientific education; MVE+PMVE, middle vocational education and preparatory middle vocational education; LVE+ES+NFE, lower vocational education, elementary schooling and no formal education

Multivariate prediction of electronic questionnaire preference showed that lower age ($p < 0.001$), male gender ($p < 0.001$), higher completed level of schooling ($p < 0.001$) and higher BMI ($p = 0.004$) independently predicted preference of electronic questionnaires in THR patients (Table V). In TKR patients, multivariate prediction of electronic questionnaire preference showed that lower age ($p < 0.001$) and higher completed level of schooling ($p < 0.001$) independently predicted preference of electronic questionnaires (Table V). The prediction model for preference of electronic questionnaires in THR patients had an r^2 of 0.31 with an optimism estimate of 0.04, yielding an optimism-corrected r^2 estimate of 0.27. The AUC was 0.81, with an optimism estimate of -0.02, indicating absence of optimism. The prediction model for preference

of electronic questionnaires in TKR patients had an r^2 of 0.41 with an optimism estimate of -0.24, indicating absence of optimism. The AUC was 0.88, with an optimism estimate of -0.004, indicating absence of optimism.

Discussion

The vast majority of THR and TKR patients prefer pen-and-paper questionnaires. THR patients who prefer electronic questionnaires are younger, more often male, have completed higher levels of schooling and are more often obese. TKR patients who prefer electronic questionnaires are younger and have completed higher levels of schooling.

A limitation of our study is the mode of questionnaire used to capture the data. In this study, we invited patients to participate by conventional mail.

Additionally, all questionnaires consisted of pen- and-paper questionnaires. Patients willing to participate in this study might be more inclined to prefer pen-and-paper questionnaires than THR and TKR patients in general, thus leading to an overestimation of the proportion of patients preferring pen-and-paper questionnaires. However, we consider it unlikely that the entire preference for pen-and-paper questionnaires is based on such selection bias. Additionally, the identified predictors for electronic questionnaire preference, such as age and completed level of schooling, are plausible, thereby indirectly validating our results.

Strengths of our study include the large sample size, allowing precise estimation and multivariate prediction of patient preference. Although the low r^2 values indicate that not all variance is explained by the predictors, the high AUC values indicate that the prediction models have a high discriminatory ability. The limited extent of optimism in r^2 and AUC estimates indicate that overfitting did not play a role in our study.²⁴ In other words, it is unlikely that the prediction models in this study have captured the peculiarities in this data set; conversely, it is likely that predictions, based on this data, will be generalisable to other, similar populations.

Unfortunately, we do not have any information on the availability of internet access of our patients. Although The Netherlands is rated as one of the most mature internet markets,²⁶ recent evidence suggests that non-users of the internet are more likely to be elderly,²⁷ which could explain pen-and-paper questionnaire preference. Practical advantages of electronic questionnaires are stressed in the current orthopaedic literature.^{6,28} Patients are sometimes considered to prefer electronic questionnaires, without any evidence supporting this claim.²⁸ Although electronic questionnaires certainly appear more efficient, our results reveal limitations in line with the findings of Rolfson et al.¹⁰ Future studies, which only measure PROMs using electronic questionnaires, might suffer from limited generalisability, as elderly and less educated patients are less likely to participate. Moreover, selection bias might occur if the association of interest is related to age or social class.

When planning a study in which PROMs will be completed by THR and TKR patients at home, we recommend using pen-and-paper questionnaires, despite their logistic limitations. Such studies should at least provide the option of pen-and-paper questionnaires, in order to prevent selection bias by questionnaire mode.

References

1. Keurentjes JC, Fiocco M, Schreurs BW, et al. Revision surgery is overestimated in hip replacement. *Bone Joint Res* 2012;1:258–262.
2. Nelissen RG, Brand R, Rozing PM. Survivorship analysis in total condylar knee arthroplasty: a statistical review. *J Bone Joint Surg [Am]* 1992;74-A:383–389.
3. Fitzpatrick R, Davey C, Buxton MJ, Jones DR. Evaluating patient-based outcome measures for use in clinical trials. *Health Technol Assess* 1998;2:1–74.
4. Rolfson O, Kärrholm J, Dahlberg LE, Garellick G. Patient-reported outcomes in the Swedish Hip Arthroplasty Register: results of a nationwide prospective observational study. *J Bone Joint Surg [Br]* 2011;93-B:867–875.
5. Kvien TK, Mowinckel P, Heiberg T, et al. Performance of health status measures with a pen based personal digital assistant. *Ann Rheum Dis* 2005;64:1480–1484.
6. Saleh KJ, Radosevich DM, Kassim RA, et al. Comparison of commonly used orthopaedic outcome measures using palm-top computers and paper surveys. *J Orthop Res* 2002;20:1146–1151.
7. Bellamy N, Wilson C, Hendrick J, et al. Osteoarthritis Index delivered by mobile phone (m-WOMAC) is valid, reliable, and responsive. *J Clin Epidemiol* 2011;64:182–190.
8. van Gelder MMHJ, Pijpe A. E-epidemiology: a comprehensive update. *OA Epidemiology* 2013;1:5.
9. Manfreda K, Bosnjak M, Berzelak J, Haas I, Vehovar V. Web surveys versus other survey modes: a meta-analysis comparing response rates. *Int J Market Res* 2008;50:79–104.
10. Rolfson O, Salomonsson R, Dahlberg LE, Garellick G. Internet-based follow-up questionnaire for measuring patient-reported outcome after total hip replacement surgery—reliability and response rate. *Value Health* 2011;14:316–321.
11. Ryan JM, Corry JR, Attewell R, Smithson MJ. A comparison of an electronic version of the SF-36 General Health Questionnaire to the standard paper version. *Qual Life Res* 2002;11:19–26.
12. Keurentjes JC, Fiocco M, So-Osman C, et al. Patients with severe radiographic osteoarthritis have a better prognosis in physical functioning after hip and knee replacement: a cohort-study. *PLoS One* 2013;8:59500.
13. Keurentjes JC, Blane D, Bartley M, et al. Socio-economic position has no effect on improvement in health-related quality of life and patient satisfaction in total hip and knee replacement: a cohort study. *PLoS One* 2013;8:56785.
14. Keurentjes JC, Fiocco M, Nelissen RG. Willingness to undergo surgery again validated clinically important differences in health-related quality of life after total hip replacement or total knee replacement surgery. *J Clin Epidemiol* 2013:Epub.
15. Evangelou E, Kerkhof HJ, Styrkarsdottir U, et al. A meta-analysis of genome-wide association studies identifies novel variants associated with osteoarthritis of the hip. *Ann Rheum Dis* 2013:Epub.
16. Charnley J. The long-term results of low-friction arthroplasty of the hip performed as a primary intervention. *J Bone Joint Surg [Br]* 1972;54-B:61–76.
17. Dunbar MJ, Robertsson O, Ryd L. What's all that noise?: the effect of co-morbidity on health outcome questionnaire results after knee arthroplasty. *Acta Orthop Scand* 2004;75:119–126.
18. Aaronson NK, Muller M, Cohen PD, et al. Translation, validation, and norming of the Dutch language version of the SF-36 Health Survey in community and chronic disease populations. *J Clin Epidemiol* 1998;51:1055–1068.
19. Ware JE, Snow KK, Kosinski M, Gandek B. *SF-36 health survey: manual and interpretation guide*. Boston: New England Medical Center, 1993.
20. Keurentjes JC, Van Tol FR, Fiocco M, Schoones JW, Nelissen RG. Minimal clinically important differences in health-related quality of life after total hip or knee replacement: a systematic review. *Bone Joint Res* 2012;1:71–77.
21. Nagelkerke NJD. A note on a general definition of the coefficient of determination. *Biometrika* 1991;78:691–692.
22. Steyerberg EW, Eijkemans MJ, Harrell FE, Habbema JD. Prognostic modeling with logistic regression analysis: a comparison of selection and estimation methods in small data sets. *Stat Med* 2000;19:1059–1079.
23. Steyerberg EW, Bleeker SE, Moll HA, Grobbee DE, Moons KG. Internal and external validation of predictive models: a simulation study of bias and precision in small samples. *J Clin Epidemiol* 2003;56:441–447.
24. Steyerberg EW. *Clinical prediction models: a practical approach to development, validation and updating*. New York: Springer, 2009.
25. No authors listed. *R: a language and environment for statistical computing*. Vol. 1. Vienna: R Foundation for Statistical Computing, 2008.
26. Deutskens E, de Ruyter K, Wetzels M, Oosterveld P. Response rate and response quality of internet-based surveys: an experimental study. *Marketing Letters* 2004;15:21–36.
27. Brandtzæg PB, Heim J, Karahasanovic A. Understanding the new digital divide: a typology of internet users in Europe. *Int J Human-Computer Studies* 2011;69:123–138.
28. Lee SJ, Kavanaugh A, Lenert L. Electronic and computer-generated patient questionnaires in standard care. *Best Pract Res Clin Rheumatol* 2007;21:637–647.

Funding statement:

- Ongoing research grant of the Dutch Arthritis Association to Professor R. G. H. H Nelissen, Department of Orthopaedic Surgery, Leiden University Medical Center: grant number LLP-013 (<http://www.reumafonds.nl/informatie-voor-doelgroepen/professionals/ongoing-research>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions:

- J. C. Keurentjes: Study conception and design, Data collection, Data analysis, Drafting of the article
- M. Fiocco: Study conception and design, Data analysis, Critical revision of the article
- C. So-Osman: Study conception and design, Data collection, Drafting of the article, Critical revision of the article
- R. Onstenk: Data collection, Drafting of the article, Critical revision of the article

- A. W. M. M. Koopman-Van Gemert: Data collection, Drafting of the article, Critical revision of the article
- R. G. Pöll: Data collection, Drafting of the article, Critical revision of the article
- R. G. H. H. Nelissen: Study conception and design, Data collection, Drafting of the article, Critical revision of the article

ICMJE Conflict of Interest:

- None declared

©2013 The British Editorial Society of Bone & Joint Surgery. This is an open-access article distributed under the terms of the Creative Commons Attributions licence, which permits unrestricted use, distribution, and reproduction in any medium, but not for commercial gain, provided the original author and source are credited.