

The influence of newspaper coverage and a media campaign on smokers' support for smoke-free bars and restaurants and on secondhand smoke harm awareness: findings from the International Tobacco Control (ITC) Netherlands Survey

Gera E Nagelhout,^{1,2} Bas van den Putte,³ Hein de Vries,¹ Matty Crone,⁴ Geoffrey T Fong,^{5,6} Marc C Willemsen^{1,2}

¹Department of Health Promotion, CAPHRI, Maastricht University, Maastricht, The Netherlands

²STIVORO for a smoke free future, The Hague, The Netherlands

³Amsterdam School of Communications Research, ASCoR, University of Amsterdam, Amsterdam, The Netherlands

⁴Department of Public Health and Primary Care, Leiden University Medical Centre, Leiden, The Netherlands

⁵Department of Psychology, University of Waterloo, Waterloo, Canada

⁶Ontario Institute for Cancer Research, Toronto, Canada

Correspondence to

Gera E Nagelhout, STIVORO, PO Box 16070, 2500 BB The Hague, The Netherlands; gnaghelout@stivoro.nl

Received 29 September 2010

Accepted 5 April 2011

Published Online First

17 May 2011

ABSTRACT

Objective To assess the influence of newspaper coverage and a media campaign about Dutch smoke-free legislation on smokers' support for smoke-free bars and restaurants and on secondhand smoke (SHS) harm awareness.

Design and main outcome measures A content analysis was conducted of 1041 newspaper articles on the smoke-free legislation published in six Dutch newspapers from March 2008 to April 2009. Smokers who were regular readers of at least one of these newspapers ($n=677$) were selected from the pre-ban and post-ban waves of the International Tobacco Control Netherlands Survey. Exposure to newspaper coverage and the implementation campaign was correlated with changes in smokers' support for smoke-free bars and restaurants and SHS harm awareness.

Results Most newspaper coverage was found to be negative towards the smoking ban (57%) and focused on economic aspects (59%) rather than health aspects (22%). Exposure to this coverage had a small but significantly negative effect on support for smoke-free bars and restaurants ($\beta=-0.09$, $p=0.013$). Among higher educated smokers, exposure to positive newspaper coverage had a more positive effect on support for smoke-free bars and restaurants. In addition, exposure to the implementation campaign had a small but significantly positive effect on SHS harm awareness ($\beta=0.11$, $p=0.001$).

Conclusions Media attention on smoke-free legislation can influence smokers' support for the legislation and SHS harm awareness. Tobacco control advocates should aim to establish positive media attention that puts forward the health arguments for the legislation.

INTRODUCTION

In many countries around the world, smoke-free legislation is being implemented to protect non-smokers from the health dangers of secondhand smoke (SHS).¹ Optimal protection is achieved when compliance with smoking bans is high. Several studies have shown that compliance is positively related to higher levels of support for the legislation and to greater SHS harm awareness.²⁻⁴ Both can possibly be influenced by means of well-designed media campaigns and advocacy. In this study, we assess the influence of newspaper

coverage and a campaign about smoke-free legislation on changes in smokers' support for the legislation and SHS harm awareness.

Both opponents and proponents of tobacco control believe that media attention can influence public support for tobacco control policies. This is evidenced by the fact that influencing media attention on smoking is a known strategy of tobacco control advocates⁵⁻⁶ and the tobacco industry⁷⁻⁸ alike. However, research on media attention on tobacco has mainly focused on describing media coverage, and few studies have linked this with individual outcomes.⁵ Two recent studies from the USA found an association between media coverage of tobacco control policy in a region and support for the policy in that region.⁹⁻¹⁰ Another recent US study found no association between self-reported media coverage and support for tobacco control policies.¹¹ These studies provided no conclusive evidence on the matter, as their cross-sectional design made it difficult to reach firm conclusions about causal relationships. Furthermore, these studies did not report on differential effects of media coverage between people from higher and lower educational levels. This is important, because smoking is increasingly a problem for lower educated groups.¹ The available research suggests that media campaigns are more effective among higher educated groups¹² and that these groups learn more from the media than lower educated groups.¹³

The theoretical concept of framing explains how media attention can influence attitudes. Frames give context to an issue, which influences how the issue is perceived and understood.¹⁴ For example, smoke-free legislation is often framed by tobacco control advocates as a measure to protect the health of non-smokers but by the tobacco industry as a measure that causes economic losses to the hospitality industry.¹⁵⁻¹⁶ A study on the effects of newspaper coverage about a tobacco tax increase found that coverage of economic aspects was associated with less support for the tax increase, while coverage of health aspects was associated with more support.⁹

The case that we present in this study is that of the implementation of the smoke-free hospitality industry legislation in The Netherlands in July 2008. The implementation of this legislation did

not proceed as planned. A small minority of bars failed to comply with the smoking ban at first,¹⁷ but this small group received a disproportionate high amount of media attention. This may have fuelled resistance to the legislation and increased the number of non-complying bars. While support for legislation typically increases considerably after its implementation,^{18–20} support for the Dutch legislation was low before and stayed low after its implementation.²¹ In fact, of all European countries, The Netherlands had the lowest support for smoke-free bars: only 44% of the population.²² The Dutch government ran a mass media implementation campaign for the smoke-free legislation on television (849 GRPsⁱ), radio (1636 GRPs) and posters (144 GRPs) from May to September 2008. Although the WHO guidelines recommend an emphasis on the public health arguments for implementing smoke-free legislation,²³ the Dutch implementation campaign communicated only the date of implementation. The television commercial showed a man dressed as a large cigarette being thrown out of hospitality industry venues onto the street.

The present study examined the effects of the implementation campaign and newspaper coverage about the smoke-free legislation on support for smoke-free bars and restaurants and on SHS harm awareness. The following hypotheses were tested: (1) positive newspaper coverage has a more positive effect on support and harm awareness than negative coverage; (2) newspaper coverage of health aspects has a more positive effect on support and harm awareness than coverage of economic aspects; (3) the implementation campaign has no effect on support or harm awareness; and (4) there are more positive effects from media attention among smokers with higher rather than lower educational levels. To study this, we used longitudinal data from the International Tobacco Control (ITC) Netherlands Survey and a content analysis of newspaper coverage about the smoke-free legislation.

METHODS

ITC Netherlands Survey

In this study, we used the pre-ban (April 2008) and post-ban (April–May 2009) internet surveys of the ITC Netherlands Survey. All respondents to the pre-ban survey were aged 15 years or older, smoked manufactured or roll-your-own cigarettes at least once-a-month and had smoked at least 100 cigarettes. The pre-ban survey was sent to 2331 smoking panel members of the TNS NIPObase and was returned by 1820 respondents (78%). Of the 1820 pre-ban respondents, 1447 participated again in the post-ban survey (80%). Additional details about the methods of the ITC Netherlands Survey can be found on the ITC Project website.²⁴

The first aspect of media attention about the smoke-free legislation that we assessed in this study is newspaper coverage. All post-ban respondents were presented with a list of 38 newspapers, with a request to indicate which newspapers they read and how often. This method of asking respondents of a population survey about their newspaper reading behaviour to assess the population effects of newspaper coverage was also recently used by Dunlop and Romer.²⁵ Because this method requires a content analysis of every included newspaper, we did not include readers of all 38 newspapers, but only respondents who read at least one of the six largest newspapers (n=677

newspaper readers). A smaller number of the post-ban respondents read one of the 32 other newspapers (n=351), read no newspaper at all (n=415), or did not answer the question (n=4). Table 1 shows the number of respondents per newspaper.

The second aspect of media attention about the smoke-free legislation was the implementation campaign. Recall of exposure to this campaign was assessed at the post-ban survey by showing pictures and texts of the three different parts of the campaign to the post-ban respondents: posters, television and radio commercials. On each part of the campaign, we asked the respondents how often they have seen or heard something about this. Response categories were *never*, *once or twice*, *sometimes*, *often* and *very often*. One exposure variable was created by computing a mean score of exposure to the three different parts of the campaign for every respondent, ranging from 0 to 4 (M=1.12, SD=0.83).

Support for smoke-free bars and restaurants was assessed at the pre-ban and post-ban surveys using the question 'Do you support or oppose a complete Dutch smoking ban in drinking establishments: cafés, bars and pubs?' and 'Do you support or oppose a complete Dutch smoking ban in restaurants?' Response categories were *strongly support*, *support*, *oppose* and *strongly oppose*. The mean of these two variables was used as an indicator of support for smoke-free bars and restaurants.

SHS harm awareness was assessed at the pre-ban and post-ban survey using the question 'In the last month, how often, if at all, did you think about the harm your smoking might be doing to other people?' Response categories were *never*, *rarely*, *sometimes*, *often* and *very often*.

Furthermore, age, gender and educational level of respondents were assessed at the pre-ban survey. Education was categorised into three levels: low (primary education and lower pre-vocational secondary education), moderate (middle pre-vocational secondary education and secondary vocational education) and high (senior general secondary education, (pre-) university education and higher professional education).

Content analysis

A content analysis was performed on the six Dutch newspapers that were most often read by smoking respondents of the ITC Netherlands Survey. Two of the newspapers were the largest regular national newspapers in The Netherlands (*De Telegraaf* and *De Volkskrant*), two were the largest free national newspapers (*Metro* and *Sp!ts*) and two were large regular regional newspapers (*AD Rotterdams Dagblad* and *De Gelderlander*).

Articles that appeared in one of these six newspapers from March 2008 to April 2009 were selected from the LexisNexis digital database, which contains all articles that appeared in these newspapers. A search with words related to smoking (eg, smoking, smoker, cigarette) and words related to the smoke-free hospitality industry legislation (eg, hospitality industry, bar, smoking ban, smoke-free) resulted in a selection of articles with a high probability of being about the smoke-free hospitality industry legislation. Of the selected articles, only the 1086 articles that were about the smoke-free hospitality industry legislation and that had a tobacco focus (ie, the article had smoking or tobacco in the heading, or at least more than half of one paragraph of the article dealt with smoking or tobacco) were included in the study.

All 1086 included articles were coded on reference to health and economic aspects and on slant towards the smoking ban (positive, negative, mixed or neutral). The coding on slant was done from the perspective of tobacco control according to the method of Smith *et al*¹⁰ The coding was done by two coders and

ⁱGRP stands for gross rating point, which is the percentage of the target audience that is reached by the campaign multiplied by the frequency of exposure. Normally, government campaigns in The Netherlands have 300 GRPs on television and 640 GRPs on radio.

Table 1 Number of respondents and articles per newspaper

	<i>Metro</i>	<i>De Telegraaf</i>	<i>Sp!ts</i>	<i>De Volkskrant</i>	<i>AD Rotterdams Dagblad</i>	<i>De Gelderlander</i>	Total
Number of respondents	327	310	302	85	60	54	677
Number of articles							
Smoke-free legislation	145	183	94	142	118	359	1041
Economic aspects	76	114	58	81	60	229	618
Health aspects	33	42	20	51	21	59	226

had a high level of reliability with respect to health aspects (Cohen's $\kappa=0.85$) and economic aspects (Cohen's $\kappa=0.85$) and a satisfactory level of reliability with respect to slant (Cohen's $\kappa=0.62$). A third coder coded the articles for which there was no agreement between the first two coders and made the decision about the coding. If the third coder disagreed with each of the first two coders (only possible for codings on slant), that article was excluded ($n=45$), which left 1041 articles for the analyses.

Analyses

Individual exposure to articles about the smoke-free legislation was estimated by combining the content analysis of newspaper coverage with information about the newspaper reading behaviour of the ITC respondents. This method was also used by Dunlop and Romer,²⁵ with the difference that the current study does not use newspaper reading frequency in general but per newspaper. For example, a respondent who read the newspaper *De Telegraaf* three times a week and the newspaper *Metro* every day could be exposed to half of the 49 articles that were positive about the smoking ban in *De Telegraaf* ($49/2 \approx 25$) and all 52 articles that were positive about the smoking ban in *Metro*. Maximum exposure of this respondent to positive articles on the smoking ban would then be estimated to be 77 ($25+52$) articles. We do not assume that this respondent did read all 77 articles, but we assume that the relative difference in maximum exposure between respondents indicated the relative differences in actual exposure.

Pearson correlation coefficients were reported between exposure to the implementation campaign and newspaper coverage and support for smoke-free bars and restaurants and SHS harm awareness. Linear regression analysis was used to determine the effect of exposure to the implementation campaign and newspaper coverage on support for smoke-free bars and restaurants and on harm awareness. The dependent variables were post-ban support for smoke-free bars and restaurants and post-ban SHS harm awareness. The predictor variables were gender, age, educational level, recall of exposure to the implementation campaign, estimation of exposure to newspaper coverage about the smoke-free legislation and the value of the dependent variable on the pre-ban survey. Estimation of exposure to newspaper coverage consisted of four predictors in the regression analyses: (1) estimated total number of articles a smoker was exposed to; (2) ratio of coverage about health aspects to economic aspects, with higher ratios meaning relatively more coverage about health aspects than economic aspects; (3) estimated percentage of articles that were positive towards the smoking ban and (4) estimated percentage of articles that were mixed or neutral. The percentage of articles that were negative towards the ban had a strong negative correlation with the percentage of positive articles and were therefore not included in the analyses. When, in additional analyses, the percentage of positive articles was replaced by that of negative articles, the regression coefficients of negative newspaper coverage were nearly the same as

those of positive newspaper coverage, with the difference that the coefficients were of different signs. The regression analyses were repeated with the interaction between educational level and recall of exposure to the implementation campaign and the interaction between educational level and estimation of exposure to newspaper coverage.

RESULTS

Characteristics of smoking newspaper readers

Smokers in the ITC Netherlands Survey who were newspaper readers had a mean age of 39.6 years ($SD=15.5$), most of the respondents were men (54%) and had a moderate educational level (low 30%, moderate 46%, high 23%). Readers of the six newspapers differed from each other on age and educational level, but not on gender. For example, readers of *De Telegraaf* were older (mean age 42.4, $SD=15.3$) and had a lower educational level (low 32%, moderate 50%, high 18%).

Newspaper coverage of smoke-free legislation

As can be seen in table 1, economic aspects of the legislation were mentioned in 618 (59%) articles and health aspects in 226 (22%) articles. Both economic and health aspects were mentioned in 688 articles. Articles that were not about economic or health aspects ($n=353$) mostly dealt with resistance to the ban, with rules about smoking rooms in bars and smoking in coffee shops. There were differences in coverage between newspapers. The total number of articles per newspaper ranged from 94 to 359 articles.

Most articles were negative towards the smoking ban (57%), while less than a third were positive (29%) and only a few articles were mixed (5%) or neutral (9%) (table 2). This was also true for articles in which the economic aspects of the legislation were mentioned (62% was negative). However, articles in which health aspects were mentioned were about equally positive about the smoking ban (42%) as negative (43%).

Differences were found in the newspaper coverage over time (figure 1). There was an increase in the number of articles in July 2008, when the smoke-free legislation was implemented. This was followed by a larger increase in the number of articles in the autumn of 2008. These articles were mostly negative about the smoking ban and mainly concerned with economic aspects.


Effects of exposure to media attention

The correlation between exposure to newspaper coverage and the implementation campaign and support for smoke-free bars

Table 2 Percentage of newspaper coverage that was positive, negative, mixed and neutral

	Smoke-free legislation	Economic aspects	Health aspects
Positive (%)	28.8	29.4	42.0
Negative (%)	57.4	61.8	42.9
Mixed (%)	4.5	5.5	8.8
Neutral (%)	9.2	3.2	6.2

Figure 1 Newspaper coverage of the smoke-free legislation from March 2008 to April 2009.



and restaurants and SHS harm awareness were generally low (table 3). There was a negative correlation between exposure to newspaper coverage and support for smoke-free bars and restaurants. There was a positive correlation between the ratio of coverage about health aspects to economic aspects and support for smoke-free bars and restaurants. There was a positive correlation between exposure to mixed or neutral newspaper coverage and support for smoke-free bars and restaurants. In addition, there was a positive correlation between exposure to the implementation campaign and SHS harm awareness.

The regression analyses showed that exposure to newspaper coverage of the smoke-free legislation had a small negative effect on support for smoke-free bars and restaurants (table 4). Exposure to the implementation campaign had a small positive effect on SHS harm awareness. There were no significant effects of theme and slant of newspaper coverage.

The regression analyses from table 4 were repeated with interactions with educational level. The only significant interaction was between educational level and positive newspaper coverage on support for smoke-free bars and restaurants ($\beta=0.10$, $p=0.012$). Among higher educated smokers, exposure to positive newspaper coverage had a more positive effect on support for smoke-free bars and restaurants than among lower educated smokers.

DISCUSSION

Our first hypothesis was that exposure to positive newspaper coverage had a more positive effect on support for smoke-free bars and restaurants and on SHS harm awareness than exposure

Table 3 Associations between media attention (implementation campaign and newspaper coverage) and post-ban support for smoke-free bars and restaurants and post-ban secondhand smoke (SHS) harm awareness (correlation coefficients)

	Support for smoke-free bars and restaurants	SHS harm awareness
Exposure to implementation campaign	0.00	0.12**
Exposure to newspaper coverage about the ban	-0.09*	0.01
Ratio of coverage about health aspects and economic aspects†	0.11**	0.03
Positive newspaper coverage about the ban	-0.07	-0.02
Negative newspaper coverage about the ban	0.02	0.00
Mixed or neutral newspaper coverage about the ban	0.09*	0.04

* $p<0.05$; ** $p<0.01$.

†A higher ratio means relatively more coverage about health aspects than economic aspects.

to negative newspaper coverage. However, no effects were found from positive or negative slant of newspaper coverage. We did find that smokers who were exposed to a larger amount of newspaper coverage about the smoke-free legislation were somewhat less likely to support smoke-free bars and restaurants. This might be explained by the fact that newspaper coverage was twice as often negative than positive about the ban.

Our second hypothesis was that exposure to newspaper coverage of health aspects of the smoke-free legislation had a more positive effect on support and harm awareness than coverage of economic aspects. We did indeed find a small positive correlation with support for smoke-free bars and restaurants, but no significant effect in the multivariate analysis. There was, however, very little newspaper coverage about the health aspects of the smoke-free legislation. Newspaper coverage contained almost three times more references to economic aspects than health aspects of the legislation.

Since the implementation campaign of the smoke-free legislation communicated only the date of implementation, our third hypothesis was that the campaign had no effect on support and harm awareness. We found no effects on support for smoke-free bars and restaurants, but we did find a small positive effect on SHS harm awareness. A possible explanation for this unexpected finding is that the campaign stimulated thinking about the reason for the smoking ban, and smokers concluded that it was implemented to protect people from the harm of SHS. However, the fact that support for smoke-free bars and restaurants was not positively influenced by the campaign is a missed opportunity. A clear campaign in which the government explained the health arguments for the smoke-free legislation could have positively influenced smokers' support for the smoke-free legislation. This may also have prevented many of the problems involving compliance with the ban.

Our last hypothesis was that there were more positive effects from media attention among smokers with a higher rather than lower educational level. We indeed found that newspaper coverage that was positive about the ban had a more positive effect on support for smoke-free bars and restaurants among higher educated smokers than among lower educated smokers. Since more support for smoke-free legislation can increase intentions to quit smoking,²⁶ this could contribute to a widening of health inequalities.

Studies about newspaper coverage of the smoke-free legislation in California¹⁵ and Ireland¹⁶ show that there was a lot of newspaper coverage about the legislation around the implementation date. Our study shows that this is also true for The Netherlands. However, while newspaper coverage of the

Research paper

Table 4 Influence of media attention (implementation campaign and newspaper coverage) on post-ban support for smoke-free bars and restaurants and on post-ban secondhand smoke (SHS) harm awareness (linear regression analyses)

	Support for smoke-free bars and restaurants n = 646 β (95% CI)	SHS harm awareness n = 668 β (95% CI)
Pre-ban value of dependent variable	0.59 (0.52 to 0.65)**	0.43 (0.36 to 0.50)**
Gender (men vs women)	0.04 (−0.02 to 0.11)	−0.12 (−0.19 to −0.05)**
Age	0.02 (−0.04 to 0.09)	0.03 (−0.05 to 0.10)
Low versus high educational level	−0.10 (−0.19 to −0.02)*	0.07 (−0.02 to 0.18)
Moderate versus high educational level	−0.08 (−0.16 to 0.01)	0.10 (0.01 to 0.19)*
Exposure to implementation campaign	0.02 (−0.04 to 0.08)	0.11 (0.04 to 0.18)**
Exposure to newspaper coverage about the ban	−0.09 (−0.14 to −0.02)*	0.04 (−0.03 to 0.11)
Ratio of coverage about health aspects and economic aspects†	−0.02 (−0.10 to 0.07)	0.04 (−0.05 to 0.13)
Positive newspaper coverage about the ban	−0.03 (−0.10 to 0.03)	−0.02 (−0.09 to 0.06)
Mixed or neutral newspaper coverage about the ban	0.03 (−0.05 to 0.11)	0.04 (−0.05 to 0.12)
Adjusted R ²	0.37	0.23

*p<0.05; **p<0.01.

†A higher ratio means relatively more coverage about health aspects than economic aspects.

smoke-free legislation decreased immediately after implementation in California and Ireland, this did not happen in The Netherlands. A few months after the implementation of the Dutch smoke-free legislation, there was a large increase in the number of articles about the legislation. This newspaper coverage was mostly about the resistance of small bars to the legislation, which was organised by a newly formed organisation of small bar owners. Investigative journalism revealed that there were ties between this organisation and the tobacco industry.²⁷ This suggests that the tobacco industry influenced public opinion about the Dutch smoke-free legislation through media attention.

The first limitation of this study is that we estimated the exposure to articles on smoking by combining individual data on the general amount of newspaper reading per newspaper with a content analysis of the articles that appeared in each newspaper. We cannot be sure that all readers of a newspaper with many articles on smoking also read these particular articles. Therefore, although we used a longitudinal design, causal relationships could not be established with absolute certainty. Causal effects of exposure to individual articles can be more reliably established in an experimental design, but this severely limits the time period that can be studied (most often, one experimental session) and reduces the ecological validity of the study. The second limitation is that we looked at coverage of smoking only in newspapers and not in other media. This may not be a problem, since the content of newspapers is highly associated with the content of other media.⁵

A year after the implementation of the smoke-free hospitality industry legislation in The Netherlands, Greece implemented similar legislation. As in The Netherlands, compliance with the ban was less than optimal. It is suggested that the unsupportive and pessimistic newspaper coverage in Greece had an important role in this.²⁸ In contrast to Greece and The Netherlands, the smoke-free legislation in Ireland is known for its immediate success.²⁹ A qualitative study suggests that this success is the result of the efforts of tobacco control advocates who consistently repeated the health arguments for the legislation in the media and who prevented anti-ban advocates from shifting the debate to the economic arguments against the legislation.¹⁶ This suggests that media attention on the health aspects of smoke-free legislation can have an important influence on the success of the legislation.

Acknowledgements We would like to thank Dagmar Feenstra for her help with creating search strings for the selection of articles about smoke-free hospitality industry legislation and with coding the articles. Furthermore, we thank Michel Vlasblom, Marleen Klaassen, Geert de Waal, Nadine Bol, Grietje Bakker and Annegre van den Bos for coding the articles. We also thank Rens Vliegenthart and Jan de Ridder for their helpful advice on selecting articles with search strings.

Funding This study was supported by a grant from the Netherlands Organisation for Health Research and Development (ZonMw), the Amsterdam School of Communication Research of the University of Amsterdam and STIVORO.

Competing interests None.

Contributors GEN conducted the statistical analyses and wrote the manuscript. BvdP advised on the design of the study, the statistical analyses and the writing of the manuscript. All other authors advised on the design of the study and the writing of the manuscript. All authors read and approved the final manuscript.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. **World Health Organization.** *Report on the Global Tobacco Epidemic, 2008: the MPOWER Package.* Geneva: World Health Organization, 2008.
2. **Borland R,** Yong HH, Siahpush M, *et al.* Support for and reported compliance with smoke-free restaurants and bars by smokers in four countries: findings from the International Tobacco Control (ITC) Four Country Survey. *Tob Control* 2006;**15** (Suppl 3):34–41.
3. **Thrasher JF,** Besley JC, González W. Perceived justice and popular support for public health laws: a case study around comprehensive smoke-free legislation in Mexico City. *Soc Sci Med* 2010;**70**:787–93.
4. **Li X,** Li Q, Dong L, *et al.* Risk factors associated with smoking behavior in recreational venues: findings from the International Tobacco Control (ITC) China Survey. *Tob Control* 2010;**19**(Suppl 2):i30–9.
5. **National Cancer Institute.** *The Role of the Media in Promoting and Reducing Tobacco Use. Tobacco Control Monograph No. 19.* Bethesda, MD: U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute, 2008.

What this paper adds

- ▶ In The Netherlands, news coverage of the smoke-free legislation was mostly negative towards the ban and focused on economic aspects rather than health aspects.
- ▶ The news coverage of the smoke-free legislation had a negative effect on support for the legislation.
- ▶ Tobacco control advocates should aim to establish positive media attention during the implementation of smoke-free legislation, focusing on the health arguments for the legislation.

6. **American Cancer Society.** *Tobacco Control Strategy Planning Guide #1. Strategy Planning for Tobacco Control Advocacy.* American Cancer Society & International Union Against Cancer, 2003.
7. **National Cancer Institute.** *Evaluating ASSIST: A Blueprint for Understanding State-level Tobacco Control. Tobacco Control Monograph No. 17.* Bethesda, MD: U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute, 2006.
8. **WHO.** *Tobacco Industry Interference with Tobacco Control.* Geneva: World Health Organization, 2008.
9. **Harris JK,** Shelton SC, Moreland-Russell S, *et al.* Tobacco coverage in print media: the use of timing and themes by tobacco control supporters and opposition before a failed tobacco tax initiative. *Tob Control* 2010;**19**:37–43.
10. **Smith KC,** Siebel C, Pham L, *et al.* News on tobacco and public attitudes towards smokefree air policies in the United States. *Health Policy* 2008;**86**:42–52.
11. **Blake KD,** Viswanath K, Blendon RJ, *et al.* The role of tobacco-specific media exposure, knowledge, and smoking status on selected attitudes toward tobacco control. *Nicotine Tob Res* 2010;**12**:117–26.
12. **Niederdeppe J,** Kuang X, Crock B, *et al.* Media campaigns to promote smoking cessation among socioeconomically disadvantaged populations: what do we know, what do we need to learn, and what should we do now? *Soc Sci Med* 2008;**67**:1343–55.
13. **Tichenor PJ,** Donohue GA, Olien CN. Mass media flow and differential growth in knowledge. *Public Opin Q* 1970;**34**:159–70.
14. **Goffman E.** *Frame Analysis: An Essay on the Organization of Experience.* New York, NY: Harper University Press, 1974.
15. **Magzamen S,** Charlesworth A, Glantz SA. Print media coverage of California's smokefree bar law. *Tob Control* 2001;**10**:154–60.
16. **Fahy D,** Trench B, Clancy L. Communicating Contentious Health Policy: lessons from Ireland's Workplace Smoking Ban. *Health Promot Pract.* Published Online First: 8 October 2009. doi:10.1177/1524839909341554.
17. **VVA.** *Inventarisatie naleefniveau rookvrije horeca winter 2009/2010 [Inventarisation level of compliance smoke-free hospitality industry legislation winter 2009/2010].* The Hague, Groningen: Voedsel en Waren Autoriteit & IntraVal, 2010.
18. **Wakefield M,** Roberts L, Owen N. Trends in prevalence and acceptance of workplace smoking bans among indoor workers in South Australia. *Tob Control* 1996;**5**:205–8.
19. **Fong GT,** Hyland A, Borland R, *et al.* Reductions in tobacco smoke pollution and increases in support for smoke-free public places following the implementation of comprehensive smoke-free workplace legislation in the Republic of Ireland: findings from the ITC Ireland/UK Survey. *Tob Control* 2006;**15**(Suppl 3):51–8.
20. **Hyland A,** Hassan LM, Higbee C, *et al.* The impact of smokefree legislation in Scotland: results from the Scottish ITC Scotland/UK longitudinal surveys. *Eur J Public Health* 2009;**19**:198–205.
21. **ITC Project.** *ITC Netherlands National Report.* Waterloo, The Hague: University of Waterloo, STIVORO, 2010. <http://www.itcproject.org>.
22. **European Commission.** *Survey on Tobacco: Analytical Report. Eurobarometer #253.* Hungary: The Gallup Organisation, 2009.
23. **World Health Organization.** *Guidelines for Controlling and Monitoring the Tobacco Epidemic.* Geneva: WHO, 1998.
24. **ITC Project.** *ITC Netherlands Survey Waves 1-3 (2008–2009) Technical Report.* Waterloo, Ontario, Canada: University of Waterloo, 2009. <http://www.itcproject.org>.
25. **Dunlop SM,** Romer D. Relation between newspaper coverage of 'light' cigarette litigation and beliefs about 'lights' among American adolescents and young adults: the impact on risk perceptions and quitting intentions. *Tob Control* 2010;**19**:267–73.
26. **Brown A,** Moodie C, Hastings G. A longitudinal study of policy effect (smoke-free legislation) on smoking norms: ITC Scotland/United Kingdom. *Nicotine Tob Res* 2009;**11**:924–32.
27. **Baltesen F,** Rosenberg E. *Tabakssector Betaalt Verzet Rookverbod [Big Tobacco Pays Dutch Opposition to Smoking Ban].* Rotterdam: NRC Handelsblad, 2009:1.
28. **Andreou G,** Gourgoulianis K, Galantomos I. The "language" of the unsuccessful anti-smoking campaign in Greece: examples from Greek newspaper headlines. *Prev Med* 2010;**51**:336–7.
29. **Smoke Free Partnership.** *Lifting the Smokescreen. 10 Reasons for a Smoke Free Europe.* Brussels: European Respiratory Society Brussels Office, 2006.



The influence of newspaper coverage and a media campaign on smokers' support for smoke-free bars and restaurants and on secondhand smoke harm awareness: findings from the International Tobacco Control (ITC) Netherlands Survey

Gera E Nagelhout, Bas van den Putte, Hein de Vries, et al.

Tob Control 2012 21: 24-29 originally published online May 17, 2011
doi: 10.1136/tc.2010.040477

Updated information and services can be found at:
<http://tobaccocontrol.bmj.com/content/21/1/24.full.html>

References

These include:

This article cites 15 articles, 10 of which can be accessed free at:
<http://tobaccocontrol.bmj.com/content/21/1/24.full.html#ref-list-1>

Article cited in:

<http://tobaccocontrol.bmj.com/content/21/1/24.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>