



# Cushing's Syndrome

## Hormonal secretion patterns, treatment and outcome



Omslag: Harvey Cushing. Portret afkomstig van 1e dag envelop, bij uitgifte van postzegel met afbeelding van H. Cushing, in de serie "Great Americans" (1988).

Cover, lay-out and printed by Optima Grafische Communicatie, Rotterdam

**Cushing's Syndrome**  
**Hormonal secretion patterns, treatment and outcome**

**Proefschrift**

ter verkrijging van de graad van Doctor aan de  
Universiteit Leiden,  
op gezag van de  
Rector Magnificus Dr. D.D. Breimer,  
hoogleraar in de faculteit der Wiskunde en  
Natuurwetenschappen en die der Geneeskunde,  
volgens besluit van het College voor Promoties

te verdedigen op donderdag  
17 maart 2005 klokke 14.15 uur

door

**Maarten Otto van Aken**

Geboren te Delft in 1968

## **PROMOTIECOMMISSIE**

Promotores: Prof. dr. J.A. Romijn  
Prof. dr. S.W.J. Lamberts, Erasmus Universiteit Rotterdam

Copromotor: dr. W.W. de Herder, Erasmus Universiteit Rotterdam

Referent: Prof. dr. A.R.M.M. Hermus, Universitair Medisch Centrum  
St. Radboud

Overige leden: Prof. dr. E.R. de Kloet  
Prof. dr. S.E. Papapoulos  
Prof. dr. H.A.P. Pols, Erasmus Universiteit Rotterdam  
dr. F. Roelfsema  
Prof. dr. J.M. Wit

Aan Manon  
Aan Pepijn, Jasmine en Friso



## CONTENTS

<b>Chapter 1</b>	9
Introduction	
<b>Chapter 2</b>	31
Irregular and frequent cortisol secretory episodes with preserved diurnal rhythmicity in primary adrenal Cushing's syndrome. <i>Journal of Clinical Endocrinology and Metabolism, online publication dec. 2004</i>	
<b>Chapter 3</b>	49
Growth hormone secretion in primary adrenal Cushing's syndrome is disorderly and inversely correlated with body mass index. <i>American Journal of Physiology: Endocrinology and Metabolism 2005; 288(1); E63-70</i>	
<b>Chapter 4</b>	67
Profound amplification of secretory-burst mass and anomalous regularity of ACTH secretory process in patients with Nelson's syndrome compared with Cushing's disease. <i>Clinical Endocrinology 2004;60;765-772</i>	
<b>Chapter 5</b>	83
Salivary cortisol in the diagnosis of Cushing's syndrome. <i>Published in part in Clinical Chemistry 2003;49(8);1408-1409</i>	
<b>Chapter 6</b>	101
Risk factors for meningitis after transsphenoidal surgery. <i>Clinical Infectious Diseases 1997;25:852-6</i>	
<b>Chapter 7</b>	111
Cerebrospinal fluid leakage during transsphenoidal surgery: postoperative external lumbar drainage reduces the risk for meningitis. <i>Pituitary 2004;7: 93-97</i>	
<b>Chapter 8</b>	121
Postoperative metyrapone test in the early assessment of outcome of pituitary surgery for Cushing's disease. <i>Clinical Endocrinology 1997;47;145-149</i>	

<b>Chapter 9</b>	131
Long-term predictive value of postsurgical cortisol concentrations for cure and risk of recurrence in Cushing's disease. <i>Journal of Clinical Endocrinology and Metabolism 2003: 88;5858-5864</i>	
<b>Chapter 10</b>	147
Decreased quality of life in patients despite long-term biochemical cure of Cushing's disease <i>Submitted</i>	
<b>Chapter 11</b>	163
Summary and conclusions.	
Samenvatting	175
Nawoord	183
Curriculum Vitae	185
Publicaties	186