

A&A 424, L17–L20 (2004)  
 DOI: 10.1051/0004-6361:200400041  
 © ESO 2004

**Astronomy  
&  
Astrophysics**

Letter to the Editor

## Discovery of six Ly $\alpha$ emitters near a radio galaxy at $z \sim 5.2$

B. P. Venemans<sup>1</sup>, H. J. A. Röttgering<sup>1</sup>, R. A. Overzier<sup>1</sup>, G. K. Miley<sup>1</sup>, C. De Breuck<sup>2</sup>, J. D. Kurk<sup>3</sup>, W. van Breugel<sup>4</sup>,  
 C. L. Carilli<sup>5</sup>, H. Ford<sup>6</sup>, T. Heckman<sup>6</sup>, P. McCarthy<sup>7</sup>, and L. Pentericci<sup>8</sup>

<sup>1</sup> Sterrewacht Leiden, PO Box 9513, 2300 RA, Leiden, The Netherlands

e-mail: [venemans@strw.leidenuniv.nl](mailto:venemans@strw.leidenuniv.nl)

<sup>2</sup> European Southern Observatory, Karl Schwarzschild Straße 2, 85748 Garching, Germany

<sup>3</sup> INAF Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, 50125, Firenze, Italy

<sup>4</sup> Lawrence Livermore National Laboratory, PO Box 808, Livermore CA, 94550, USA

<sup>5</sup> National Radio Astronomy Observatory, PO Box 0, Socorro, NM 87801, USA

<sup>6</sup> Dept. of Physics & Astronomy, The Johns Hopkins University, 3400 North Charles Street, Baltimore MD, 21218–2686, USA

<sup>7</sup> The Observatories of the Carnegie Institution of Washington, 813 Santa Barbara Street, Pasadena CA, 91101, USA

<sup>8</sup> Dipartimento di Fisica, Università degli studi Roma Tre, via della Vasca Navale 84, Roma, 00146, Italy

Received 10 May 2004 / Accepted 17 July 2004

**Abstract.** We present the results of narrow-band and broad-band imaging with the Very Large Telescope\* of the field surrounding the radio galaxy TN J0924–2201 at  $z = 5.2$ . Fourteen candidate Ly $\alpha$  emitters with a rest-frame equivalent width of  $>20 \text{ \AA}$  were detected. Spectroscopy of 8 of these objects showed that 6 have redshifts similar to that of the radio galaxy. The density of emitters at the redshift of the radio galaxy is estimated to be a factor 1.5–6.2 higher than in the field, and comparable to the density of Ly $\alpha$  emitters in radio galaxy protoclusters at  $z = 4.1, 3.1$  and  $2.2$ . The Ly $\alpha$  emitters near TN J0924–2201 could therefore be part of a structure that will evolve into a massive cluster. These observations confirm that substantial clustering of Ly $\alpha$  emitters occurs at  $z > 5$  and support the idea that radio galaxies pinpoint high density regions in the early Universe.

**Key words.** galaxies: active – galaxies: clusters: general – galaxies: evolution – cosmology: observations – cosmology: early Universe

### 1. Introduction

One of the most intriguing questions in modern astrophysics concerns the formation of structure in the early Universe (e.g. Bahcall et al. 1997). The narrow-band imaging technique can efficiently select objects with a strong Ly $\alpha$  line in a narrow redshift range, and is therefore ideal for finding and investigating overdense regions at high redshift (Steidel et al. 2000; Möller & Fynbo 2001; Shimasaku et al. 2003; Palunas et al. 2004). For example, Steidel et al. (2000) used narrow-band imaging to map the extent of a large-scale structure at  $z \sim 3.09$ , discovered in a survey for continuum-selected Lyman-break galaxies. Shimasaku et al. (2003) serendipitously found a large-scale structure at  $z \sim 4.9$  while searching for Ly $\alpha$  emitters in the Subaru Deep Field. Their results demonstrate that Mpc-scaled structures have already formed by  $z \sim 4.9$  and that Ly $\alpha$  emitters must be very biased tracers of mass in the early Universe.

Narrow-band imaging of distant powerful radio galaxies at  $z = 2\text{--}4$  has shown that these objects are often located in rich environments, possibly the early stages in the formation of massive clusters (Pascarelle et al. 1996; Le Fèvre et al. 1996;

Pentericci et al. 2000; Venemans et al. 2002, 2003; Kurk et al. 2004). An interesting question is out to which redshift such large-scale structures (protoclusters) can be detected. The most distant known radio galaxy is TN J0924–2201, with a redshift of  $z = 5.2$  (van Breugel et al. 1999). In this letter, we describe broad- and narrow-band observations of this radio galaxy, and report the discovery of 6 Ly $\alpha$  emitters in the field whose redshifts are close to that of the radio galaxy<sup>1</sup>.

### 2. Observations and candidate selection

#### 2.1. Imaging observations and candidate selection

To search for candidate Ly $\alpha$  emitters near TN J0924–2201, narrow-band and broad-band (*I*- and *V*-band) imaging of the field were carried out during two separate observing sessions in 2002 March and April with the VLT Yepun (UT4), using the FOCal Reducer/low dispersion Spectrograph 2 (FOR2). The custom made narrow-band filter had a *FWHM* of  $89 \text{ \AA}$  and a central wavelength of  $7528 \text{ \AA}$ , which encompasses the

<sup>1</sup> Throughout this Letter, magnitudes are in the AB system and a  $\Lambda$ -dominated cosmology with  $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ,  $\Omega_M = 0.3$ , and  $\Omega_\Lambda = 0.7$  is assumed.

\* Based on observations carried out at the European Southern Observatory, Paranal, Chile, programs LP167.A-0409 and 70.A-0589.

wavelength of the Ly $\alpha$  emission line at  $z \sim 5.2$ . The effective exposure times are 36 860 s (narrow-band), 9750 s ( $I$ -band) and 3600 s ( $V$ -band). The seeing in the narrow-band,  $I$ -band and  $V$ -band images is  $0''.8$ ,  $0''.8$  and  $1''.05$  respectively. The  $3\sigma$  limiting magnitudes in an aperture with a  $2''.0$  diameter are 26.29, 26.65 and 26.80 for the narrow-band,  $I$  and  $V$ -band respectively. For a Ly $\alpha$  emitter at  $z \sim 5.2$  with negligible continuum, the  $5\sigma$  limiting line luminosity is  $L_{\text{lim}}(\text{Ly}\alpha) = 3 \times 10^{42} \text{ erg s}^{-1}$ . The images have an area useful for detecting Ly $\alpha$  emitters of  $46.8 \text{ arcmin}^2$ .

A total of 3471 objects were detected in the narrow-band image with a signal-to-noise greater than 5 using the program SExtractor (Bertin & Arnouts 1996). For each object, the observed equivalent width was calculated using a method that will be described in a future paper (Venemans et al., in preparation). Ly $\alpha$  emitters at  $z \sim 5.19$  with a rest-frame equivalent width of  $EW_0 > 20 \text{ \AA}$  would have an observed equivalent width ( $EW_\lambda$ ) of  $124 \text{ \AA}$ . We find 24 such objects in the field. The  $V$ -band image was used to identify low redshift interlopers with an emission line falling in the narrow-band filter. Ten of the 24 objects with  $EW_\lambda > 124 \text{ \AA}$  were also detected in the  $V$ -band with a signal-to-noise greater than 2, and had  $V - I$  colors that were much bluer ( $V - I < 1.2$ ) than a  $V - I$  color of  $\sim 2.75$  as expected for a galaxy at  $z \sim 5.2$  (e.g. Songaila 2004). The remaining 14 candidates were our high priority candidates for follow-up spectroscopy.


## 2.2. Spectroscopy

For the spectroscopy, a mask was constructed which included the radio galaxy and 8 of the 14 high priority candidate Ly $\alpha$  emitters. This was the maximum number that could be fitted on the mask. The rest of the mask was filled with objects having an excess flux in the narrow-band, but with a lower equivalent width than our selection criterion and/or with a blue  $V - I$  color. The observations were carried out on 2003 March 3 and 4 using FORS2 on the VLT Yepun. The mask was observed through the 600RI grism (with a peak efficiency of 87%) with  $1''.4$  slits. The pixels were  $2 \times 2$  binned to decrease the readout time and noise, giving a spatial scale of  $0''.25 \text{ pixel}^{-1}$  and a dispersion of  $1.66 \text{ \AA pixel}^{-1}$ . The total exposure time was 20 676 s. The mean airmass was 1.23 and the seeing in the individual frames varied between  $0''.7$  and  $1''.0$ , giving a spectral resolution of  $185\text{--}265 \text{ km s}^{-1}$  for point sources. For the wavelength calibration, exposures were taken of He, HgCd, Ar and Ne lamps. The rms of the wavelength calibration was always better than  $0.25 \text{ \AA}$  ( $\sim 10 \text{ km s}^{-1}$ ).

## 3. Results

The radio galaxy and all of the 8 observed candidate Ly $\alpha$  emitters showed an emission line near  $7500 \text{ \AA}$ . The redshift of the radio galaxy of  $z = 5.1989 \pm 0.0006$  is consistent with the redshift of  $z = 5.2$ , reported by van Breugel et al. (1999).

Two of the 8 candidate Ly $\alpha$  objects (emitters #463 and #559) are identified with [O III]  $\lambda 5007$  at  $z \sim 0.5$ , confirmed by the accompanying lines [O III]  $\lambda 4959$  and H $\beta$  (Fig. 1 and Table 1). The other six spectra (Fig. 1) did not show any



**Fig. 1.** Part of the spectra of the eight spectroscopically observed high priority emitters and the radio galaxy. For clarity the spectra are offset by  $1.5 \times 10^{-18} \text{ erg s}^{-1} \text{ cm}^{-2}$ . The solid lines indicate the zero-point of the spectra, the dotted lines the  $1\sigma$  uncertainty in the data, and the dashed lines are the scaled transmission curves of the narrow-band filter. The regions in the spectrum where strong telluric skylines dominate are indicated with hashed lines.

other emission line in a wavelength range covering more than  $3300 \text{ \AA}$  (see Table 1).

To distinguish high redshift Ly $\alpha$  emitters from low redshift interlopers various tests can be applied (see Stern et al. 2000, for a review).

**Asymmetric line profile:** A characteristic feature of a high redshift Ly $\alpha$  line is the flux decrement on the blue wing of the Ly $\alpha$  emission (e.g. Dawson et al. 2002). Following Rhoads et al. (2003), the asymmetry of an emission line can be described by the parameters  $a_\lambda$  and  $a_f$ . These parameters measure the ratio of the line width and line flux redward and blueward of the line peak and depend both on the characteristics of the line (line width, amount of absorption, merged doublet) and on the resolution of the spectrum (Rhoads et al. 2003, and reference therein). Simulations of observed spectra indicate that Gaussian Ly $\alpha$  lines with a  $FWHM$  of  $150\text{--}800 \text{ km s}^{-1}$  and with the blue side fully absorbed have  $a_\lambda = 1.0\text{--}1.6$  and  $a_f = 1.0\text{--}1.4$ , while [O II]  $\lambda 3727$  emitters have  $a_\lambda \approx 0.9$  and  $a_f = 0.8\text{--}0.9$ . This is consistent with values found by Rhoads et al. (2003), who measure typical values of  $0.9 < a_f < 1.9$  and  $0.9 < a_\lambda < 3.1$  for a sample of high redshift Ly $\alpha$  emitters and for [O II] emitters at  $z \sim 1$   $a_f \approx 0.8$  and  $a_\lambda \approx 0.9$ . Only two of the emission lines (of emitters #1388 and #2881) have a signal-to-noise that is high enough to measure the asymmetry. These two lines are (marginally) asymmetric (with  $a_\lambda = 2.0 \pm 0.9(2.2 \pm 0.6)$  and  $a_f = 1.7 \pm 0.8(1.4 \pm 0.6)$  for emitter #1388 (2881)), an indication that emitters #1388 and #2881 are Ly $\alpha$  emitters at  $z \sim 5.2$ .

**Continuum break:** A high redshift Ly $\alpha$  emitter must have a continuum break across the Ly $\alpha$  line, caused by the Ly $\alpha$  forest between the galaxy and the observer. Madau (1995) predict a break of a factor  $\sim 5$  across the Ly $\alpha$  line at  $z \sim 5$ . To measure continuum in our spectra, regions that were not effected by strong telluric lines were chosen redward and blueward of

**Table 1.** Properties of the eight spectroscopically observed high priority candidates and the radio galaxy.

Object	Position		$z$	Flux erg s $^{-1}$ cm $^{-2}$	$EW_0$ Å	$FWHM$ km s $^{-1}$	$SFR_{UV}$ $M_{\odot}$ yr $^{-1}$	$SFR_{Ly\alpha}$ $M_{\odot}$ yr $^{-1}$
	$\alpha_{J2000}$	$\delta_{J2000}$						
259	09 24 07.07	-22 02 09.2	$5.1834 \pm 0.0002$	$8.8 \pm 0.8 \times 10^{-18}$	$>103$	$208 \pm 39$	$<3.2$	$3.9 \pm 0.7$
524	09 24 09.41	-22 02 00.5	$5.1683 \pm 0.0003$	$1.2 \pm 0.1 \times 10^{-17}$	$97^{+866}_{-21}$	$392 \pm 25$	$9.1 \pm 3.6$	$10.4 \pm 1.2$
1388	09 24 16.68	-22 01 16.9	$5.1772 \pm 0.0003$	$4.1 \pm 0.5 \times 10^{-18}$	$59^{+476}_{-14}$	$295 \pm 38$	$5.1 \pm 2.0$	$3.5 \pm 0.5$
2688	09 24 25.67	-22 03 01.1	$5.1731 \pm 0.0003$	$3.1 \pm 0.4 \times 10^{-18}$	$>88$	$167 \pm 63$	$<2.9$	$3.0 \pm 0.6$
2849	09 24 24.30	-22 02 30.9	$5.1765 \pm 0.0003$	$8.0 \pm 0.9 \times 10^{-18}$	$47^{+785}_{-13}$	$249 \pm 32$	$6.1 \pm 2.7$	$3.3 \pm 0.7$
2881	09 24 23.88	-22 03 44.8	$5.1683 \pm 0.0005$	$1.4 \pm 0.2 \times 10^{-17}$	$42^{+45}_{-9}$	$479 \pm 28$	$13.7 \pm 3.4$	$6.8 \pm 1.1$
463 <sup>a</sup>	09 24 08.48	-22 00 04.0	$0.4983 \pm 0.0001$	$4.7 \pm 0.7 \times 10^{-18}$	$339^{+1966}_{-91}$	$<200$	–	–
559 <sup>a</sup>	09 24 09.51	-22 00 18.3	$0.51515 \pm 0.00003$	$1.3 \pm 0.1 \times 10^{-17}$	$207^{+160}_{-47}$	$<70$	–	–
RG	09 24 19.90	-22 01 42.0	$5.1989 \pm 0.0006$	$2.1 \pm 0.2 \times 10^{-17}$	$83^{+148}_{-14}$	$1161 \pm 55$	$11.6 \pm 3.5$	$11.4 \pm 0.7$

<sup>a</sup> [O III]  $\lambda 5007$  emitter.

the emission line. Four spectra had a significant ( $>3\sigma$ ) detection of continuum emission redward of the emission line, resulting in  $2\sigma$  lower limits on their flux decrements of 3.6–5.3. Such large breaks in the optical are exclusively found in high redshift objects (e.g. Stern et al. 2000). Hammer et al. (1997) showed that observed [O II] emitters at  $0.5 < z < 1.0$  have a total 4000 Å and Balmer break of factor  $<3$ . Therefore, the continuum break measured in four of the emitters is most likely caused by neutral H I absorption, and hence these emitters can be identified with Ly $\alpha$  emitters at  $z \sim 5.2$ .

*Equivalent width:* The emission line objects have observed equivalent widths in excess of  $\sim 250$  Å. The two emitters which do not show a convincing line asymmetry and do not show continuum both redward and blueward of the emission line, have observed equivalent widths of  $EW_{\lambda} > 540$  Å. This would correspond to a rest-frame equivalent width of  $>269$  Å if the emission line is [O II]  $\lambda 3727$  at  $z \sim 1.0$ . Such high [O II] equivalent width emitters are rare. The total number of  $z \sim 1$  [O II] emitters expected in our field, derived from Teplitz et al. (2003), is  $\sim 1$ . However, the fraction of [O II] emitters with a rest-frame  $EW > 200$  Å is  $<2.5\%$  (Teplitz et al. 2003), which indicates that these two emission line objects are probably not [O II] emitters at  $z \sim 1$ , but Ly $\alpha$  emitters at  $z = 5.2$ .

*Emission line ratios:* As mentioned above, no other emission lines were found in the spectra of the emitters. To estimate the likelihood that the emitters are [O II] emitters, we can derive an upper limit on the flux of the [Ne III]  $\lambda 3869$  line and compare that to local emission line galaxies (Fynbo et al. 2001). Stacking the six spectra to increase the signal-to-noise, we find an upper limit on the ratio of [Ne III] line flux over the [O II] flux of flux ([Ne III])/flux([O II])  $< 0.07$  ( $2\sigma$ ). Using the spectrophotometric catalogue of local emission line galaxies of Terlevich et al. (1991), we found that only 5 out of a sample of 151 of the galaxies with both [O II] and [Ne III] lines detected ( $5/151 \approx 3\%$ ) have such a weak neon line. With the estimated number of [O II] emitters (see above), we expect that  $<1$  of our 6 emission line galaxies is an [O II] emitter.

On the basis of these four lines of arguments, we conclude that these 6 line emitters are almost certainly Ly $\alpha$  emitters at  $z = 5.2$ .

The extracted Ly $\alpha$  lines were fitted with a Gaussian function to estimate the redshift, line flux and widths ( $FWHMs$ ). In Table 1 the properties of the Ly $\alpha$  emitters are summarised. The IDs correspond to the object's number in the catalogue. To correct for the instrumental broadening, the observed  $FWHM$  was deconvolved with the resolution. The star formation rates ( $SFR_{UV}$  and  $SFR_{Ly\alpha}$ ) were calculated from the measured UV continuum fluxes and line fluxes in the images assuming a flat  $f_{\nu}$  spectrum and UV flux density to SFR conversion of Madau et al. (1998).

The velocities of the six confirmed Ly $\alpha$  emitters cluster within a range of 900 km s $^{-1}$  in the rest-frame, while the narrow-band filter is  $\sim 3500$  km s $^{-1}$  wide. The peak of the Ly $\alpha$  emission of the radio galaxy is roughly 1000 km s $^{-1}$  away from the central velocity of the emitters. This is different from other  $z > 2$  radio galaxy protoclusters, where the radio galaxy has a velocity close to the average velocity of the Ly $\alpha$  emitters (Pentericci et al. 2000; Kurk et al. 2004; Venemans et al. 2002). This could be due to H I absorption on the Ly $\alpha$  emission line of the radio galaxy. It has been shown that in radio galaxies this absorption can cause a velocity shift of the Ly $\alpha$  line up to 1000 km s $^{-1}$  as compared to other UV emission lines (e.g., Röttgering et al. 1997).

Of the remaining objects covered by the mask, one is identified as a [O II]  $\lambda 3727$  emitter, also showing [Ne III]  $\lambda 3869$  emission and nine were identified as [O III]  $\lambda 5007$  emitters, confirmed by various lines such as [O III]  $\lambda 4959$ , H $\beta$ , H $\gamma$ , H $\delta$ , [Ne III]  $\lambda 3869$  and [O II]  $\lambda 3727$ . In total 11 [O III] emitters were confirmed in the field, all having a redshift of  $z \sim 0.5$ .

#### 4. Discussion and conclusions

The fraction of foreground contaminants in our sample is estimated to be  $2/8 \sim 25\%$ . There are 6 additional unconfirmed high priority candidate Ly $\alpha$  emitters in the field. Based on the fraction of contaminants in our sample,  $\sim 4$  of those are expected to be  $z \sim 5.2$  Ly $\alpha$  emitters.

Is there an overdensity of Ly $\alpha$  emitters near TN J0924–2201? To investigate this question, we have to compare the density of Ly $\alpha$  emitters in our field with the density in blank fields. The largest survey near  $z \sim 5$

for Ly $\alpha$  emitters is the search for Ly $\alpha$  emitters at  $z \sim 4.79$  in the Subaru Deep Field (SDF, Shimasaku et al. 2004). This survey is comparable in depth to our observations ( $L_{\text{lim}}(\text{Ly}\alpha) = 3 \times 10^{42} \text{ erg s}^{-1}$  for an emitter at  $z = 4.79$  with no continuum) and the selection criteria applied to identify Ly $\alpha$  emitters are very similar to ours ( $EW_{\lambda} > 80 \text{ \AA}$ , Shimasaku et al. 2004). In the SDF, Shimasaku et al. find 51 candidate Ly $\alpha$  emitters in an area of  $25' \times 45'$ . However, there is no spectroscopic confirmation of these candidates. We therefore conservatively assume that all their candidates are Ly $\alpha$  emitters at  $z \sim 4.8$ , resulting in a number density of Ly $\alpha$  emitters in the SDF of  $2.1 \pm 0.3 \times 10^{-4} \text{ Mpc}^{-3}$  (Shimasaku et al. 2004). Excluding the radio galaxy, the density of confirmed Ly $\alpha$  emitters in our field is  $5.3^{+3.2}_{-2.1} \times 10^{-4} \text{ Mpc}^{-3}$ , which is a factor  $2.5^{+1.6}_{-1.0}$  higher than in the SDF. If the four unconfirmed candidate Ly $\alpha$  emitters are included, this factor rises to  $4.2^{+2.0}_{-1.4}$ . We used the data from the SDF to estimate the chance of finding 6 or more Ly $\alpha$  emitters in within a single  $6'8 \times 6'8$  FORS2 field by counting the number of emitters in randomly placed  $6'8 \times 6'8$  apertures. In only 7% of the cases, more than 6 Ly $\alpha$  emitters were found. This further indicates that the TN J0924–2201 field is overdense in Ly $\alpha$  emitters.

Ly $\alpha$  emitters at high redshift show large cosmic variance in their clustering properties (e.g. Shimasaku et al. 2004). Various authors have found that the distribution of Ly $\alpha$  emitters on the sky and/or in redshift space can be very inhomogeneous (e.g. Ouchi et al. 2003; Fynbo et al. 2003; Shimasaku et al. 2003; Palunas et al. 2004; Hu et al. 2004). For example, most of the Ly $\alpha$  emitters found at  $z = 4.86$  in the SDF are concentrated within a large-scale structure with a radius of  $\sim 6'$  ( $\sim 2.5 \text{ Mpc}$ , Shimasaku et al. 2003). It is therefore possible that the Ly $\alpha$  emitters around TN J0924–2201 in the  $\sim 6'8 \times 6'8$  field of view of FORS2 are located inside such a large-scale structure.

It is interesting to compare the (over)density in this field with the protoclusters that were found around radio galaxies at  $z = 4.1, 3.1$  and  $2.2$ , each with at least 20 confirmed protocluster members and estimated masses of  $\sim 10^{14} - 10^{15} M_{\odot}$  (Pentericci et al. 2000; Kurk et al. 2004; Venemans et al. 2002, Venemans et al. in prep.). In the TN 0924–2201 field objects were selected with a (Ly $\alpha$ ) line luminosity of  $> 3 \times 10^{42} \text{ erg s}^{-1}$ . At  $z = 4.1, 3.1$  and  $2.2$ , this luminosity limit corresponds to a limit of  $> 1.5, 3.1$  and  $7.0 \times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$ . The number of candidate (confirmed) emitters with a line brighter than the luminosity limit in the  $z = 4.1, 3.1$  and  $2.2$  protoclusters is 10 (10), 12 (12) and 8 (6) respectively. This is roughly the same number of Ly $\alpha$  emitters as in the TN J0924–2201 field, which contains six confirmed and four possible Ly $\alpha$  emitters. The Ly $\alpha$  emitters at  $z = 5.2$  might therefore be the bright end of a population of star forming galaxies in a protocluster at  $z = 5.2$ , making it the most distant known protocluster. Deep multi-color observations should confirm this by detecting other populations of galaxies (e.g. Lyman break galaxies) in the protocluster.

**Acknowledgements.** We thank the staff on Paranal, Chile for their splendid support, and William Grenier of Andover Corporation for his help in our purchase of the customised narrow-band filter. We also thank the referee, J. Fynbo, for his comments that improved this manuscript. GKM acknowledges funding by an Academy Professorship of the Royal Netherlands Academy of Arts and Sciences (KNAW). The work by WvB was performed under the auspices of the US Department of Energy, National Nuclear Security Administration by the University of California, Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48. The NRAO is operated by associated universities Inc., under cooperative agreement with the NSF. This work was supported by the European Community Research and Training Network “The Physics of the Intergalactic Medium”.

## References

- Bahcall, N. A., Fan, X., & Cen, R. 1997, ApJ, 485, L53  
 Bertin, E., & Arnouts, S. 1996, A&AS, 117, 393  
 Dawson, S., Spinrad, H., Stern, D., et al. 2002, ApJ, 570, 92  
 Fynbo, J. P. U., Ledoux, C., Möller, P., Thomsen, B., & Burud, I. 2003, A&A, 407, 147  
 Fynbo, J. U., Möller, P., & Thomsen, B. 2001, A&A, 374, 443  
 Hammer, F., Flores, H., Lilly, S. J., et al. 1997, ApJ, 481, 49  
 Hu, E. M., Cowie, L. L., Capak, P., et al. 2004, AJ, 127, 563  
 Kurk, J. D., Pentericci, L., Röttgering, H. J. A., & Miley, G. K. 2004, A&A, accepted  
 Le Fèvre, O., Deltorn, J. M., Crampton, D., & Dickinson, M. 1996, ApJ, 471, L11  
 Madau, P. 1995, ApJ, 441, 18  
 Madau, P., Pozzetti, L., & Dickinson, M. 1998, ApJ, 498, 106  
 Möller, P., & Fynbo, J. U. 2001, A&A, 372, L57  
 Ouchi, M., Shimasaku, K., Furusawa, H., et al. 2003, ApJ, 582, 60  
 Palunas, P., Teplitz, H. I., Francis, P. J., Williger, G. M., & Woodgate, B. E. 2004, ApJ, 602, 545  
 Pascarelle, S. M., Windhorst, R. A., Driver, S. P., Ostrander, E. J., & Keel, W. C. 1996, ApJ, 456, L21  
 Pentericci, L., Kurk, J. D., Röttgering, H. J. A., et al. 2000, A&A, 361, L25  
 Rhoads, J. E., Dey, A., Malhotra, S., et al. 2003, AJ, 125, 1006  
 Röttgering, H. J. A., van Ojik, R., Miley, G. K., et al. 1997, A&A, 326, 505  
 Shimasaku, K., Ouchi, M., Okamura, S., et al. 2003, ApJ, 586, L111  
 Shimasaku, K., Hayashino, T., Matsuda, Y., et al. 2004, ApJ, 605, L93  
 Songaila, A. 2004, AJ, 127, 2598  
 Steidel, C. C., Adelberger, K. L., Shapley, A. E., et al. 2000, ApJ, 532, 170  
 Stern, D., Bunker, A., Spinrad, H., & Dey, A. 2000, ApJ, 537, 73  
 Teplitz, H. I., Collins, N. R., Gardner, J. P., et al. 2003, ApJS, 146, 209  
 Terlevich, R., Melnick, J., Masegosa, J., Moles, M., & Copetti, M. V. F. 1991, A&AS, 91, 285  
 van Breugel, W., De Breuck, C., Stanford, S. A., et al. 1999, ApJ, 518, L61  
 Venemans, B. P., Kurk, J. D., Miley, G. K., et al. 2002, ApJ, 569, L11  
 Venemans, B. P., Kurk, J. D., Miley, G. K., & Röttgering, H. J. A. 2003, New Astron. Rev., 47, 353