

The Annals of Probability
2003, Vol. 31, No. 3, 1205–1227
© Institute of Mathematical Statistics, 2003

DETERMINATE MULTIDIMENSIONAL MEASURES, THE EXTENDED CARLEMAN THEOREM AND QUASI-ANALYTIC WEIGHTS¹

BY MARCEL DE JEU

Leiden University

We prove in a direct fashion that a multidimensional probability measure μ is determinate if the higher-dimensional analogue of Carleman's condition is satisfied. In that case, the polynomials, as well as certain proper subspaces of the trigonometric functions, are dense in all associated L_p -spaces for $1 \leq p < \infty$. In particular these three statements hold if the reciprocal of a quasi-analytic weight has finite integral under μ . We give practical examples of such weights, based on their classification.

As in the one-dimensional case, the results on determinacy of measures supported on \mathbb{R}^n lead to sufficient conditions for determinacy of measures supported in a positive convex cone, that is, the higher-dimensional analogue of determinacy in the sense of Stieltjes.

1. Introduction and overview. We will be concerned with determinacy and density results for probability measures on \mathbb{R}^n for a fixed n . Establishing notation, let (\cdot, \cdot) be the standard inner product on \mathbb{R}^n with corresponding norm $\|\cdot\|$. For $\lambda \in \mathbb{R}^n$ define $e_{i\lambda} : \mathbb{R}^n \mapsto \mathbb{C}$ by $e_{i\lambda}(x) = \exp i(\lambda, x)$ ($x \in \mathbb{R}^n$). We let \mathcal{M}^* be the set of all positive Borel measures μ on \mathbb{R}^n such that

$$\int_{\mathbb{R}^n} \|x\|^d d\mu(x) < \infty$$

for all $d \geq 0$. A measure μ is said to be determinate if $\mu \in \mathcal{M}^*$ and if μ is uniquely determined in \mathcal{M}^* by the set of integrals

$$\int_{\mathbb{R}^n} P(x) d\mu(x)$$

of all polynomials P on \mathbb{R}^n .

There are several results in the literature concerning the determinacy of elements of \mathcal{M}^* and the related matter of the density of the polynomials in the associated L_p -spaces. Connections are furthermore known between these

Received October 2001; revised May 2002.

¹Supported in part by a PIONIER grant of the Netherlands Organisation for Scientific Research (NWO).

AMS 2000 subject classifications. Primary 44A60; secondary 41A63, 41A10, 42A10, 46E30, 26E10.

Key words and phrases. Determinate multidimensional measures, Carleman criterion, L_p -spaces, multidimensional approximation, polynomials, trigonometric functions, multidimensional quasi-analytic classes, quasi-analytic weights.

properties for a multidimensional measure and the corresponding properties for its one-dimensional marginal distributions. We refer to [1, 2] for an overview of the field.

These results in the literature yield sufficient conditions for a measure to be determinate. The resulting criteria are however not always easy to apply, since they tend to ultimately involve the computation of moment sequences. Given a particular measure such a computation need not be an attractive task.

In this paper, on the contrary, we establish an *integral* criterion of some generality to conclude that a measure is determinate. A criterion of this type is evidently easier to apply. Moreover, if this criterion is satisfied, then the polynomials are dense in the associated L_p -spaces for finite p , and the same holds for $\text{Span}_{\mathbb{C}}\{e_{i\lambda} \mid \lambda \in S\}$ for any subset S of \mathbb{R}^n which is somewhere dense, that is, is such that its closure \bar{S} has nonempty interior. Our criterion is established along the following lines.

We first prove that a multidimensional probability measure is determinate and that the density results hold as described above if the higher dimensional analogue of Carleman's condition is satisfied. This should, analogously to [4], be compared with the classical one-dimensional Carleman theorem, which asserts determinacy but is not concerned with density. We will in arbitrary dimension refer to the total conclusion of the determinacy and the density as described above as the *extended* Carleman theorem.

Our proof of the extended Carleman theorem is based on a result on multidimensional quasi-analytic classes. It is a "direct" proof and close in spirit to the classical proof of the one-dimensional Carleman theorem as in, for example, [11]. We will also indicate an alternative derivation based on the recent literature. This alternative derivation, however, is considerably less direct than our approach.

Having established the extended Carleman theorem, we subsequently note that a measure satisfies the necessary hypotheses if the reciprocal of a so called quasi-analytic weight has finite integral. Such (multidimensional) weights are defined and studied systematically in [6]. The sufficiency of the aforementioned integral condition on the measure is then in fact almost trivial, given the definition of these weights in terms of divergent series as in Section 3. Verifying this divergence for a particular weight is, however, in general not an easy computation, but—and this is the crucial point—quasi-analytic weights can alternatively be characterized by the divergence of certain integrals, which is on the contrary usually a rather straightforward condition to verify. Using these two equivalent characterizations, we are thus finally led to results in the vein of the following theorem.

THEOREM 1.1. *Suppose $R > 0$ and a nondecreasing function $\rho : (R, \infty) \mapsto \mathbb{R}_{\geq 0}$ of class C^1 are such that*

$$\int_R^\infty \frac{\rho(s)}{s^2} ds = \infty.$$

If μ is a positive Borel measure on \mathbb{R}^n such that

$$\int_{\|x\|>R} \exp\left(\int_R^{\|x\|} \frac{\rho(s)}{s} ds\right) d\mu(x) < \infty,$$

then μ is determinate. Furthermore, the polynomials and $\text{Span}_{\mathbb{C}}\{e_{i\lambda} \mid \lambda \in S\}$ are then dense in $L_p(\mathbb{R}^n, \mu)$ for all $1 \leq p < \infty$ and for every subset S of \mathbb{R}^n which is somewhere dense.

A particular case is obtained by choosing $\rho(s) = \varepsilon s$ for some $\varepsilon > 0$. Then one sees that

$$\int_{\mathbb{R}^n} \exp(\varepsilon\|x\|) d\mu(x) < \infty$$

is a sufficient condition; this is a classical type of result. However, a measure $\mu \in \mathcal{M}^*$ is now also seen to be determinate, and the polynomials and spaces $\text{Span}_{\mathbb{C}}\{e_{i\lambda} \mid \lambda \in S\}$ for somewhere dense S are dense in the associated L_p -spaces for finite p if, for example,

$$(1) \quad \int_{a_2\|x\|>2} \exp\left(\frac{a_1\|x\|}{\log a_2\|x\|}\right) d\mu(x) < \infty$$

for some $a_1, a_2 > 0$. This is a substantially weaker condition. In Section 4 we will give some additional and even more lenient sufficient conditions, formulated in terms of elementary functions as above. It will also become apparent that the integrand need not be radial as in Theorem 1.1. Although such radial integrands may be sufficient for most applications, this is not the most general situation in which our results apply. We return to the possible consequences of this observation in Section 6.

In the discussion so far we considered what can be called determinacy in the sense of Hamburger in arbitrary dimension, that is, the question whether a measure on \mathbb{R}^n is determined by its integrals of the polynomials, without any restriction on its support. Naturally, in the one-dimensional case the question of determinacy has also been studied under the condition that the support of the measure is contained in the interval $[0, \infty)$. This determinacy in the sense of Stieltjes has an analogue in arbitrary dimension, by asking whether a measure on \mathbb{R}^n is determined by its integrals of the polynomials, under the assumption that its support is contained in a given positive convex cone with the origin as vertex. The simultaneous distribution of nonnegative random variables provides an obvious practical example. To facilitate the formulation, we adapt the following terminology.

DEFINITION 1.2. Let $\{v_1, \dots, v_n\}$ be a basis of \mathbb{R}^n and let $C = \mathbb{R}_{\geq 0} \cdot v_1 + \dots + \mathbb{R}_{\geq 0} \cdot v_n$ be the corresponding positive convex cone. Let $\mu \in \mathcal{M}^*$ be supported

in C . Then μ is C -determinate if a measure $\nu \in \mathcal{M}^*$, which is also supported in C and is such that

$$\int_C P(x) d\nu(x) = \int_C P(x) d\mu(x)$$

for all polynomials P on \mathbb{R}^n , is necessarily equal to μ .

As in the one-dimensional case, sufficient conditions for determinacy in the sense of Hamburger imply sufficient conditions for C -determinacy. The density results do not transfer in general. Concentrating on radial integrands again, we thus obtain the following result.

THEOREM 1.3. *Let $\{v_1, \dots, v_n\}$ be a basis of \mathbb{R}^n and let $C = \mathbb{R}_{\geq 0} \cdot v_1 + \dots + \mathbb{R}_{\geq 0} \cdot v_n$ be the corresponding positive convex cone. Let μ be a positive Borel measure on \mathbb{R}^n which is supported in C .*

Suppose $R > 0$ and a nondecreasing function $\rho : (R, \infty) \mapsto \mathbb{R}_{\geq 0}$ of class C^1 are such that

$$\int_R^\infty \frac{\rho(s)}{s^2} ds = \infty$$

and

$$\int_{\sqrt{\|x\|} > R} \exp\left(\int_R^{\sqrt{\|x\|}} \frac{\rho(s)}{s} ds\right) d\mu(x) < \infty.$$

Then μ is C -determinate.

Aside, we mention that under an additional condition one can conclude that μ is actually determinate, as will be discussed in Section 5.

As a consequence of the theorem, if $\mu \in \mathcal{M}^*$ is supported in C , and if, for example,

$$\int_C \exp(\varepsilon\sqrt{\|x\|}) d\mu(x) < \infty$$

for some $\varepsilon > 0$, or if

$$(2) \quad \int_{a_2\sqrt{\|x\|} > 2} \exp\left(\frac{a_1\sqrt{\|x\|}}{\log a_2\|x\|}\right) d\mu(x) < \infty$$

for some $a_1, a_2 > 0$, then μ is C -determinate.

To conclude this introductory discussion, we first of all mention that for quite a few (one-dimensional) common distributions occurring in practice the determinacy or nondeterminacy is known; see, for example, [14] for a number of examples. It seems that many of the known positive results on determinacy in one dimension follow from condition (1) in the Hamburger case or (2) in the

Stieltjes case, the latter possibly combined with the result as discussed at the end of Section 5.

Secondly, let us note that the typical practical sufficient condition from which we conclude determinacy in this paper is the integrability of a function of a suitable type. The “underlying” reason for this determinacy is a Carleman-type criterion, which is satisfied as a consequence of this integrability. It is an interesting problem to determine a set of functions with the property that a measure satisfies such a Carleman-type criterion *precisely* if a function in this set is integrable. These matters are addressed in [9] for the one-dimensional case.

This paper is organized as follows.

In Section 2 we establish the extended Carleman theorem.

Section 3 is a preparation for Sections 4 and 5. It contains the definition of quasi-analytic weights, their classification and main properties, referring to [6] for proofs.

In Section 4 the results of Sections 2 and 3 are put together, resulting in integral criteria for determinacy (without restrictions on the support) and the density results.

Section 5 is concerned with determinacy in the sense of Stieltjes, that is, with C -determinacy. Integral criteria are obtained and a condition is discussed under which one can conclude determinacy, rather than just C -determinacy.

Section 6 contains a tentative remark on the possibility of the existence of distinguished marginal distributions.

2. The extended Carleman theorem. In this section we establish the extended Carleman theorem. The determinacy of the measure, the density of the polynomials and the density of the spaces $\text{Span}_{\mathbb{C}}\{e_{i;\lambda} \mid \lambda \in S\}$ are all seen to be closely related, since they all ultimately rest on the following theorem on multidimensional quasi-analytic classes.

THEOREM 2.1. *For $j = 1, \dots, n$ let $\{M_j(m)\}_{m=0}^\infty$ be a sequence of nonnegative real numbers such that*

$$\sum_{m=1}^\infty \frac{1}{M_j(m)^{1/m}} = \infty.$$

Assume that $f : \mathbb{R}^n \mapsto \mathbb{C}$ is of class C^∞ and that there exists $C \geq 0$ such that

$$\left| \frac{\partial^\alpha f}{\partial \lambda^\alpha}(\lambda) \right| \leq C \prod_{j=1}^n M_j(\alpha_j)$$

for all $\alpha \in \mathbb{N}^n$ and all $\lambda \in \mathbb{R}^n$. Then, if $\frac{\partial^\alpha f}{\partial \lambda^\alpha}(0) = 0$ for all $\alpha \in \mathbb{N}^n$, f is actually identically zero on \mathbb{R}^n .

A proof by induction, starting from the Denjoy–Carleman theorem in one dimension, can be found in [6]. The result in [6] is in fact somewhat stronger than the statement above. A slightly weaker version, on the other hand, which is however not entirely sufficient in our situation, can already be found in [10]. The proof in [10] is more complicated than the proof in [6], but in [10] the necessity of the hypotheses is investigated as well.

For reference purposes we state the following elementary fact, the verification of which is omitted.

LEMMA 2.2. *Let $\{a(m)\}_{m=1}^\infty$ be a nonnegative nonincreasing sequence of real numbers. If k and l are strictly positive integers, then $\sum_{m=1}^\infty a(km) = \infty$ if and only if $\sum_{m=1}^\infty a(lm) = \infty$.*

THEOREM 2.3 (Extended Carleman theorem). *Let $\mu \in \mathcal{M}^*$ and suppose $\{v_1, \dots, v_n\}$ is a basis of \mathbb{R}^n . For $j = 1, \dots, n$ and $m = 0, 1, 2, \dots$ define*

$$s_j(m) = \int_{\mathbb{R}^n} (v_j, x)^m d\mu(x).$$

If each of the sequences $\{s_j(m)\}_{m=1}^\infty$ ($j = 1, \dots, n$) satisfies Carleman’s condition

$$(3) \quad \sum_{m=1}^\infty \frac{1}{s_j(2m)^{1/2m}} = \infty,$$

then μ is determinate. Furthermore, the polynomials and $\text{Span}_{\mathbb{C}}\{e_{i\lambda} \mid \lambda \in S\}$ are then dense in $L_p(\mathbb{R}^n, \mu)$ for all $1 \leq p < \infty$ and for every subset S of \mathbb{R}^n which is somewhere dense.

PROOF. Using the obvious fact that a linear automorphism of \mathbb{R}^n induces an automorphism of the polynomials and a permutation of the spaces of trigonometric functions $\text{Span}_{\mathbb{C}}\{e_{i\lambda} \mid \lambda \in S\}$ for somewhere dense S , one sees easily that we may assume that $\{v_1, \dots, v_n\}$ is the standard basis of \mathbb{R}^n . One also verifies that we may assume in addition that $\mu(\mathbb{R}^n) = 1$. Under these two assumptions we turn to the proof.

Treating the determinacy of μ first, we write $\mu_1 = \mu$ and we suppose that $\mu_2 \in \mathcal{M}^*$ is a probability measure on \mathbb{R}^n with the same integrals of the polynomials as μ_1 . Let $\nu = \frac{1}{2}(\mu_1 + \mu_2)$ and introduce, in the usual multi-index notation:

$$t(\alpha) = \int_{\mathbb{R}^n} |x|^\alpha d\nu, \quad \alpha \in \mathbb{N}^n.$$

Let

$$(4) \quad t_j(s) = \int_{\mathbb{R}^n} |x_j|^s d\nu, \quad j = 1, \dots, n, s \geq 0.$$

As is well known, the Hölder inequality and the fact that $\nu(\mathbb{R}^n) = 1$ imply that

$$(5) \quad t_j(s_1)^{1/s_1} \leq t_j(s_2)^{1/s_2}$$

for $j = 1, \dots, n$ and $1 \leq s_1 \leq s_2 < \infty$. In addition, regarding $|x|^\alpha = \prod_{j=1}^n |x_j|^{\alpha_j}$ as a product of n elements of $L_n(\mathbb{R}^n, \nu)$, the generalized Hölder inequality [7], Section VI.11.1, yields

$$(6) \quad t(\alpha) \leq \prod_{j=1}^n t_j(\alpha_j n)^{1/n}, \quad j = 1, \dots, n, \alpha \in \mathbb{N}^n.$$

Consider the Fourier transforms

$$\widehat{\mu}_k(\lambda) = \int_{\mathbb{R}^n} e^{i(\lambda, x)} d\mu_k(x), \quad k = 1, 2, \lambda \in \mathbb{R}^n.$$

Then $\widehat{\mu}_1$ and $\widehat{\mu}_2$ are of class C^∞ on \mathbb{R}^n with derivatives

$$(7) \quad \frac{\partial^\alpha \widehat{\mu}_k}{\partial \lambda^\alpha}(\lambda) = \int_{\mathbb{R}^n} i^{|\alpha|} x^\alpha e^{i(\lambda, x)} d\mu_k(x), \quad k = 1, 2, \alpha \in \mathbb{N}^n, \lambda \in \mathbb{R}^n.$$

By assumption we therefore have

$$(8) \quad \frac{\partial^\alpha \widehat{\mu}_1}{\partial \lambda^\alpha}(0) = \frac{\partial^\alpha \widehat{\mu}_2}{\partial \lambda^\alpha}(0), \quad \alpha \in \mathbb{N}^n.$$

From (7) we see that

$$\frac{1}{2} \left| \frac{\partial^\alpha (\widehat{\mu}_1 - \widehat{\mu}_2)}{\partial \lambda^\alpha}(\lambda) \right| \leq t(\alpha), \quad \alpha \in \mathbb{N}^n, \lambda \in \mathbb{R}^n,$$

and then combination with (6) yields

$$\frac{1}{2} \left| \frac{\partial^\alpha (\widehat{\mu}_1 - \widehat{\mu}_2)}{\partial \lambda^\alpha}(\lambda) \right| \leq \prod_{j=1}^n t_j(\alpha_j n)^{1/n}, \quad \alpha \in \mathbb{N}^n, \lambda \in \mathbb{R}^n.$$

We claim that the nonnegative sequences $\{t_j(mn)^{1/mn}\}_{m=0}^\infty$ ($j = 1, \dots, n$) satisfy the hypotheses of Theorem 2.1, that is, we claim that

$$(9) \quad \sum_{m=1}^\infty \frac{1}{t_j(mn)^{1/mn}} = \infty.$$

To see this we fix j . If $|x_j| = 0$ almost everywhere (ν) then (9) is obvious. If $|x_j|$ is not ν -almost everywhere equal to 0, we define the (then finite valued) nonnegative sequence $\{h_j(m)\}_{m=1}^\infty$ by

$$h_j(m) = t_j(m)^{-1/m}, \quad m = 1, 2, \dots$$

By (5) the sequence $\{h_j(m)\}_{m=1}^\infty$ is nonincreasing. For m even, we have $s_j(m) = t_j(m)$; the hypothesis (3) therefore translates as $\sum_{m=1}^\infty h_j(2m) = \infty$. Lemma 2.2 then implies that $\sum_{m=1}^\infty h_j(mn) = \infty$, which is (9). This establishes the claim.

To conclude the proof of the determinacy we note that by (8) all derivatives of $\frac{1}{2}(\widehat{\mu}_1 - \widehat{\mu}_2)$ vanish at 0. Therefore, Theorem 2.1 now shows that $\widehat{\mu}_1 = \widehat{\mu}_2$, implying that $\mu_1 = \mu_2$, as was to be proved.

We turn to the density statements in $L_p(\mathbb{R}^n, \mu)$ for $1 \leq p < \infty$. Fix such p and let $1 < q \leq \infty$ be the conjugate exponent.

We treat the polynomials first. Suppose $f \in L_q(\mathbb{R}^n, \mu)$ is such that

$$(10) \quad \int_{\mathbb{R}^n} P(x) f(x) d\mu(x) = 0$$

for all polynomials P . We need to prove that $f = 0$ a.e. (μ) . Define the complex Borel measure ξ_f on \mathbb{R}^n by

$$(11) \quad \xi_f(E) = \int_E f(x) d\mu(x)$$

for Borel sets E . Consider the Fourier transform

$$(12) \quad \widehat{\xi}_f(\lambda) = \int_{\mathbb{R}^n} e^{i(\lambda, x)} d\xi_f(x) = \int_{\mathbb{R}^n} e^{i(\lambda, x)} f(x) d\mu(x).$$

Then $\widehat{\xi}_f$ is of class C^∞ on \mathbb{R}^n with derivatives

$$(13) \quad \frac{\partial^\alpha \widehat{\xi}_f}{\partial \lambda^\alpha}(\lambda) = \int_{\mathbb{R}^n} i^{|\alpha|} x^\alpha e^{i(\lambda, x)} f(x) d\mu(x), \quad \alpha \in \mathbb{N}^n, \lambda \in \mathbb{R}^n.$$

By assumption we therefore have

$$(14) \quad \frac{\partial^\alpha \widehat{\xi}_f}{\partial \lambda^\alpha}(0) = 0, \quad \alpha \in \mathbb{N}^n.$$

For $\alpha \in \mathbb{N}^n$ and $\lambda \in \mathbb{R}^n$, we have the following estimate, as a consequence of (13) and the generalized Hölder inequality (the norms refer to μ).

$$\begin{aligned} \left| \frac{\partial^\alpha \widehat{\xi}_f}{\partial \lambda^\alpha}(\lambda) \right| &\leq \| |x|^\alpha |f| \|_1 \\ &\leq \|f\|_q \| |x|^\alpha \|_p \\ &= \|f\|_q \left\| \prod_{j=1}^n |x_j|^{\alpha_j p} \right\|_1^{1/p} \\ &\leq \|f\|_q \prod_{j=1}^n \| |x_j|^{\alpha_j p} \|_n^{1/p} \\ &= \|f\|_q \prod_{j=1}^n t_j(\alpha_j p n)^{1/n p}. \end{aligned}$$

Here we have used (4) for the definition of $t_j(s)$ ($j = 1, \dots, n, s \geq 0$), which is correct since we already know that $\nu = \mu$. We claim that the nonnegative sequences $\{t_j(mpn)^{1/np}\}_{m=0}^\infty$ ($j = 1, \dots, n$) satisfy the hypotheses of Theorem 2.1, that is, we claim that

$$(15) \quad \sum_{m=1}^\infty \frac{1}{t_j(mpn)^{1/mnp}} = \infty.$$

To see this we again fix j . If $|x_j| = 0$ almost everywhere (μ) then (15) is again obvious. If $|x_j|$ is not μ -almost everywhere equal to 0, then we note that (5) implies that

$$(16) \quad \sum_{m=1}^{\infty} \frac{1}{t_j(mpn)^{1/mnp}} \geq \sum_{m=1}^{\infty} \frac{1}{t_j(m([p] + 1)n)^{1/m([p]+1)n}},$$

where $[p]$ is the largest integer not exceeding p . In the notation as in the proof of the determinacy, the right-hand side of (16) is $\sum_{m=1}^{\infty} h_j(m([p] + 1)n)$. Again, Lemma 2.2 then implies that this series is divergent since $\sum_{m=1}^{\infty} h_j(2m)$ diverges, thus establishing the claim.

In view of (14), we now conclude from Theorem 2.1 that $\widehat{\xi}_f = 0$, implying $\xi_f = 0$ and finally that $f = 0$ a.e. (μ), as was to be proved.

Finally, let us prove the density of $\text{Span}_{\mathbb{C}}\{e_{i\lambda} \mid \lambda \in S\}$ for a subset S of \mathbb{R}^n such that \overline{S} has nonempty interior. Assume that $a \in \mathbb{R}^n$ is an interior point of \overline{S} and suppose $f \in L_p(\mathbb{R}^n, \mu)$ vanishes on $\text{Span}_{\mathbb{C}}\{e_{i\lambda} \mid \lambda \in S\}$. Consider the Fourier transform

$$(17) \quad \widehat{e_{ia}f}(\lambda) = \int_{\mathbb{R}^n} e^{i(\lambda+a \cdot x)} f(x) d\mu(x), \quad \lambda \in \mathbb{R}^n.$$

Then $\widehat{e_{ia}f}$ is of class C^∞ and the assumption on f implies that $\widehat{e_{ia}f}$ is identically zero on a neighborhood of $0 \in \mathbb{R}^n$. Evidently all derivatives of $\widehat{e_{ia}f}$ then vanish at 0, which shows that $e_{ia}f$ vanishes on the polynomials. Since we had already shown that these are dense in $L_p(\mathbb{R}^n, \mu)$, we conclude that $f = 0$ a.e. (μ), as was to be proved. \square

We comment on the relation between Theorem 2.3 and the literature.

The fact that the divergence of the series in Theorem 2.3 is sufficient for the determinacy of the measure can already be found in [13], where a combination of quasi-analytic methods and a Hilbert space approach is used.

Theorem 2.3 is also related to the following result ([16], page 21): if $\mu \in \mathcal{M}^*$ and if we have $\sum_{m=1}^{\infty} 1/\sqrt[2m]{\lambda(2m)} = \infty$, where

$$(18) \quad \lambda(m) = \int_{\mathbb{R}^n} \sum_{j=1}^n x_j^m d\mu(x), \quad m = 0, 1, 2, \dots,$$

then μ is determinate. It is under this condition on the $\lambda(2m)$ even true [1, 2] that the polynomials are dense in $L_p(\mathbb{R}^n, \mu)$ for all $1 \leq p < \infty$, a property which is stronger than determinacy of the measure. As to this last implication, it was first proved in [3] in the one-dimensional case that, for $\mu \in \mathcal{M}^*$, the density of the polynomials in $L_p(\mathbb{R}, \mu)$ for some finite $p > 2$ implies that μ is determinate, and this result was later generalized to arbitrary dimension in [8].

The determinacy and polynomial density under the condition on the $\lambda(2m)$ also follow from Theorem 2.3. Indeed, taking the standard basis in Theorem 2.3, one

obviously has $s_j(2m) \leq \lambda(2m)$. The divergence of the series for $\lambda(2m)$ therefore implies the divergence of the series for all $s_j(m)$, so that the conclusions of Theorem 2.3 on determinacy and density hold.

Conversely, the special case $\mathbb{R}^n = \mathbb{R}$ (see [4]) of the results on determinacy and polynomial density as quoted above can be taken as a starting point to derive Theorem 2.3, albeit in a more indirect fashion than in the present paper. Indeed, assuming that the basis in Theorem 2.3 is the standard basis, one concludes from this one-dimensional starting point that all marginal distributions of the measure in Theorem 2.3 are determinate and that the polynomials are dense in all L_p -spaces ($1 \leq p < \infty$) associated with these marginal distributions. The results in [15] then imply that the analogous two statements hold for the measure itself. The additional density of the trigonometric functions then also follows. Indeed, the conclusion of the proof of Theorem 2.3 shows, in fact, that for $\mu \in \mathcal{M}^*$ and $1 \leq p < \infty$, the density of the polynomials in $L_p(\mathbb{R}^n, \mu)$ implies the density in $L_p(\mathbb{R}^n, \mu)$ of $\text{Span}_{\mathbb{C}}\{e_{i\lambda} \mid \lambda \in S\}$ for all somewhere dense subsets S of \mathbb{R}^n . The author is indebted to Christian Berg for communicating this last result and its proof.

3. Quasi-analytic weights. This section is a preparation for Sections 4 and 5. We define quasi-analytic weights, mention the relevant properties and the classification and give practical examples.

The notion of quasi-analytic weight is a delicate one, which is studied systematically in [6]. Simple and intuitive properties are sometimes not immediately obvious and may require an argument. We therefore refer for proofs to [6], from which all results in this section are taken.

A weight on \mathbb{R}^n is an arbitrary bounded nonnegative function on \mathbb{R}^n . We emphasize that we assume no regularity.

DEFINITION 3.1. Let w be a weight on \mathbb{R}^n . Suppose $\{v_1, \dots, v_n\}$ is a basis of \mathbb{R}^n . If

$$\sum_{m=1}^{\infty} \frac{1}{\|(v_j, x)^m w(x)\|_{\infty}^{1/m}} = \infty$$

for $j = 1, \dots, n$ then w is *quasi-analytic with respect to* $\{v_1, \dots, v_n\}$. A weight is *standard quasi-analytic* if it is quasi-analytic with respect to the standard basis of \mathbb{R}^n . A weight is *quasi-analytic* if it is quasi-analytic with respect to some basis.

The terminology “quasi-analytic” refers not to regularity of the weight itself, which might, for example, even fail to be Lebesgue measurable. The reason then for this terminology lies—as has become customary—in the fact that certain crucial associated functions have the quasi-analytic property, by which is meant that one can conclude that such an associated function is actually identically zero once one has established that the function and all its derivatives vanish at one fixed

point. These associated functions thus share this property with analytic functions in function theory, which explains the nomenclature. In our case, the reader may verify that it is indeed the divergence of the series in Definition 3.1 that validates the application of Theorem 2.1 on the quasi-analytic property in the proof of the basic Theorem 2.3.

A weight which vanishes outside a compact set is quasi-analytic with respect to all bases. By a small computation, the same holds for a weight of type $\exp(-\varepsilon \|x\|)$ with $\varepsilon > 0$. The set of quasi-analytic weights is invariant under the group of affine automorphisms of \mathbb{R}^n and under multiplication with nonnegative constants.

Let w be a weight, quasi-analytic with respect to the basis $\{v_1, \dots, v_n\}$. Then $\lim_{\|x\| \rightarrow \infty} \|x\|^d w(x) = 0$ for all $d \geq 0$, that is, w is rapidly decreasing. A minorant of w outside a compact set is again quasi-analytic with respect to $\{v_1, \dots, v_n\}$. One can prove that w always has a pointwise majorant of class C^∞ which is strictly positive and quasi-analytic with respect to $\{v_1, \dots, v_n\}$. In one dimension, such a majorant can, in addition, be required to be even and strictly decreasing on $[0, \infty)$.

There are several closely related ways of characterizing quasi-analytic weights other than by Definition 3.1, which is a technically convenient characterization but not a very practical one to verify. The formulation in the following paragraphs seems to fit most applications. For additional material the reader is referred to [6].

A weight w on \mathbb{R}^n is quasi-analytic if and only if there exists an affine automorphism A of \mathbb{R}^n and quasi-analytic weights w_j ($j = 1, \dots, n$) on \mathbb{R} such that

$$(19) \quad w(Ax) \leq \prod_{j=1}^m w_j(x_j)$$

for all $x \in \mathbb{R}^n$. More precisely, if w is quasi-analytic with respect to the basis $\{v_1, \dots, v_n\}$ of \mathbb{R}^n , then quasi-analytic weights w_j on \mathbb{R} satisfying (19) exist for any A with linear component A_0 defined by $A_0^t v_j = e_j$ ($j = 1, \dots, n$); here A_0^t is the transpose of A_0 with respect to the standard inner product on \mathbb{R}^n . Conversely, if (19) holds for some A and quasi-analytic weights w_j on \mathbb{R} , and if A_0 is the linear component of A , then w is quasi-analytic with respect to the basis $\{v_1, \dots, v_n\}$ of \mathbb{R}^n defined by $A_0^t v_j = e_j$ ($j = 1, \dots, n$).

The matter has now been reduced to \mathbb{R} . As a first equivalent characterization on the real line, a weight w on \mathbb{R} is quasi-analytic if and only if there exist $R > 0$, $C \geq 0$ and a nondecreasing function $\rho : (R, \infty) \mapsto \mathbb{R}_{\geq 0}$ of class C^1 such that

$$(20) \quad \int_R^\infty \frac{\rho(s)}{s^2} ds = \infty$$

and

$$w(x) \leq C \exp\left(-\int_R^{|x|} \frac{\rho(s)}{s} ds\right)$$

if $|x| \geq R$.

As a second and closely related equivalent characterization on the real line, a weight on \mathbb{R} is quasi-analytic if and only if there exists a weight \tilde{w} on \mathbb{R} and $R > 0$ such that $w(t) \leq \tilde{w}(t)$ and $\tilde{w}(t) = \tilde{w}(-t) > 0$ both hold for $|t| > R$, such that $s \mapsto -\log \tilde{w}(e^s)$ is convex on $(\log R, \infty)$ and such that

$$\int_R^\infty \frac{\log \tilde{w}(t)}{1+t^2} = -\infty.$$

Weights such as \tilde{w} are classical and figure, for example, in the Bernstein problem [11]. The connection between these classical weights and the one-dimensional version of Definition 3.1 seems to have gone largely unnoticed, although some ingredients can be found in [12], proof of Theorem 2, under additional regularity conditions on the weight.

If w is a quasi-analytic weight on \mathbb{R} , then one obtains a quasi-analytic weight w' on \mathbb{R}^n by putting $w'(x) = w(\|x\|)$ for $x \in \mathbb{R}^n$. Such w' is then quasi-analytic with respect to all bases of \mathbb{R}^n . All minorants of w' outside a compact set are then again quasi-analytic with respect to all bases of \mathbb{R}^n . The first alternative characterization of quasi-analytic weights on \mathbb{R} , combined with this radial extension procedure thus yields the following.

PROPOSITION 3.2. *Suppose $R > 0$ and a nondecreasing function $\rho : (R, \infty) \mapsto \mathbb{R}_{\geq 0}$ of class C^1 are such that*

$$\int_R^\infty \frac{\rho(s)}{s^2} ds = \infty.$$

If w is a weight such that

$$w(x) \leq C \exp\left(-\int_R^{\|x\|} \frac{\rho(s)}{s} ds\right)$$

whenever $\|x\| \geq R$, then w is a weight on \mathbb{R}^n which is quasi-analytic with respect to all bases of \mathbb{R}^n .

The following result in terms of elementary functions is based on the second alternative characterization of quasi-analytic weights on \mathbb{R} , combined with the radial extension procedure.

PROPOSITION 3.3. *Define repeated logarithms by $\log_0 t = t$ and, inductively, for $j \geq 1$, by $\log_j t = \log(\log_{j-1} t)$, where t is assumed to be sufficiently large for the definition to be meaningful in the real context. For $j = 0, 1, 2, \dots$ let $a_j > 0$ and let $p_j \in \mathbb{R}$ be such that $p_j = 0$ for all sufficiently large j . Put $j_0 = \min\{j = 0, 1, 2, \dots \mid p_j \neq 1\}$. Let $C > 0$ and suppose $w : \mathbb{R}^n \mapsto \mathbb{R}_{\geq 0}$ is bounded.*

Then, if $p_{j_0} < 1$ and if

$$w(x) \leq C \exp\left(-\|x\|^2 \left(\prod_{j=0}^{\infty} \log_j^{p_j} a_j \|x\|\right)^{-1}\right)$$

for all sufficiently large $\|x\|$, w is a weight on \mathbb{R}^n which is quasi-analytic with respect to all bases of \mathbb{R}^n .

Note the occurrence of \log_0 (i.e., of the identity) in the proposition, which permits a uniform formulation.

Thus, to give explicit examples, a weight on \mathbb{R}^n is quasi-analytic if it is for all sufficiently large $\|x\|$ majorized by one of the expressions

$$\begin{aligned} & C \exp\left(-\frac{\|x\|^{1-\nu}}{a_0}\right), \\ & C \exp\left(-\frac{\|x\|}{a_0(\log a_1 \|x\|)^{1+\nu}}\right), \\ & C \exp\left(-\frac{\|x\|}{a_0 \log a_1 \|x\| (\log \log a_2 \|x\|)^{1+\nu}}\right), \\ & \dots \end{aligned}$$

for some $C, a_0, a_1, a_2, \dots > 0$ and $\nu \leq 0$. The case $\nu = 0$ yields a sequence of families of quasi-analytic weights, each consisting of weights that are negligible at infinity compared with any member of the succeeding family.

Explicit nonradial examples of standard quasi-analytic weights on \mathbb{R}^n in terms of elementary functions can be obtained as tensor products of quasi-analytic weights on \mathbb{R} taken from Proposition 3.3. All minorants of such tensor products outside a compact set are then again standard quasi-analytic weights on \mathbb{R}^n . Insertion of an affine automorphism of \mathbb{R}^n in the argument of the weight yields additional quasi-analytic weights.

For the sake of completeness, we mention that we have the following negative criterion for a weight to be quasi-analytic: if w is a quasi-analytic weight on \mathbb{R}^n and if $x, y \in \mathbb{R}^n$ are such that $y \neq 0$ and such that $t \mapsto w(x + ty)$ is Lebesgue measurable on \mathbb{R} , then for all $R > 0$, we have

$$\int_R^{\infty} \frac{\log w(x + ty)}{1 + t^2} dt = \int_{-\infty}^{-R} \frac{\log w(x + ty)}{1 + t^2} dt = -\infty.$$

This shows that Proposition 3.3 is sharp in the sense that the corresponding statement for $p_{j_0} > 1$ does not hold.

The set of quasi-analytic weights has some interesting characteristics. Contrary to what the explicit examples above suggest, this set is not closed under addition. More precisely, one can construct weights w_1 and w_2 on \mathbb{R}^n , each of which is quasi-analytic with respect to all bases, but such that $w_1 + w_2$ is not quasi-analytic

with respect to any basis. One can also construct weights which are quasi-analytic with respect to just *one* basis (up to scaling); in Section 6 we will make some tentative remarks on a possible parallel of this phenomenon for measures. For $n \geq 2$ it implies that such quasi-analytic weights on \mathbb{R}^n are not minorants outside a compact set of quasi-analytic weights as obtained from the radial extension procedure.

4. Integral criteria for determinacy. We will now combine the results in Sections 2 and 3.

THEOREM 4.1 (First main theorem). *Let μ be a positive Borel measure on \mathbb{R}^n such that*

$$\int_{\mathbb{R}^n} w(x)^{-1} d\mu < \infty$$

for some measurable quasi-analytic weight. Then μ is determinate. Furthermore, the polynomials and $\text{Span}_{\mathbb{C}}\{e_{i\lambda} \mid \lambda \in S\}$ are then dense in $L_p(\mathbb{R}^n, \mu)$ for all $1 \leq p < \infty$ and for every subset S of \mathbb{R}^n which is somewhere dense.

Note that since quasi-analytic weights are rapidly decreasing, the measure in the theorem is automatically in \mathcal{M}^* .

PROOF OF THEOREM 4.1. Suppose w is quasi-analytic with respect to the basis $\{v_1, \dots, v_n\}$. We may assume that w is strictly positive: if necessary we can replace w by a strictly positive measurable (say, smooth) majorant which is quasi-analytic with respect to $\{v_1, \dots, v_n\}$. We may also assume that $\|w\|_{\infty} = 1$.

In the notation of Theorem 2.3 we then have, for $j = 1, \dots, n$ and $m = 0, 1, 2, \dots$,

$$\begin{aligned} s_j(2m) &= \int_{\mathbb{R}^n} (v_j, x)^{2m} d\mu(x) \\ (21) \qquad &= \int_{\mathbb{R}^n} (v_j, x)^{2m} w(x) w(x)^{-1} d\mu(x) \\ &\leq \|(v_j, x)^{2m} w(x)\|_{\infty} \int_{\mathbb{R}^n} w(x)^{-1} d\mu(x). \end{aligned}$$

Now the sequences $\{\|(v_j, x)^m w(x)\|_{\infty}^{1/m}\}_{m=1}^{\infty}$ ($j = 1, \dots, n$) are easily seen to be nondecreasing, as a consequence of the normalization $\|w\|_{\infty} = 1$. The quasi-analyticity of w with respect to $\{v_1, \dots, v_n\}$ therefore implies by Lemma 2.2 that

$$\sum_{m=1}^{\infty} \frac{1}{\|(v_j, x)^{2m} w(x)\|_{\infty}^{1/2m}} = \infty, \qquad j = 1, \dots, n.$$

This divergence implies, in view of the estimate in (21), that the hypotheses of Theorem 2.3 are satisfied, which concludes the proof. \square

Theorem 4.1 can also be found in [6], where the density part is seen to be a consequence of more general considerations on the closure of modules over the polynomials and trigonometric functions in topological vector spaces. In [6] the determinacy of the measure is then concluded from [8] since there exists $p > 2$ such that the polynomials are dense in the associated L_p -space. This way of deriving Theorem 4.1 is considerably more involved than the present proof.

The combination of Theorem 4.1 with the results on quasi-analytic weights in Section 3 now yields various integral criteria for determinacy and density, as mentioned in the Introduction. Variation in these criteria, in particular, variation in the degree of regularity of the weights involved, is possible in view of the various ways in which quasi-analytic weights can be characterized. In the criteria as described in this section, all integrands are of class C^2 outside a compact set. For cases where this is too stringent the reader is referred to [6].

A nonradial criterion is the following:

THEOREM 4.2. *For $j = 1, \dots, n$ let $R_j > 0$ and a nondecreasing function $\rho_j : (R_j, \infty) \mapsto \mathbb{R}_{\geq 0}$ of class C^1 be such that*

$$\int_{R_j}^{\infty} \frac{\rho_j(s)}{s^2} ds = \infty.$$

Define $f_j : \mathbb{R} \mapsto \mathbb{R}_{\geq 0}$ by

$$f_j(x) = \exp\left(\int_{R_j}^{|x|} \frac{\rho_j(s)}{s} ds\right)$$

for $|x| > R_j$ and by $f_j(x) = 1$ for $|x| \leq R_j$. Let A be an affine automorphism of \mathbb{R}^n . If μ is a positive Borel measure on \mathbb{R}^n such that

$$\int_{\mathbb{R}^n} \prod_{j=1}^n f_j((Ax)_j) d\mu(x) < \infty,$$

then μ is determinate. Furthermore, the polynomials and $\text{Span}_{\mathbb{C}}\{e_{i\lambda} \mid \lambda \in S\}$ are then dense in $L_p(\mathbb{R}^n, \mu)$ for all $1 \leq p < \infty$ and for every subset S of \mathbb{R}^n which is somewhere dense.

PROOF. From the first alternative characterization of quasi-analytic weights on \mathbb{R} , as given in Section 3, we see that the weights $1/f_j$ are all quasi-analytic weights on \mathbb{R} . Their tensor product is then a quasi-analytic weight on \mathbb{R}^n and then the same holds for the image of this tensor product under an element of the affine group. We now apply Theorem 4.1. \square

We now specialize to the case of radial integrands. Theorem 1.1 evidently follows from Theorem 4.1 and Proposition 3.2. In addition, the combination of Proposition 3.3 and Theorem 4.1 implies the following. As with Proposition 3.3, note the occurrence of \log_0 , that is, of the identity.

THEOREM 4.3. *Define repeated logarithms \log_j ($j = 0, 1, 2, \dots$) as in Proposition 3.3. For $j = 0, 1, 2, \dots$ let $a_j > 0$ and let $p_j \in \mathbb{R}$ be such that $p_j = 0$ for all sufficiently large j . Put $j_0 = \min\{j = 0, 1, 2, \dots \mid p_j \neq 1\}$ and assume $p_{j_0} < 1$.*

Let μ be a positive Borel measure on \mathbb{R}^n . If

$$\int_{\|x\| \geq R} \exp\left(\|x\|^2 \left(\prod_{j=0}^{\infty} \log_j^{p_j} a_j \|x\|\right)^{-1}\right) d\mu < \infty,$$

for some $R \geq 0$ which is sufficiently large to ensure that the integrand is defined, then μ is determinate. Furthermore, the polynomials and $\text{Span}_{\mathbb{C}}\{e_{i\lambda} \mid \lambda \in S\}$ are then dense in $L_p(\mathbb{R}^n, \mu)$ for all $1 \leq p < \infty$ and for every subset S of \mathbb{R}^n which is somewhere dense.

As explicit examples, if one of the functions (tacitly assumed to be equal to 1 on a sufficiently large compact set)

$$\begin{aligned} &\exp\left(\frac{\|x\|^{1-\nu}}{a_0}\right), \\ &\exp\left(\frac{\|x\|}{a_0(\log a_1 \|x\|)^{1+\nu}}\right), \\ &\exp\left(\frac{\|x\|}{a_0 \log a_1 \|x\| (\log \log a_2 \|x\|)^{1+\nu}}\right), \\ &\dots \end{aligned}$$

has finite integral under μ for some $a_0, a_1, a_2, \dots > 0$ and $\nu \leq 0$, then the conclusions in Theorem 4.3 hold. The classical condition of the integrability of $\exp(\varepsilon\|x\|)$ for some $\varepsilon > 0$ can be weakened quite substantially.

To conclude we mention that explicit nonradial reciprocals of quasi-analytic weights can be obtained in terms of elementary functions by taking the tensor product of the reciprocals of the one-dimensional versions of the majorants in Proposition 3.3, when these majorants are in addition defined to be equal to 1 on a sufficiently large compact subset of \mathbb{R} .

5. Determinacy in the sense of Stieltjes. In this section we are concerned with determinacy in the sense of Stieltjes, that is, with C -determinacy as in Definition 1.2. Analogously to the one-dimensional case, The Carleman criterion in Theorem 2.3 implies a similar sufficient condition for C -determinacy. When combined with the results on quasi-analytic weights again, we obtain integral criteria for C -determinacy. At the end of the section we discuss a condition (which is satisfied for absolutely continuous measures) enabling one to conclude that the measure is not just C -determinate, but in fact determinate.

THEOREM 5.1 (Carleman criterion for C -determinacy). *Let $\{v_1, \dots, v_n\}$ be a basis of \mathbb{R}^n and let $C = \mathbb{R}_{\geq 0} \cdot v_1 + \dots + \mathbb{R}_{\geq 0} \cdot v_n$ be the corresponding positive convex cone. Define the dual basis $\{v'_1, \dots, v'_n\}$ by $(v'_i, v_j) = \delta_{ij}$ ($i, j = 1, \dots, n$).*

Let $\mu \in \mathcal{M}^$ be supported in C . For $j = 1, \dots, n$ and $m = 0, 1, 2, \dots$, define*

$$s_j(m) = \int_C (v'_j, x)^m d\mu(x),$$

and suppose that each of the sequences $\{s_j(m)\}_{m=1}^\infty$ ($j = 1, \dots, n$) satisfies

$$(22) \quad \sum_{m=1}^\infty \frac{1}{s_j(m)^{1/2m}} = \infty.$$

Then μ is C -determinate.

Note that the $s_j(m)$ are defined in terms of distinguished coordinates on C , namely those corresponding to extremal generators of C .

PROOF OF THEOREM 5.1. The proof generalizes the well-known proof in one dimension. Let \mathcal{M}_C^* be the measures in \mathcal{M}^* which are supported in C . As a first preparation, define $\phi : \mathbb{R}^n \mapsto \mathbb{R}^n$ by

$$\phi(x) = \begin{cases} x, & \text{if } x \notin C, \\ \sum_{j=1}^n \sqrt{x_j} v_j, & \text{if } x = \sum_{j=1}^n x_j v_j, x_j \geq 0, j = 1, \dots, n. \end{cases}$$

For $\xi \in \mathcal{M}^*$, define $\xi_\phi \in \mathcal{M}^*$ by putting $\xi_\phi(A) = \xi(\phi^{-1}(A))$ for a Borel set A . The assignment $\xi \mapsto \xi_\phi$ defines an injective map from \mathcal{M}^* to \mathcal{M}^* which maps \mathcal{M}_C^* into itself, and

$$(23) \quad \int_{\mathbb{R}^n} P(x) d\xi_\phi(x) = \int_{\mathbb{R}^n} (P \circ \phi)(x) d\xi(x)$$

for all polynomials P and all $\xi \in \mathcal{M}^*$.

As a second preparation, let G be the group of linear isomorphisms of \mathbb{R}^n having 2^n elements, corresponding to all possible sign changes in the coordinates with respect to the basis $\{v_1, \dots, v_n\}$. For $\xi \in \mathcal{M}^*$ and $g \in G$, define $g \cdot \xi \in \mathcal{M}^*$ by putting $(g \cdot \xi)(A) = \xi(g^{-1}(A))$ for a Borel set A and let $\bar{\xi} = 2^{-n} \sum_{g \in G} g \cdot \xi$. The averaging map $\xi \mapsto \bar{\xi}$ is not injective as a map from \mathcal{M}^* to itself, but it is injective as a map from \mathcal{M}_C^* to \mathcal{M}^* . To see this, let $J \subset \{1, \dots, n\}$ have cardinality $|J|$ and define $C_J = \{x \in C \mid x_j = 0 \Leftrightarrow j \in J\}$. Then, for $\xi \in \mathcal{M}_C^*$, we have $\bar{\xi}(A_J) = 2^{|J|-n} \xi(A_J)$ for any Borel subset A_J of C_J . Thus the restriction of ξ to C_J can be retrieved from $\bar{\xi}$. Since the C_J form a disjoint covering of C , we see that $\xi \in \mathcal{M}_C^*$ can be reconstructed from $\bar{\xi}$, as claimed.

Furthermore, if $\xi \in \mathcal{M}^*$, then

$$(24) \quad \int_{\mathbb{R}^n} (P \circ g)(x) d\bar{\xi}(x) = \int_{\mathbb{R}^n} P(x) d\bar{\xi}(x)$$

for all polynomials P , so that

$$(25) \quad \int_{\mathbb{R}^n} \prod_{j=1}^n (v'_j, x)^{e_j} d\bar{\xi}(x) = 0$$

if the e_j are nonnegative integers, at least one of which is odd. This follows from (24) by choosing an element in G which sends the integrand in the left-hand side of (25) to its negative. On the other hand, if $\xi \in \mathcal{M}^*$ and if the e_j are all even nonnegative integers, then

$$(26) \quad \int_{\mathbb{R}^n} \prod_{j=1}^n (v'_j, x)^{e_j} d\bar{\xi}(x) = \int_{\mathbb{R}^n} \prod_{j=1}^n (v'_j, x)^{e_j} d\xi(x)$$

as a consequence of the invariance of the integrand under G .

Combining (23), (25) and (26), we conclude that, for $\xi \in \mathcal{M}_C^*$,

$$(27) \quad \int_{\mathbb{R}^n} \prod_{j=1}^n (v'_j, x)^{e_j} d\bar{\xi}_\phi(x) = \int_C \prod_{j=1}^n (v'_j, x)^{e_j/2} d\xi(x)$$

if the e_j are all even nonnegative integers, whereas the integral on the left-hand side is 0 if the e_j are nonnegative integers, at least one of which is odd.

Turning to the theorem, we first of all note that $\overline{\mu_\phi}$ satisfies the conditions of Theorem 2.3 as a consequence of (27), so $\overline{\mu_\phi}$ is determinate as a measure on \mathbb{R}^n . Suppose then that $\nu \in \mathcal{M}_C^*$ yields the same integrals for all polynomials as μ . Then (27) implies that $\overline{\mu_\phi}$ and $\overline{\nu_\phi}$ also have the same integrals for all polynomials and we conclude that $\overline{\mu_\phi} = \overline{\nu_\phi}$. By the injectivity of the maps as observed above, it first follows that $\mu_\phi = \nu_\phi$ and subsequently that $\mu = \nu$. \square

We will now combine this with the results in Section 3.

THEOREM 5.2 (Second main theorem). *Let $\{v_1, \dots, v_n\}$ be a basis of \mathbb{R}^n and let $C = \mathbb{R}_{\geq 0} \cdot v_1 + \dots + \mathbb{R}_{\geq 0} \cdot v_n$ be the corresponding positive convex cone. Define the dual basis $\{v'_1, \dots, v'_n\}$ by $(v'_i, v_j) = \delta_{ij}$ ($i, j = 1, \dots, n$). Let w be a measurable weight on \mathbb{R}^n , quasi-analytic with respect to $\{v'_1, \dots, v'_n\}$. For $x = \sum_{j=1}^n x_j v_j \in C$ define $\phi(x) = \sum_{j=1}^n \sqrt{x_j} v_j$. Let μ be a positive Borel measure on \mathbb{R}^n which is supported in C and suppose that*

$$\int_C (w \circ \phi)(x)^{-1} d\mu(x) < \infty.$$

Then μ is C -determinate.

PROOF. The argument parallels the proof of Theorem 4.1. We may assume that w is strictly positive: if necessary we can replace w by a strictly positive

measurable (say, smooth) majorant which is quasi-analytic with respect to $\{v'_1, \dots, v'_n\}$. We may also assume that $\|w\|_\infty = 1$.

In the notation of Theorem 5.1 we then have, for $j = 1, \dots, n$ and $m = 0, 1, 2, \dots$,

$$s_j(m) \leq \left[\sup_{x \in C} (v'_j, x)^m (w \circ \phi)(x) \right] \int_C (w \circ \phi)(x)^{-1} d\mu(x).$$

Now

$$\begin{aligned} \sup_{x \in C} (v'_j, x)^m (w \circ \phi)(x) &= \sup_{x_1, \dots, x_n \geq 0} \left(v'_j, \sum_{j=1}^n x_j v_j \right)^m w \left(\sum_{j=1}^n \sqrt{x_j} v_j \right) \\ &= \sup_{t_1, \dots, t_n \geq 0} \left(v'_j, \sum_{j=1}^n t_j^2 v_j \right)^m w \left(\sum_{j=1}^n t_j v_j \right) \\ &= \sup_{x \in C} (v'_j, x)^{2m} w(x) \\ &\leq \|(v'_j, x)^{2m} w\|_\infty. \end{aligned}$$

As in the proof of Theorem 4.1, we conclude from these estimates and Lemma 2.2 that the hypotheses of Theorem 5.1 are satisfied. \square

The proof of the following theorem is left to the reader. It follows from Theorem 5.2, using the results in Section 3 on quasi-analytic weights, in a way similar to the proof of Theorem 4.2.

THEOREM 5.3. *Let $\{v_1, \dots, v_n\}$ be a basis of \mathbb{R}^n and let $C = \mathbb{R}_{\geq 0} \cdot v_1 + \dots + \mathbb{R}_{\geq 0} \cdot v_n$ be the corresponding positive convex cone. Define the dual basis $\{v'_1, \dots, v'_n\}$ by $(v'_i, v_j) = \delta_{ij}$ ($i, j = 1, \dots, n$). For $x = \sum_{j=1}^n x_j v_j \in C$ define $\phi(x) = \sum_{j=1}^n \sqrt{x_j} v_j$.*

For $j = 1, \dots, n$ let $R_j > 0$ and a nondecreasing function $\rho_j : (R_j, \infty) \mapsto \mathbb{R}_{\geq 0}$ of class C^1 be such that

$$\int_{R_j}^\infty \frac{\rho_j(s)}{s^2} ds = \infty.$$

Define $f_j : \mathbb{R} \mapsto \mathbb{R}_{\geq 0}$ by

$$f_j(x) = \exp \left(\int_{R_j}^{|x|} \frac{\rho_j(s)}{s} ds \right)$$

for $|x| > R_j$ and by $f_j(x) = 1$ for $|x| \leq R_j$.

If μ is a positive Borel measure on \mathbb{R}^n which is supported in C and if

$$\int_C \prod_{j=1}^n f_j((v'_j, \phi(x))) d\mu(x) < \infty,$$

then μ is C -determinate.

We turn to radial integrands, for which we will use the following practical result as a starting point.

THEOREM 5.4. *Let $\{v_1, \dots, v_n\}$ be a basis of \mathbb{R}^n and let $C = \mathbb{R}_{\geq 0} \cdot v_1 + \dots + \mathbb{R}_{\geq 0} \cdot v_n$ be the corresponding positive convex cone. Let w be a measurable quasi-analytic weight on the real line. Let μ be a positive Borel measure on \mathbb{R}^n which is supported in C and suppose that*

$$\int_C w(\sqrt{\|x\|})^{-1} d\mu(x) < \infty.$$

Then μ is C -determinate.

PROOF. As with Theorem 5.2, the proof parallels that of Theorem 4.1. We may assume that w is strictly positive, by replacing w with a quasi-analytic majorant with this property if necessary. We may also assume that $\|w\|_\infty = 1$. In the notation of Theorem 5.1 we then have, for $j = 1, \dots, n$ and $m = 0, 1, 2, \dots$,

$$s_j(m) \leq \left[\sup_{x \in C} (v'_j, x)^m w(\sqrt{\|x\|}) \right] \int_C w(\sqrt{\|x\|})^{-1} d\mu(x).$$

Now

$$\begin{aligned} \sup_{x \in C} (v'_j, x)^m w(\sqrt{\|x\|}) &\leq \|v'_j\|^m \sup_{x \in C} \|x\|^m w(\sqrt{\|x\|}) \\ &\leq \|v'_j\|^m \sup_{t \in \mathbb{R}} |t^{2m} w(t)|. \end{aligned}$$

As in the proof of Theorem 4.1, we conclude from these estimates and Lemma 2.2 that the hypotheses of Theorem 5.1 are satisfied. \square

Theorem 1.3 is now obvious, given Theorem 5.4 and Proposition 3.2. The combination of Proposition 3.3 and Theorem 5.4 implies the following (as with Proposition 3.3, note the occurrence of \log_0 , i.e., of the identity).

THEOREM 5.5. *Define repeated logarithms \log_j ($j = 0, 1, 2, \dots$) as in Proposition 3.3. For $j = 0, 1, 2, \dots$ let $a_j > 0$ and let $p_j \in \mathbb{R}$ be such that $p_j = 0$ for all sufficiently large j . Put $j_0 = \min\{j = 0, 1, 2, \dots \mid p_j \neq 1\}$ and assume $p_{j_0} < 1$.*

Let $\{v_1, \dots, v_n\}$ be a basis of \mathbb{R}^n and let $C = \mathbb{R}_{\geq 0} \cdot v_1 + \dots + \mathbb{R}_{\geq 0} \cdot v_n$ be the corresponding positive convex cone. Suppose μ is a positive Borel measure on \mathbb{R}^n which is supported in C and such that

$$\int_{\|x\| \geq R} \exp\left(\|x\|^{3/2} \left(\prod_{j=0}^{\infty} \log_j^{p_j} a_j \sqrt{\|x\|}\right)^{-1}\right) d\mu < \infty$$

for some $R \geq 0$ which is sufficiently large to ensure that the integrand is defined. Then μ is C -determinate.

As a consequence, if $\mu \in \mathcal{M}^*$ is supported in a positive convex cone C as above and if one of the functions (tacitly assumed to be equal to 1 on a sufficiently large compact set)

$$\begin{aligned} &\exp\left(\frac{\|x\|^{1/2-\nu}}{a_0}\right), \\ &\exp\left(\frac{\sqrt{\|x\|}}{a_0(\log a_1\|x\|)^{1+\nu}}\right), \\ &\exp\left(\frac{\sqrt{\|x\|}}{a_0 \log a_1\|x\|(\log \log a_2\|x\|)^{1+\nu}}\right), \\ &\dots \end{aligned}$$

has finite integral under μ for some $a_0, a_1, a_2, \dots > 0$ and $\nu \leq 0$, then μ is C -determinate.

We end this section by showing that in many cases—for example, if the measure is absolutely continuous with respect to Lebesgue measure—the conclusion of C -determinacy in the Theorems 5.1–5.5 can be strengthened to determinacy. It is sufficient to discuss strengthening Theorem 5.1, since this result implies the others. We assume that the basis $\{v_1, \dots, v_n\}$ generating the cone C is the standard basis; this simplifies the discussion and the general case follows from this by a linear transformation.

To start with, note that the hypotheses in Theorem 5.1 imply that the marginal distributions of μ are determinate in the sense of Stieltjes. Recall (see [5], page 481) that in one dimension a measure, which is supported in $[0, \infty)$ and which is determinate in the sense of Stieltjes, is actually determinate if its support does not contain 0 and/or its support is not equal to a discrete unbounded set. Therefore, if the support of each marginal distribution satisfies this condition, all marginal distributions are actually determinate. The results in [15] then imply that μ itself is determinate. To summarize:

Let $\{v_1, \dots, v_n\}$, C and μ be as in Theorems 5.1–5.4 or 5.5. Define marginal distributions of μ in terms of the projections corresponding to the basis $\{v_1, \dots, v_n\}$. If the support of each of these marginal distributions does not contain 0 and/or is not equal to a discrete unbounded set, then μ is determinate.

6. Closing remark. As mentioned in Section 3 there exist quasi-analytic weights on \mathbb{R}^n which are quasi-analytic with respect to a unique basis (up to scaling). For $n \geq 2$ the demonstration of this phenomenon in [6] is based on the

construction of n strictly positive logarithmically convex sequences $\{M_j(m)\}_{m=1}^\infty$ ($j = 1, \dots, n$) such that

$$\sum_{m=1}^{\infty} M_j(m)^{-1/m} = \infty, \quad j = 1, \dots, n,$$

but

$$\sum_{m=1}^{\infty} (\max(M_{j_1}(m), M_{j_2}(m)))^{-1/m} < \infty, \quad 1 \leq j_1 \neq j_2 \leq n.$$

Sequences satisfying the first of these equations also figure in Theorem 2.1

Now Theorem 2.1 can evidently be formulated with respect to any basis of \mathbb{R}^n , leading naturally to the notion of quasi-analytic classes with respect to bases. It is an interesting question whether there then exists an analogue of the aforementioned phenomenon for quasi-analytic weights. More precisely, can one, perhaps using sequences as above satisfying *both* equations, establish the existence of smooth functions that are in a quasi-analytic class with respect to a *unique* basis, up to scaling? If so, then in view of the proof of the extended Carleman theorem, additional argumentation could conceivably lead to the construction of multidimensional measures to which the extended Carleman theorem applies, but applies with *only one* basis, again up to scaling. Such measures would then have a distinguished set of marginal distributions.

Acknowledgment. It is a pleasure to thank Christian Berg for a helpful exposition on the subject and useful comments on a previous version of the paper.

REFERENCES

- [1] BERG, C. (1995). Recent results about moment problems. In *Probability Measures on Groups and Related Structures XI* (H. Heyer, ed.) 1–13. World Scientific, River Edge, NJ.
- [2] BERG, C. (1996). Moment problems and polynomial approximation. 100 ans après Th. J. Stieltjes. *Ann. Fac. Sci. Toulouse Math.* **6** (special issue) 9–32.
- [3] BERG, C. and CHRISTENSEN, J. P. R. (1981). Density questions in the classical theory of moments. *Ann. Inst. Fourier (Grenoble)* **31** 99–114.
- [4] BERG, C. and CHRISTENSEN, J. P. R. (1983). Exposants critiques dans le problème des moments. *C. R. Acad. Sci. Paris Sér. I Math.* **296** 661–663.
- [5] CHIHARA, T. S. (1968). On indeterminate Hamburger moment problems. *Pacific J. Math.* **27** 475–484.
- [6] DE JEU, M. F. E. (2001). Subspaces with equal closure. *Constr. Approx.* To appear.
- [7] DUNFORD, N. and SCHWARTZ, J. T. (1957). *Linear Operators. Part I.* Wiley, New York.
- [8] FUGLEDE, B. (1983). The multidimensional moment problem. *Expo. Math.* **1** 47–65.
- [9] HOFFMAN-JORGENSEN, J. (2002). The moment problem. Preprint.
- [10] HRYPTUN, V. G. (1976). An addition to a theorem of S. Mandelbrojt. *Ukrain. Mat. Zh.* **28** 841–844. (English translation: *Ukrainian Math. J.* **28** 655–658.)
- [11] KOOSIS, P. (1988). *The Logarithmic Integral. I.* Cambridge Univ. Press.
- [12] LIN, G. D. (1997). On the moment problems. *Statist. Probab. Lett.* **35** 85–90.

- [13] NUSSBAUM, A. E. (1966). Quasi-analytic vectors. *Ark. Mat.* **6** 179–191.
- [14] PAKES, A. G., HUNG, W. L. and WU, J. W. (2001). Criteria for the unique determination of probability distributions by moments. *Aust. N. Z. J. Stat.* **43** 101–111.
- [15] PETERSEN, L. C. (1982). On the relation between the multidimensional moment problem and the one-dimensional moment problem. *Math. Scand.* **51** 361–366.
- [16] SHOHAT, J. and TAMARKIN, J. D. (1943). *The Problem of Moments*. Amer. Math. Soc., Providence, RI.

MATHEMATICAL INSTITUTE
LEIDEN UNIVERSITY
P. O. BOX 9512
2300 RA LEIDEN
THE NETHERLANDS
E-MAIL: mdejeu@math.leidenuniv.nl
WEB: www.math.leidenuniv.nl/~mdejeu/