

RESEARCH ARTICLE

Selective Blockade of Interferon- α and - β Reveals Their Non-Redundant Functions in a Mouse Model of West Nile Virus Infection

Kathleen C. F. Sheehan^{1,4}, Helen M. Lazear², Michael S. Diamond^{1,2,3,4}, Robert D. Schreiber^{1,4*}

1 Department of Pathology and Immunology, Washington University School of Medicine, St. Louis, Missouri, United States of America, **2** Department of Medicine, Washington University School of Medicine, St. Louis, Missouri, United States of America, **3** Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, Missouri, United States of America, **4** Center for Human Immunology and Immunotherapy Programs, Washington University School of Medicine, St. Louis, Missouri, United States of America

* Schreiber@immunology.wustl.edu



OPEN ACCESS

Citation: Sheehan KCF, Lazear HM, Diamond MS, Schreiber RD (2015) Selective Blockade of Interferon- α and - β Reveals Their Non-Redundant Functions in a Mouse Model of West Nile Virus Infection. PLoS ONE 10(5): e0128636. doi:10.1371/journal.pone.0128636

Academic Editor: Rafael Aldabe, Centro de Investigación en Medicina Aplicada (CIMA), SPAIN

Received: October 6, 2014

Accepted: April 30, 2015

Published: May 26, 2015

Copyright: © 2015 Sheehan et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by U54 AI057160 (RDS) <http://www.niaid.nih.gov/labsandresources/resources/rce/Pages/default.aspx>, R01 CA043059 (RDS) <http://www.cancer.gov>, U19 AI083019 (MSD) <http://www.niaid.nih.gov>, U19 AI106772 (MSD) <http://www.niaid.nih.gov> and the R01 AI104972 (MSD) <http://www.niaid.nih.gov>. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abstract

Although type I interferons (IFNs) were first described almost 60 years ago, the ability to monitor and modulate the functional activities of the individual IFN subtypes that comprise this family has been hindered by a lack of reagents. The major type I IFNs, IFN- β and the multiple subtypes of IFN- α , are expressed widely and induce their effects on cells by interacting with a shared heterodimeric receptor (IFNAR). In the mouse, the physiologic actions of IFN- α and IFN- β have been defined using polyclonal anti-type I IFN sera, by targeting IFNAR using monoclonal antibodies or knockout mice, or using *Irfnb*^{-/-} mice. However, the corresponding analysis of IFN- α has been difficult because of its polygenic nature. Herein, we describe two monoclonal antibodies (mAbs) that differentially neutralize murine IFN- β or multiple subtypes of murine IFN- α . Using these mAbs, we distinguish specific contributions of IFN- β versus IFN- α in restricting viral pathogenesis and identify IFN- α as the key mediator of the antiviral response in mice infected with West Nile virus. This study thus suggests the utility of these new reagents in dissecting the antiviral and immunomodulatory roles of IFN- β versus IFN- α in murine models of infection, immunity, and autoimmunity.

Introduction

The type I interferons (IFNs), first identified by their ability to control viral infection [1, 2], are now known to contribute broadly to innate and adaptive immunity [3]. In mice, the type I IFN family includes IFN- β (encoded by a single gene), multiple IFN- α subtypes (14 genes and 3 pseudogenes), IFN- ζ (limitin) [4], IFN- ϵ [5, 6] and IFN- κ [7, 8]. The type I IFNs are encoded by single exon genes (with the exception of IFN- κ , which contains 1 intron) of similar structure, size, and conservation of protein sequence [6, 9, 10] but with divergent regulatory elements [11, 12]. Type I IFNs are induced after microbial products are sensed via pattern-

Competing Interests: The authors have declared that no competing interests exist.

recognition receptors (PRRs), which triggers activation and nuclear translocation of IRF-family transcription factors (IRF-1, -3, -5 and -7) [13–16]. While most cell types can produce IFN- β , the predominant source of IFN- α is hematopoietic cells, particularly plasmacytoid dendritic cells. [7, 17, 18].

All type I IFNs bind to the same receptor, IFNAR, a ubiquitously expressed heterodimer consisting of two subunits IFNAR1 and IFNAR2 [19–22]. Type I IFN binding to IFNAR activates the receptor associated tyrosine kinases, JAK1 and TYK2, which phosphorylate the latent transcription factors STAT1 and STAT2 to bind IRF-9. These form the ISGF3 complex, which then enters the nucleus, binds to the IFN response element in the promoters of hundreds of IFN stimulated genes (ISGs), and initiates their transcription. These ISGs promote antiviral, anti-proliferative, anti-tumor, and immunomodulatory functions [17, 23, 24]. More specifically, type I IFNs inhibit viral entry, transcription, translation, and assembly in host cells; augment host adaptive immune responses [25, 26]; and activate several key innate immune cell types including natural killer cells [27, 28], dendritic cells (DC) [29], CD8⁺ T cells [30] and B cells [31, 32]. Beyond the canonical STAT1-STAT2 signaling pathway, type I IFN-dependent activation of STAT1 and STAT3 homo- and heterodimers results in variable, context-specific, shifts in the balance of downstream signaling pathways, altering priming and induction of inflammatory responses [14, 33]. Moreover, individual type I IFN subtypes bind IFNAR with different affinities that may influence downstream gene activation [34]. Although type I IFNs trigger expression of an array of effector genes that limit viral infection [35, 36], their downstream effects also can be deleterious, leading to inflammatory immunopathology, cellular toxicity [37], cellular dysfunction [38–42] and suppression of antibacterial responses [43, 44]. Type I IFNs also have critical roles in promoting autoimmune disease [45–50] and in inducing host-protective responses to cancer [23, 51–54].

A deeper understanding of the physiologic functions of type I IFN has come from experiments using gene-targeted mice lacking *Ifnar1*, *Ifnar2* [21, 55–59], or *Ifnb* [60–62] or studies in which the various ligands or receptors are engineered to express point mutations that alter downstream signaling [63, 64]. However, distinguishing the specific *in vivo* effects of IFN- β and IFN- α has remained challenging. Better tools are needed to understand the mechanisms by which this family of cytokines functions in health and disease and how to balance protective versus harmful responses. Herein, we describe the generation and characterization of two monoclonal antibodies (mAbs), HD β -4A7 and TIF-3C5, which selectively bind and neutralize murine IFN- β or many IFN- α subtypes, respectively. Using a mouse model of West Nile virus (WNV) infection, we demonstrate the efficacy of these neutralizing mAbs *in vivo* and identify distinct roles for IFN- β and IFN- α in controlling WNV pathogenesis. Thus, these mAbs provide a new opportunity to investigate the physiological actions of IFN- α and IFN- β *in vivo* in many biological processes.

Materials and Methods

Ethics Statement

These studies were carried out in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institute of Health. All protocols are approved by the Washington University School of Medicine Animal Studies Committee (Animal Welfare Assurance # A3381-01). All efforts were made to minimize suffering of animals.

Animals

Specific pathogen free C57BL/6 mice were obtained from either Jackson Laboratories (Bar Harbor, ME) or Taconic Farms (Hudson, NY). *Ifnar1*^{-/-} [65], *Ifnb*^{-/-} [60], and *Irf7*^{-/-} [66] mice, all on

a C57BL/6 background, were bred in the barrier animal facilities of the Washington University School of Medicine and genotyped prior to experimentation. Male and female mice from 6–18 weeks of age were used in these studies. Armenian hamsters (F1) were purchased from Cytogen Research and Development (Cambridge, MA) and housed individually with enrichment.

Cytokines and Reagents

Recombinant murine IFN- β was purchased from BioLegend (San Diego, CA), additionally *E. coli*-derived IFN- β and IFN- $\alpha 5$ were provided by Daved Fremont (Washington University School of Medicine). Recombinant murine IFN- $\alpha 1$, - $\alpha 4$, - $\alpha 11$ and - $\alpha 13$ were provided by Anthony Coyle and Ricardo Cibotti (MedImmune, Inc., Gaithersburg, MD). Murine IFN- γ was obtained from Genentech (South San Francisco, CA) and IFN- $\alpha A/D$ was a gift from Hoffmann-LaRoche (Nutley, NJ). IFN- αA was purchased from PBL Assay Sciences (Piscataway, NJ). The following mAbs were produced by Leinco Technologies, Inc. (St. Louis, MO): MAR1-5A3 (neutralizing anti-murine IFNAR1) [67], GIR-208 (control murine IgG1) [68], H22 (neutralizing anti-murine IFN- γ) [69], PIP (control Armenian hamster mAb specific for bacterial glutathione-S-transferase) [70], and YTS-169 (control rat IgG). Control mouse IgG2a mAb (OKT3) was purchased from BioXCell (West Lebanon, NH). Type I IFN antibodies 7F-D3 (rat anti-IFN- β) and 4E-A1 (rat anti-IFN- α) were purchased from Abcam (Cambridge, MA). Secondary antibody reagents were purchased from Jackson ImmunoResearch (West Grove, PA).

Production of Type I IFN specific mAbs

Mouse mAbs specific for murine IFN- β were developed using *Ifnb*^{-/-} mice immunized by seven rounds of hydrodynamic injections of plasmid DNA encoding murine IFN- β at intervals of two weeks or greater using methods previously described [67]. Murine *Ifnb* cDNA was cloned from a 129SvMa cDNA library derived from lipopolysaccharide and polyinosinic:polycytidylic acid (pI:C) stimulated bone marrow macrophages and subcloned into *EcoRI* and *XbaI* sites of pEF4/myc-His A vector (Invitrogen, Grand Island, NY), which adds an in-frame 3' myc epitope and polyhistidine tag (5' sense primer: TAGATTTACCACATGAACAACAGGTGGATC; 3' antisense primer: TATCTAGAGTTTTGGAAGTTTCTGGT). The *Ifnb* sequence was excised from this vector using *EcoRI* and *PmeI* sites and ligated into pRK5 vector via *EcoRI* and *SmaI* sites and verified by Sanger sequencing. To produce IFN- β specific mAbs, immune splenocytes from genetically immunized *Ifnb*^{-/-} mice were fused to the murine P3X63Ag8.653 myeloma cell line according to published procedures [67]. Supernatants from growth-positive wells were screened by ELISA for binding using plate-bound recombinant murine IFN- β compared to murine IFN- γ or IFN- α , and then for their capacity to block IFN- β effects on target cells (see below). One hybridoma cell line with neutralizing activity was identified: HD β -4A7. This hybridoma produced a murine IgG2a mAb, which was purified by protein A affinity chromatography [67].

Antibodies reactive to IFN- α species were developed following immunization of Armenian hamsters with recombinant murine IFN- $\alpha 5$ alone, or with multiple IFN- α subtypes (IFN- $\alpha 1$, - $\alpha 4$, - $\alpha 5$, and - $\alpha 11$) in Freund's adjuvant using conventional techniques [71]. Hybridomas were identified by ELISA screening where reactivity to IFN- $\alpha 5$, murine IFN- $\alpha 1$, murine IFN- $\alpha 13$, and human IFN- $\alpha A/D$ but not to murine IFN- β was used as the selection criteria. TIF-3C5 and TIF-1D6 were purified by protein A affinity chromatography as described [71].

In vitro IFN activity measurement

Type I IFN activity was measured using three *in vitro* functional assays: (1) induction of STAT1 phosphorylation (P-STAT1); (2) upregulation of MHC class I (MHC-I) expression; and (3) antiviral activity. P-STAT1 was measured by flow cytometry of L929 cells after

incubation with mAbs and various doses of type I IFNs for 20 min at 37°C. Cells were fixed using 2% paraformaldehyde, permeabilized with 90% methanol and stained with a mAb to P-STAT1 (BD Phosflow anti-Stat1, BD Biosciences, San Jose, CA) for 1 h on ice. Enhancement of MHC-I expression was assessed by flow cytometry following treatment of a murine fibrosarcoma cell line (F510 or 1969) [72, 73] with different subtypes of IFN for 72 h in the absence or presence of mAbs as described [67]. IFN-induced antiviral effects were assessed using a cytopathic effect (CPE) assay with L929 cells and vesicular stomatitis virus (VSV, Indiana strain 5 x 10⁴ TCID₅₀) as previously described [67]. In each assay, mAb was titrated as indicated in the text or Figure legends. One unit of IFN activity is defined as the volume of serum containing half-maximal stimulating activity.

Pharmacokinetics

Groups of wild-type mice were injected i.p. with 0.25 mg HD β -4A7, TIF-3C5, or control hamster IgG (PIP) mAbs in 0.5 ml of PBS. Blood was collected from two to five mice on days 0 to 20 following mAb administration. TIF-3C5 and HD β -4A7 levels in the serum were determined by direct ELISA using plates coated with murine IFN- α 4 or IFN- β , respectively. Control Ig (PIP) was measured by indirect ELISA on anti-Armenian hamster Ig coated plates. Values were determined based on a standard curve of purified mAb diluted in normal mouse serum.

In vivo pI:C stimulation

Groups of two to five mice were injected with 0.1 mg pI:C (GE Healthcare Life Science, Piscataway, NJ) and blood was collected at multiple time points as described in the text or figure legends.

West Nile virus infection

The WNV-NY strain (3000.0259) was isolated in New York in 2000 and passaged once in C6/36 *Aedes albopictus* cells to generate a virus stock that was used in all experiments [74, 75]. WNV was diluted in Hank's Balanced Salt Solution supplemented with 1% heat-inactivated fetal bovine serum. Groups of eight to 12 week-old age and sex matched mice were inoculated by footpad injection with 10² plaque forming units (PFU) of WNV in a volume of 50 μ l. Mice were anesthetized with ketamine (~1.7 mg/mouse) and xylazine (~50 μ g/mouse) prior to footpad inoculation. Antibodies were delivered via intraperitoneal injection in PBS. GIR-208, PIP, or an IgG2a mAb (2H2) specific to dengue virus prM protein [76] were used as isotype controls for MARI-5A3, TIF-3C5 and HD β -4A7, respectively. Survival was monitored over 21 days. Mice were monitored daily for disease signs and were euthanized by controlled CO₂ administration when they exhibited severe morbidity, including hindlimb paralysis or non-responsiveness.

Serum analysis from WNV-infected mice

Wild-type, *Ifnar1*^{-/-}, *Ifnb*^{-/-} or *Irf7*^{-/-} mice were inoculated with WNV and serum was collected at one to six days following infection. Serum was treated at pH 3.0 for 15 min at 37°C, followed by neutralization in HEPES (final concentration 250 mM), prior to assay. This treatment inactivates WNV while preserving type I IFN activity [16].

Statistical analysis

For serum bioassay, differences were analyzed with the Mann-Whitney test. Kaplan-Meier survival curves were analyzed by the log rank test. All data were analyzed using Prism software (GraphPad, San Diego, CA).

Results and Discussion

Development of mAbs against specific type I IFNs

The goal of this project was to develop mAbs that neutralized the functional activity of IFN- β or all forms of IFN- α . We employed a multi-tiered screening strategy to identify functional, subtype-specific mAbs. Having previously developed a blocking mAb specific for IFNAR1 (MAR1-5A3) using hydrodynamic genetic immunization of *Ifnar1*^{-/-} mice [67], we used this same method to immunize *Ifnb*^{-/-} mice with plasmid DNA encoding full-length murine IFN- β tagged with myc and hexahistidine. This immunization method allows for the secretion of native, glycosylated IFN- β and presentation of this ‘foreign’ antigen in an *Ifnb*^{-/-} mouse with a largely intact immune system. Animals injected with plasmid DNA displayed serum titers greater than 1:30,000 as determined by ELISA using recombinant murine IFN- β . Immune sera did not detect recombinant murine IFN- α 1, - α 4, - α 5, - α 13 or IFN- γ and completely blocked IFN- β -induced STAT1 phosphorylation in L929 cells (data not shown). Fusion of immune splenocytes generated five stable hybridoma lines as assessed by selective binding to murine IFN- β by ELISA and lack of reactivity to murine IFN- γ and multiple IFN- α subtypes. Two of these IFN- β specific hybridomas secreted mAbs of the IgG2a isotype, one of which, HD β -4A7, exhibited neutralizing activity and was selected for further characterization. A second mAb, HD β -5F5, bound IFN- β specifically but did not block its function. The three other IFN- β -specific hybridomas secreted IgM and were not analyzed further.

We next sought to produce a ‘pan-IFN- α ’ mAb that would neutralize all 14 murine IFN- α subtypes (75–99% amino acid identity) [6, 9, 10] but not IFN- β (~30% identity) [77]. We could not use the same approach that generated the IFN- β mAbs because mice lacking the multi-gene IFN- α complex have not been reported. Instead, we immunized Armenian hamsters with several different combinations of murine IFN- α proteins including injection with a single IFN- α subtype (IFN- α 5) or with multiple IFN- α subtypes (IFN- α 1, - α 4, - α 5, and - α 11) either mixed together or following sequential immunizations. Each of the immunization strategies generated polyclonal neutralizing antisera with similar titers against IFN- α 5. Immune serum derived from the hamster immunized repeatedly with purified recombinant murine IFN- α 5 displayed the highest titer (>1:25,000), as assessed with an IFN- α 5 ELISA, and showed no binding to IFN- β . This immune serum also blocked IFN- α 5-induced STAT1 phosphorylation in L929 fibroblasts (data not shown). Following fusion, many hybridomas were identified that showed ELISA reactivity with multiple subtypes of recombinant murine IFN- α . However, only mAb derived from the TIF-3C5 hybridoma neutralized all of the IFN- α subtypes tested. In comparison, mAbs derived from other hybridomas bound or blocked different subsets of the IFN- α s used (S1 Table).

In vitro neutralization of type I IFN activity

Three biological assays were used to identify mAbs that neutralized specific type I IFNs: (1) induction of STAT1 phosphorylation; (2) upregulation of cell surface MHC-I expression; and (3) inhibition of CPE caused by VSV infection. We initially measured the ability of HD β -4A7 and TIF-3C5 to block IFN- β - or IFN- α -induced STAT1 phosphorylation, as this represents a rapid response following ligand binding to IFNAR1/IFNAR2 that occurs within minutes of receptor engagement. HD β -4A7 inhibited STAT1 phosphorylation in a dose-dependent manner, comparable to that achieved using an IFNAR1 blocking mAb, MAR1-5A3 (Fig 1A and 1B). A second IFN- β -specific IgG2a mAb, HD β -5F5, as well as several IFN- α -specific mAbs, failed to block STAT1 phosphorylation induced by IFN- β (data not shown). To examine inhibition of IFN- α activity *in vitro*, we tested STAT1 phosphorylation in response to IFN- α 4 stimulation.




Fig 1. Blockade of type I IFN activation of P-STAT1 *in vitro*. Titrations of HD β -4A7 and TIF-3C5 (0.01–10 μ g) or MAR1-5A3 (10 μ g) were preincubated for 60 min with 3 ng of IFN- β (A, B) or 3.3 ng of IFN- α 4 (C, D) and then incubated with L929 cells for 20 min, followed by staining for P-STAT1 and processing by flow cytometry. E. Individual IFN- α subtypes (3.3 ng) were preincubated with 10 μ g of mAb and then incubated with L929 cells for 20 min, followed by staining for P-STAT1 and flow cytometric analysis. Data are representative of three or more independent experiments.

doi:10.1371/journal.pone.0128636.g001

TIF-3C5 inhibited IFN- α 4-mediated phosphorylation in a dose-dependent manner, similar to that achieved using anti-IFNAR1 mAb, whereas incubation of IFN- α 4 with the anti-IFN- β mAb HD β -4A7 did not alter phospho-STAT1 levels (Fig 1C and 1D). TIF-3C5 also blocked induction of STAT1 phosphorylation in response to recombinant IFN- α A, IFN- α 1, IFN- α 5, IFN- α 11 (partial blockade), and IFN- α 13 (Fig 1E), with no inhibition of IFN- β (data not shown). Thus, TIF-3C5 exhibited neutralizing activity for all six of the recombinant IFN- α subtypes tested.




Fig 2. Inhibition of type I IFN MHC I upregulation. Type I IFNs (IFN- β , 800 pg; IFN- α 1, 40,000 pg; IFN- α 4, 4000 pg; IFN- α 5, 400 pg; IFN- α 11, 1000 pg; or IFN- α 13, 1000 pg) were preincubated with 1 to 10 μ g of HD β -4A7 (A, B) or 0.1 to 10 μ g of TIF-3C5 (C, D), MAR1-5A3 (10 μ g) or control hamster IgG (10 μ g) for 30 min then added to fibrosarcoma cells for 72 h and stained for H2-K^b class I MHC antigens by flow cytometry. Plots and histograms are representative of three or more independent assays using either F510 (A, B) or 1969 (C, D) fibrosarcoma cells.

doi:10.1371/journal.pone.0128636.g002

We next examined type I IFN induction of cell surface expression of MHC-I. We used the mouse sarcoma lines F510 or 1969 that constitutively express low levels of MHC class I, but upregulate cell surface expression of H-2K^b and H-2D^b following IFN treatment. HD β -4A7 efficiently blocked IFN- β -mediated H2-K^b expression whereas TIF-3C5 had no effect (Fig 2A and 2B). In parallel experiments, TIF-3C5 inhibited MHC-I upregulation in a dose dependent manner upon stimulation with IFN- α 4 (Fig 2C and 2D) or additional IFN- α subtypes (IFN- α A, - α 1, - α 5, - α 11 and - α 13), consistent with TIF-3C5 reactivity against all species of IFN- α tested. HD β -4A7 and TIF-3C5 blocked IFN- β and IFN- α activity, respectively, in a manner that was similar to IFNAR blockade by MAR1-5A3.

We also assessed the ability of HD β -4A7 and TIF-3C5 to neutralize type I IFN antiviral activity using a cytopathic effect bioassay. HD β -4A7 neutralized 10 U (16 pg) of IFN- β antiviral activity with a 50% inhibitory dose (ID₅₀) of 600 ng/well, compared to 30 ng/well for the IFNAR1-binding mAb, MAR1-5A3 (Fig 3A). As expected, TIF-3C5 did not inhibit IFN- β antiviral activity. Although TIF-3C5 completely blocked the antiviral activity of each IFN- α subtypes tested, it did so with differing efficiencies (Fig 3B). IFN- α 1 was blocked at low concentrations of mAb (ID₅₀ = 9 ng/well) whereas more TIF-3C5 was needed to inhibit IFN- α 4 (ID₅₀ = 2000 ng/well). It is unclear in these studies if this spectrum of activity reflects the unique binding affinity of TIF-3C5 for each IFN- α subtype, the specific activity of each of the IFN- α subtype used, or if particular IFN- α subtypes are more effective at triggering specific biological functions. Hence, we have identified two mAbs, HD β -4A7 and TIF-3C5, capable of selectively neutralizing the *in vitro* activities of recombinant IFN- β and IFN- α , respectively.

We compared the *in vitro* neutralizing activity of HD β -4A7 and TIF-3C5 to previously reported rat mAbs targeting IFN- β and IFN- α (7F-D3 and 4E-A1, respectively) [78]. HD β -4A7




Fig 3. Neutralization of type I IFN induced antiviral activity *in vitro*. Ten units IFN- β (A) or IFN- α (B) were preincubated for 1 h with indicated mAbs and then added to L929 cells overnight. Ten units/well type I IFN activity used in these assays are as follows: IFN- β , 16 pg; IFN- α A, 22 pg; IFN- α 1, 15 pg; IFN- α 4, 70 pg; IFN- α 5, 20 pg; IFN- α 13, 13 pg. IFN-treated cells subsequently were infected with VSV for 48 h and cell viability determined by crystal violet staining and optical density measurements. Results shown are representative of three or more independent experiments.

doi:10.1371/journal.pone.0128636.g003

exhibited less potent neutralizing activity than 7F-D3, and the relative potencies of TIF-3C5 and 4E-A1 depended on the subtype of IFN- α tested (Fig 4). While 7F-D3 and 4E-A1 exhibit neutralizing activity against murine IFN- β and IFN- α , anti-globin responses can effect mAb pharmacokinetics, making these existing rat IgG reagents problematic for repeated *in vivo* administration [79]. The new mAbs reported in this work are more amenable to *in vivo* use, as HD β -4A7 is of murine origin and TIF-3C5 is derived from Armenian hamster, a species whose IgG is not immunogenic in mice [70,78]

In vivo production of type I IFN

We next tested the ability of HD β -4A7 and TIF-3C5 to neutralize type I IFNs generated *in vivo* in response to inflammatory stimuli or infection. *Ifnar1*^{-/-}, *Ifnb*^{-/-} or *Irf7*^{-/-} mice were used to generate distinct mixtures of IFN compared to those generated using wild-type animals. *Ifnar1*^{-/-} mice produce high levels of circulating IFNs as they lack a receptor to bind and internalize type I IFNs. *Ifnb*^{-/-} mice do not produce IFN- β , but generate IFN- α and have normal responses to type I IFN [60, 61]. *Irf7*^{-/-} mice produce IFN- β , however IFN- α production is largely ablated [66, 80]. In the first set of experiments, mice were treated with the TLR3/MDA5 agonist pI:C to induce IFN production, and serum was collected as a source of naturally produced IFN. Antiviral activity in the serum was quantitated using the VSV CPE bioassay (Fig 5A–5D). Wild-type mice injected with pI:C produced peak type I IFN activity at 3 h post injection, with




Fig 4. *In vitro* neutralizing activity of Type I IFN mAbs. Approximately 20 units of recombinant IFN- β (A) or IFN- α 1 (B), IFN- α 4 (C) or IFN- α 13 (D) were preincubated for 1 h with indicated mAbs and then added to L929 cells overnight. Constant levels of Type I IFN used in these assays are as follows: IFN- β , 4.4 ng; IFN- α 1, 0.15 ng; IFN- α 4, 0.05 ng; IFN- α 13, 34 ng. IFN-treated cells subsequently were infected with VSV for 48 h and cell viability determined by crystal violet staining and optical density measurements. Results shown are representative of two independent experiments.

doi:10.1371/journal.pone.0128636.g004




Fig 5. Kinetics of native type I IFN production *in vivo*. Wild-type (A, E), *Ifnar1*^{-/-} (B, F), *Ifnb*^{-/-} (C, G) or *Irf7*^{-/-} (D, H) mice were treated with 100 μ g of pl:C (A-D) or infected with 10² PFU of WNV (E-H) and serum was assayed for IFN activity by VSV CPE bioassay. Samples from three independent sets of pl:C treated mice (3 to 5 per group) were collected at 1, 3 and 6 h post injection. Serum from groups of 3 to 8 mice of each genotype was collected from naïve and WNV infected mice at days 1 to 6 post infection. Each genotype was sampled from at least two (*Ifnar1*^{-/-} and *Irf7*^{-/-}) or more (wild-type and *Ifnb*^{-/-}) independent infections. IFN levels in all samples were determined at least three times. As shown herein, symbols represent samples derived from individual animals at the times indicated and assayed at the same time.

doi:10.1371/journal.pone.0128636.g005

levels declining rapidly by 12 h (data not shown). The kinetics of production were identical for each of the gene-targeted strains examined with peak levels similar to those described using pl:C treated wild-type or reporter mice [81, 82]. *Ifnb*^{-/-} mice produced slightly elevated peak levels of type I IFN, whereas *Irf7*^{-/-} animals generated lower levels overall.

We infected mice with WNV, an encephalitic flavivirus, and collected sera one to six days after infection. Again, we quantitated serum IFN activity using the VSV CPE bioassay (Fig 5E–5H). Serum levels of IFN- α and IFN- β rose rapidly in *Ifnar1*^{-/-} mice, which sustain extremely high levels of viral replication and viral nucleic acids, further driving IFN production [16]. As previously observed, serum antiviral activity in *Ifnb*^{-/-} mice was similar to that of wild-type mice, implying that a substantial component of the antiviral activity in wild-type serum comes from IFN- α [61]. As expected, only minimal antiviral activity was detected in the serum from *Irf7*^{-/-} mice [80].

Neutralization of *in vivo* generated IFN- β and IFN- α

We used our neutralizing mAbs to characterize the composition of natural IFN produced by wild-type, *Ifnar1*^{-/-}, *Ifnb*^{-/-} or *Irf7*^{-/-} mice in response to pl:C treatment (Fig 6) or WNV infection (Fig 7). IFN-containing sera were incubated with control or neutralizing mAbs and the antiviral activity measured by the VSV CPE bioassay. All of the antiviral activity present in the serum samples was attributable to type I IFN because it was abrogated by addition of the IFNAR1-blocking mAb MAR1-5A3 (Figs 6 and 7, panels F, L, R and X). Consistent with this observation, treatment with an IFN- γ neutralizing mAb, H22, did not affect the antiviral activity (Figs 6 and 7, panels B, H, N and T), which was identical to cultures treated with buffer or control IgG. In all cases, the combination of both HD β -4A7 and TIF-3C5 ablated all antiviral




Fig 6. Detection of IFN- β and IFN- α in serum following pI:C treatment. Serum was collected from wild-type (A-F), *Ifnar1*^{-/-} (G-L), *Ifnb*^{-/-} (M-R) or *Irf7*^{-/-} (S-X) mice injected with 100 μ g of pI:C at 3 to 6 hrs. Volumes of serum containing approximately 5–10 units of IFN activity (as determined in Fig 4) were incubated with 10 μ g of mAb for 1 h and then titrations were added to L929 cells overnight. IFN-treated cells were infected with VSV and cellular viability assessed after 48 h using crystal violet staining and optical density measurements. Each panel is representative of six independent samples.

doi:10.1371/journal.pone.0128636.g006

activity in the bioassay (Figs 6 and 7, panels E, K, Q and W), similar to that seen with MAR1-5A3, revealing that other type I IFNs (e.g., IFN- κ or IFN- ζ) or type III IFNs (IFN- λ) did not contribute to antiviral activity under these conditions. In sera from wild-type mice stimulated with pI:C for 6 h, the type I IFN activity was blocked completely by TIF-3C5 with no effect of neutralization with HD β -4A7, indicating that IFN- α was the only type I IFN produced at this




Fig 7. Detection of IFN- β and IFN- α in serum following WNV infection. Serum was collected from wild-type (A-F), *Ifnar1*^{-/-} (G-L), *Ifnb*^{-/-} (M-R) or *Irf7*^{-/-} (S-X) mice infected with 10² PFU of WNV at day 3 (wild-type and *Ifnar1*^{-/-}) or day 4 (*Ifnb*^{-/-} and *Irf7*^{-/-}). Volumes of serum containing approximately 5–10 units of IFN activity (as determined in Fig 4) were incubated with 10 μ g of mAb for 1 h and then titrations were added to L929 cells overnight. IFN-treated cells were infected with VSV and cellular viability assessed after 48 h using crystal violet staining and optical density measurements. Each panel is representative of six independent samples.

doi:10.1371/journal.pone.0128636.g007

time point (Fig 6C and 6D). Identical results were obtained using serum samples obtained at 3 h post injection of pI:C (data not shown). Analogously, in serum obtained 3 days after WNV infection of wild-type mice, TIF-3C5 fully blocked antiviral activity, with HD β -4A7 having virtually no impact, indicating that IFN- α is the dominant type I IFN in serum following WNV

infection at that time point (Fig 7C and 7D). Neither TIF-3C5 nor HD β -4A7 alone could completely neutralize the antiviral activity present in serum from *Ifnar1*^{-/-} mice following either pI:C treatment or WNV challenge, demonstrating that both IFN- β and IFN- α are produced in these immunodeficient mice in response to these stimuli although IFN- α predominates (Figs 6 and 7, panels I and J). As expected, HD β -4A7 had no impact on the antiviral activity found in serum from *Ifnb*^{-/-} mice and all of the IFN activity in these samples was neutralized by TIF-3C5 (Figs 6 and 7, panels O and P). With serum from pI:C-treated *Irf7*^{-/-} mice, HD β -4A7 neutralized antiviral activity whereas TIF-3C5 had no effect (Fig 6V and 6U). In contrast, TIF-3C5 neutralized all of the type I IFN activity from WNV-infected *Irf7*^{-/-} mice, with no neutralization by HD β -4A7 (Fig 7U and 7V). Thus, IFN- α was generated in *Irf7*^{-/-} mice in response to WNV infection but not after pI:C treatment. This IFN- α activity may be due to the IFN- α 4 subtype, which is an unusual IFN- α that can be induced directly in response to pattern recognition receptor signaling and does not require IFN- β production or IRF-7-dependent transcriptional activity for its expression [83]. Although *Irf7*^{-/-} mice would be expected to induce some IFN- β in response to WNV infection, this low level of IFN- β may be induced at early time points and cleared from the circulation by IFNAR1 or fall below the sensitivity of this assay. Collectively, these data establish that HD β -4A7 and TIF-3C5 can bind and neutralize naturally-produced IFN- β and IFN- α , respectively, and provide a sensitive tool to interrogate the appearance and quantity of either IFN in complex biological samples such as serum.

Type I IFN mAbs enhance lethality of WNV infection

Having identified mAbs that selectively neutralize IFN- β and IFN- α *in vitro*, we assessed whether they could neutralize IFN activity *in vivo*. Pharmacokinetic analysis revealed circulatory half-lives of 14 days for HD β -4A7 and 3.5 days for TIF-3C5 as determined by ELISA using IFN- β and IFN- α 5, respectively, and 4 days for PIP control Armenian hamster IgG (data not shown). To evaluate the ability of HD β -4A7 and TIF-3C5 to neutralize IFNs *in vivo*, we used a well-characterized model of WNV infection in which type I IFNs control viral replication and pathogenesis [61, 84, 85].

In vivo mAb efficacy was assessed by monitoring survival following WNV infection. We observed increased lethality in wild-type mice treated with 500 μ g of TIF-3C5 one day prior and two days following WNV infection, compared to mice receiving isotype control IgG (Fig 8A), but this difference did not achieve statistical significance ($P > 0.05$). We tried an alternate dosing regimen and found that three administrations of TIF-3C5 (250 μ g one day prior, one day following, and three days following WNV infection) produced a greater and statistically significant increase in lethality compared to two doses of 500 μ g (Fig 8B). TIF-3C5 treatment enhanced lethality compared to isotype control mAb in both wild-type and *Ifnb*^{-/-} mice, indicating a key antiviral role for IFN- α in the presence or absence of IFN- β [86]. Wild-type mice treated with 250 μ g of HD β -4A7 one day prior and two days following WNV infection succumbed with kinetics identical to *Ifnb*^{-/-} mice, indicating that HD β -4A7 completely neutralizes IFN- β activity *in vivo* (Fig 8C). Combined administration of both HD β -4A7 and TIF-3C5 in wild-type mice, or TIF-3C5 treatment in *Ifnb*^{-/-} mice, phenocopied treatment with MARI-5A3 (Fig 8D). Since IFN- β and IFN- α fully accounted for the type I IFN response observed, these data suggest that other type I IFN subtypes do not contribute significantly to the antiviral response to WNV in this system. TIF-3C5 treatment produced a small increase lethality in *Irf7*^{-/-} mice (Fig 8A), consistent with the ability of this mAb to neutralize antiviral activity in serum from WNV-infected *Irf7*^{-/-} mice (Fig 7U). We conclude that while some residual IFN- α is induced in the absence of IRF-7, it is insufficient to support an antiviral response, perhaps because IRF-7 is necessary for the expression of antiviral effector molecules in response to




Fig 8. Type I IFN blocking mAbs enhance lethality of WNV infection. Wild-type or *Irf7*^{-/-} mice were administered 500 μ g of TIF-3C5 or isotype control mAb by intraperitoneal injection one day prior, and two days following subcutaneous infection with 10² PFU of WNV. **B.** Wild-type or *Ifnb*^{-/-} mice received 250 μ g of TIF-3C5 or isotype control mAb one day prior, one day following, and three days following WNV infection. **C.** Wild-type or *Ifnb*^{-/-} mice received 250 μ g of HD β -4A7 or isotype control antibody one day prior and two days following WNV infection. **D.** Wild-type or *Ifnb*^{-/-} mice received 250 μ g of HD β -4A7 antibody or isotype control one day prior and three days following WNV infection, as well as 500 μ g of TIF-3C5 antibody (or control) one day prior, one day following, and three days following infection. Wild-type and *Ifnar*^{-/-} mice were given 500 μ g of MAR1-5A3 or isotype control mAb one day prior, one day following, and three days following infection. Survival was monitored for 21 days. Data represent 9 to 20 mice per group, from 2 or more independent experiments. *, $P < 0.05$; ***, $P < 0.001$; ****, $P < 0.0001$ (log-rank test).

doi:10.1371/journal.pone.0128636.g008

IFN- α signaling. We presume that the IRF-7-independent IFN- α produced is IFN- α 4. Together, our observations provide new insights into the specific roles of IFN- β and IFN- α in the response to WNV infection. Specifically, we detected very little IFN- β in serum of WNV infected mice (days 2 to 6 after infection) and found that the type I IFN response was dominated by IFN- α ; it is likely that small quantities of IFN- β are sufficient to initiate positive feedback amplification driving IFN- α production. Since TIF-3C5 neutralizes all murine IFN- α subtypes tested, the present studies do not distinguish unique functions for different IFN- α subtypes *in vivo*; neutralizing mAbs targeting individual or defined sets of IFN- α subtypes will be needed to determine their specific properties. Although our observations define the specific contributions of IFN- α versus IFN- β in serum, other IFN subtypes may act in a localized manner in particular cells or tissues. Furthermore, the relative contributions of IFN subtypes to the antiviral response for different infections may depend on the tropism and evasion strategies of the virus studied.

In summary, we generated two new mAbs, HD β -4A7 and TIF-3C5, that are highly specific for murine IFN- β and most, if not all, IFN- α subtypes, respectively, and can block type I IFN activity *in vitro* and *in vivo*. We used these novel reagents to distinguish specific contributions of IFN- β and IFN- α to the antiviral response to WNV infection. Our observations reveal a complex interplay between IFN production and host survival from infection. These unique reagents provided us with an ability to monitor and/or selectively ablate the functional activity of individual type I IFN subtypes in complex models of immune and inflammatory stimulation.

Unlike studies in gene-targeted animals, these reagents will permit modulation of IFN activity at particular time points relative to infectious or inflammatory stimuli in the context of an immune system that has developed in a normal IFN milieu. These mAbs will be valuable tools to interrogate the specific functional roles of IFN- β and IFN- α in inflammatory, infectious, or autoimmune models of disease.

Supporting Information

S1 ARRIVE Checklist.

(PDF)

S1 Table. ELISA Binding Specificity of Type I IFN Antibodies. Binding of mAbs to adsorbed recombinant IFNs was measured by ELISA.

(DOCX)

Acknowledgments

The authors thank Daved Fremont and Chung Lee for providing recombinant IFN- α 5 and IFN- β and are grateful to Anthony Coyle and Richard Cibotti for the gift of recombinant IFN- α 1, - α 4, - α 11, and - α 13. The IFN- β plasmid DNA was constructed by Mark Diamond. We thank Marina Cella for reagents and helpful discussions.

Author Contributions

Conceived and designed the experiments: KCFS HML MSD RDS. Performed the experiments: KCFS HML. Analyzed the data: KCFS HML MSD RDS. Contributed reagents/materials/analysis tools: KCFS HML MSD RDS. Wrote the paper: KCFS HML MSD RDS.

References

1. Isaacs A, Lindenmann J. Virus interference. I. The interferon. *Proc R Soc Lond B Biol Sci.* 1957; 147 (927):258–67. Epub 1957/09/12. PMID: [13465720](#).
2. Nagano Y, Kojima Y. [Inhibition of vaccinia infection by a liquid factor in tissues infected by homologous virus]. *Comptes rendus des seances de la Societe de biologie et de ses filiales.* 1958; 152(11):1627–9. Epub 1958/01/01. PMID: [13639454](#).
3. Gresser I, Tovey MG, Maury C, Bandu MT. Role of interferon in the pathogenesis of virus diseases in mice as demonstrated by the use of anti-interferon serum. II. Studies with herpes simplex, Moloney sarcoma, vesicular stomatitis, Newcastle disease, and influenza viruses. *The Journal of experimental medicine.* 1976; 144(5):1316–23. Epub 1976/11/02. PMID: [186555](#); PubMed Central PMCID: PMC2190450.
4. Oritani K, Medina KL, Tomiyama Y, Ishikawa J, Okajima Y, Ogawa M, et al. Limitin: An interferon-like cytokine that preferentially influences B-lymphocyte precursors. *Nature medicine.* 2000; 6(6):659–66. Epub 2000/06/03. doi: [10.1038/76233](#) PMID: [10835682](#).
5. Fung KY, Mangan NE, Cumming H, Horvat JC, Mayall JR, Stifter SA, et al. Interferon-epsilon protects the female reproductive tract from viral and bacterial infection. *Science.* 2013; 339(6123):1088–92. Epub 2013/03/02. doi: [10.1126/science.1233321](#) PMID: [23449591](#); PubMed Central PMCID: PMC3617553.
6. Hardy MP, Owczarek CM, Jermini LS, Ejdeback M, Hertzog PJ. Characterization of the type I interferon locus and identification of novel genes. *Genomics.* 2004; 84(2):331–45. Epub 2004/07/06. doi: [10.1016/j.ygeno.2004.03.003](#) PMID: [15233997](#).
7. LaFleur DW, Nardelli B, Tsareva T, Mather D, Feng P, Semenuk M, et al. Interferon-kappa, a novel type I interferon expressed in human keratinocytes. *The Journal of biological chemistry.* 2001; 276 (43):39765–71. Epub 2001/08/22. doi: [10.1074/jbc.M102502200](#) PMID: [11514542](#).
8. Buontempo PJ, Jubin RG, Buontempo CA, Wagner NE, Reyes GR, Baroudy BM. Antiviral activity of transiently expressed IFN-kappa is cell-associated. *Journal of interferon & cytokine research: the official journal of the International Society for Interferon and Cytokine Research.* 2006; 26(1):40–52. Epub 2006/01/24. doi: [10.1089/jir.2006.26.40](#) PMID: [16426147](#).

9. van Pesch V, Lanaya H, Renauld JC, Michiels T. Characterization of the murine alpha interferon gene family. *Journal of virology*. 2004; 78(15):8219–28. Epub 2004/07/16. doi: [10.1128/JVI.78.15.8219-8228.2004](https://doi.org/10.1128/JVI.78.15.8219-8228.2004) PMID: [15254193](https://pubmed.ncbi.nlm.nih.gov/15254193/); PubMed Central PMCID: PMC446145.
10. Zwarthoff EC, Mooren AT, Trapman J. Organization, structure and expression of murine interferon alpha genes. *Nucleic acids research*. 1985; 13(3):791–804. Epub 1985/02/11. PMID: [2987810](https://pubmed.ncbi.nlm.nih.gov/2987810/); PubMed Central PMCID: PMC341035.
11. Khabar KS, Young HA. Post-transcriptional control of the interferon system. *Biochimie*. 2007; 89(6–7):761–9. Epub 2007/04/06. doi: [10.1016/j.biochi.2007.02.008](https://doi.org/10.1016/j.biochi.2007.02.008) PMID: [17408842](https://pubmed.ncbi.nlm.nih.gov/17408842/); PubMed Central PMCID: PMC1994070.
12. Donlin LT, Jayatilleke A, Giannopoulou EG, Kalliolias GD, Ivashkiv LB. Modulation of TNF-Induced Macrophage Polarization by Synovial Fibroblasts. *J Immunol*. 2014; 193(5):2373–83. Epub 2014/07/25. doi: [10.4049/jimmunol.1400486](https://doi.org/10.4049/jimmunol.1400486) PMID: [25057003](https://pubmed.ncbi.nlm.nih.gov/25057003/); PubMed Central PMCID: PMC4135020.
13. Akira S, Takeda K. Toll-like receptor signalling. *Nature reviews Immunology*. 2004; 4(7):499–511. Epub 2004/07/02. doi: [10.1038/nri1391](https://doi.org/10.1038/nri1391) PMID: [15229469](https://pubmed.ncbi.nlm.nih.gov/15229469/).
14. Ivashkiv LB, Donlin LT. Regulation of type I interferon responses. *Nature reviews Immunology*. 2014; 14(1):36–49. Epub 2013/12/24. doi: [10.1038/nri3581](https://doi.org/10.1038/nri3581) PMID: [24362405](https://pubmed.ncbi.nlm.nih.gov/24362405/); PubMed Central PMCID: PMC4084561.
15. Tamura T, Yanai H, Savitsky D, Taniguchi T. The IRF family transcription factors in immunity and oncogenesis. *Annual review of immunology*. 2008; 26:535–84. Epub 2008/02/29. doi: [10.1146/annurev.immunol.26.021607.090400](https://doi.org/10.1146/annurev.immunol.26.021607.090400) PMID: [18303999](https://pubmed.ncbi.nlm.nih.gov/18303999/).
16. Lazear HM, Lancaster A, Wilkins C, Suthar MS, Huang A, Vick SC, et al. IRF-3, IRF-5, and IRF-7 coordinately regulate the type I IFN response in myeloid dendritic cells downstream of MAVS signaling. *PLoS pathogens*. 2013; 9(1):e1003118. Epub 2013/01/10. doi: [10.1371/journal.ppat.1003118](https://doi.org/10.1371/journal.ppat.1003118) PMID: [23300459](https://pubmed.ncbi.nlm.nih.gov/23300459/); PubMed Central PMCID: PMC3536698.
17. Prakash A, Smith E, Lee CK, Levy DE. Tissue-specific positive feedback requirements for production of type I interferon following virus infection. *The Journal of biological chemistry*. 2005; 280(19):18651–7. Epub 2005/03/16. doi: [10.1074/jbc.M501289200](https://doi.org/10.1074/jbc.M501289200) PMID: [15767254](https://pubmed.ncbi.nlm.nih.gov/15767254/); PubMed Central PMCID: PMC1167601.
18. Barchet W, Cella M, Odermatt B, Asselin-Paturel C, Colonna M, Kalinke U. Virus-induced interferon alpha production by a dendritic cell subset in the absence of feedback signaling in vivo. *The Journal of experimental medicine*. 2002; 195(4):507–16. Epub 2002/02/21. PMID: [11854363](https://pubmed.ncbi.nlm.nih.gov/11854363/); PubMed Central PMCID: PMC2193622.
19. Uze G, Schreiber G, Piehler J, Pellegrini S. The receptor of the type I interferon family. *Current topics in microbiology and immunology*. 2007; 316:71–95. Epub 2007/11/01. PMID: [17969444](https://pubmed.ncbi.nlm.nih.gov/17969444/).
20. Pestka S, Krause CD, Walter MR. Interferons, interferon-like cytokines, and their receptors. *Immunological reviews*. 2004; 202:8–32. Epub 2004/11/18. doi: [10.1111/j.0105-2896.2004.00204.x](https://doi.org/10.1111/j.0105-2896.2004.00204.x) PMID: [15546383](https://pubmed.ncbi.nlm.nih.gov/15546383/).
21. de Weerd NA, Samarajiwa SA, Hertzog PJ. Type I interferon receptors: biochemistry and biological functions. *The Journal of biological chemistry*. 2007; 282(28):20053–7. Epub 2007/05/16. doi: [10.1074/jbc.R700006200](https://doi.org/10.1074/jbc.R700006200) PMID: [17502368](https://pubmed.ncbi.nlm.nih.gov/17502368/).
22. de Weerd NA, Nguyen T. The interferons and their receptors—distribution and regulation. *Immunology and cell biology*. 2012; 90(5):483–91. Epub 2012/03/14. doi: [10.1038/icb.2012.9](https://doi.org/10.1038/icb.2012.9) PMID: [22410872](https://pubmed.ncbi.nlm.nih.gov/22410872/).
23. Diamond MS, Kinder M, Matsushita H, Mashayekhi M, Dunn GP, Archambault JM, et al. Type I interferon is selectively required by dendritic cells for immune rejection of tumors. *The Journal of experimental medicine*. 2011; 208(10):1989–2003. Epub 2011/09/21. doi: [10.1084/jem.20101158](https://doi.org/10.1084/jem.20101158) PMID: [21930769](https://pubmed.ncbi.nlm.nih.gov/21930769/); PubMed Central PMCID: PMC3182061.
24. Fuertes MB, Woo SR, Burnett B, Fu YX, Gajewski TF. Type I interferon response and innate immune sensing of cancer. *Trends in immunology*. 2013; 34(2):67–73. Epub 2012/11/06. doi: [10.1016/j.it.2012.10.004](https://doi.org/10.1016/j.it.2012.10.004) PMID: [23122052](https://pubmed.ncbi.nlm.nih.gov/23122052/); PubMed Central PMCID: PMC3565059.
25. Clemens MJ, Elia A. The double-stranded RNA-dependent protein kinase PKR: structure and function. *Journal of interferon & cytokine research: the official journal of the International Society for Interferon and Cytokine Research*. 1997; 17(9):503–24. Epub 1997/10/23. PMID: [9335428](https://pubmed.ncbi.nlm.nih.gov/9335428/).
26. Stark GR, Kerr IM, Williams BR, Silverman RH, Schreiber RD. How cells respond to interferons. *Annual review of biochemistry*. 1998; 67:227–64. Epub 1998/10/06. doi: [10.1146/annurev.biochem.67.1.227](https://doi.org/10.1146/annurev.biochem.67.1.227) PMID: [9759489](https://pubmed.ncbi.nlm.nih.gov/9759489/).
27. Trinchieri G. Type I interferon: friend or foe? *The Journal of experimental medicine*. 2010; 207(10):2053–63. Epub 2010/09/15. doi: [10.1084/jem.20101664](https://doi.org/10.1084/jem.20101664) PMID: [20837696](https://pubmed.ncbi.nlm.nih.gov/20837696/); PubMed Central PMCID: PMC2947062.

28. Salazar-Mather TP, Ishikawa R, Biron CA. NK cell trafficking and cytokine expression in splenic compartments after IFN induction and viral infection. *J Immunol*. 1996; 157(7):3054–64. Epub 1996/10/01. PMID: [8816415](#).
29. Le Bon A, Etchart N, Rossmann C, Ashton M, Hou S, Gewert D, et al. Cross-priming of CD8+ T cells stimulated by virus-induced type I interferon. *Nature immunology*. 2003; 4(10):1009–15. Epub 2003/09/23. doi: [10.1038/ni978](#) PMID: [14502286](#).
30. Le Bon A, Durand V, Kamphuis E, Thompson C, Bulfone-Paus S, Rossmann C, et al. Direct stimulation of T cells by type I IFN enhances the CD8+ T cell response during cross-priming. *J Immunol*. 2006; 176(8):4682–9. Epub 2006/04/06. PMID: [16585561](#).
31. Le Bon A, Schiavoni G, D'Agostino G, Gresser I, Belardelli F, Tough DF. Type I interferons potently enhance humoral immunity and can promote isotype switching by stimulating dendritic cells in vivo. *Immunity*. 2001; 14(4):461–70. Epub 2001/05/05. PMID: [11336691](#).
32. Le Bon A, Thompson C, Kamphuis E, Durand V, Rossmann C, Kalinke U, et al. Cutting edge: enhancement of antibody responses through direct stimulation of B and T cells by type I IFN. *J Immunol*. 2006; 176(4):2074–8. Epub 2006/02/04. PMID: [16455962](#).
33. Hertzog PJ, Williams BR. Fine tuning type I interferon responses. *Cytokine & growth factor reviews*. 2013; 24(3):217–25. Epub 2013/05/29. doi: [10.1016/j.cytogfr.2013.04.002](#) PMID: [23711406](#).
34. Schreiber G, Piehler J. The molecular basis for functional plasticity in type I interferon signaling. *Trends in immunology*. 2015; 36(3):139–49. Epub 2015/02/18. doi: [10.1016/j.it.2015.01.002](#) PMID: [25687684](#).
35. Schoggins JW, Wilson SJ, Panis M, Murphy MY, Jones CT, Bieniasz P, et al. A diverse range of gene products are effectors of the type I interferon antiviral response. *Nature*. 2011; 472(7344):481–5. Epub 2011/04/12. doi: [10.1038/nature09907](#) PMID: [21478870](#); PubMed Central PMCID: PMC3409588.
36. Schoggins JW, MacDuff DA, Imanaka N, Gainey MD, Shrestha B, Eitson JL, et al. Pan-viral specificity of IFN-induced genes reveals new roles for cGAS in innate immunity. *Nature*. 2014; 505(7485):691–5. Epub 2013/11/29. doi: [10.1038/nature12862](#) PMID: [24284630](#); PubMed Central PMCID: PMC4077721.
37. Pascual V, Chaussabel D, Banchereau J. A genomic approach to human autoimmune diseases. *Annual review of immunology*. 2010; 28:535–71. Epub 2010/03/03. doi: [10.1146/annurev-immunol-030409-101221](#) PMID: [20192809](#); PubMed Central PMCID: PMC2847838.
38. Boasso A, Hardy AW, Anderson SA, Dolan MJ, Shearer GM. HIV-induced type I interferon and tryptophan catabolism drive T cell dysfunction despite phenotypic activation. *PLoS one*. 2008; 3(8):e2961. Epub 2008/08/14. doi: [10.1371/journal.pone.0002961](#) PMID: [18698365](#); PubMed Central PMCID: PMC2491901.
39. Henry T, Kirimanjeswara GS, Ruby T, Jones JW, Peng K, Perret M, et al. Type I IFN signaling constrains IL-17A/F secretion by gammadelta T cells during bacterial infections. *J Immunol*. 2010; 184(7):3755–67. Epub 2010/02/24. doi: [10.4049/jimmunol.0902065](#) PMID: [20176744](#); PubMed Central PMCID: PMC2879132.
40. Herbeuval JP, Shearer GM. HIV-1 immunopathogenesis: how good interferon turns bad. *Clin Immunol*. 2007; 123(2):121–8. Epub 2006/11/23. doi: [10.1016/j.clim.2006.09.016](#) PMID: [17112786](#); PubMed Central PMCID: PMC1930161.
41. Boasso A, Royle CM, Doumazos S, Aquino VN, Biasin M, Piacentini L, et al. Overactivation of plasmacytoid dendritic cells inhibits antiviral T-cell responses: a model for HIV immunopathogenesis. *Blood*. 2011; 118(19):5152–62. Epub 2011/09/21. doi: [10.1182/blood-2011-03-344218](#) PMID: [21931112](#); PubMed Central PMCID: PMC3217402.
42. Swiecki M, Wang Y, Vermi W, Gilfillan S, Schreiber RD, Colonna M. Type I interferon negatively controls plasmacytoid dendritic cell numbers in vivo. *The Journal of experimental medicine*. 2011; 208(12):2367–74. Epub 2011/11/16. doi: [10.1084/jem.20110654](#) PMID: [22084408](#); PubMed Central PMCID: PMC3256963.
43. Carrero JA, Calderon B, Unanue ER. Type I interferon sensitizes lymphocytes to apoptosis and reduces resistance to *Listeria* infection. *The Journal of experimental medicine*. 2004; 200(4):535–40. Epub 2004/08/11. doi: [10.1084/jem.20040769](#) PMID: [15302900](#); PubMed Central PMCID: PMC2211931.
44. Carrero JA. Confounding roles for type I interferons during bacterial and viral pathogenesis. *International immunology*. 2013; 25(12):663–9. Epub 2013/10/26. doi: [10.1093/intimm/dxt050](#) PMID: [24158954](#); PubMed Central PMCID: PMC3839063.
45. Miller JC, Ma Y, Bian J, Sheehan KC, Zachary JF, Weis JH, et al. A critical role for type I IFN in arthritis development following *Borrelia burgdorferi* infection of mice. *J Immunol*. 2008; 181(12):8492–503. Epub 2008/12/04. PMID: [19050267](#); PubMed Central PMCID: PMC3024833.
46. Biggioggero M, Gabbriellini L, Meroni PL. Type I interferon therapy and its role in autoimmunity. *Autoimmunity*. 2010; 43(3):248–54. Epub 2010/03/20. doi: [10.3109/08916930903510971](#) PMID: [20298125](#).

47. Rizza P, Moretti F, Belardelli F. Recent advances on the immunomodulatory effects of IFN-alpha: implications for cancer immunotherapy and autoimmunity. *Autoimmunity*. 2010; 43(3):204–9. Epub 2010/03/02. doi: [10.3109/08916930903510880](https://doi.org/10.3109/08916930903510880) PMID: [20187707](https://pubmed.ncbi.nlm.nih.gov/20187707/).
48. Nestle FO, Conrad C, Tun-Kyi A, Homey B, Gombert M, Boyman O, et al. Plasmacytoid dendritic cells initiate psoriasis through interferon-alpha production. *The Journal of experimental medicine*. 2005; 202(1):135–43. Epub 2005/07/07. doi: [10.1084/jem.20050500](https://doi.org/10.1084/jem.20050500) PMID: [15998792](https://pubmed.ncbi.nlm.nih.gov/15998792/); PubMed Central PMCID: PMC2212894.
49. Theofilopoulos AN, Baccala R, Beutler B, Kono DH. Type I interferons (alpha/beta) in immunity and autoimmunity. *Annual review of immunology*. 2005; 23:307–36. Epub 2005/03/18. doi: [10.1146/annurev.immunol.23.021704.115843](https://doi.org/10.1146/annurev.immunol.23.021704.115843) PMID: [15771573](https://pubmed.ncbi.nlm.nih.gov/15771573/).
50. Axtell RC, Raman C, Steinman L. Type I interferons: beneficial in Th1 and detrimental in Th17 autoimmunity. *Clinical reviews in allergy & immunology*. 2013; 44(2):114–20. Epub 2012/01/11. doi: [10.1007/s12016-011-8296-5](https://doi.org/10.1007/s12016-011-8296-5) PMID: [22231516](https://pubmed.ncbi.nlm.nih.gov/22231516/).
51. Trinchieri G, Santoli D, Knowles BB. Tumour cell lines induce interferon in human lymphocytes. *Nature*. 1977; 270(5638):611–3. Epub 1977/12/15. PMID: [593387](https://pubmed.ncbi.nlm.nih.gov/593387/).
52. Dunn GP, Bruce AT, Sheehan KC, Shankaran V, Uppaluri R, Bui JD, et al. A critical function for type I interferons in cancer immunoediting. *Nature immunology*. 2005; 6(7):722–9. Epub 2005/06/14. doi: [10.1038/ni1213](https://doi.org/10.1038/ni1213) PMID: [15951814](https://pubmed.ncbi.nlm.nih.gov/15951814/).
53. Swann JB, Hayakawa Y, Zerafa N, Sheehan KC, Scott B, Schreiber RD, et al. Type I IFN contributes to NK cell homeostasis, activation, and antitumor function. *J Immunol*. 2007; 178(12):7540–9. Epub 2007/06/06. PMID: [17548588](https://pubmed.ncbi.nlm.nih.gov/17548588/).
54. Jaks E, Gavutis M, Uze G, Martal J, Piehler J. Differential receptor subunit affinities of type I interferons govern differential signal activation. *Journal of molecular biology*. 2007; 366(2):525–39. Epub 2006/12/19. doi: [10.1016/j.jmb.2006.11.053](https://doi.org/10.1016/j.jmb.2006.11.053) PMID: [17174979](https://pubmed.ncbi.nlm.nih.gov/17174979/).
55. Hwang SY, Hertzog PJ, Holland KA, Sumarsono SH, Tymms MJ, Hamilton JA, et al. A null mutation in the gene encoding a type I interferon receptor component eliminates antiproliferative and antiviral responses to interferons alpha and beta and alters macrophage responses. *Proceedings of the National Academy of Sciences of the United States of America*. 1995; 92(24):11284–8. Epub 1995/11/21. PMID: [7479980](https://pubmed.ncbi.nlm.nih.gov/7479980/); PubMed Central PMCID: PMC40616.
56. Mueller C, Kornilova L, Wiest G, Steinhoff N. Psychophysical studies of visuo-vestibular interaction in microgravity. *Acta astronautica*. 1994; 33:9–13. Epub 1994/07/01. PMID: [11539543](https://pubmed.ncbi.nlm.nih.gov/11539543/).
57. Kamphuis E, Junt T, Waibler Z, Forster R, Kalinke U. Type I interferons directly regulate lymphocyte recirculation and cause transient blood lymphopenia. *Blood*. 2006; 108(10):3253–61. Epub 2006/07/27. doi: [10.1182/blood-2006-06-027599](https://doi.org/10.1182/blood-2006-06-027599) PMID: [16868248](https://pubmed.ncbi.nlm.nih.gov/16868248/).
58. Fenner JE, Starr R, Cornish AL, Zhang JG, Metcalf D, Schreiber RD, et al. Suppressor of cytokine signaling 1 regulates the immune response to infection by a unique inhibition of type I interferon activity. *Nature immunology*. 2006; 7(1):33–9. Epub 2005/11/29. doi: [10.1038/ni1287](https://doi.org/10.1038/ni1287) PMID: [16311601](https://pubmed.ncbi.nlm.nih.gov/16311601/).
59. Kawai T, Akira S. Innate immune recognition of viral infection. *Nature immunology*. 2006; 7(2):131–7. Epub 2006/01/21. doi: [10.1038/ni1303](https://doi.org/10.1038/ni1303) PMID: [16424890](https://pubmed.ncbi.nlm.nih.gov/16424890/).
60. Takaoka A, Mitani Y, Suemori H, Sato M, Yokochi T, Noguchi S, et al. Cross talk between interferon-gamma and—alpha/beta signaling components in caveolar membrane domains. *Science*. 2000; 288(5475):2357–60. Epub 2000/07/06. PMID: [10875919](https://pubmed.ncbi.nlm.nih.gov/10875919/).
61. Lazear HM, Pinto AK, Vogt MR, Gale M Jr., Diamond MS. Beta interferon controls West Nile virus infection and pathogenesis in mice. *Journal of virology*. 2011; 85(14):7186–94. Epub 2011/05/06. doi: [10.1128/JVI.00396-11](https://doi.org/10.1128/JVI.00396-11) PMID: [21543483](https://pubmed.ncbi.nlm.nih.gov/21543483/); PubMed Central PMCID: PMC3126609.
62. Deonarain R, Alcamì A, Alexiou M, Dallman MJ, Gewert DR, Porter AC. Impaired antiviral response and alpha/beta interferon induction in mice lacking beta interferon. *Journal of virology*. 2000; 74(7):3404–9. Epub 2000/03/09. PMID: [10708458](https://pubmed.ncbi.nlm.nih.gov/10708458/); PubMed Central PMCID: PMC111842.
63. Runkel L, Pfeffer L, Lewerenz M, Monneron D, Yang CH, Murti A, et al. Differences in activity between alpha and beta type I interferons explored by mutational analysis. *The Journal of biological chemistry*. 1998; 273(14):8003–8. Epub 1998/05/09. PMID: [9525899](https://pubmed.ncbi.nlm.nih.gov/9525899/).
64. Lewerenz M, Mogensen KE, Uze G. Shared receptor components but distinct complexes for alpha and beta interferons. *Journal of molecular biology*. 1998; 282(3):585–99. Epub 1998/09/17. doi: [10.1006/jmbi.1998.2026](https://doi.org/10.1006/jmbi.1998.2026) PMID: [9737924](https://pubmed.ncbi.nlm.nih.gov/9737924/).
65. Muller U, Steinhoff U, Reis LF, Hemmi S, Pavlovic J, Zinkernagel RM, et al. Functional role of type I and type II interferons in antiviral defense. *Science*. 1994; 264(5167):1918–21. Epub 1994/06/24. PMID: [8009221](https://pubmed.ncbi.nlm.nih.gov/8009221/).

66. Honda K, Yanai H, Negishi H, Asagiri M, Sato M, Mizutani T, et al. IRF-7 is the master regulator of type-I interferon-dependent immune responses. *Nature*. 2005; 434(7034):772–7. Epub 2005/04/01. doi: [10.1038/nature03464](https://doi.org/10.1038/nature03464) PMID: [15800576](https://pubmed.ncbi.nlm.nih.gov/15800576/).
67. Sheehan KC, Lai KS, Dunn GP, Bruce AT, Diamond MS, Heutel JD, et al. Blocking monoclonal antibodies specific for mouse IFN-alpha/beta receptor subunit 1 (IFNAR-1) from mice immunized by in vivo hydrodynamic transfection. *Journal of interferon & cytokine research: the official journal of the International Society for Interferon and Cytokine Research*. 2006; 26(11):804–19. Epub 2006/11/23. doi: [10.1089/jir.2006.26.804](https://doi.org/10.1089/jir.2006.26.804) PMID: [17115899](https://pubmed.ncbi.nlm.nih.gov/17115899/).
68. Sheehan KC, Calderon J, Schreiber RD. Generation and characterization of monoclonal antibodies specific for the human IFN-gamma receptor. *J Immunol*. 1988; 140(12):4231–7. Epub 1988/06/15. PMID: [2453573](https://pubmed.ncbi.nlm.nih.gov/2453573/).
69. Schreiber RD. Identification of gamma-interferon as a murine macrophage-activating factor for tumor cytotoxicity. *Contemporary topics in immunobiology*. 1984; 13:171–98. Epub 1984/01/01. PMID: [6428805](https://pubmed.ncbi.nlm.nih.gov/6428805/).
70. Bach EA, Szabo SJ, Dighe AS, Ashkenazi A, Aguet M, Murphy KM, et al. Ligand-induced autoregulation of IFN-gamma receptor beta chain expression in T helper cell subsets. *Science*. 1995; 270(5239):1215–8. Epub 1995/11/17. PMID: [7502050](https://pubmed.ncbi.nlm.nih.gov/7502050/).
71. Sheehan KC, Ruddle NH, Schreiber RD. Generation and characterization of hamster monoclonal antibodies that neutralize murine tumor necrosis factors. *J Immunol*. 1989; 142(11):3884–93. Epub 1989/06/01. PMID: [2469726](https://pubmed.ncbi.nlm.nih.gov/2469726/).
72. Shankaran V, Ikeda H, Bruce AT, White JM, Swanson PE, Old LJ, et al. IFN-gamma and lymphocytes prevent primary tumour development and shape tumour immunogenicity. *Nature*. 2001; 410(6832):1107–11. Epub 2001/04/27. doi: [10.1038/35074122](https://doi.org/10.1038/35074122) PMID: [11323675](https://pubmed.ncbi.nlm.nih.gov/11323675/).
73. Koebel CM, Vermi W, Swann JB, Zerafa N, Rodig SJ, Old LJ, et al. Adaptive immunity maintains occult cancer in an equilibrium state. *Nature*. 2007; 450(7171):903–7. Epub 2007/11/21. doi: [10.1038/nature06309](https://doi.org/10.1038/nature06309) PMID: [18026089](https://pubmed.ncbi.nlm.nih.gov/18026089/).
74. Ebel GD, Dupuis AP 2nd, Ngo K, Nicholas D, Kauffman E, Jones SA, et al. Partial genetic characterization of West Nile virus strains, New York State, 2000. *Emerging infectious diseases*. 2001; 7(4):650–3. Epub 2001/10/05. doi: [10.3201/eid0704.010408](https://doi.org/10.3201/eid0704.010408) PMID: [11585527](https://pubmed.ncbi.nlm.nih.gov/11585527/); PubMed Central PMCID: PMC2631742.
75. Diamond MS, Shrestha B, Marri A, Mahan D, Engle M. B Cells and Antibody Play Critical Roles in the Immediate Defense of Disseminated Infection by West Nile Encephalitis Virus. *Journal of Virology*. 2003; 77(4):2578–86. doi: [10.1128/jvi.77.4.2578-2586.2003](https://doi.org/10.1128/jvi.77.4.2578-2586.2003) PMID: [12551996](https://pubmed.ncbi.nlm.nih.gov/12551996/)
76. Gentry MK, Henchal EA, McCown JM, Brandt WE, Dalrymple JM. Identification of distinct antigenic determinants on dengue-2 virus using monoclonal antibodies. *The American journal of tropical medicine and hygiene*. 1982; 31(3 Pt 1):548–55. Epub 1982/05/01. PMID: [6177259](https://pubmed.ncbi.nlm.nih.gov/6177259/).
77. Taniguchi T, Mantei N, Schwarzstein M, Nagata S, Muramatsu M, Weissmann C. Human leukocyte and fibroblast interferons are structurally related. *Nature*. 1980; 285(5766):547–9. Epub 1980/06/19. PMID: [6157095](https://pubmed.ncbi.nlm.nih.gov/6157095/).
78. Belardelli F, Gessani S, Proietti E, Locardi C, Borghi P, Watanabe Y, et al. Studies on the expression of spontaneous and induced interferons in mouse peritoneal macrophages by means of monoclonal antibodies to mouse interferons. *The Journal of general virology*. 1987; 68 (Pt 8):2203–12. Epub 1987/08/01. PMID: [3039046](https://pubmed.ncbi.nlm.nih.gov/3039046/).
79. Isaacs JD. The antiglobulin response to therapeutic antibodies. *Seminars in immunology*. 1990; 2(6):449–56. Epub 1990/11/01. PMID: [2104281](https://pubmed.ncbi.nlm.nih.gov/2104281/).
80. Daffis S, Samuel MA, Suthar MS, Keller BC, Gale M Jr., Diamond MS. Interferon regulatory factor IRF-7 induces the antiviral alpha interferon response and protects against lethal West Nile virus infection. *Journal of virology*. 2008; 82(17):8465–75. Epub 2008/06/20. doi: [10.1128/JVI.00918-08](https://doi.org/10.1128/JVI.00918-08) PMID: [18562536](https://pubmed.ncbi.nlm.nih.gov/18562536/); PubMed Central PMCID: PMC2519659.
81. Santiago-Raber ML, Baccala R, Haraldsson KM, Choubey D, Stewart TA, Kono DH, et al. Type-I interferon receptor deficiency reduces lupus-like disease in NZB mice. *The Journal of experimental medicine*. 2003; 197(6):777–88. Epub 2003/03/19. doi: [10.1084/jem.20021996](https://doi.org/10.1084/jem.20021996) PMID: [12642605](https://pubmed.ncbi.nlm.nih.gov/12642605/); PubMed Central PMCID: PMC2193854.
82. Pulverer JE, Rand U, Lienenklaus S, Kugel D, Zietara N, Kochs G, et al. Temporal and spatial resolution of type I and III interferon responses in vivo. *Journal of virology*. 2010; 84(17):8626–38. Epub 2010/06/25. doi: [10.1128/JVI.00303-10](https://doi.org/10.1128/JVI.00303-10) PMID: [20573823](https://pubmed.ncbi.nlm.nih.gov/20573823/); PubMed Central PMCID: PMC2919002.
83. Marie I, Durbin JE, Levy DE. Differential viral induction of distinct interferon-alpha genes by positive feedback through interferon regulatory factor-7. *The EMBO journal*. 1998; 17(22):6660–9. Epub 1998/11/21. doi: [10.1093/emboj/17.22.6660](https://doi.org/10.1093/emboj/17.22.6660) PMID: [9822609](https://pubmed.ncbi.nlm.nih.gov/9822609/); PubMed Central PMCID: PMC1171011.

84. Samuel MA, Diamond MS. Alpha/beta interferon protects against lethal West Nile virus infection by restricting cellular tropism and enhancing neuronal survival. *Journal of virology*. 2005; 79(21):13350–61. Epub 2005/10/18. doi: [10.1128/JVI.79.21.13350-13361.2005](https://doi.org/10.1128/JVI.79.21.13350-13361.2005) PMID: [16227257](https://pubmed.ncbi.nlm.nih.gov/16227257/); PubMed Central PMCID: PMC1262587.
85. Daffis S, Lazear HM, Liu WJ, Audsley M, Engle M, Khromykh AA, et al. The naturally attenuated Kunjin strain of West Nile virus shows enhanced sensitivity to the host type I interferon response. *Journal of virology*. 2011; 85(11):5664–8. Epub 2011/03/18. doi: [10.1128/JVI.00232-11](https://doi.org/10.1128/JVI.00232-11) PMID: [21411525](https://pubmed.ncbi.nlm.nih.gov/21411525/); PubMed Central PMCID: PMC3094947.
86. Erlandsson L, Blumenthal R, Eloranta ML, Engel H, Alm G, Weiss S, et al. Interferon-beta is required for interferon-alpha production in mouse fibroblasts. *Current biology: CB*. 1998; 8(4):223–6. Epub 1998/03/21. PMID: [9501984](https://pubmed.ncbi.nlm.nih.gov/9501984/).