

Synthesis and crystal structures of 2-methyl-4-aryl-5-oxo-5H-indeno [1,2-b] pyridine carboxylate derivatives

Kumar, Vinodh; Pandian, Ramesh; Naushad , Edayadukka; Peters, Günther H.J.; Nanjappagounder, P. M.

Published in:
Chemistry Central Journal

DOI:
[10.1186/1752-153X-8-34](https://doi.org/10.1186/1752-153X-8-34)

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Kumar, V., Pandian, R., Naushad , E., Peters, G. H. J., & Nanjappagounder, P. M. (2014). Synthesis and crystal structures of 2-methyl-4-aryl-5-oxo-5H-indeno [1,2-b] pyridine carboxylate derivatives. *Chemistry Central Journal*, 8(1), [34]. <https://doi.org/10.1186/1752-153X-8-34>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

RESEARCH ARTICLE

Open Access

Synthesis and crystal structures of 2-methyl-4-aryl-5-oxo-5*H*-indeno [1,2-*b*] pyridine carboxylate derivatives

Ramesh Pandian^{1,2†}, Edayadulla Naushad^{3†}, Vinodhkumar Vijayakumar⁴, Günther H Peters⁵ and Ponnuswamy Mondikalipudur Nanjappagounder^{1*}

Abstract

Background: Hantzsch 1,4-dihydropyridines (Hantzsch1,4-DHP) have been extensively utilized as the analogs of nicotinamide adenine dinucleotide (NADH) coenzyme to study the mechanism and various redox processes. During the redox processes 1,4-DHP systems undergo transformation into the corresponding pyridine derivatives through oxidation. Consequently, the interest in this aromatization reaction, investigation of a wide range of 1, 4-DHPs continues to attract the attention of researchers. Herein, we report the preparation of pyridine derivatives and the crystal structures determined by X-ray crystallographic methods.

Results: The crystal structures and conformational studies of two organic compounds, namely ethyl 2-methyl-4-phenyl-5-oxo-5*H*-indeno [1,2-*b*] pyridine-3-carboxylate (I) and ethyl 2-methyl-4-(4 chlorophenyl)-5-oxo-5*H*-indeno [1,2-*b*] pyridine-3-carboxylate (II) are reported. The terminal ethyl group of the compound I is disordered over two positions with the refined occupancies of 0.645 & 0.355 and C8 one dimensional zig-zag chain running along 101 direction through C-H...O type of intermolecular interactions. In the compound II, C-H...O interactions connect the molecules to form an R₂² (16) dimer running along 011 direction.

Conclusion: The crystal structures ethyl 2-methyl-4-phenyl-5-oxo-5*H*-indeno [1,2-*b*] pyridine-3-carboxylate and ethyl 2-methyl-4-(4 chlorophenyl)-5-oxo-5*H*-indeno [1,2-*b*] pyridine-3-carboxylate have been investigated in detail. The terminal ethyl group of compound I is disordered. In compound II, the substitution of Cl atom in the phenyl ring alters the configuration of carboxylate group with respect to the pyridine indane ring.

Background

Hantzsch 1,4-dihydropyridines (Hantzsch1,4-DHP) have been extensively utilized as the analogs of nicotinamide adenine dinucleotide (NADH) coenzyme to study the mechanism and the synthetic potential of various redox processes [1,2]. Hantzsch 1,4-DHP based drugs such as nifedipine and niguldipine are widely used as calcium channel blockers for the treatment of cardiovascular disorders including angina, hypertension and cardiac arrhythmias [3]. During the redox processes and in the course of drug metabolism [4], 1,4-DHP systems are oxidatively transformed into the corresponding pyridine

derivatives. Consequently, this aromatization reaction continues to attract the attention of researchers to establish a general protocol applicable to a wide range of 1,4-dihydropyridines. A number of methods and reagents have been reported recently in the literature for this purpose [5-14].

Some of these methods suffer from disadvantages such as the use of strong or toxic oxidants, the requirement of severe conditions or need excess of the oxidants. Other drawbacks are the long reaction times, production of by-products, the lower yields of products and/or the requirement of tedious work-up procedures.


N-Bromosuccinimide (NBS) is a versatile reagent for the oxidation of primary and secondary alcohols, α -hydroxycarboxylic acids [15], α -hydroxycarboxylic esters [16], hydrazines and hydrazones [15]. In addition, NBS is preferred for allylic bromination. While hydroxy acids like

* Correspondence: mnpsy2004@yahoo.com

†Equal contributors

¹Centre of Advanced Study in Crystallography and Biophysics, University of Madras, Guindy Campus, Chennai 600 025, India

Full list of author information is available at the end of the article



Scheme 1 Synthesis scheme of the dihydropyridines.

malic acid, tartaric acid, citric acid etc. are converted to aldehydes and ketones, polyhydric alcohols (glycol, glycerol and hexitols) are quantitatively decomposed to carbon dioxide and water [17] with NBS. NBS also promotes reactions of sterically hindered cresols *via* *p*-benzoquinone methide [18].

Having synthesized a number of 1, 4-dihydropyridines derived from indane-1,3-dione, we have dehydrogenated them to the corresponding pyridines. The reagent of the choice for effecting dehydrogenation is NBS in methanol (Schemes 1 and 2). This reagent was earlier employed to effect dehydrogenation of simple dihydropyridines [19].

Experimental

The title compounds reported in the present work were prepared by the following procedure [19,20].

Preparation of 4a-b

To an alcoholic solution (50 mL) of indane-1,3-dione **2** (0.01 mol), appropriate aromatic aldehydes **1a-b** (0.01 mol), ethyl acetoacetate **3** (0.01 mol), ammonium acetate (0.02 mol) and a drop of piperidine were added and the mixture was refluxed for 1 hr. The reaction


mixture was concentrated to half of its original volume and allowed to cool in an ice-chest. The solid **4a-b** thus separated was filtered, washed with ice cold aqueous ethanol and crystallized from petroleum ether (60–80°C)-chloroform (1: 1) (Scheme 1).

Preparation of 5a-b

To a solution of ethyl 2-methyl-4-aryl-5-oxo-1*H*,4*H*-indeno [1,2-*b*] dihydropyridine-3-carboxylate **4a-b** (0.5 g, 1.87 mmol) in methanol (10.0 mL), N-bromosuccinimide (0.33 g, 1.87 mmol) was added and the reaction mixture was stirred at room temperature. The colour of the solution changes immediately and the reaction proceeds instantaneously within five minutes. The course of the reaction was monitored by TLC. The reaction mixture was diluted with water (50 mL) and extracted with chloroform (3 × 20 mL). The organic layer was separated, dried over anhydrous sodium sulfate and filtered (Scheme 2). Evaporation of the solvent afforded the products ethyl 2-methyl-4-phenyl-5-oxo-5*H*-indeno [1,2-*b*] pyridine-3-carboxylate (Scheme 3) or ethyl 2-methyl-4-(4-chlorophenyl)-5-oxo-5*H*-indeno [1,2-*b*] pyridine-3-carboxylate respectively in excellent yields (Scheme 4). For compound (**5a**): Yield 96%; M.p. 212°C. For compound (**5b**): Yield 89%; M.p. 198°C.




Scheme 2 Synthesis scheme of the compounds I and II.



Results and Discussion


In both the compounds, the indenopyridine ring is almost planar, with r.m.s deviation of 0.035(2) Å [C3] and 0.087(2) Å [C11] for compounds I and II, respectively. The keto atom O substituted in the indenopyridine in both the molecules are slightly out of plane [0.048(2) & 0.217(1) Å for I & II]. The substitution of the Cl atom in the phenyl ring plays a vital role while packing the molecules in the unit cell and promotes the change of conformation of the carboxylate group. This is evidenced from the torsion angle values of [C10-C11-15-O2] and [C12-C11-C15-O2] 114.5(2)° & -63.5(2)° for (I) and -74.9(2)° & 114.0(1)° for (II), respectively. The terminal ethyl group in compound I is disordered over two positions with refined occupancies of 0.645 & 0.355. The phenyl ring and indenopyridine



are oriented by an angle of 67.8(1)° in compound (I) which is almost similar in compound (II) amounting the value of 55.2(1)°. The overall conformations in both the molecules are similar as can be seen from the superimposed rmsd value 0.154 Å (Figure 1). Both the structures are stabilized by C-H...O type of intra and intermolecular interactions. In compound I, molecules at (x, y, z) and (x + 1/2, -y - 1/2, z + 1/2) are linked through intermolecular C20-H20...O1 hydrogen bond to form a C8 zig-zag chain (Figure 2) running along 101 direction [21]. The combination of C5-H5...O3 and C22-H22...O1 intermolecular hydrogen bonds, lead to the formation of a R²₂ (16) ring motif chain running along [0 1 1] direction (Figure 3), observed in compound II.

X-ray Crystallography

Single crystal X-ray intensity data for the compounds (I) and (II) were collected using a Bruker Kappa APEX II area-detector diffractometer with MoK_α (0.71073 Å) radiation at room temperature (293 K). The data reduction was carried out using the program SAINT [22]. The absorption corrections were applied using the Multi-scan method using SADABS program [23]. The structures of both the compounds were solved by direct methods using SHELXS97 [24] and all the non-hydrogen atoms were refined anisotropically by full-matrix least-squares on F² taking all the unique reflections using SHELXL97 [24]. The hydrogen atoms attached with carbon atoms were placed in their calculated positions and included in the isotropic refinement using the riding model with C-H = 0.93 Å (-CH) or 0.97 Å (-CH₂) Å or






Figure 2 Figure showing the intermolecular hydrogen bonds resulting in C8 zig-zag motif in compound (I).




Figure 3 Figure showing the intermolecular hydrogen bonds resulting in R^2_2 (16) ring motifs chain running along 0 1 1 direction in compound (II).

Table 1 The crystal data, experimental conditions and structure refinement parameters for the compounds (I) and (II)

Parameters	Compound (I)	Compound (II)
Empirical formula	C ₂₂ H ₁₇ NO ₃	C ₂₂ H ₁₆ ClNO ₃
Formula weight	343.37	377.81
Wavelength		0.71073 Å
Crystal system, space group	Monoclinic, P2 ₁ /n	Triclinic, P-1
Unit cell dimensions	a = 7.5078(5) Å b = 21.0935(15) Å c = 11.5058(3) Å β = 104.876(2)°	a = 9.7750(8) Å; α = 113.199(2)° b = 9.8262(4) Å; β = 102.572(3)° c = 10.8687(5) Å; γ = 99.791(3)°
Volume	1761.1(2) Å ³	897.65(9) Å ³
Z, Calculated density	4, 1.295 g/cm ³	2, 1.398 g/cm ³
Absorption coefficient	0.086 mm ⁻¹	0.236 mm ⁻¹
F (000)	720	392
Crystal size	0.23 × 0.20 × 0.19 mm ³	0.22 × 0.18 × 0.17 mm ³
Theta range for data collection	1.93 to 30.48°	2.15 to 30.99°
Limiting indices	-10 ≤ h ≤ 10, -30 ≤ k ≤ 30, -15 ≤ l ≤ 16	-13 ≤ h ≤ 14, -14 ≤ k ≤ 14, -15 ≤ l ≤ 15
Reflections collected/unique	5323/2998 [R (int) = 0.032]	5580/4173 [R (int) = 0.0261]
Completeness	99.4%	97.5%
Absorption correction		Multi-scan
Refinement method		Full-matrix least-squares on F ²
Data/restraints/parameters	2998/0/238	5580/0/244
Goodness-of-fit on F ²	1.008	1.051
Final R indices [I > 2σ (I)]	R ₁ = 0.0574, wR ₂ = 0.1494	R ₁ = 0.0464, wR ₂ = 0.1334
R indices (all data)	R ₁ = 0.1047, wR ₂ = 0.1819	R ₁ = 0.0638, wR ₂ = 0.1470
Extinction coefficient	0.0098(18)	0
Largest diff. peak and hole	0.333 and -0.240 e.Å ⁻³	0.393 and -0.332 e.Å ⁻³

0.96 Å (-CH₃) Å with Uiso (H) = 1.2Ueq (parent C atom). The crystal data, experimental conditions and structure refinement parameters for the compounds (I) and (II) are presented in Table 1. Table 2 gives the geometry of the intra and intermolecular interactions. The

molecular structure of compounds (I) and (II) with the atom numbering scheme using ORTEP3 [25] are given in Figure 4 and Figure 5, respectively. The least-squares plane, geometrical and puckering parameters of both the compounds were calculated using PLATON software package [26-28].

Table 2 The geometry of the hydrogen bonds (Å, °)

D-H...A	D (D-H)	D (H...A)	D (D...A)	<(DHA)
Compound (I)				
C (20)-H (20)...O (1) ⁱ	0.93	2.40	3.232(3)	149
Compound (II)				
C (14)-H (14A)...O (2)	0.96	2.44	3.132(2)	128
C (5)-H (5)...O (3) ⁱⁱ	0.93	2.60	3.472(2)	157
C (22)-H (22)...O (1) ⁱⁱⁱ	0.93	2.58	3.458(2)	157
C (16)-H (16A)...Cg(3) ^{iv}	0.97	2.72	3.566(2)	146

Symmetry transformations used: (i) x + 1/2, -y - 1/2, z + 1/2; (ii) x, y - 1, z - 1; (iii) 1 - x, -y, -z; (iv) -x, -y, -z; Cg3 centroid atom of the ring (C2-C7).

Conclusions

The title compounds were synthesized, crystallized and the crystal structures have been determined by single-crystal X-ray diffraction methods. The terminal ethyl group of the compound I is disordered over two positions with the refined occupancies of 0.645 & 0.355. C-H...O intermolecular hydrogen bond builds up a one dimensional zig-zag chain running along 101 directions. In compound II, C-H...O hydrogen bonds connect the molecules to form a R²₂ (16) dimer chain running along 011 direction.




Figure 4 ORTEP plot of compound (I) showing with atoms ellipsoids are drawn at 40% probability level.




Figure 5 ORTEP plot of compound (II) showing with atoms ellipsoids drawn at 40% probability level.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

PR collected the X-ray data and solved the crystal structures under the guidance of MNP. NE synthesized the title compounds under the guidance of Prof. PR (Late). PR and NE contributed equally to this work. All authors have read and approved the final manuscript.

Acknowledgements

One of the authors NE is grateful to Mother Teresa Women's University, Kodaikanal, Tamilnadu-India and DST-CURIE programme for their encouragement and providing facilities for doing this research work. NE also put forth his heartfelt thanks to his guide Dr. P. Ramesh (Late) for the encouragement and motivation of the research work. PR thanks Prof. Kyeong Kyu Kim, Laboratory of Structural Biology, Department of Molecular Cell Biology, Samsung Biomedical Research Institute, Sungkyunkwan University School of Medicine, Suwon 440-746, South Korea and the National Research Foundation of Korea for the financial support in the form of postdoctoral fellowship (2012K2A4A1034867 and 2011-0030915).

Additional material

Crystallographic data (excluding structure factors) for the structures of compounds (I) and (II) reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication numbers, CCDC 996464 and CCDC 996465, respectively. Copies of the data can be obtained free of charge, on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK. (fax: +44-(0)1223-336033 or email: deposit@ccdc.cam.ac.uk).

Author details

¹Centre of Advanced Study in Crystallography and Biophysics, University of Madras, Guindy Campus, Chennai 600 025, India. ²Laboratory of Structural Biology, Department of Molecular Cell Biology, Samsung Biomedical Research Institute, Sungkyunkwan University School of Medicine, Suwon 440-746, South Korea. ³Department of Chemistry, Mother Teresa Women's University, Kodaikanal 624 101, India. ⁴Department of Life Sciences and Chemistry, Roskilde University, Roskilde DK-4000, Denmark. ⁵Department of Chemistry, Technical University of Denmark, Lyngby DK-2800 Kgs, Denmark.

Received: 20 February 2014 Accepted: 9 May 2014
Published: 29 May 2014

References

1. Stout DM, Meyers AI: Recent advances in the chemistry of dihydropyridines. *Chem Rev* 1982, **82**:223-243.
2. Kill RJ, Widdowson DA: In *Bioorganic Chemistry*. Edited by Van Tamelen EE. New York: Academic; 1978:239.
3. Triggle DJ: In *Comprehensive Medicinal Chemistry*. Edited by Emmett JC. Oxford: Pergamon; 1990. Vol. 3, Ch. 14.1.
4. Janis RA, Triggle DJ: New developments in Ca²⁺ channel antagonists. *J Med Chem* 1983, **26**:775-785.
5. Nasr-Esfahani M, Moghadam M, Tangestaninejad S, Mirkhani V: Biomimetic oxidation of Hantzsch 1,4-dihydropyridines with tetra-n-butylammoniumperiodate catalyzed by tetraphenylporphyrinatomanganese (III) chloride [Mn (TPP) Cl]. *Bioorg Med Chem Lett* 2005, **13**:3276-3278.
6. Moghadam M, Nasr-Esfahani M, Tangestaninejad S, Mirkhani V: Mild and efficient oxidation of Hantzsch 1,4-dihydropyridines with sodium periodate catalyzed by a new polystyrene-bound Mn (TPP) Cl. *Bioorg Med Chem Lett* 2006, **16**:2026-2030.
7. Balogh M, Hermezc I, Meszaros Z, Laszlo P: Aromatization of 1, 4-Dihydropyridines by Clay-Supported Metal Nitrates. *Helv Chim Acta* 1984, **67**:2270-2272.
8. Eynde JJV, Orazio RD, Haverabeke YV: Potassium permanganate, a versatile reagent for the aromatization of Hantzsch 1,4-dihydropyridines. *Tetrahedron* 1994, **50**:2479-2484.
9. Eynde JJV, Mayence A, Maquestiau A: A novel application of the oxidizing properties of pyridiniumchlorochromate: aromatization of Hantzsch 1,4-dihydropyridines. *Tetrahedron* 1992, **48**:463-468.
10. Delgado F, Alvarez C, Garcia O, Penieris G, Marques C: Unusual oxidative dealkylation of certain 4-alkyl-1, 4-dihydropyridines with MnO₂/bentonite

using microwave irradiation in the absence of solvent (II). *Synth Commun* 1991, **21**:2137-2141.

11. Mashraqui SH, Karnik MA: Bismuth nitrate pentahydrate. A convenient reagent for the oxidation of Hantzsch 1, 4-dihydropyridines. *Synthesis* 1998, **5**:713-714.
12. Wang B, Hu Y: The Aromatization of Hantzsch 1,4-Dihydropyridines by Tetrakis-Pyridine Cobalt (II) Dichromate (TPCD). *Synth Commun* 1999, **29**:4193-4199.
13. Maquestiau A, Mayence A, Eynde JJV: Ultrasound-promoted aromatization of hantzsch 1,4-Dihydropyridines by clay-supported cupric nitrate. *Tetrahedron Lett* 1991, **32**:3839-3840.
14. Khadikar B, Borkat S: Silica Gel Supported Ferric Nitrate: A Convenient Oxidizing Reagent. *Synth Commun* 1998, **28**:207-212.
15. Barakat MZ, Wahap MFA, Sadr MM: Some reactions with N-Bromosuccinimide. *J Am Chem Soc* 1955, **77**:1670-1672.
16. Kurse PF, Geurkink JN, Grist KL: Studies with N-Halo Reagents. II. New Syntheses of β -Bromo- α -keto Esters, Ethyl Phenylglyoxylate and Phenacyl Bromide Using N-Bromosuccinimide. *J Am Chem Soc* 1954, **76**:5796-5797.
17. Abdel-wahab MF, Barakat MZ: Die Verwendung von N-Bromosuccinimid bei Bromierungs-und Abbaureaktionen. *Monatsh Chem* 1957, **88**:692-701.
18. Duar S, Turk J, Speigle J, Cobin J, Masnovi J, Baker JR: NBS-Promoted Reactions of Symmetrically Hindered Methylphenols via p-Benzoquinone Methide. *J Org Chem* 2000, **65**:108-115.
19. Nagarajan R, Anthonyraj JCA, Muralidharan D, Saikumar C, Perumal PT: A convenient method for the oxidation of Hantzsch 1,4-dihydropyridines with N-bromosuccinimide. *Indian J Chem* 2006, **45B**:826-828.
20. Edayadulla N, Ramesh P: Synthesis of 2-methyl-4-aryl-5-oxo-1H, 4H-indeno [1,2-b] dihydropyridine-3-carboxylates. *Indian J Heterocyclic Chem* 2007, **16**:371-374.
21. Bernstein J, Davis RE, Shimon L, Chang NL: Patterns in Hydrogen Bonding: Functionality and Graph Set Analysis in Crystals. *Angew Chem Int Ed Engl* 1995, **34**:1555-1573.
22. Bruker: APEX2 and SAINT. Madison, Wisconsin, USA: Bruker AXS Inc; 2004.
23. Sheldrick GM: SADABS. Version 2.03. Germany: University of Göttingen; 2001.
24. Sheldrick GM: A short history of SHELX. *Acta Crystallogr A* 2008, **A64**:112-122.
25. Farrugia LJ: ORTEP-3 for Windows - a version of ORTEP-III with a Graphical User Interface (GUI). *J Appl Crystallogr* 1997, **30**:565.
26. Cremer D, Pople JA: A General Definition of Ring Puckering Coordinates. *J Am Chem Soc* 1975, **97**:1354-1358.
27. Nardelli M: Ring asymmetry parameters from out-of-plane atomic displacements. *Acta Appl Crystallogr* 1983, **C39**:1141-1142.
28. Spek AL: PLATON. *J Appl Crystallogr A* 2003, **36**:7-13.

doi:10.1186/1752-153X-8-34

Cite this article as: Pandian et al.: Synthesis and crystal structures of 2-methyl-4-aryl-5-oxo-5H-indeno [1,2-b] pyridine carboxylate derivatives. *Chemistry Central Journal* 2014 **8**:34.

Publish with **ChemistryCentral** and every scientist can read your work free of charge

"Open access provides opportunities to our colleagues in other parts of the globe, by allowing anyone to view the content free of charge."

W. Jeffery Hurst, The Hershey Company.

- available free of charge to the entire scientific community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.chemistrycentral.com/manuscript/



ChemistryCentral