

Nutrient (N and P) dynamics of the invasive macroalga *Gracilaria vermiculophylla*
nutrient uptake kinetics and nutrient release through decomposition

Pedersen, Morten Foldager; Lørup Johnsen, Kathrine

Published in:
Marine Biology

DOI:
[10.1007/s00227-017-3197-7](https://doi.org/10.1007/s00227-017-3197-7)

Publication date:
2017

Document Version
Peer reviewed version

Citation for published version (APA):
Pedersen, M. F., & Lørup Johnsen, K. (2017). Nutrient (N and P) dynamics of the invasive macroalga *Gracilaria vermiculophylla*: nutrient uptake kinetics and nutrient release through decomposition. *Marine Biology*, 164, 172. <https://doi.org/10.1007/s00227-017-3197-7>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@ruc.dk providing details, and we will remove access to the work immediately and investigate your claim.

1 **Nutrient (N and P) dynamics of the invasive macroalga *Gracilaria***
2 ***vermiculophylla*: Nutrient uptake kinetics and nutrient release through**
3 **decomposition**

4
5 **Morten F. Pedersen* and Kathrine L. Johnsen**

6
7 Department of Science and Environment (DSE)
8 Roskilde University, Box 260, DK-4000 Roskilde, Denmark.

9
10 * Corresponding author: Morten Foldager Pedersen, email: mfp@ruc.dk, phone: +45
11 46742088.

12
13
14
15 **Acknowledgments**

16 We thank technician Anne B. Faarborg for assistance in the laboratory and two anonymous
17 referees for valuable comments to an earlier version of the manuscript.

18
19 **Compliance with Ethical Standards:**

20 **Funding:** This study was supported by grant no. 272-08-0577 (MARINVA) from the Danish
21 Council for Independent Research.

22 **Conflict of interest:** The authors declare that they have no conflict of interest.

23 **Ethical approval:** All applicable international, national, and/or institutional guidelines for the
24 care and use of animals were followed. This article does not contain any studies with human
25 participants performed by any of the authors.

26 **Abstract**

27 The invasive alga *Gracilaria vermiculophylla* was introduced to Europe two decades ago and
28 has since become dominant in many shallow estuaries. *G. vermiculophylla* is a relatively fast-
29 growing alga that thrives well at low nutrient availability in summer, suggesting that it uses
30 nutrients efficiently, which might give it a competitive advantage over many native species.
31 We studied therefore the nutrient dynamics of *G. vermiculophylla* and compared it to those of
32 a range of native macroalgal species. Nutrient uptake rates (NH_4^+ , NO_3^- and PO_4^{3-}) were
33 compared to growth related requirements and we found that *G. vermiculophylla* needs
34 relatively high nutrient concentrations to sustain fast and non-limited growth. This compares
35 to the nutrient dynamics of many fast-growing algae and we found thus no indication that *G.*
36 *vermiculophylla* should have any particularly advantage relative to other, sympatric species.
37 The nutrient storage capacity of *G. vermiculophylla* was, in contrast, relatively large and
38 comparable to that of more slow-growing algae, which, when combined with the low nutrient
39 uptake experienced in summer, could explain how *G. vermiculophylla* can sustain non-
40 nutrient limited growth through most of the growth season. The biomass of *G.*
41 *vermiculophylla* can be massive and estimates showed that gross nutrient uptake could exceed
42 the amount of nutrients received from land. The turnover of *Gracilaria* biomass is however
43 fast and nutrients bound in the resulting detritus are quickly mineralized during
44 decomposition, which is especially important during late summer when water temperatures
45 are high. Invasion and subsequent dominance by *G. vermiculophylla* may thus affect local
46 nutrient cycling significantly.

47

48 **Key words:** Nitrate, ammonium, phosphate, uptake kinetics, mineralization, seaweed.

49 **Introduction**

50 The red alga *Gracilaria vermiculophylla* originates from NE Asia (Tseng and Xia 1999), but
51 was introduced to North America and Europa with imported oysters intended for aquaculture
52 in the 1990's (Mollet et al. 1998). *G. vermiculophylla* is invasive (Nyberg et al. 2009) and has
53 spread along the Atlantic coast of Europe and is now found from Venice lagoon in the
54 Mediterranean to southern Norway in Scandinavia (Thomsen et al. 2007; Sfriso et al. 2012).
55 *G. vermiculophylla* was first observed in Danish waters in 2003 and is now common in many
56 estuaries of the western Baltic Sea (Thomsen et al. 2007; Weinberger et al. 2008). *G.*
57 *vermiculophylla* is mostly abundant in sheltered, soft-bottom areas where it may become
58 dominant and sometimes replaces previously common seaweeds such as *Ulva* sp. (Nejrup and
59 Pedersen 2010) or *Fucus vesiculosus* (Weinberger et al. 2008). *G. vermiculophylla* is
60 considered an ecosystem engineer (Wallentinus and Nyberg 2007; Byers et al. 2012) and
61 where abundant it may not only affect the composition of the algal assemblage, but also affect
62 local biogeochemistry, nutrient cycling including the transfer of nutrients between abiotic and
63 biotic components (Tyler and McGlathery 2006; Hardison et al. 2010; Gulbransen &
64 McGlathery 2013) and alter trophic relations in the food web (Wallentinus and Nyberg 2007).

65 *Gracilaria vermiculophylla* has a number of traits that are typical for invasive
66 algae; it is relatively fast-growing ($22\% \text{ d}^{-1}$; Raikar et al. 2001), recruits from both spores and
67 fragments (Rueness 2005; Thomsen et al. 2007; Nyberg et al. 2009), it is tolerant to
68 desiccation and extreme levels of light, temperature and salinity (Yokoya et al. 1999; Raikar
69 et al. 2001; Nejrup and Pedersen 2012; Nejrup et al. 2013). Recent studies have further shown
70 that *G. vermiculophylla* is avoided by many native herbivores within its invaded range
71 (Thomsen and McGlathery 2007; Weinberger et al. 2008; Nejrup et al. 2012; Hammann et al.
72 2013) due to inducible chemical defenses (Nylund et al. 2011; Hammann et al. 2016).

73 Low nutrient availability (permanently or periodically) restricts growth of fast-

74 growing species more than that of more slow-growing species because the latter are better
75 adapted to cope with low nutrient availability (Pedersen and Borum 1996; Pedersen and
76 Borum 1997; Pedersen et al. 2010) and nutrient richness may therefore affect species
77 composition of algal assemblages (Bokn et al. 2003; Karez et al. 2004; Kraufvelin et al. 2006;
78 Kraufvelin et al. 2010). *G. vermiculophylla* is relatively fast growing and should therefore
79 have high nutrient demands per unit biomass and time and be susceptible to nutrient
80 limitation during late spring and summer where insolation and water temperature is high
81 enough to support rapid growth, but where the availability of nutrients is low. Nejrup and
82 Pedersen (2010) studied seasonal variations in biomass and growth of *G. vermiculophylla* in
83 two Danish estuaries with low nutrient availability in summer, but were unable to detect any
84 significant increase in growth following experimental nutrient enrichment. This result
85 indicates that *G. vermiculophylla* has the capacity to acquire dissolved nutrients efficiently
86 even when these are present at low concentrations or, that it is able to sustain growth in
87 summer by using internal nutrient reserves obtained during winter and early spring where
88 nutrient availability is high. Both strategies are comparable to those of more slow-growing
89 species that can sustain near maximum growth rates during extended periods of low nutrient
90 availability (Pedersen and Borum 1996; Pedersen and Borum 1997; Pedersen et al. 2010).

91 Fast growth combined with an efficient uptake capacity (relative to its demands)
92 and/or high nutrient storage capacity may leave *G. vermiculophylla* competitively superior
93 under low nutrient availability, which could add to explain its recent success in European and
94 north American estuarine waters. Few studies have investigated the nutrient dynamics of *G.*
95 *vermiculophylla* and most of these have only assessed one or a few aspects of the nutrient
96 dynamics, i.e. either uptake kinetics (Tyler et al. 2005; Tyler and McGlathery 2006; Abreu et
97 al. 2011) or the role of *in situ* nutrient limitation (e.g. Thomsen and McGlathery 2007; Nejrup
98 and Pedersen 2010). No studies have yet evaluated the full set of dynamics including uptake

99 kinetics, requirements for growth and the role of stored nutrients.

100 Seaweeds may acquire and temporarily immobilize a large proportion of the
101 nutrients received from land, especially in shallow estuaries and coastal lagoons where algal
102 biomass can be substantial (Tyler and McGlathery 2003; Pedersen et al. 2004). Nutrients
103 incorporated into macroalgal biomass become temporarily unavailable for other primary
104 producers until they are released through grazing or decomposition. The turnover rate of algal
105 biomass and, hence, the release of nutrients, differs systematically among seaweeds with
106 different life strategies; slow-growing macroalgae tend generally to be grazed less and
107 decompose more slowly than fast-growing species (e.g. Buchsbaum et al. 1991; Enriquez et al.
108 1993; Banta et al. 2004; Conover et al. 2016). Algal assemblages dominated by fast-growing
109 and bloom-forming macroalgae do therefore have a faster and more variable turnover of
110 nutrients than those dominated by slow-growing, perennial macrophytes (Duarte and Cebrián
111 1996; Banta et al. 2004) and release of nutrients from dense populations of opportunistic
112 seaweeds may periodically exceed land-derived inputs and the efflux of nutrients from
113 sediments (Tyler et al. 2003). Grazing on *G. vermiculophylla* is insignificant (e.g. Weinberger
114 et al. 2008; Nejrup and Pedersen 2010) so most of the nutrients contained in the biomass must
115 consequently be released through decomposition. Invasion and subsequent dominance by *G.*
116 *vermiculophylla* may thus potentially influence nutrient dynamics at the ecosystem level if the
117 decomposition rate of *G. vermiculophylla* differs markedly from the species it has replaced.

118 The aim of this study was 2-fold. The major aim was to evaluate all aspects of the
119 nutrient dynamics of *G. vermiculophylla* and to compare it to those of other common,
120 indigenous algal species to assess whether *G. vermiculophylla* uses nutrients more efficiently
121 than the algae it is potentially competing with. We wanted next to study the release of major
122 nutrients (N and P) from decomposing *G. vermiculophylla* and compare it to that of other
123 common, indigenous algal species to assess how *G. vermiculophylla* may affect the turn over

124 of nutrients in estuarine systems once it has become dominant. We measured nutrient uptake
125 kinetics for ammonium (NH_4^+), nitrate (NO_3^-) and phosphate (PO_4^{3-}) and compared those to
126 growth related N and P requirements, the latter being determined from culturing algae under a
127 range of nutrient concentrations. These data were combined and used to model how growth
128 relates to nutrient availability and additionally, used to estimate the storage capacity for N and
129 P in *G. vermiculophylla*. We conducted finally a series of decomposition experiments to
130 evaluate how fast N and P bound in biomass was released from dead and decaying *G.*
131 *vermiculophylla* under different temperature regimes

132

133 **Methods**

134 *Gracilaria vermiculophylla* was collected at Fyns Hoved, Denmark (55° 36.9' N, 10° 36.7' E)
135 in October 2013. The algae were cleaned and transported to the laboratory where they were
136 kept in 80 L storage tanks until being used in the experiments. The storage tanks and the
137 experimental chambers used for the uptake and growth experiments were kept at constant
138 temperature (15°C) and salinity 25 (PSU) and were illuminated by lamps equipped with
139 halogen spots (OSRAM 12V, 35W) providing a light intensity of ca. 90 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$
140 PAR in a 16:8 hr light:dark cycle. The chosen temperature and salinity is optimal for *G.*
141 *vermiculophylla* (Nejrup and Pedersen 2012) while the light intensity is sufficient to saturate
142 growth of this species (ca. 60 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$; Nejrup et al. 2013). The initial N and P
143 content in the algae were 1.7 % DW and 0.19 % DW, respectively.

144 **Nutrient uptake kinetics.** Uptake kinetics for NH_4^+ , NO_3^- and PO_4^{3-} were determined in three
145 separate experiments using a combination of the multiple-flask and the depletion methods as
146 described by Harrison et al. (1989) and Pedersen (1994). Eight PVC-beakers (Vol = 1.6 L)
147 were filled with a known volume (1.0-1.4 L) of seawater (salinity 25) with different added
148 concentrations of the nutrient species in question (NO_3^- range: 15-76 μM ; NH_4^+ range: 1-76

149 μM ; PO_4^{3-} range: 1-14 μM). Each beaker was bubbled with air to ensure circulation and
150 reduce the thickness of boundary layers. Measurements of nutrient uptake were initiated by
151 adding an algal sample (5-10 g FW) to each beaker where it was attached to a PVC-net to
152 keep it submerged during the incubation. Three replicate water samples (each 5.0 mL) were
153 taken from each beaker during the first hour of the incubation (at time = 0, 15, 30, 45 and 60
154 minutes) and then subsequently at every 30 min for the remaining part of the experiment.
155 Water samples were immediately frozen at -20°C for later analysis of nutrients. The
156 incubations lasted for 120 – 300 minutes and all algae were subsequently harvested and dried
157 to constant dry weight (DW) at 85°C and weighted. Concentrations of NO_3^- in the water
158 samples were analyzed using a Lachat (QuickChem FIA+ 8000 Series) autoanalyzer while
159 concentrations of NH_4^+ and PO_4^{3-} were analyzed manually; the concentration of NH_4^+ was
160 analyzed using the salicylate-hypochlorite method (Bower and Holm-Hansen 1980) while that
161 of PO_4^{3-} was analyzed spectrophotometrically following Strickland and Parsons (1968).

162 Biomass specific nutrient uptake rates (V) were estimated from changes in substrate
163 concentration (S) over the course of the experiment:

164
$$V = \frac{(S_0 \times \text{Vol}_0) - (S_T \times \text{Vol}_T)}{t \times B} \quad (\text{Eq. 1})$$

165 where V is the uptake rate (in $\mu\text{mol g}^{-1} \text{DW h}^{-1}$), S_0 and Vol_0 are the substrate concentration (in
166 μM) and volume (in L) at the beginning of a time interval, while S_T and Vol_T are the substrate
167 concentration and volume at the end of a time interval. t is the time elapsed between two
168 successive samplings and B is the DW biomass. Uptake rates determined on algae from
169 different beakers, but obtained during identical time intervals, were plotted against the mean
170 substrate concentrations obtained in each beaker during that specific time interval. The
171 Michaelis-Menten function (Eq. 2) was fitted to data by least square non-linear regression
172 using SYSTAT v. 13:

173
$$V = \frac{V_{\text{max}} \times S}{K_m + S} \quad (\text{Eq. 2})$$

174 where V is the uptake rate (in $\mu\text{mol g}^{-1} \text{DW h}^{-1}$), V_{max} is the maximum uptake rate, K_m is the
175 half-saturation constant (in μM) and S is the substrate concentration (in μM). Uptake rates of
176 NH_4^+ and PO_4^{3-} were initially enhanced when the algae were exposed to nutrients (figure 1)
177 while no such transiently enhanced uptake was observed in the case of NO_3^- . Parameter
178 estimates of V_{max} and K_m (table 1) were therefore represented by 2 sets of uptake kinetics: (1)
179 one representing transiently enhanced uptake rates (i.e. surge uptake), measured over the
180 initial 30 minutes after exposure to nutrients and, (2) one representing rates obtained after 90
181 minutes in the case of NH_4^+ and later than 30 minutes after exposure in the case of NO_3^- and
182 PO_4^{3-} (hereafter called assimilation).

183 **Nutrient requirements.** Algae were first pre-conditioned for 4-6 weeks in 10 aquaria
184 (volume = 20 L) receiving different quantities of dissolved N or P, to obtain specimens with
185 different tissue nutrient concentrations. Nutrients were added from stock-solutions of
186 NH_4NO_3 and KH_2PO_3 . Algae receiving different levels of NH_4NO_3 received KH_2PO_3 in
187 excess while algae receiving different levels of KH_2PO_3 received NH_4NO_3 in excess to ensure
188 that only one nutrient was limiting at the time. The aquaria were exposed to low light (40
189 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ PAR in a 16:8 hr light:dark cycle) using shade screens to ensure slow
190 growth, which eased the accumulation of nutrients in the tissues. Tissue N-concentrations in
191 algae intended for the N-growth experiment ranged from 1.0 to 3.7% N of DW at the end of
192 the pre-condition period while the P-content in these algae averaged 0.23 % of DW. Final
193 tissue P-concentrations in algae intended for the P-growth experiment ranged from 0.04 to
194 0.27 % of DW while the N-content in these algae averaged 2.4 % of DW .

195 Growth measurements were subsequently conducted in PVC-beakers (volume =
196 1.6L), which were filled with GF/C-filtered seawater (salinity 25) that had been stripped for
197 inorganic nutrients by letting *Ulva lactuca* grow in it for 2-3 days prior to use. The beakers
198 were placed under saturating light ($90 \mu\text{mol m}^{-2} \text{s}^{-1}$ PAR) and each beaker was bubbled with

199 atmospheric air to create circulation and ensure exchange of O₂ and CO₂. The initial FW
200 biomass was determined on all pre-conditioned specimens prior to the growth measurements
201 and the DW:FW ratio (and initial tissue N or P concentrations) was determined on a number
202 of sacrificed subsamples after drying them to constant weight at 85°C. The algae were left to
203 grow for 5-7 days after which they were harvested, dried to constant weight and analyzed for
204 final tissue N or P content. Growth rates were estimated from changes in biomass over time
205 assuming exponential growth:

$$206 \quad \mu = \frac{\ln B_T - \ln B_0}{t} \quad \text{Eq. 3}$$

207 where μ is the relative growth rate (d⁻¹), B_0 and B_T are the initial and final DW biomass and t
208 is the incubation time. Growth rates were plotted against tissue nutrient concentrations and
209 the Droop equation was fitted to data using least square, non-linear regression (SYSTAT v.
210 13):

$$211 \quad \mu = \mu_{max} \times \left(1 - \frac{Q_S}{Q}\right) \quad \text{Eq. 4}$$

212 where μ is the relative growth rate (d⁻¹), μ_{max} is the maximum growth rate, Q_S is the
213 subsistence cell quota (in % of DW), i.e. the lowest tissue nutrient concentration that allows
214 growth and Q is the average tissue N or P concentration (i.e. = $(Q_{Init} + Q_{Final})/2$). The critical
215 cell quota (Q_C) was defined as the tissue nutrient concentration above which growth is not
216 limited by lack of nutrients. The Droop function is a continuous function so we arbitrarily
217 determined Q_C as the tissue nutrient concentration where the corresponding growth rate
218 equaled 67% of the estimated μ_{max} (Pedersen and Borum 1996).

219 **Modeling substrate dependent growth.** The kinetics of substrate dependent growth were
220 estimated by combining data for substrate dependent uptake kinetics and tissue nutrient
221 dependent growth. Substrate dependent growth of macroalgae is often described by Monod
222 kinetics (e.g. Rosenberg et al. 1984) with parameters μ^*_{max} and K_{μ} , which can be estimated

223 from equations 5 and 6 assuming that nutrient uptake and use of nutrients for growth is in
224 steady state (Turpin 1988):

$$225 \quad \mu_{max}^* = \frac{(\mu_{max} \times V_{max})}{[(\mu_{max} \times Q_S) + V_{max}]} \quad \text{Eq. 5}$$

$$226 \quad K_{\mu} = \frac{(K_m \times \mu_{max} \times Q_{min})}{[(\mu_{max} \times Q_S) + V_{max}]} \quad \text{Eq. 6}$$

227 where V_{max} and K_m are the maximum uptake rate (in μ mol g^{-1} DW h^{-1}) and the Michaelis-
228 Menten constant (in μ M), respectively, obtained from the nutrient uptake experiments (Eq. 2)
229 while μ_{max} and Q_S are the maximum growth rate (d^{-1}) and subsistence cell quota (in % DW)
230 obtained from the growth experiments (Eq. 4).

231 **Storage capacity.** The amount of N or P being stored in excess of what is needed to obtain
232 maximum growth was estimated as the difference between the critical quota (Q_C) and the
233 highest observed quota (Q_{max}) in algae from the growth experiment. The excess amount of
234 cellular N or P, respectively, can support growth at maximum rates until Q_C is reached after
235 which growth ceases as the cell quota approaches Q_S . We defined storage capacity as the
236 number of days ($T_{Storage}$) that this pool of N or P could support growth (at maximum and
237 reduced rates) without additional acquisition of N or P from the medium. The storage capacity
238 was estimated from a numerical solution of equation 4 for Q . We used a time step of 0.1 day
239 for each iteration and started out with an initial cell quota equal to the observed Q_{max} . A new
240 cell quota was estimated for each time step using the estimated growth rate from the previous
241 time step. The storage capacity that could support non-nutrient limited growth was then
242 defined as the time passing until the cell quota reached Q_C while the storage capacity that
243 could support growth at reduced rates was defined as the time it took to reduce the quota from
244 Q_C to Q_S . Potential time lags to mobilize nutrient reserves were not accounted for.

245 **Decomposition and mineralization.** Decomposition of *G. vermiculophylla* and
246 mineralization of N and P bound in its tissue was studied using litterbags. 105 algal samples

247 (each 0.6 – 2.5 g FW) were cleaned and their initial FW biomass determined. Fifteen samples
248 were initially sacrificed and dried for determination of the initial DW:FW ratio. The
249 remaining ninety samples were placed in separate litterbags with a mesh size of 1 mm, which
250 were distributed equally among nine aquaria. Each aquarium had a volume of 45 L and was
251 filled with ca. 20 L of sediment from the sampling site and 20 L of seawater (salinity 25). The
252 aquaria were kept dark in 3 climate chambers with temperatures 5, 15 and 25°C, respectively,
253 thus simulating a typical winter, spring/fall or summer situation. The water was bubbled with
254 atmospheric air to keep it aerated and exchanged with freshly collected seawater monthly.
255 The litterbags with live algae were covered by ca. 1 cm of sediment in an attempt to simulate
256 slight burial as observed in the field. Three litterbags (one per replicate aquarium at each
257 temperature) were retrieved periodically, and the algal remains were rinsed for sand and mud,
258 dried to constant DW at 85°C, weighed and stored for later analysis of tissue nutrients (C, N
259 and P). Decomposition rates were estimated by fitting a multiple-G model (Eq.; Westrich and
260 Berner 1984) to the data (i.e. remaining biomass vs. time). This model assumes that detritus
261 may be made up by several fractions, each decomposing at a specific rate (k_i) plus a refractory
262 fraction (G_R) that does not decompose within the time-scale studied. If only one actively
263 decomposing pool of matter and a refractory pool can be deduced from data, the model
264 simplifies to:

$$265 \quad B_T = B_0 e^{-kt} + G_R \quad \text{Eq. 7}$$

266 where B_T and B_0 are the final and initial biomass (measured in units of C, N or P), k is the
267 decay rate, t is the number of days elapsed since initiating the experiment and G_R is the size of
268 the residual (i.e. non-reactive) fraction of the detritus.

269 **Tissue nutrient analyses.** Tissue concentrations of C and N were determined on dried and
270 ground samples using an EA 1110 CHNS elemental analyzer (CE Instruments). Tissue P was
271 determined on dried and ground algae after oxidation with boiling H_2SO_4 followed by

272 spectrophotometric analysis (Strickland and Parsons 1968).

273

274 **Results**

275 **Nutrient uptake kinetics.** Uptake rates of NH_4^+ , NO_3^- and PO_4^{3-} increased with increasing
276 substrate concentration and saturated at high concentrations (figure 1) why uptake kinetics
277 were described by the Michaelis-Menten function (R^2 -values ranged from 0.629 to 0.930; all
278 p-values <0.001). The maximum surge uptake rate of NH_4^+ was 2-fold higher than the
279 maximum assimilation rate, while the half-saturation constant for uptake (K_m) during surge
280 uptake was 3-fold higher than that for assimilation. These differences resulted in a higher
281 affinity (α) for NH_4^+ at low substrate concentrations during assimilation than during surge
282 uptake. The initial maximum uptake rate for NO_3^- was not different from that observed later in
283 the experiment (i.e. no surge uptake), but it was 3 to 4-fold lower than both V_{max}^{Surge} and V_{max}^{Ass}
284 for NH_4^+ . K_m for NO_3^- averaged 9 μM and the affinity at low substrate concentration ca. 1.7
285 μM . Initial V_{max} and K_m for PO_4^{3-} were ca. 2-fold higher than for P-uptake occurring after 30
286 minutes whereas the affinity for P at low PO_4^{3-} concentrations (α) was approximately the
287 same for surge uptake and assimilation.

288 **Growth rate and nutrient requirements.** Growth rate increased as a function of N or P
289 quotas (figure 2; R^2 for N = 0.876, $p < 0.001$; R^2 for P = 0.876, $p < 0.001$). Estimated maximum
290 growth rate at infinite N or P quota varied slightly between the two experiments (0.088 –
291 0.105 d^{-1}), but did not differ significantly according to the 95% CL's (table 2). The N
292 subsistence quota (Q_S) was 0.71% of DW while the critical N quota (Q_C) was 2.14% of DW,
293 resulting in a N-requirement of 169 $\mu\text{mol N g}^{-1} \text{DW d}^{-1}$ at maximum growth rate. The
294 subsistence quota for P was 0.05% of DW while the critical P quota was 0.14% of DW,
295 resulting in a P-demand of 4.1 $\mu\text{mol P g}^{-1} \text{DW d}^{-1}$.

296 **Substrate dependent growth.** Substrate dependent growth was modeled from data on uptake
297 kinetics and quota dependent growth and was represented by the derived Monod parameters
298 (table 3). Estimated maximum growth rate based on uptake of NH_4^+ (0.098 d^{-1}) or NO_3^- (0.091
299 d^{-1}) as the N source did not differ much. In contrast, the half-saturation constant for growth
300 (K_μ) based on NO_3^- was 2-fold larger than that for growth on NH_4^+ indicating a higher affinity
301 for NH_4^+ . Substrate concentrations needed to saturate growth (S_{Sat}) ranged from $5 \mu\text{M}$ for NH_4^+
302 to $10 \mu\text{M}$ for NO_3^- . Estimated maximum growth rate based on uptake of PO_4^{3-} (0.084 d^{-1}) was
303 slightly lower than for NH_4^+ or NO_3^- and K_μ was $0.12 \mu\text{M PO}_4^{3-}$. The substrate concentration
304 needed to support maximum growth rate was ca. $1 \mu\text{M PO}_4^{3-}$.

305 **Nutrient storage capacity.** The amount of N stored in excess of the critical quota (i.e. $Q_{max} -$
306 Q_C) attained $14.0 \text{ mg N g}^{-1} \text{ DW}$, while the internal quantity of N that could support sustained
307 growth at reduced rates (i.e. $Q_C - Q_S$) was $14.3 \text{ mg N g}^{-1} \text{ DW}$ (table 2). These internal stocks
308 of N could support continual growth at maximum or reduced rates for a total of 33 days (6.5
309 days at maximum rate and 26.6 days at reduced rates) without acquisition of dissolved N from
310 the surroundings (table 4). The amount of P in excess of the critical limit was 1.3 mg P g^{-1}
311 DW and the amount of P that could support sustained growth at reduced rates was 1.0 mg P
312 $\text{g}^{-1} \text{ DW}$ (table 2). The internal stocks of P were could support growth for a total period of 43
313 days (10.5 days at maximum rate and 32.5 days at reduced rates; table 4) without having to
314 acquire dissolved P from the medium.

315 **Decomposition and mineralization.** Decomposition of *G. vermiculophylla* followed a simple
316 1-G model without a significant refractory pool (i.e. a simple exponential decline; figure 3,
317 table 5; R^2 ranged from 0.774 to 0.908, all p-values <0.001) since this model provided better
318 fits (higher R^2 -values) than multiple G-models with or without a refractory pool. All biomass
319 disappeared within 120 days in the 15 and 25°C treatments, while about 30% of the original
320 biomass was left in the 5°C treatment at the end of the experiment. Decay rates based on loss

321 of C biomass ranged from 0.013 to 0.065 d⁻¹ depending on temperature and increased almost
322 5-fold across the temperature range from 5 to 25°C. Temperature had the same effect on loss
323 of tissue-bound N and P. Decomposition rates ranged from 0.016 to 0.079 d⁻¹ when expressed
324 in units of N and were marginally higher than when expressed in units of C. Decay rates
325 expressed in units of P ranged from 0.023 to 0.1 d⁻¹ depending on temperature and were
326 substantially higher than those expressed in units of C or N, indicating that P was mineralized
327 faster than C and N during decomposition.

328

329 **Discussion**

330 The balance between uptake capacity and nutrient requirements and, thus, the risk of suffering
331 nutrient limitation during periods of low nutrient availability is related to the maximum
332 growth rate of algae. Fast-growing algae have a large relative surface area (i.e. high SA:V
333 ratio) and a high capacity for nutrient uptake (Wallentinus 1984; Hein et al. 1995), but also
334 large nutrient requirements per unit biomass and time dictated by their fast growth (Pedersen
335 and Borum 1997; Pedersen et al. 2010). In contrast, slow-growing species have lower uptake
336 capacities and lower requirements, but uptake rate and nutrient demands are better scaled at
337 low nutrient availability in these algae (Pedersen and Borum 1997; Pedersen et al. 2010), why
338 slow-growing algae generally can grow at near maximum rates when concentrations of
339 dissolved inorganic N (DIN) and P (DIP) in the water are low.

340 Our findings failed to support the hypothesis presented by Nejrup and Pedersen
341 (2010); *G. vermiculophylla* did not have an exceptionally high affinity for inorganic nutrients
342 relative to its requirements for growth. Nutrient demands are largely determined by apparent
343 growth rate and critical cell quotas. *G. vermiculophylla* attained maximum growth rates
344 between 0.09 and 0.11 d⁻¹, which is comparable to those reported from other studies (Yokoya
345 et al. 1999; Raikar et al. 2001; Nejrup and Pedersen 2010; Nejrup et al. 2013), but lower than

346 for truly fast-growing, sheet-like and filamentous species and higher than for most fucoids
347 and Laminarians (see table 6 for a comparison with algae indigenous to Scandinavian waters).
348 *G. vermiculophylla* can thus be ranked as an algal species with an intermediate growth rate.

349 The experimentally determined critical N quota of *G. vermiculophylla* is rather high
350 and comparable to those of fast-growing species such as *U. lactuca* and *C. virgatum* (table 6).
351 Red algae have generally higher N quotas than green and brown algae due to the prevalence
352 of protein rich pigments in red algae (Hurd et al. 2014). The critical P quota of *G.*
353 *vermiculophylla* was, in contrast, low and corresponded to those of more slow-growing
354 species such as *F. vesiculosus* and *A. nodosum* (table 6). The observed maximum growth rates
355 and critical N and P quotas resulted in maximum N and P requirements corresponding to ca.
356 169 and 4 $\mu\text{mol N or P g}^{-1} \text{DW d}^{-1}$, respectively, which is lower than for truly fast-growing
357 algae such as *U. lactuca* and *C. virgatum*, but higher than those for more slow-growing
358 species (Pedersen and Borum 1997; Pedersen et al. 2010).
359 High nutrient requirements do not necessarily represent a problem if the affinity for these
360 nutrients is high enough. The obtained nutrient uptake kinetics of *G. vermiculophylla*
361 corresponded to those found by other authors (e.g. Tyler et al. 2005; Abreu et al. 2011) and
362 show that maximum uptake rates for dissolved inorganic N and P by far exceed the
363 requirements even when *G. vermiculophylla* is growing at maximum rate. Such uptake rates
364 can, however, only be obtained at relatively high and often ecologically irrelevant substrate
365 concentrations. The main question was therefore whether the capacity of *G. vermiculophylla*
366 to acquire nutrients is better scaled to growth related nutrient demands than in native algae
367 that it potentially competes with. This is best evaluated by comparing species-specific
368 substrate dependent growth and its related parameters across species. The half-saturation
369 constants (K_{μ}) for growth and the substrate concentrations needed to saturate growth (S_{SAT})
370 on either NH_4^+ or NO_3^- tended to be higher than those of more slow-growing algae and

371 comparable to those of faster growing algae showing that *G. vermiculophylla* requires
372 relatively high DIN concentrations to saturate growth while the opposite is true for DIP. The
373 half-saturation constant (K_{μ}) for growth on PO_4^{3-} and the concentration of DIP needed to
374 saturate growth were low and comparable to those of more slow-growing algae such as *Fucus*
375 *vesiculosus* and *F. serratus*, which could give *G. vermiculophylla* an advantage under low P
376 availability. P-limitation is however rare in most temperate estuaries with terrigenous
377 sediments, where N-limitation is more common (e.g. Howarth and Marino 2006). Nejrup and
378 Pedersen (2010) showed that the tissue N:P ratio in *G. vermiculophylla* from Holckenhavn
379 Fjord and Fyns Hoved reached 40-45 for a short period in early spring, thus inferring P-
380 limitation. The absolute P-content in these algae was however relatively high (>0.2 % of DW)
381 and experimental fertilization did not stimulate growth. Slow growth in early spring seems
382 thus to be dictated by low water temperature and insolation rather than by P-limitation and
383 low P-requirements do therefore not seem to represent a clear advantage for *G.*
384 *vermiculophylla*. In summary, we found no clear evidence that the nutrient kinetics of *G.*
385 *vermiculophylla* result in higher nutrient use efficiency than in comparable, sympatric
386 indigenous algal species and therefore, no indication that *G. vermiculophylla* should have a
387 specific advantage relative to native species under nutrient replete conditions.

388 Nutrient storage plays an essential role for seaweeds in areas where nutrient
389 availability undergoes large seasonal variations. Little is known about the forms in which
390 excess P is stored in macroalgae, but N is mainly stored as proteins and amino acids while
391 inorganic N and N bound in pigments may constitute smaller and less important N reserves
392 (Bird et al. 1982; McGlathery et al. 1996). Although some amino acids play essential roles for
393 osmoregulation (e.g. proline; Kirst 1989) and photoprotection (e.g. mycosporine-like amino
394 acids; Karsten et al. 2000) there is a strong correlation between total N content and the pool
395 sizes of proteins, amino acids, inorganic N and pigment bound N, which shows that all these

396 pools take part in N storage and can be mobilized when necessary (Bird et al. 1982;
397 McGlathery et al. 1996).

398 The amount of N and P stored in excess of that needed to support the maximum
399 growth rate in *G. vermiculophylla* was ca. 14 mg N g⁻¹ DW and 1.3 mg P g⁻¹ DW,
400 respectively. The observed N-reserve was somewhat less than for a range of indigenous algal
401 species (Pedersen and Borum 1996; Pedersen et al. 2010), which suggests that the algae used
402 in our experiments may not have been saturated with nutrients. The storage capacity for
403 nutrients (defined as the time where growth could be sustained without additional nutrient
404 uptake from the medium) was ca. 33 days for N and 43 days for P, respectively. The storage
405 capacity for N was larger for a number of fast-growing indigenous algae (*U. lactuca*, *C.*
406 *virgatum*, *Cladophora* sp. and *Chaetomorpha linum*: range from 9.6 to 17.5 days), but smaller
407 than for slow-growing *F. vesiculosus* (45.9 days; Pedersen and Borum 1996).

408 The N and P-reserves obtained through winter and spring can obviously not sustain
409 maximum growth of *G. vermiculophylla* throughout extended periods of low nutrient
410 availability (i.e. mid May to mid September). However, even though *in situ* nutrient
411 concentrations in Holckenhavn Fjord and Fyns Hoved were low (Nejrup and Pedersen 2010),
412 they were not low enough to prevent nutrient uptake from the water completely. The observed
413 concentrations of DIN ranged between 1 and 3 µM from mid May to mid September and
414 should be high enough to cover about 50% of the N-demand for maximum growth, meaning
415 that the N reserves could last for about twice as long as estimated assuming no uptake (i.e. ca.
416 2 months). Equally important, average *in situ* growth rates in Holckenhavn Fjord during the
417 growing season were some what lower than those attained in this study (ca. 0.05 d⁻¹, range
418 0.032 – 0.065 d⁻¹; Nejrup and Pedersen 2010), meaning that ‘dilution’ of the internal nutrient
419 reserves would occur more slowly and the stores last longer than predicted when assuming
420 growth at maximum attainable rates. In other words, the combined effect of continuous

421 nutrient uptake (albeit at reduced rates) and slower growth (i.e. lower nutrient demands per
422 unit biomass and time) increases the storage capacity of the algae by a factor of ca. 4, i.e. to
423 130-160 days, which should be enough to sustain non-nutrient limited growth during most of
424 the major growth season. This conclusion is supported by the fact that Nejrup and Pedersen
425 (2010) were unable to stimulate *in situ* growth significantly by experimental nutrient
426 enrichment and by the fact that the N content in algae from the control treatment in
427 Holckenhavn Fjord and Fyns Hoved remained close to ca. 2.5% of DW (i.e. above the critical
428 N content) throughout summer. Relatively fast and non-nutrient limited growth by *G.*
429 *vermiculophylla* during summer can thus be explained by the use of reserve N (and P) rather
430 than by an extraordinarily high affinity for inorganic nutrients.

431 The large biomass of *G. vermiculophylla* found in some estuaries, e.g. >1000 g DW
432 m⁻² in some samples from Hog Island Bay, Virginia, USA (Thomsen et al. 2006), 464 g DW
433 m⁻² on average in Holckenhavn Fjord, Denmark (Nejrup and Pedersen 2010) and 6-700 g DW
434 m⁻² in parts of Venice lagoon, Italy (Sfriso et al. 2012) suggests that large amounts of DIN
435 and DIP are assimilated and become bound in living or dead biomass and, thus, that *G.*
436 *vermiculophylla* has the potential to affect nutrient cycling significantly in such systems (e.g.
437 Tyler et al. 2003; Tyler and McGlathery 2006). Like for other bloom-forming algae, the
438 biomass of *G. vermiculophylla* might undergo large and quick temporal variations where
439 algae suffer high mortality and the biomass is turned into detritus that decomposes under
440 release of nutrients. Whether detritus originating from *G. vermiculophylla* serves as a
441 temporary sink for nutrients during summer or functions as an internal source of nutrients
442 within the system depends on: 1) the turn-over of biomass and, thus, the production of detritus
443 and, 2) the rate at which that detritus decays and the bound nutrients become mineralized.

444 We have no direct estimates of the biomass turnover of *G. vermiculophylla*, but data
445 on seasonal changes in biomass and *in situ* growth rates from Holckenhavn Fjord, from

446 Nejrup and Pedersen (2010) show that the potential production of *G. vermiculophylla* may
447 reach ca. 5200 g DW m⁻² during the growing season (from March to October) corresponding
448 to a net incorporation of ca. 78 g N and ca. 10 g P m⁻² (assuming a mean N and P content of
449 1.5% of DW and 0.2% of DW, respectively). The average load of total nitrogen (TN) and
450 total phosphorus (TP) per unit of estuarine area across 47 Danish estuaries is ca. 20 g TN m⁻²
451 y⁻¹ and 0.8 g TP m⁻² y⁻¹ (Conley et al. 2000) so *G. vermiculophylla* may have a substantial
452 effect on nutrient cycling in Holckenhavn Fjord. The biomass of *G. vermiculophylla* changed
453 little over the season (ranging from 464 g DW m⁻² in March to 176 g DW m⁻² in October), so
454 biomass losses and, hence, the production of detritus must have been in the same order of
455 magnitude as the production of biomass (estimated to ca. 5400 g DW m⁻² in 7 months) since
456 grazing on *G. vermiculophylla* is insignificant (Nejrup and Pedersen 2010). Large amounts of
457 nutrient are thus being acquired and bound into living and dead algal biomass and will
458 become released when the detritus decomposes. Decomposition rates of *G. vermiculophylla*
459 ranged from 0.013 to 0.065 d⁻¹ depending on temperature and were markedly lower than those
460 reported by Conover et al. (2016). Conover et al. (2016) used dead tissues killed by freezing
461 which were placed on top of the sediment (i.e. under aerobic conditions) whereas we used live
462 material that was slightly covered by sediment (i.e. partly hypoxic conditions). Freezing may
463 break the cell membrane and wall and cause a rapid loss of soluble compounds from the
464 detritus and, thus, speed up decomposition. The use of live material may, in contrast, delay
465 initial decay and the initial loss of biomass may be due to respiration rather than decay. Dark
466 respiration rate in *G. vermiculophylla* corresponds, however, to a much smaller biomass loss
467 than that observed in our decomposition study (e.g. <0.003 d⁻¹; Nejrup et al. 2013) suggesting
468 that the biomass losses we observed were caused by decomposition mainly.

469 Decomposition of *G. vermiculophylla* was strongly temperature dependent;
470 decomposition was 4-5 fold faster at 25°C than at 5°C, corresponding to a half-time of about

471 10 days under late summer conditions suggesting that nutrients bound in detritus generated
472 during summer and early autumn are released almost completely within the time-scale of
473 weeks whereas it takes much longer during winter and early spring. An increasing algal
474 biomass in early spring and subsequent production of detritus during early summer may thus
475 immobilize bio-available nutrients whereas the opposite will be true in late summer and early
476 autumn where the production of detritus increases and dead *G. vermiculophylla* decomposes
477 fast and, thus, will act as a major source of nutrients. Nutrients released from living and dead
478 *G. vermiculophylla* may not only be recycled back to live and nutrient deplete *Gracilaria* in
479 the upper layers of the mats, but may also be taken up by other primary producers including
480 phytoplankton and microphytobenthos or diffuse into the sediment where it can take part in
481 biogeochemical processes (Hardison et al. 2010; Gulbransen and McGlathery 2013).

482 Decomposition rate for *G. vermiculophylla* at 15°C were obtained under the same
483 conditions as rates for other estuarine macroalgae reported by Banta et al. (2004) and
484 Pedersen et al. (2005), which allows for a direct comparison with these data. Detritus from *G.*
485 *vermiculophylla* decomposed completely within 120 days at 15°C. The decay rate (0.032 d^{-1})
486 was lower than those of *Ulva lactuca* and *Ceramium virgatum* ($0.038 - 0.040\text{ d}^{-1}$) and similar
487 to that of *Fucus vesiculosus* (0.028 d^{-1} ; Banta et al. 2004). *F. vesiculosus* contains however
488 refractory compounds, which will leave ca. 7% of the initial biomass after 340 days (Banta et
489 al. 2004). Detritus from *G. vermiculophylla* decomposed markedly faster than that from
490 *Sargassum muticum* (0.016 d^{-1}) and *Halidrys siliquosa* (0.019 d^{-1}), respectively (Pedersen et
491 al. 2005). The release of nutrients through decomposition and, thus, the potential impact of *G.*
492 *vermiculophylla* on local nutrient cycling depends therefore partly on which species it has
493 succeeded. Nutrient cycling may be slowed down when fast-growing species like *Ulva* sp.
494 and *Ceramium* sp. are replaced by *Gracilaria* because the former are grazed more and their
495 detritus decomposes faster than for *G. vermiculophylla*. Dominance by *G. vermiculophylla*

496 will thus tend cause a slower turnover of nutrients and the biomass of living and dead
497 *Gracilaria* may act as a temporary sink for nutrients. The opposite is expected to be the case
498 if *G. vermiculophylla* succeeds slow-growing species such as *Fucus* sp. *Halidrys siliquosa*
499 etc., which are less susceptible to grazing and produces detritus that decomposes slowly
500 and/or incompletely. Here dominance of *G. vermiculophylla* will tend to speed up the
501 turnover of nutrients due to its relatively fast and complete decomposition.
502

503 **References**

- 504 Abreu MH, Pereira R, Buschmann AH, Sousa-Pinto I, Yarish C (2011) Nitrogen uptake
505 responses of *Gracilaria vermiculophylla* (Ohmi) Papenfuss under combined and single
506 addition of nitrate and ammonium. *J Exp Mar Biol Ecol* 407: 190–199.
- 507 Banta GT, Pedersen MF, Nielsen SL (2004) Decomposition of marine primary producers:
508 Consequences for nutrient recycling and retention in coastal ecosystems. In: GT Banta,
509 MF Pedersen, SL Nielsen (eds.), *Estuarine nutrient cycling: The influence of primary*
510 *producers: The fate of nutrients and biomass* (pp. 187-216). Kluwer Academic Publishers,
511 Dordrecht/Boston/London.
- 512 Bird KT, Habig C, DeBusk T (1982) Nitrogen allocation and storage patterns in *Gracilaria*
513 *tikvahiae* (Rhodophyta). *J Phycol* 18: 344-348.
- 514 Bokn TL, Duarte CM, Pedersen MF, Marba N, Moy FE, Barrón C, Bjerkeng B, Borum J,
515 Christie H, Fotel FL, Hoell EE, Karez R, Kersting K, Kraufvelin P, Lindblad C, Olsen M,
516 Sommer U, Sørensen K (2003) The response of experimental rocky shore communities to
517 nutrient additions: A synthesis of the EULIT experiment. *Ecosystems* 6: 577-594.
- 518 Bower CE, Holm-Hansen T (1980) A salicylate-hypochlorite method for determining
519 ammonia in seawater. *Can J Fish Aquat Sci* 37: 794-798.
- 520 Buchsbaum R, Valiela I, Swain T, Dzierzeski M, Allen S (1991) Available and refractory
521 nitrogen in detritus of coastal vascular plants and macroalgae. *Mar Ecol Prog Ser* 72: 131–
522 143.
- 523 Byers JE, Gribben PE, Yeager C, Sotka EE (2012) Impacts of an abundant introduced
524 ecosystem engineer within mudflats in the southeastern US coast. *Biol Invasions* 14:
525 2587- 2600.
- 526 Conley DJ, Kaas H, Møhlenberg F, Rasmussen B, Windolf J (2000) Characteristics of Danish
527 Estuaries. *Estuaries* 23: 820-837.

528 Duarte CM, Cebrian J (1996) The fate of marine autotrophic production. *Limnol Oceanogr*
529 41: 1758-1766.

530 Enriquez S, Duarte CM, Sand-Jensen K (1993) Patterns in decomposition rates among
531 photosynthetic organisms: the importance of detritus C:N:P content. *Oecologia* 94: 457–
532 471.

533 Gulbransen D, McGlathery K (2013) Nitrogen transfers mediated by perennial, non-native
534 macroalga: a ¹⁵N tracer study. *Mar Ecol Prog Ser* 482: 299-304.

535 Hammann M, Wang G, Rickert E, Boo SM, Weinberger F (2013) Invasion success of the
536 seaweed *Gracilaria vermiculophylla* correlates with low palatability. *Mar Ecol Prog Ser*
537 486: 93-103.

538 Hammann M, Rempt M, Pohnert G, Wang G, Boo SM, Weinberger F (2016) Increased
539 potential for wound activated production of Prostaglandin E-2 and related toxic
540 compounds in non-native populations of *Gracilaria vermiculophylla*. *Harmful Algae* 51:
541 81-88.

542 Hardison A, Canuel E, Anderson I, Veuger B (2010) Fate of macroalgae in benthic systems:
543 carbon and nitrogen cycling within the microbial community. *Mar Ecol Prog Ser* 414: 41–
544 55.

545 Harrison PJ, Parslow JS, Conway HL (1989) Determination of nutrient uptake kinetic
546 parameters: a comparison of methods. *Mar Ecol Prog Ser* 52: 301-312.

547 Healey FP (1980) Slope of the Monod equation as an indicator of advantage in nutrient
548 competition. *Microb Ecol* 5: 281-286.

549 Hein M, Pedersen MF, Sand-Jensen K (1995). Size-dependent nitrogen uptake in micro-and
550 macroalgae. *Mar Ecol Prog Ser* 118: 247-253.

551 Howarth RW, Marino R (2006) Nitrogen is the limiting nutrient for eutrophication in coastal
552 marine ecosystems: evolving views over three decades. *Limnol Oceanogr* 51: 364–376.

553 Hurd CL, Harrison PJ, Bischof K, Lobban CS (2014) Seaweed ecology and physiology.
554 Cambridge University Press. Cambridge, UK.

555 Karez R, Engelbert S, Kraufvelin P, Pedersen MF, Sommer U (2004) Biomass response and
556 changes in composition of ephemeral macroalgal assemblages along an experimental
557 gradient of nutrient enrichment. *Aquat Bot* 78: 103-117.

558 Karsten U, Sawall T, West J, Wiencke C (2000) Ultraviolet sunscreen compounds in
559 epiphytic red algae from mangroves. *Hydrobiologia* 432: 159-171.

560 Kirst GO (1989) Salinity tolerance of eukaryotic marine algae. *Annu Rev Plant Phys* 41: 21–
561 53.

562 Kraufvelin P, Moy FE, Christie H, Bokn TL (2006) Nutrient addition to experimental rocky
563 shore communities revisited: delayed responses, rapid recovery. *Ecosystems* 9: 1076–
564 1093.

565 Kraufvelin P, Lindholm A, Pedersen MF, Kirkerud LA, Bonsdorff E (2010) Biomass,
566 diversity and production of rocky shore macroalgae at two nutrient enrichment and wave
567 action levels. *Mar Biol* 157: 29-47.

568 McGlathery KJ, Pedersen MF, Borum J (1996) Changes in intracellular nitrogen pools and
569 feedback controls on nitrogen uptake in *Chaetomorpha linum* (Chlorophyta). *J Phycol* 32:
570 393-401.

571 Mollet J, Rahaoui A, Lemoine Y (1998) Yield, chemical composition and gel strength of
572 agarocolloids of *Gracilaria gracilis*, *Gracilariopsis longissima* and the newly reported
573 *Gracilaria cf. vermiculophylla* from Roscoff (Brittany, France). *J Appl Phycol* 10: 59–66.

574 Nejrup LB, Pedersen MF (2010) Growth and biomass development of the introduced red alga
575 *Gracilaria vermiculophylla* is unaffected by nutrient limitation and grazing. *Aquat Biol*
576 10: 249-259.

577 Nejrup LB, Pedersen MF (2012) The effect of temporal variability in salinity on the invasive
578 red alga *Gracilaria vermiculophylla*. *Europ J Phycol* 47: 254-263.

579 Nejrup LB, Pedersen MF, Vinzent J (2012) Grazer avoidance may explain the invasiveness of
580 the red alga *Gracilaria vermiculophylla* in Scandinavian waters. *Mar Biol* 159: 1703–
581 1712.

582 Nejrup LB, Staehr PA, Thomsen MS (2013) Temperature- and light-dependent growth and
583 metabolism of the invasive red algae *Gracilaria vermiculophylla* – a comparison with two
584 native macroalgae. *Europ J Phycol* 48: 295–308.

585 Nyberg CD, Thomsen MS, Wallentinus I (2009) Flora and fauna associated with the
586 introduced red alga *Gracilaria vermiculophylla*. *Europ J Phycol* 44: 395–403.

587 Nylund GM, Weinberger F, Rempt M, Pohnert G (2011) Metabolomic assessment of induced
588 and activated chemical defence in the invasive red alga *Gracilaria vermiculophylla*. *PLoS*
589 *One* 6: e29359.

590 Pedersen MF (1994) Transient ammonium uptake in the macroalga *Ulva lactuca*
591 (Chlorophyta): Nature, regulation, and the consequences for choice of measuring
592 technique. *J Phycol* 30: 980-986.

593 Pedersen MF, Borum J (1996) Nutrient control of algal growth in estuarine waters: Nutrient
594 limitation and the importance of nitrogen requirements and nitrogen storage among
595 phytoplankton and species of macroalgae. *Mar Ecol Prog Ser* 142: 261-272.

596 Pedersen MF, Borum J (1997) Nutrient control of estuarine macroalgae: growth strategy and
597 the balance between nitrogen requirements and uptake. *Mar Ecol Prog Ser* 161: 155-163.

598 Pedersen MF, Nielsen SL, Banta G (2004) Interactions between vegetation and nutrient
599 dynamics in coastal marine ecosystems: an introduction. In: GT Banta, MF Pedersen, SL
600 Nielsen (eds.), *Estuarine nutrient cycling: The influence of primary producers: The fate of*
601 *nutrients and biomass* (pp. 1-15). Kluwer Academic Publishers, Dordrecht/Boston/London.

602 Pedersen MF, Borum J, Fotel FL (2010) Phosphorus dynamics and limitation of fast-and
603 slow-growing temperate seaweeds in Oslofjord, Norway. *Mar Ecol Prog Ser* 399: 103-115.

604 Pedersen MF, Staehr PA, Wernberg T, Thomsen MS (2005) Biomass dynamics of exotic
605 *Sargassum muticum* and native *Halidrys siliquosa* in Limfjorden, Denmark-Implications
606 of species replacements on turnover rates. *Aquat Bot* 83: 31-47.

607 Raikar SV, Iima M, Fujita Y (2001) Effect of temperature, salinity and light intensity on the
608 growth of *Gracilaria* spp. (Gracilariales, Rhodophyta) from Japan, Malaysia and India.
609 *Indian J Mar Sci* 30: 98-104.

610 Rosenberg G, Probyn TA, Mann KH (1984) Nutrient uptake and growth kinetics in brown
611 seaweeds: Response to continuous and single additions of ammonium. *J Exp Mar Biol Ecol*
612 80: 125-146.

613 Rueness, J. (2005). Life history and molecular sequences of *Gracilaria vermiculophylla*
614 (Gracilariales, Rhodophyta), a new introduction to European waters. *Phycologia* 44: 120-
615 128.

616 Sfriso A, Wolf MA, Maistro S, Sciuto K, Moro I (2012) Spreading and autoecology of the
617 invasive species *Gracilaria vermiculophylla* (Gracilariales, Rhodophyta) in the lagoons of
618 the north-western Adriatic Sea (Mediterranean Sea, Italy). *Est Coast Shelf Sci* 114: 192-
619 198.

620 Strickland JD, Parsons TR (1968) A practical handbook of seawater analysis. *Bull Fish Res*
621 *Board Can.* No 167.

622 Thomsen MS, McGlathery KJ, Tyler AC (2006) Macroalgal distribution in a shallow, soft
623 bottom lagoon, with emphasis on the non-native *Gracilaria vermiculophylla* and *Codium*
624 *fragile*. *Est Coast* 29: 465-473.

625 Thomsen MS, Staehr PA, Nyberg CD, Schwärter S, Krause-Jensen D, Silliman BR (2007)
626 *Gracilaria vermiculophylla* (Ohmi) Papenfuss, 1967 (Rhodophyta, Gracilariaceae) in

627 northern Europe, with emphasis on Danish conditions, and what to expect in the future.
628 Aquat Invasions 2: 83-94.

629 Thomsen MS, McGlathery KJ (2007) Stress tolerance of the invasive macroalgae *Codium*
630 *fragile* and *Gracilaria vermiculophylla* in a soft-bottom turbid lagoon. Biol Invasions 9:
631 499–513.

632 Tseng CK, Xia BM (1999) On the *Gracilaria* in the Western Pacific and the Southeastern Asia
633 Region. Bot Mar 42: 209-217.

634 Turpin DH (1988) Physiological mechanisms in phytoplankton resource competition. Chapter
635 8 in Sandgren CD (ed) Growth and reproductive strategies of freshwater phytoplankton.
636 Cambridge University Press. Cambridge, UK. pp. 316-368.

637 Tyler AC, McGlathery KJ, Macko SA (2005) Uptake of urea and amino acids by the
638 macroalgae *Ulva lactuca* (Chlorophyta) and *Gracilaria vermiculophylla* (Rhodophyta).
639 Mar Ecol Prog Ser 294: 161-172.

640 Tyler AC, McGlathery KJ (2003) Benthic algae control sediment-water column fluxes of
641 organic and inorganic nitrogen compounds in a temperate lagoon. Limnol Oceanogr 48:
642 2125-2137.

643 Tyler AC, McGlathery KJ (2006) Uptake and release of nitrogen by the macroalgae
644 *Gracilaria vermiculophylla* (Rhodophyta). J Phycol 42: 515–525.

645 Wallentinus I (1984) Comparisons of nutrient uptake rates for Baltic macroalgae with
646 different thallus morphologies. Mar Biol 80:215–225.

647 Wallentinus I, Nyberg CD (2007) Introduced marine organisms as habitat modifiers. Mar.
648 Pollut Bull 55: 323–32.

649 Weinberger F, Buchholz B, Karez R, Wahl M (2008) The invasive red alga *Gracilaria*
650 *vermiculophylla* in the Baltic Sea: adaptation to brackish water may compensate for light
651 limitation. Aquat Biol 3: 251-264.

652 Westrich JT, Berner RA (1984) The role of sedimentary organic matter in bacterial sulfate
653 reduction: The G model tested. *Limnol Oceanogr* 29: 236–249.
654 Yokoya NS, Kakita H, Obika H, Kitamura T (1999) Effects of environmental factors and
655 plant growth regulators on growth of the red alga *Gracilaria vermiculophylla* from
656 Shikoku Island, Japan. *Hydrobiologia*, 398/399, 339-347.

657 **Figure legends**

658 **Fig. 1** *Gracilaria vermiculophylla*. Uptake rates (V) of ammonium, nitrate and phosphate as a
659 function of substrate concentration (S). Surge uptake (●) was measured over the initial 30
660 minutes after exposure to nutrients, while assimilation rates (○) were measured later than 30
661 minutes after exposure. Lines represent the best fits of the Michalis-Menten function to the
662 data

663


664 **Fig. 2** *Gracilaria vermiculophylla*. Relationship between relative growth rate (μ) and N
665 quota (figure A) or P quota (figure B) as determined in laboratory experiments. Lines
666 represent the best fits of the Droop equation to the data

667



668 **Fig. 3** *Gracilaria vermiculophylla*. Changes in C, N and P biomass during decomposition of
669 dead *Gracilaria vermiculophylla* at three different temperatures. A: 5°C, B: 15°C and C:
670 25°C. Lines represent the best fits to an exponential function. Values are means \pm 95%
671 confidence limits (n=3)

672

Figure 1.




675 **Figure 2.**



676

677

678 **Figure 3.**



679

680

Table 1. Nutrient (NH_4^+ , NO_3^- and PO_4^{3-}) uptake kinetics. Parameters V_{max} ($\mu\text{mol N or P g}^{-1} \text{ DW h}^{-1}$), K_m (μM) and affinity for uptake at low substrate concentrations (V_{max}/K_m ; Healy 1980) are given for both initial surge uptake measured over the first 30 minutes of exposure to nutrients and for uptake and assimilation measured later than 30 minutes after exposure to nutrients. Numbers are means $\pm 95\%$ confidence limits.

	Surge uptake (0-30 min.)			Assimilation (30-300 min.)		
	V_{max} ($\mu\text{mol g}^{-1} \text{ DW h}^{-1}$)	K_m (μM)	V_{max}/K_m	V_{max} ($\mu\text{mol g}^{-1} \text{ DW h}^{-1}$)	K_m (μM)	V_{max}/K_m
NH_4^+	76.8 \pm 24.5	28.7 \pm 19.7	2.68	32.3 \pm 4.4	8.9 \pm 3.2	3.63
NO_3^-	18.3 \pm 5.6	10.4 \pm 12.4	1.76	13.4 \pm 1.3	8.3 \pm 2.3	1.61
PO_4^{3-}	2.47 \pm 1.01	5.5 \pm 5.1	0.45	1.0 \pm 0.2	2.4 \pm 1.0	0.42

681

682

683

684

Table 2. Parameter estimates of maximum growth rate (μ_{max}) and subsistence cell quota (Q_S) as determined from fitting the Droop function to data (i.e. growth rates vs. cell quota). Estimated means $\pm 95\%$ confidence limits. Critical cell quotas (Q_C) were estimated as the quota where growth rate equalled 66.7% of μ_{max} while maximum cell quotas (Q_{max}) correspond to the highest quotas obtained in the experiment.

	μ_{max} (d^{-1})	Q_S (% of DW)	Q_C (% of DW)	Q_{max} (% of DW)	Requirement ($\mu\text{mol g}^{-1} \text{ DW d}^{-1}$)
Nitrogen	0.105 \pm 0.016	0.706 \pm 0.098	2.14	3.54	169.2
Phosphorus	0.088 \pm 0.013	0.045 \pm 0.007	0.14	0.27	4.1

685

Table 3. Modelled substrate dependent growth kinetics. Monod parameters (μ_{max}^* and K_μ) and the substrate concentration at which 0.9 of the maximum growth rate is obtained (S_{sat}).

	μ_{max}^* (d ⁻¹)	K_μ (μ M)	μ_{max}/K_μ	S_{sat} (μ M)
Ammonium	0.098	0.57	0.17	5.1
Nitrate	0.091	1.17	0.08	10.4
Phosphorus	0.084	0.12	0.70	1.03

Table 4. Storage capacity ($T_{Storage}$) for N and P, respectively, in *Gracilaria vermiculophylla*.

	$T_{Storage}$ at max. growth rate (days)	$T_{Storage}$ at reduced growth rate (days)	$T_{Storage}$ total (days)
Storage capacity for N	6.5	26.5	33.0
Storage capacity for P	10.5	32.5	43.0

687

688

689

690

691

692

693

694

695

Table 5. Parameter estimates (Y_0 : intercept with Y-axis; k : decay rate and, $T_{0.5}$: half time) for decomposition of C biomass and mineralization of tissue bound N and P in *Gracilaria vermiculophylla*. Estimated means \pm 95% confidence limits. Decomposition experiments were conducted at 5°C (A), 15°C (B) and 25°C (C), respectively.

	Y_0 (%)	k (d ⁻¹)	$T_{0.5}$ (days)	R^2
C at 5°C	103.4 \pm 3.4	0.013 \pm 0.002	52.9 \pm 8.9	0.895
C at 15°C	107.2 \pm 14.0	0.032 \pm 0.007	21.5 \pm 4.1	0.857
C at 25°C	94.6 \pm 17.3	0.065 \pm 0.023	10.7 \pm 2.8	0.774
N at 5°C	99.5 \pm 9.5	0.016 \pm 0.004	44.6 \pm 7.9	0.877
N at 15°C	109.5 \pm 15.1	0.037 \pm 0.009	18.8 \pm 3.8	0.842
N at 25°C	99.9 \pm 9.8	0.079 \pm 0.015	12.7 \pm 5.3	0.926
P at 5°C	94.4 \pm 14.5	0.023 \pm 0.008	30.5 \pm 7.8	0.780
P at 15°C	109.8 \pm 17.8	0.043 \pm 0.012	16.2 \pm 3.7	0.805
P at 25°C	98.8 \pm 11.2	0.100 \pm 0.022	6.9 \pm 1.2	0.908

Table 6. Maximum relative growth rate (μ_{max}), critical N and P quotas (N_C and P_C), half-saturation constants for growth (K_μ) and saturating substrate concentrations (S_{sat}) for *Gracilaria vermiculophylla* and eleven common algal species native to Scandinavian waters.

Species	μ_{max} (d ⁻¹)	Critical quota		NH ₄ ⁺		NO ₃ ⁻		PO ₄ ³⁻	
		N _C (%DW)	P _C	K _μ (μM)	S _{sat}	K _μ (μM)	S _{sat}	K _μ (μM)	S _{sat}
<i>Ulva lactuca</i> ^{1,2,3}	0.196 - 0.452	2.17	0.20	0.75	6.71	1.45	13.02	0.27	2.39
<i>Ceramium virgatum</i> ^{1,2,3}	0.136 - 0.299	3.10	0.44	0.63	5.65	-	-	0.22	1.94
<i>Cladophora sp.</i> ^{1,2}	0.188 - 0.208	2.05	-	0.36	3.28	0.83	7.45	-	-
<i>Chaetomorpha linum</i> ^{1,2}	0.139 - 0.142	1.15	-	0.11	1.00	0.13	1.18	-	-
<i>Gracilaria vermiculophylla</i>	0.084 - 0.098	2.14	0.14	0.57	5.10	1.17	10.40	0.12	1.03
<i>Chordaria flagelliformis</i> ⁴	0.091 - 0.126			0.31					
<i>Codium fragile</i> ^{1,2}	0.074 - 0.083	1.58	-	0.25	2.22	0.57	5.11	-	-
<i>Fucus disticus</i> ⁴	0.067 - 0.081			0.24					
<i>Fucus vesiculosus</i> ^{1,2,3}	0.038 - 0.040	1.71	0.12	0.29	2.66	0.81	7.32	0.09	0.78
<i>Fucus serratus</i> ³	0.040	-	0.22	-	-	-	-	0.16	0.88
<i>Ascophyllum nodosum</i> ³	0.014	-	0.15	-	-	-	-	0.06	0.35
<i>Laminaria digitata</i> ³	0.006	-	0.22	-	-	-	-	0.03	0.15

Data sources:
¹⁾ Pedersen and Borum 1996
²⁾ Pedersen and Borum 1997
³⁾ Pedersen et al. 2010
⁴⁾ Rosenberg et al. 1984