

Anti-staphylococcal activity of *Syagrus coronata* essential oil

Biofilm eradication and in vivo action on *Galleria mellonella* infection model

dos Santos, Bruno Souza; Macedo Bezerra Filho, Clóvis; Adelson Alves do Nascimento Junior, José; Brust, Flávia Roberta; Bezerra-Silva, Patrícia Cristina; da Rocha, Suyana Karoline Lino; Krogfelt, K. A.; do Amaral Ferraz Navarro, Daniela Maria; dos Santos Correia, Maria Tereza; Napoleão, Thiago Henrique; da Silva, Luís Claudio Nascimento; Macedo, Alexandre José; da Silva, Márcia Vanusa; Guedes Paiva, Patrícia Maria

Published in:
Microbial Pathogenesis

DOI:
[10.1016/j.micpath.2019.04.009](https://doi.org/10.1016/j.micpath.2019.04.009)

Publication date:
2019

Document Version
Peer reviewed version

Citation for published version (APA):

dos Santos, B. S., Macedo Bezerra Filho, C., Adelson Alves do Nascimento Junior, J., Brust, F. R., Bezerra-Silva, P. C., da Rocha, S. K. L., Krogfelt, K. A., do Amaral Ferraz Navarro, D. M., dos Santos Correia, M. T., Napoleão, T. H., da Silva, L. C. N., Macedo, A. J., da Silva, M. V., & Guedes Paiva, P. M. (2019). Anti-staphylococcal activity of *Syagrus coronata* essential oil: Biofilm eradication and *in vivo* action on *Galleria mellonella* infection model. *Microbial Pathogenesis*, 131, 150-157. <https://doi.org/10.1016/j.micpath.2019.04.009>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@ruc.dk providing details, and we will remove access to the work immediately and investigate your claim.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/332754122>

Anti-staphylococcal activity of *Syagrus coronata* essential oil: Biofilm eradication and in vivo action on *Galleria mellonella* infection model

Article in *Microbial Pathogenesis* · April 2019

DOI: 10.1016/j.micpath.2019.04.009

CITATIONS

4

READS

96

14 authors, including:



Clovis Macêdo

Universidade Federal de São Paulo

15 PUBLICATIONS 67 CITATIONS

[SEE PROFILE](#)



José Adelson Alves do Nascimento Junior

2 PUBLICATIONS 9 CITATIONS

[SEE PROFILE](#)



KA Krogfelt

Roskilde University & Statens Serum Institut

543 PUBLICATIONS 14,088 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Project

Ecofisiologia de cultivares de *Opuntia ficus-indica* Mill (Cactaceae) de tolerância contrastante à cochonilha-do-carmim *Dactylopius opuntiae* [View project](#)



Project

Diagnostic Methods in Leptospirosis [View project](#)

Accepted Manuscript



Anti-staphylococcal activity of *Syagrus coronata* essential oil: Biofilm eradication and *in vivo* action on *Galleria mellonella* infection model

Bruno Souza dos Santos, Clóvis Macedo Bezerra Filho, José Adelson Alves do Nascimento Junior, Flávia Roberta Brust, Patrícia Cristina Bezerra-Silva, Suyana Karoline Lino da Rocha, Karen Angeliki Krogfelt, Daniela Maria do Amaral Ferraz Navarro, Maria Tereza dos Santos Correia, Thiago Henrique Napoleão, Luís Claudio Nascimento da Silva, Alexandre José Macedo, Márcia Vanusa da Silva, Patrícia Maria Guedes Paiva

PII: S0882-4010(18)31395-0

DOI: <https://doi.org/10.1016/j.micpath.2019.04.009>

Reference: YMPAT 3481

To appear in: *Microbial Pathogenesis*

Received Date: 7 August 2018

Revised Date: 4 April 2019

Accepted Date: 5 April 2019

Please cite this article as: Souza dos Santos B, Bezerra Filho ClóMacedo, Alves do Nascimento Junior JoséAdelson, Brust FláRoberta, Bezerra-Silva PatríCristina, Lino da Rocha SK, Krogfelt KA, Maria do Amaral Ferraz Navarro D, Tereza dos Santos Correia M, Napoleão TH, Nascimento da Silva LuíClaudio, Macedo AlexandreJosé, Vanusa da Silva Má, Guedes Paiva PatríMaria, Anti-staphylococcal activity of *Syagrus coronata* essential oil: Biofilm eradication and *in vivo* action on *Galleria mellonella* infection model, *Microbial Pathogenesis* (2019), doi: <https://doi.org/10.1016/j.micpath.2019.04.009>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Anti-staphylococcal activity of *Syagrus coronata* essential oil: biofilm eradication and *in***
2 ***vivo* action on *Galleria mellonella* infection model**

3

4 Bruno Souza dos Santos^a, Clóvis Macedo Bezerra Filho^a, José Adelson Alves do Nascimento
5 Junior^a, Flávia Roberta Brust^b, Patrícia Cristina Bezerra-Silva^c, Suyana Karoline Lino da
6 Rocha^c, Karen Angeliki Krogfelt^d, Daniela Maria do Amaral Ferraz Navarro^c; Maria Tereza
7 dos Santos Correia^a, Thiago Henrique Napoleão^a, Luís Claudio Nascimento da Silva^c,
8 Alexandre José Macedo^b, Márcia Vanusa da Silva^a, Patrícia Maria Guedes Paiva^{a,*}

9

10 ^a*Departamento de Bioquímica, Universidade Federal de Pernambuco, Recife, Pernambuco,*
11 *Brazil.*

12 ^b*Faculdade de Farmácia, Universidade Federal do Rio Grande do Sul, Porto Alegre, Rio*
13 *Grande do Sul, Brazil.*

14 ^c*Departamento de Química Fundamental, Universidade Federal de Pernambuco, Recife,*
15 *Pernambuco, Brazil.*

16 ^d*Department of Virology and Microbiological Diagnostics, Statens Serum Institut, Denmark.*

17 ^e*Programa de Pós-Graduação em Biologia Parasitária, Universidade CEUMA, Maranhão,*
18 *Brazil.*

19

20 *Corresponding author. E-mail address: ppaivaufpe@yahoo.com.br

21

22 **Abstract**

23

24 In this study, essential oil extracted from *Syagrus coronata* seeds (SCEO) was evaluated for
25 antibacterial and antibiofilm activities against *Staphylococcus aureus*; in addition, *Galleria*
26 *mellonella* model was used as an *in vivo* infection model. SCEO was mainly composed by
27 fatty acids (89.79%) and sesquiterpenes (8.5%). The major components were octanoic acid,
28 dodecanoic acid, decanoic acid and γ -eudesmol. SCEO showed bactericidal activity (minimal
29 bactericidal concentration from 312 to 1250 $\mu\text{g/mL}$) against all tested *S. aureus* clinical
30 isolates, which showed distinct biofilm-forming and multiple drug resistance phenotypes.
31 SCEO weakly reduced biomass but remarkably decreased cell viability in pre-formed
32 biofilms of *S. aureus* isolate UFPEDA-02 (ATCC-6538). Electron microscopy analysis
33 showed that SCEO treatments decreased the number of bacterial cells (causing structural
34 alterations) and lead to loss of the roughness in the multiple layers of the three-dimensional
35 biofilm structure. In addition, overproduction of exopolymeric matrix was observed. SCEO
36 at 31.2 mg/kg improved the survival of *G. mellonella* larvae inoculated with UFPEDA-02
37 isolate and reduced the bacterial load in hemolymph and melanization. In conclusion, SCEO
38 is an antibacterial agent against *S. aureus* strains with different resistance phenotypes and able
39 to disturb biofilm architecture. Our results show SCEO as a potential candidate to drug
40 development.

41

42 **Keywords:** *Staphylococcus aureus*; antibiofilm; antibacterial activity; volatile oil.

43

44 1. Introduction

45

46 Antimicrobial resistance is one of the most serious public health problems, especially
47 in developing countries where infectious diseases still represent a major cause of human
48 mortality [1]. *Staphylococcus aureus* is highlighted as one of the major human pathogens due
49 to its high ability to produce virulence factors that mediate evasion of immune system and
50 host tissue damage [2–4]. Diseases caused by *S. aureus* involve skin infections (boils,
51 folliculitis, and abscesses) and diseases with greater severity such as pneumonia, meningitis,
52 osteomyelitis, endocarditis, bacteremia, and sepsis [5, 6]. In addition, the widespread and
53 indiscriminate use of antibiotics has caused selective pressure favoring the development of
54 resistant strains, such as methicillin-resistant *S. aureus* (MRSA) and other multidrug resistant
55 phenotypes. MRSA is associated with high rates of morbidity and mortality [5–7].

56 As other bacteria, *S. aureus* often survive by adhering to surfaces on which they form
57 complex structures called biofilms [2]. Biofilms are conglomerates of microbial cells
58 protected by self-synthesized extracellular polysaccharide matrices. Bacterial biofilms are one
59 of the most common causes of persistent infection and represents a major health problem, as it
60 plays an important role in nosocomial infections when formed in internal medical devices
61 such as implanted catheters, artificial heart valves, or bone and joint prostheses [8, 9]. The
62 ability of *S. aureus* to form biofilms in implanted medical devices is an important virulence
63 factor for this pathogen [9]. Biofilm producer strains usually exhibit increased resistance to
64 antibiotics and are responsible for persistent infections [8, 10].

65 The failure of the antibiotics currently used in treating infections caused by multidrug
66 resistant microorganisms has driven the search for new compounds and alternative treatments,
67 particularly those involving plant-derived products such as essential oils, flavonoids and other
68 secondary metabolites [11–13]. Essential oils are mixtures of odoriferous and volatile

69 compounds that have been widely reported as antimicrobial agents [12, 14]. One example of
70 essential oil bearing plant is *Syagrus coronata* (Martius) Beccari (Arecaceae, Arecoideae),
71 popularly known as “licuri” or “ouricuri” [15].

72 *Syagrus coronata* is an edible oil crop known to produce high amount of oils, with
73 potential use for various purposes [16, 17]. In addition, it has a number of applications in folk
74 medicine including snakebites, ocular inflammations, mycoses, wound healing, and spinal
75 pain treatment [18]. Various biological activities have been reported for *S. coronata* seed oil,
76 including antibacterial and insecticidal properties [16, 19]. In addition, it has shown
77 moisturizing property [20].

78 *Galleria mellonella* larvae (waxmoth) is an alternative model that has attracted
79 attention due the methodological simplicity and reliability in the evaluation of infections
80 caused by different human pathogens, in the discovery of new virulence genes, as well as in
81 the evaluation of toxicity and efficacy of antimicrobial agents [21–23].

82 In the present study, *S. coronata* seed essential oil (SCEO) was evaluated for antibiotic
83 and antibiofilm activity against *S. aureus* and *G. mellonella* was used as an *in vivo* infection
84 model.

85

86 **2. Material and methods**

87

88 *2.1. Plant material*

89

90 Seeds of *S. coronata* were collected at the Catimbau National Park region (PARNA do
91 Catimbau, Pernambuco, Brazil - 8° 30' 57" S, 37° 20' 59" W) in December 2015. They were
92 dried at 30 °C in an open and airy area for three weeks. The taxonomic identification of the
93 plant was performed by Dr. Alexandre Gomes da Silva in the herbarium of the *Instituto*

94 *Agronômico de Pernambuco* (IPA). The voucher specimen was deposited under the number
95 86,950. The access was recorded (AFD8A80) in the *Sistema Nacional de Gestão do*
96 *Patrimônio Genético e do Conhecimento Tradicional. Associado* (SisGen).

97

98 2.2. Extraction of *S. coronata* essential oil (SCEO)

99

100 The dried seeds (200 g) were powdered and essential oil was obtained by
101 hydrodistillation method for 4 h in a modified Clevenger-type apparatus. The SCEO layer was
102 separated from the hydrolate (aqueous layer), dried over anhydrous sodium sulfate, and stored
103 in a hermetically sealed amber-glass vial at -20 °C until required for analysis. The percentage
104 yield of essential oil was taken as the ratio between the weight of oil obtained and the weight
105 of seed powder. The whole procedure was repeated 3 times.

106

107 2.3. Gas chromatographic (GC) analyses

108

109 SCEO was esterified by acid catalysis with boron trifluoride (BF₃) [24]. GC was
110 performed using a Thermo Fisher Scientific (Waltham, MA, USA) Trace GC Ultra gas
111 chromatograph equipped with a flame ionization detector (FID), a split/splitless injector and a
112 Hamilton Bonaduz (Switzerland) HB-5 fused silica capillary column (30 m × 0.25 mm; film
113 thickness 0.25 μm). The oven temperature was held at 40 °C for 2 min and then increased at 4
114 °C/min to 230 °C. The injector and detector were both maintained at 250 °C, and the essential
115 oil solution and esterified fractions were injected in the splitless mode. Each analysis was
116 carried out in triplicate.

117

118 GC coupled to mass spectrometry (GC-MS) was carried out using an Agilent
Technologies (Palo Alto, CA, USA) series 5975C quadrupole analyzer equipped with an

119 Agilent J & W nonpolar DB-5 fused silica capillary column (60 m x 0.25 μm i.d.; film
120 thickness 0.25 μm). The oven temperature was held at 60 $^{\circ}\text{C}$ for 3 min, then increased at 2.5
121 $^{\circ}\text{C}/\text{min}$ to 240 $^{\circ}\text{C}$ and subsequently held for 10 min. The helium carrier gas flow was
122 maintained with a constant pressure of 100 kPa, and the injector was operated at 250 $^{\circ}\text{C}$ in the
123 split mode (1:20). The detector temperature was 280 $^{\circ}\text{C}$, the ionization potential was 70 eV,
124 and mass spectra were scanned in the range 20-350 m/z at a rate of 0.5 scans/s [25].

125 Individual components of SCEO were initially identified according their Retention
126 Indices (RI), obtained by co-injection of oil samples and C_8 - C_{30} n-alkanes, calculated
127 according to the equation of van Den Dool and Kratz [26] and compared with the literature
128 [27]. The acquired mass spectra were matched with those stored in the library of the GC-MS
129 system (MassFinder 4 comprising NIST08 MS Library and Wiley Registry of Mass Spectral
130 Data, 9th Edition) and with other published data. The composition of essential oil was
131 expressed as percentages of total peak area as recorded by GC-FID.

132

133 2.4. Phenotypic characterization of *S. aureus* strains

134

135 2.4.1. Antibiotic susceptibility profile

136

137 Twenty *S. aureus* clinical isolates (Table 2) were obtained from the Collection of
138 Microorganisms of the *Departamento de Antibióticos* of the *Universidade Federal de*
139 *Pernambuco* (UFPEDA, WDCM0114), Brazil. *S. aureus* clinical isolates susceptibility was
140 determined according to Kirby Bauer's disk diffusion technique [28] using the antibiotics:
141 oxacillin, ciprofloxacin, nitrofurantoin, amikacin, gentamicin, clindamycin, chloramphenicol,
142 tetracycline, and trimethoprim. Antibiotic susceptibility was interpreted according the Clinical
143 and Laboratory Standards Institute [31].

144 The multiple antibiotic resistance (MAR) index was calculated using the formula
145 $MAR = x/y$, where x is the number of antibiotics to which the isolate demonstrated resistance
146 and y is the total number of antibiotics tested [29].

147

148 2.4.2. Evaluation of *S. aureus* biofilm formation

149

150 Biofilm formation was evaluated and quantified using a microtiterplate test [30].
151 Briefly, it was added 20 μ L of the bacterial suspension (1.5×10^8 CFU/mL), 20 μ L of Milli-Q
152 water and 160 μ L of brain heart infusion broth (BHI) in each well of the plate. After 24 h of
153 incubation at 37 °C, the non-adhered cells were removed, and the biofilm was washed three
154 times with saline solution (0.9% NaCl). Biofilms were heat-fixed at 60 °C for 1 h and then
155 stained with 0.4% (w/v) crystal violet for 15 min at 30 °C. Finally, the plate was washed four
156 times with water and the biofilm was resuspended with ethanol for 30 min. The optical
157 density (OD) was measured at 570 nm. The biofilm production was classified according to
158 Stepanovic et al. [30].

159

160 2.5. Determination of minimal inhibitory (MIC) and bactericidal (MBC) concentrations

161

162 Minimal inhibitory concentration (MIC) were determined by broth microdilution
163 method. Initially, overnight bacterial culture was prepared on Mueller Hinton Agar
164 (MHA) plates. The, a bacterial suspension at 1.5×10^8 CFU/mL was prepared in saline
165 solution (0.9% NaCl). SCEO (0.039–10,000 μ g/mL in 5% dimethyl sulfoxide, DMSO)
166 was serially diluted in microplates containing Mueller Hinton Broth (MHB). Each well
167 received 10 μ L of bacterial suspension, except the wells used as sterility control. In
168 negative control, it was used 5% DMSO. The plates were incubated at 37 °C and, after 24

169 h, wells received 20 μ L of 0.01% (w/v) resazurin solution to follow bacterial growth (i.e.
170 change of blue to pink color). After 24-h incubation, the MIC was defined as the lowest
171 SCEO concentration that inhibited bacterial growth. Suspension from wells before the
172 addition of resazurin were transferred to MHA plates and incubated for other 24 h. MBC
173 was determined as the lowest SCEO concentration able to prevent bacterial growth. The
174 MIC₅₀ and MIC₉₀ were determined as the MIC values that inhibits 50% and 90% of the *S.*
175 *aureus* isolates (n = 20).

176

177 *2.6. Biofilm eradication assays: Quantification of biofilm biomass and viability of biofilm*
178 *cells*

179

180 The biofilm eradication ability of SCEO was evaluated according to Zimmer et al.
181 [32]. For this assay, the *S. aureus* UFPEDA-02 (ATCC-6538) strain was selected due to its
182 source (wound) and high biofilm production ability. The biofilm was formed according
183 previously described and, after 24 h of incubation at 37 °C, planktonic cells were removed
184 and the SCEO diluted in BHI broth was added at different concentrations (156, 312, 624 and
185 1,248 μ g/mL in 5% DMSO). The plate was incubated again at 37 °C and after 24 h the wells
186 were washed three times with saline solution (0.9% NaCl). Adherent biofilms were heat-fixed
187 at 60 °C for 1 h and then stained with 0.4% (w/v) crystal violet for 15 min at 30 °C. Finally,
188 the plate was washed four times with water and the stained biofilm was solubilized in ethanol
189 for 30 min. The absorbance (570 nm) was measured. Vancomycin (1 μ g/mL) was used as
190 antibiotic control.

191 The viability of cells within biofilms exposed SCEO was assessed using MTT (3-(4,5-
192 dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide) assay [33]. Biofilms were grown
193 as described above and, after 24 h, it was exposed to the SCEO (156, 312, 624 and 1,248

194 $\mu\text{g/mL}$ in 5% DMSO). After the incubation period, the content of wells was removed and the
195 remaining biofilm was washed two times with saline. MTT solution at 0.3 mg/mL (200 μL)
196 was added to each well and incubated for 90 min at 37 °C. The wells were then washed once
197 with saline and the purple formazan crystals were dissolved with 200 μL of DMSO for 20 min
198 and then the absorbance at 540 nm was measured. Vancomycin (1 $\mu\text{g/mL}$) was used as
199 positive control.

200

201 2.7. Scanning electron microscopy (SEM)

202

203 Biofilm was grown and treated with SCEO (312 and 624 $\mu\text{g/mL}$ in 5% DMSO) in 96-
204 wells microtiter plates containing a piece of Permanox™ slide in each well (Nalge Nunc
205 International, USA). After 48 h of incubation at 37 °C, the slide pieces were fixed and stored
206 in 2.5% (v/v) glutaraldehyde at -20 °C until microscopy analysis. The samples were washed
207 with 100 mM cacodylate buffer pH 7.2 and dehydrated in increasing concentrations of
208 acetone. The slides were dried by the CO₂ critical point technique (CPD 030 Balzers,
209 Liechtenstein), fixed on aluminum stubs, covered with gold film and examined in a JEOL
210 JSM-6060 microscope. Vancomycin (1 $\mu\text{g/mL}$) was used as antibiotic control.

211

212 2.8. In vivo assays using *Galleria mellonella*

213

214 2.8.1. Survival assay

215

216 *Galleria mellonella* larvae (200 mg) were randomly distributed in groups ($n =$
217 10/group) and infected with 10 μL of *S. aureus* UFPEDA-02 (ATCC-6538) suspension (1×10^5
218 CFU/larvae) injected into the last left proleg. After 2 h incubation at 37 °C, larvae received a

219 single dose of 10 μ L of *S. coronata* essential oil solutions at MIC or 2 \times MIC (that resulted in
220 doses of 15.6 mg/kg or 31.2 mg/kg, respectively) and the plates were re-incubated at 37 $^{\circ}$ C.
221 Larvae infected with *S. aureus* and inoculated with vehicle (PBS) were used as positive
222 control, while uninfected larvae also treated with vehicle were taken as negative control.
223 Mortality rates of each group were observed daily during 5 days.

224

225 2.8.2. Early melanization assay

226

227 The effect of SCEO in the production of melanin induced by *S. aureus* infection was
228 measured as previously described by Scorzoni et al. [34] with modifications. The larvae
229 ($n=10$ /group) were infected with *S. aureus* (1×10^6 CFU/larva) and immediately treated with
230 SCEO (15.6 or 31.2 mg/kg). Larvae infected and treated with vehicle (PBS) were used as
231 positive control while larvae inoculated only with vehicle composed the negative group. After
232 1 h and 3 h of incubation, the hemolymph of four larvae from each group was collected by
233 cutting them with a scalpel blade through the cephalocaudal direction and squeezing. The
234 obtained hemolymph was diluted in cold PBS and the melanin production was detected by
235 measuring the absorbance at 405 nm.

236

237 2.8.3. Bacterial load in hemolymph

238

239 To evaluate the effect of SCEO in bacterial load in hemolymph, the larvae ($n =$
240 10/group) were infected with *S. aureus* (1×10^5 CFU) and treated with essential oil (31.2
241 mg/kg) as described in section 2.8.1. The hemolymph of five larvae was collected daily for 3
242 days, serially diluted in PBS and 4 μ L of each dilution was plated on MHA. After incubation
243 for 24 h at 37 $^{\circ}$ C, the number of CFU/mL was determined.

244

245 *2.9. Statistical analysis*

246

247 All assays were performed in triplicate in at least two independent experiments.
248 Statistical analyses were performed by one-way analysis of variance (ANOVA). All
249 analyses were carried out using GraphPrism, version 7. Differences were considered
250 significant at $p < 0.05$. Differences in the survival of *G. mellonella* larvae were determined
251 using the Kaplan-Meier method and log-rank test was used to compare survival curves.

252

253 **3. Results and discussion**

254

255 *3.1. Chemical composition of SCEO*

256

257 The hydrodistillation of *S. coronata* seeds allowed to obtain the essential oil with yield
258 of $0.41 \pm 0.1\%$. The SCEO components detected by GC/MS and GC/FID are presented in
259 Table 1. A total of 11 volatile constituents were identified, corresponding to 98.63% of the
260 total oil, being most of them fatty acids (89.79%) and sesquiterpenes (8.5%). The most
261 abundant components were octanoic acid, dodecanoic acid, decanoic acid and γ -eudesmol.
262 Previous studies have reported that *S. coronata* oils are dominated by free carboxylic acids,
263 accounting for approximately 80% of the total composition, and octanoic acid has been
264 reported as the major volatile component of *S. coronata* oil [15, 16, 19].

265

266 *3.2. Phenotypic characteristics of S. aureus isolates*

267

268 The twenty *S. aureus* clinical isolates were from several sources such as catheter tip,
269 purulent exudate, bone fragment, surgical wound and human lesions (Table 2). The results
270 showed that 13 strains were resistant to oxacillin and classified as MRSA [35]. In fact, 11
271 MRSA clinical isolates were multidrug resistant as they showed resistance to at least 3
272 antibiotics while 1 MSSA was a multidrug resistant strain (Table 3). Microtiterplate assay
273 revealed that 13 clinical isolates were strong biofilm producers, 6 strains were moderate
274 biofilm producers, while one strain was a weak biofilm producer (Table 2).

275

276 3.3. SCEO is a bactericide agent against *S. aureus*

277

278 SCEO showed antimicrobial efficacy against all selected isolates of *S. aureus*,
279 including those with biofilm-forming and multiple drug resistance phenotypes. The MIC
280 values for the oil ranged from 156 µg/mL to 625 µg/mL (Table 3). The MBC values were
281 equal to or 2-fold higher than each respective MIC, ranging from 312 to 1250 µg/mL,
282 indicating the bactericidal effect of the oil. The MIC₅₀ and MIC₉₀ corresponded to 312 and
283 625 µg/mL, respectively. Essential oils from plants such *Caryophyllus aromaticus*,
284 *Cinnamomum zeylanicum*, *Eugenia uniflora*, *Rosmarinus officinalis*, *Vernonia polyanthes*,
285 and *Baccharis dracunculifolia* have been shown to be effective against clinical isolates of *S.*
286 *aureus*, with MIC ranging from 0.25 to 56 mg/mL for MRSA and 0.25 to 50.8 mg/mL for
287 MSSA [36].

288 The main constituents of SCEO are medium chain fatty acids, which have previously
289 been identified as bioactive components against bacteria and yeasts, tending to be more active
290 against gram-positive bacteria than gram-negative [37, 38]. For example, the octanoic acid,
291 the major component of SCEO, has antibacterial properties against a range of gram-positive

292 and gram-negative pathogens; dodecanoic and decanoic acids have been also reported as
293 antimicrobial agents [39, 40].

294

295 3.4. SCEO effects viability of eradicates *S. aureus* biofilm

296

297 The effect of SCEO on the biomass and viability of preformed biofilm was evaluated
298 using the strong biofilm producer *S. aureus* strain UFPEDA 02 (ATCC-6538). SCEO showed
299 a slightly effect on biofilm matrix; significant reduction was observed only with 312 and 624
300 $\mu\text{g/mL}$ concentrations (Figure 1). These findings were similar to vancomycin results, used as
301 antibiotic control. On the other hand, SCEO was able to significantly decrease cell viability
302 inside of the biofilm structure at all tested concentrations. Bacterial cell viability decreased
303 more than 50% when the biofilm was submitted to the lowest concentration (156 $\mu\text{g/mL}$,
304 corresponding to 0.5 \times MIC), while minimal viability was detected when the biofilms were
305 exposed to the highest concentrations (624 and 1,248 $\mu\text{g/mL}$, corresponding to 2 \times MIC and
306 4 \times MIC 2) (Figure 1). Although the cell viability was strongly reduced in the treatment at
307 1248 $\mu\text{g/mL}$, there was no significant reduction in biofilm biomass, which can be due to a
308 defensive response of the bacterial cells to this high oil concentration before they became
309 inviable. Vancomycin showed low effect against the bacteria within the biofilm, confirming
310 that planktonic bacterial susceptibility to antibiotics may not correspond to a good prediction
311 for bacteria in biofilm lifestyle. This may represent a key point in the failure of antimicrobial
312 treatment in the clinical routine as well as in the evaluation and development of new
313 antimicrobial agents [41, 42].

314 The SEM analysis revealed untreated *S. aureus* biofilm as aggregates composed by
315 cells with preserved structure (Figure 2A). No remarkable alterations were observed in
316 biofilm treated with vancomycin (Figure 2B). SCEO treatments with MIC, 312 $\mu\text{g/mL}$

317 (Figure 2C), and supra-MIC, 624 $\mu\text{g/mL}$ (Figure 2D) concentrations decreased the number of
318 live cells in biofilms and led to loss of the roughness in the multiple layers of the three-
319 dimensional structure of bacterial biofilm. SCEO at 312 $\mu\text{g/mL}$ caused alteration in the
320 cellular structure of *S. aureus* (Figure 2C), which may be related to the bactericidal action of
321 the oil. Another effect induced by SCEO in *S. aureus* biofilms was the overproduction of
322 exopolymeric matrix (Figure 2D), which can be a protective mechanism against the
323 aggression caused by the treatment [43–45]. This datum corroborates with those reported in
324 Figure 1. The maintenance of the three-dimensional matrix architecture (with dense areas,
325 pores and channels) is crucial to determine the way of life in biofilm due to its influence on
326 factors such as diffusion of nutrients, oxygen, residual products, and motility [46, 47]. The
327 biofilm eradication ability of the major components of SCEO was already reported. Hogan et
328 al. [48] demonstrated that application of ML:8, an emulsion based on octanoic acid, reduced
329 *S. aureus* biofilm viability in more than 97% after 24 h treatment *in vitro*. Hess et al. [49]
330 showed that dodecanoic acid was also able to reduce the viability of biofilm cells of *S.*
331 *aureus*; however, it did not reduce the biofilm biomass.

332

333 3.5. SCEO reduces the deleterious effects of *S. aureus* infection in *G. mellonella*

334

335 Based on the MIC values, we selected two concentrations of SCEO to evaluate its
336 antimicrobial action using *G. mellonella* larvae. The inoculation of SCEO at 15.6 mg/kg or
337 31.2 mg/kg did not change the survival rate of *G. mellonella* larvae. In addition, *G. mellonella*
338 larvae exposed to SCEO developed the pupal stage in the same time period than untreated
339 larvae. These data show that SCEO showed no toxicity to this insect.

340 The survival rate of *G. mellonella* larvae was reduced by infection with *S. aureus*
341 UFPEDA-02, resulting in the death of all larvae in 3 days. This effect was inhibited when the

342 larvae was treated with a single SCEO dose of 31.2 mg/kg, which resulted in survival rate of
343 60% in 4 days after infection. The mortality rate at dose of 15.6 mg/kg was not significantly
344 different to that of the untreated group (Figure 3A). Aiming to investigate if the mortality rate
345 was related to antibacterial activity of SCEO, the bacterial survival in hemolymph was
346 evaluated. The hemolymph of larvae infected with *S. aureus* exhibited increased levels of
347 bacterial load, approximately, 6, 8, and 9 log CFU/mL in 1st, 2nd, and 3rd days post-infection,
348 respectively. These values were reduced in treatment with SCEO at 31.2 mg/kg to 4, 5, and 5
349 log CFU/mL 1, 2, and 3 days post-infection, respectively (Figure 3B).

350 We also analyzed the effect of SCEO at 31.2 mg/kg in melanin production, employing
351 a model of acute infection by inoculating the larvae with a high-density inoculum.
352 Melanogenesis is an essential component of *G. mellonella* immune response against microbial
353 infection [50]. However, the overproduction of this pigment has been associated to death
354 induced by microorganisms [51]. The melanin in hemolymph significantly increased after 1
355 and 3 h of infection with *S. aureus* but SCEO was able to reduce larvae melanization induced
356 by *S. aureus* infection in both periods (Figure 3C). These results corroborate with the benefic
357 effects of this oil in infected larvae.

358

359 4. Conclusion

360

361 This work demonstrated that SCEO is an antibacterial agent against *S. aureus* strains
362 with different resistance phenotypes. In addition, the oil was able to disturb biofilm formed by
363 a strong biofilm producer isolate, and this antibiofilm activity was probably associated to the
364 decrease of viability of cells inside the biofilm. *In vivo* antibacterial activity of SCEO against
365 *S. aureus* improved survival of *G. mellonella* larvae and this fact indicates SCEO as a
366 potential candidate to drug development for treatment of *S. aureus* infections.

367

368 **Acknowledgments**

369

370 The authors express their gratitude to the *Conselho Nacional de Desenvolvimento*
371 *Científico e Tecnológico* (CNPq) for research grants (446902/2014-4) and fellowships
372 (MTSC, AJM, THN and PMGP). They are also grateful to the *Fundação de Amparo à*
373 *Ciência e Tecnologia do Estado de Pernambuco* (FACEPE; APQ-0108-2.08/14; APQ-0661-
374 2.08/15), *Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul* (FAPERGS;
375 PRONEM 16/2551-0000244-4); *Fundação de Amparo à Pesquisa e ao Desenvolvimento*
376 *Científico e Tecnológico do Maranhão* (FAPEMA; COOPI-02860/16); and the *Coordenação*
377 *de Aperfeiçoamento de Pessoal de Nível Superior* (CAPES; AUXPE 1454/2013) for financial
378 support. KAK thanks the support of the International Network Programme, Danish Agency
379 for Science, Technology and Innovation (6144-00103B). BSS would like to thank FACEPE
380 for graduate scholarship (IBPG-0775-2.08/13). The authors are also grateful to technical
381 assistance from the CMM/UFRGS for the microscopies (SEM).

382

383 **References**

384

- 385 [1] C.L. Ventola, The antibiotic resistance crisis: part 1: causes and threats. *P T* 40 (2015)
386 277-283.
- 387 [2] A.E. Paharik, A.R. Horswill, The staphylococcal biofilm: adhesins, regulation, and host
388 response. *Microbiol Spectr* 4 (2016) doi:10.1128/microbiolspec
- 389 [3] F.E. Guerra, T.R. Borgogna, D.M. Patel, E.W. Sward, J.M. Voyich, Epic immune battles
390 of history: neutrophils vs. *Staphylococcus aureus*, *Front. Cell Infect. Microbiol.* 7
391 (2017), 286.

- 392 [4] G. Pietrocola, G. Nobile, S. Rindi, P. Speziale, *Staphylococcus aureus* manipulates innate
393 immunity through own and host-expressed proteases, *Front. Cell Infect. Microbiol.* 7
394 (2017) 166.
- 395 [5] R.A. Cosimi, N. Beik, D.W. Kubiak, J.A. Johnson, Ceftaroline for severe methicillin-
396 resistant *Staphylococcus aureus* infections: a systematic review. *Open Forum Infect.*
397 *Dis.* 4 (2017), ofx084.
- 398 [6] W.J.B. van Wamel, *Staphylococcus aureus* infections, some second thoughts, *Curr. Opin.*
399 *Infect. Dis.* 30 (2017) 303-308.
- 400 [7] J. Haaber, J.R. Penades, H. Ingmer, Transfer of antibiotic resistance in *Staphylococcus*
401 *aureus*, *Trends Microbiol.* 25 (2017) 893-905.
- 402 [8] A.M.S. Figueiredo, F.A. Ferreira, C.O. Beltrame, M.F. Cortes, The role of biofilms in
403 persistent infections and factors involved in *ica*-independent biofilm development and
404 gene regulation in *Staphylococcus aureus*, *Crit. Rev. Microbiol.* 43 (2017) 602-620.
- 405 [9] D.E. Moormeier, K.W. Bayles, *Staphylococcus aureus* biofilm: a complex developmental
406 organism, *Mol. Microbiol.* 104 (2017) 365-376.
- 407 [10] H. Mccarthy, J.K. Rudkin, N.S. Black, L. Gallagher, E. O'neill, J.P. O'gara, Methicillin
408 resistance and the biofilm phenotype in *Staphylococcus aureus*, *Front Cell Infect*
409 *Microbiol* 5 (2015) 1.
- 410 [11] B.S. Santos, L.C.N. Silva, T.D. Silva, J.F. Rodrigues, M.A. Grisotto, M.T.S. Correia,
411 T.H. Napoleão, M.V. Silva, P.M.G. Paiva, Application of omics technologies for
412 evaluation of antibacterial mechanisms of action of plant-derived products. *Front*
413 *Microbiol* 7 (2016) 1466.
- 414 [12] M.K. Swamy, M.S. Akhtar, U.R. Sinniah, Antimicrobial properties of plant essential oils
415 against human pathogens and their mode of action: an updated review. *Evid. Based*
416 *Complement. Alternat. Med.* 2016 (2016) 3012462.

- 417 [13] Silva, L.C.N., Silva, M.V., Correia, M.T.S., Editorial: New frontiers in the search of
418 antimicrobials agents from natural products, *Front Microbiol* 8 (2017) 210.
- 419 [14] M.T. Islam, A.M. Mata, R.P. Aguiar, M.F. Paz, M.V. Alencar, P.M. Ferreira, A.A.C.
420 Melo-Cavalcante, Therapeutic potential of essential oils focusing on diterpenes,
421 *Phytother Res* 30 (2016) 1420-1444.
- 422 [15] S. Belviso, D. Ghirardello, M. Giordano, G.S. Ribeiro, J.S. Alves, S. Parodi, S. Risso, G.
423 Zeppa, Phenolic composition, antioxidant capacity and volatile compounds of licuri
424 (*Syagrus coronata* (Martius) Beccari) fruits as affected by the traditional roasting
425 process, *Food Res. Int.* 51 (2013) 39-45.
- 426 [16] C.M.A. Bessa, R.S.D. Nascimento, R.C.C. Alves, J.M. Anselmo, A.P.S. Silva, A.G.
427 Silva, V.L.M. Lima, J.F. Tavares, L.C.N. Silva, M.V. Silva, M.T.S. Correia, *Syagrus*
428 *coronata* seed oils have antimicrobial action against multidrug-resistant *Staphylococcus*
429 *aureus*, *J. Med. Plants Res.* 10 (2016) 310-317.
- 430 [17] E. Toensmeier, The carbon farming solution: a global toolkit of perennial crops and
431 regenerative agriculture practices for climate change mitigation and food security.
432 Chelsea Green Publishing, Vermont (2016).
- 433 [18] W.M.D. Andrade, M.A. Ramos, W.M.S. Souto, J.S. Bento-Silva, U.P.D. Albuquerque,
434 E.D.L. Araújo, Knowledge uses and practices of the licuri palm (*Syagrus coronata*
435 (Mart.) Becc.) around protected areas in northeastern Brazil holding the endangered
436 species lear's macaw (*Anodorhynchus leari*), *Trop. Conserv. Sci.* 8 (2015) 893-911.
- 437 [19] L.M.M. Santos, J.S. Nascimento, M.G. Santos, N.B. Marriel, P.C. Bezerra-Silva, S.K.L.
438 Rocha, A.G. Silva, M.T.S. Correia, P.M.G. Paiva, G.F. Martins, M.V. Silva, T.H.
439 Napoleão, Fatty acid-rich volatile oil from *Syagrus coronata* seeds has larvicidal and
440 oviposition-deterrent activities against *Aedes aegypti*, *Physiol. Mol. Plant Pathol.* 100
441 (2017) 35-40.

- 442 [20] L.B. Leal, G.D. Sousa, K.B. Seixas, P.H.N.D. Souza, D.P.D. Santana, Determination of
443 the critical hydrophile-lipophile balance of licuri oil from *Syagrus coronata*:
444 Application for topical emulsions and evaluation of its hydrating function, Braz. J.
445 Pharm. Sci. 49 (2013) 167-173.
- 446 [21] T.A.F. Ferro, J.M.M. Araujo, B.L.S. Pinto, J.S. Santos, E.B. Souza, B.L.R. Silva, V.L.P.
447 Colares, T.M.G. Novais, C.M.B. Filho, C. Struve, J.B. Calixto, V. Monteiro-Neto,
448 L.C.N. Silva, E.S. Fernandes, Cinnamaldehyde inhibits *Staphylococcus aureus*
449 virulence factors and protects against infection in a *Galleria mellonella* model, Front
450 Microbiol 7 (2016) 2052.
- 451 [22] R. Jønsson, C. Struve, H. Jenssen, K.A. Krogh, The wax moth *Galleria mellonella* as a
452 novel model system to study enteroaggregative *Escherichia coli* pathogenesis,
453 Virulence 8 (2017) 1894-1899.
- 454 [23] K.M.T. Astvad, J. Meletiadis, S. Whalley, M.C. Arendrup, Fluconazole
455 pharmacokinetics in the *Galleria mellonella* larvae and performance evaluation of a
456 bioassay compared to LC-MS/MS for haemolymph specimens, Antimicrob Agents
457 Chemother. 61 (2017) e00895-17.
- 458 [24] L.C.M. Lima, D.M.A.F. Navarro, L.P. Souza-Santos, Effect of diet on the fatty acid
459 composition of the copepod *Tisbe biminiensis*, J. Crust. Biol. 33 (2013) 372-381.
- 460 [25] E.S. Autran, I.A. Neves, C.S.B. Silva, G.K.N. Santos, C.A.G. Câmara, D.M.A.F.,
461 Chemical composition, oviposition deterrent and larvicidal activities against *Aedes*
462 *aegypti* of essential oils from *Piper marginatum* Jacq. (Piperaceae), Bioresour Technol
463 100 (2009), 2284-2288.
- 464 [26] H. van Den Dool, P.D. Kratz, A generalization of the retention index system including
465 linear temperature programmed gas-liquid partition chromatography, J Chromatogr 11
466 (1963) 463-471.

- 467 [27] R.P. Adams, Identification of essential oil components by gas chromatography/mass
468 spectroscopy, Allured Publishing, Carol Stream, USA.
- 469 [28] A.W. Bauer, M.D.K. Kirby, J.C. Sherris, M. Truck, Antibiotic susceptibility testing by a
470 standardized single disk method, *Am. J. Clin. Pathol.* 45 (1966) 493-496.
- 471 [29] P.H. Krumperman, Multiple antibiotic resistance indexing of *Escherichia coli* to identify
472 high-risk sources of fecal contamination of foods, *Appl. Environ. Microbiol.* 46 (1983)
473 165-170.
- 474 [30] S. Stepanovic, D. Vukovic, V. Hola, G. Di Bonaventura, S. Djukic, I. Cirkovic, F.
475 Ruzicka, Quantification of biofilm in microtiter plates: overview of testing conditions
476 and practical recommendations for assessment of biofilm production by staphylococci,
477 *APMIS* 115 (2007) 891-899.
- 478 [31] Clinical and Laboratory Standards Institute, Performance standards for antimicrobial
479 susceptibility testing; 24th informational supplement. CLSI document M100-S24
480 (2014).
- 481 [32] K.R. Zimmer, A.J. Macedo, R.B. Giordani, J.M. Conceicao, G.G. Nicastro, A.L.
482 Boechat, R.L. Baldini, W.R. Abraham, C. Termignoni, A steroidal molecule present in
483 the egg wax of the tick *Rhipicephalus (Boophilus) microplus* inhibits bacterial biofilms.
484 *Environ. Microbiol.* 15 (2013) 2008-2018.
- 485 [33] T. Mosmann, Rapid colorimetric assay for cellular growth and survival: Application to
486 proliferation and cytotoxicity assays, *J. Immunol. Meth.* 65 (1983) 55-63.
- 487 [34] L. Scorzoni, M.P. Lucas, A.C. Mesa-Arango, A.M. Fusco-Almeida, E. Lozano, M.
488 Cuenca-Estrella, M.J. Mendes-Giannini, O. Zaragoza, Antifungal efficacy during
489 *Candida krusei* infection in non-conventional models correlates with the yeast in vitro
490 susceptibility profile, *PLoS One* 8 (2013) e60047.

- 491 [35] V.A. Kumar, K. Steffy, M. Chatterjee, M. Sugumar, K.R. Dinesh, A. Manoharan, S.
492 Karim, R. Biswas, Detection of oxacillin-susceptible *mecA*-positive *Staphylococcus*
493 *aureus* isolates by use of chromogenic medium MRSA ID, J. Clin. Microbiol. 51 (2013)
494 318-319.
- 495 [36] L.N. Barbosa, I.S. Probst, B.F.M.T. Andrade, F.C.B. Alves, M. Albano, L.M.C.
496 M.L.R.S. Cunha, J.T. Doyama, V.L.M. Rall, A.F. Júnior, A. In vitro antibacterial and
497 chemical properties of essential oils including native plants from Brazil against
498 pathogenic and resistant bacteria, J. Oleo Sci. 64 (2015), 289-298.
- 499 [37] A.P. Desbois, V.J. Smith, Antibacterial free fatty acids: activities, mechanisms of action
500 and biotechnological potential, Appl. Microbiol. Biotechnol. 85 (2010) 1629-1642.
501
- 502 [38] E.M.S. Lillebæk, S.L. Nielsen, R.S. Thomasen, N.J. Faergeman, B.H. Kallipolitis,
503 Antimicrobial medium- and long-chain free fatty acids prevent PrfA-dependent
504 activation of virulence genes in *Listeria monocytogenes*, Res. Microbiol. 168 (2017)
505 547-557.
- 506 [39] R. Hulankova, G. Borilova, I. Steinhauserova, Combined antimicrobial effect of oregano
507 essential oil and caprylic acid in minced beef. Meat Sci 95 (2013) 190-194.
- 508 [40] S.A. Kim, M.S. Rhee, Synergistic antimicrobial activity of caprylic acid in combination
509 with citric acid against both *Escherichia coli* O157:H7 and indigenous microflora in
510 carrot juice, Food Microbiol. 49 (2015), 166-172.
- 511 [41] J. Claessens, M. Roriz, R. Merckx, P. Baatsen, L. Van Mellaert, J. Van Eldere, Inefficacy
512 of vancomycin and teicoplanin in eradicating and killing *Staphylococcus epidermidis*
513 biofilms in vitro, Int. J. Antimicrob. Agents 45 (2015) 368-375.

- 514 [42] B. Ozturk, N. Gunay, B.M. Ertugrul, S. Sakarya, Effects of vancomycin, daptomycin,
515 and tigecycline on coagulase-negative staphylococcus biofilm and bacterial viability
516 within biofilm: an in vitro biofilm model, *Can. J. Microbiol.* 62 (2016) 735-743.
- 517 [43] D.S. Trentin, R.B. Giordani, K.R. Zimmer, A.G. Silva, M.V. Silva, M.T.S. Correia, I.J.
518 Baumvol, A.J. Macedo, Potential of medicinal plants from the Brazilian semi-arid
519 region (Caatinga) against *Staphylococcus epidermidis* planktonic and biofilm lifestyles,
520 *J. Ethnopharmacol.* 137 (2011) 327-335.
- 521 [44] Z. Sanchez, A. Tani, K. Kimbara, Extensive reduction of cell viability and enhanced
522 matrix production in *Pseudomonas aeruginosa* PAO1 flow biofilms treated with a D-
523 amino acid mixture, *Appl. Environ. Microbiol.* 79 (2013) 1396–1399.
- 524 [45] B. Kundukad, M. Schussman, K. Yang, T. Seviour, L. Yang, S.A. Rice, S. Kjelleberg,
525 P.S. Doyle, Mechanistic action of weak acid drugs on biofilms, *Sci. Rep.* 7 (2017) 4783.
- 526 [46] H.C. Flemming, J. Wingender, The biofilm matrix, *Nat. Rev. Microbiol.* 8 (2010), 623-
527 633.
- 528 [47] H.C. Flemming, J. Wingender, U. Szewzyk, P. Steinberg, S.A. Rice, S. Kjelleberg,
529 Biofilms: an emergent form of bacterial life, *Nat Rev Microbiol* 14 (2016) 563-575.
- 530 [48] S. Hogan, M. Zapotoczna, N.T. Stevens, H. Humphreys, J.P. O'gara, E. O'neill,
531 Eradication of *Staphylococcus aureus* catheter-related biofilm infections using ML:8
532 and Citrox, *Antimicrob. Agents Chemother.* 60 (2016) 5968-5975.
- 533 [49] D.J. Hess, M.J. Henry-Stanley, C.L. Wells, The natural surfactant glycerol monolaurate
534 significantly reduces development of *Staphylococcus aureus* and *Enterococcus faecalis*
535 biofilms, *Surg. Infect.* 16 (2015) 538-542.
- 536 [50] A. Zdybicka-Barabas, P. Mak, T. Jakubowicz, M. Cytrynska, Lysozyme and defense
537 peptides as suppressors of phenoloxidase activity in *Galleria mellonella*, *Arch. Insect*
538 *Biochem. Physiol.* 87 (2014) 1-12.

539 [51] H. Ciesielczuk, J. Betts, L. Phee, M. Doumith, R. Hope, N. Woodford, D.W. Wareham,
540 Comparative virulence of urinary and bloodstream isolates of extra-intestinal
541 pathogenic *Escherichia coli* in a *Galleria mellonella* model, *Virulence* 6 (2015) 145-
542 151.

543

544 **Figure captions**

545

546 **Figure 1.** Effect of the *Syagrus coronata* essential oil (SCEO) on biomass and cell viability in
547 *S. aureus* UFPEDA-02 biofilm. Biomass was quantified using the microtiterplate method (OD
548 570 nm) and viability was determined by MTT assay (OD 540 nm). (*) $p < 0.05$; (**) $p < 0.01$.

549

550 **Figure 2.** SEM images of biofilms formed by *S. aureus* UFPEDA-02. (A) Untreated biofilm.
551 (B) Biofilm treated with vancomycin at 1 $\mu\text{g}/\text{mL}$. (C) Biofilm treated with *Syagrus coronata*
552 essential oil (SCEO) at 312 $\mu\text{g}/\text{mL}$. Arrows point cells with altered structure. (D) Biofilm
553 treated with SCEO at 624 $\mu\text{g}/\text{mL}$. The asterisks indicate overproduction of exopolymeric
554 matrix.

555

556 **Figure 3.** Effects of SCEO on *Galleria mellonella* larvae infected with *S. aureus* UFPEDA-
557 02. (A) Survival curves of uninfected insects treated with PBS as well as infected insects
558 treated with PBS (control) or SCEO at 15.6 and 31.2 mg/kg. (B) Bacterial load and (C)
559 melanization in larvae uninfected treated with PBS as well as insects treated with PBS
560 (control) or SCEO at 31.2 mg/kg. (*) $p < 0.05$ (**) $p < 0.01$.

Table 1. Constituents of *Syagrus coronata* seed essential oil (SCEO).

N°	Compound ^a	RI		Content (as % of total oil)
		Determined ^b	Literature ^c	
1	Octanoic acid	1195	1167	46.77 ± 1.85
2	Decanoic acid	1378	1364	20.93 ± 0.29
3	<i>trans</i> -Caryophyllene	1421	1417	0.41 ± 0.05
4	Viridiflorene	1497	1496	0.53 ± 0.09
5	δ-Cadinene	1525	1522	0.44 ± 0.06
6	Dodecanoic acid	1573	1565	22.09 ± 3.51
7	Caryophyllene oxide	1586	1582	0.61 ± 0.23
8	Ethyl dodecanoate	1595	1594	0.34 ± 0.06
9	γ-Eudesmol	1634	1630	4.26 ± 0.41
10	β-Eudesmol	1653	1649	0.41 ± 0.04
11	α-Eudesmol	1656	1652	1.84 ± 0.61
Total				98.63

^a Constituents listed in order of elution on a non-polar DB-5 column; ^b Retention indexes (RI) calculated from retention times in relation a series of C₈-C₃₀ *n*-alkanes on a 30 m DB-5 capillary column; ^c Values taken from Adams (2007).

Isolate	Source	Phenotypic evaluation	Crystal violet assay
---------	--------	-----------------------	----------------------

Table 2. Isolation source, phenotypic evaluation and biofilm formation ability of *S. aureus* isolates used in this study.

		Colony color	Colony consistency	OD ₅₇₀	Biofilm formation
UFPEDA-02 (ATCC-6538)	Human lesion	Almost black	Dry	1.47±0.12	+++
UFPEDA-659	Catheter tip	Red	Crystalline	0.59±0.13	++
UFPEDA-662	Catheter tip	Almost black	Rough	1.22±0.08	+++
UFPEDA-670	Catheter tip	Red	Crystalline	0.52±0.03	++
UFPEDA-671	Bone Fragment	Almost black	Dry e Rough	1.03±0.15	+++
UFPEDA-672	Bone Fragment	Almost black	Rough	1.14±0.08	+++
UFPEDA-674	Purulent exudate	Bordeaux red	Crystalline	0.62±0.09	++
UFPEDA-679	Surgical wound	Black	Rough	0.77±0.07	++
UFPEDA-683	Purulent exudate	Almost black	Rough	1.06±0.19	+++
UFPEDA-689	Purulent exudate	Black	Rough	1.22±0.11	+++
UFPEDA-691	Catheter tip	Red	Rough	0.45±0.06	++
UFPEDA-699	Catheter tip	Red	Crystalline	1.01±0.15	+++
UFPEDA-700	Diabetic foot ulcer	Bordeaux red	Crystalline	1.01±0.22	+++
UFPEDA-705	Surgical wound	Black	Rough	1.49±0.18	+++
UFPEDA-709	Purulent exudate	Red	Crystalline	1.31±0.09	+++
UFPEDA-718	Tracheal secretion	Red	Crystalline	0.38±0.05	+
UFPEDA-726	Nasal secretion	Red	Crystalline	1.18±0.12	+++
UFPEDA-731	Surgical wound	Almost black	Rough	0.58±0.05	++
UFPEDA-733	Bone Fragment	Bordeaux red	Crystalline	1.28±0.16	+++
UFPEDA-802	Nasal secretion	Red	Dry	1.14±0.2	+++

(+++) **Strong biofilm forming strain. (++) Moderate biofilm forming strain. (+) Weak biofilm forming strain.**


1 **Table 3.** Antibiotic resistance profile of *S. aureus* isolates and antibacterial activity of *Syagrus coronata* essential oil (SCEO).


Clinical isolate	Susceptibility profile	MAR	SCEO activity	
			MIC($\mu\text{g/mL}$)	MBC
UFPEDA-02	Susceptible	0	312	312
UFPEDA-659	CFO,OXA, NAL	0.15	312	312
UFPEDA-662	AMP, CFO, OXA, NAL	0.2	625	625
UFPEDA-670	AMP, CFO, OXA, NAL, CIP, CLI, TRI	0.35	312	625
UFPEDA-671	AMP, CFO, OXA, NAL, CIP, AMI, GEN, CLI, CLO, TET, TRI	0.55	312	312
UFPEDA-672	AMP, CFO, OXA,NAL, CIP, NIT, CLI, TRI	0.4	156	312
UFPEDA-674	AMP, NAL, CLI, TET	0.2	312	625
UFPEDA-679	AMP, CFO, OXA, CFL, CFZ, NAL, VAN, AMI, CLI	0.45	625	625
UFPEDA-683	AMP, OXA, CFL, CFO, CFZ, CPM, CRX, CTX, NAL, CIP, VAN, AMI, GEN, CLI, CLO, TRI	0.8	625	1250
UFPEDA-689	AMP, CFZ, NAL, GEN, CLI, CLO, TET, TRI	0.4	625	625
UFPEDA-691	NAL, CIP, CLO	0.15	156	312
UFPEDA-699	AMP, NAL, CLI, CLO	0.2	156	312
UFPEDA-700	AMP, CFO, OXA, CIP, TET	0.25	312	156
UFPEDA-705	AMP, OXA, CFL, CFO, CPM, CRX, NAL, NIT, GEN	0.45	312	312
UFPEDA-709	AMP, CFO, OXA, NAL, CLI, TET	0.3	625	625
UFPEDA-718	AMP, NAL, CIP	0.15	312	312
UFPEDA-726	AMP, CFO, OXA, CIP, GEN, CLO, TRI	0.35	312	312
UFPEDA-731	AMP, CFO, OXA, CFL, CFO, CRX, NAL, CIP, GEN, CLI, CLO, TRI	0.6	312	312
UFPEDA-733	AMP, NAL, CIP, CLO	0.2	625	625
UFPEDA-802	AMP, OXA, CFL, CFO, CFZ, CPM, CRX, CTX, NAL, CIP, AMI, GEN, CLI, CLO, TET,	0.8	625	625




2 AMP: ampicillin. OXA: oxacillin. CFL: cephalothin. CFZ: cefazolin. CPM: cefepime. CFO: cefoxitin, CTX: cefotaxime. CRX: cefuroxime. IMI:
3 imipenem. MER: meropenem. NAL: nalidixic acid. CIP: ciprofloxacin. NIT: nitrofurantoin. AMI: amikacin. GEN: gentamicin. VAN:
4 vancomycin. CLI: clindamycin. CLO: chloramphenicol. TET: tetracycline. TRI: trimethoprim. MAR: multiple antibiotic resistance index. MIC:

- 5 minimum inhibitory concentration. MBC: minimum bactericidal concentration. The MIC₅₀ and MIC₉₀ of SCEO were 312 and 625 µg/mL,
6 respectively.

ACCEPTED MANUSCRIPT







Highlights

- Essential oil extracted from *Syagrus coronata* seeds (SCEO) was obtained.
- SCEO showed bactericidal activity (MBC from 312 to 1250 µg/mL) against *S. aureus*.
- SCEO decreased cell viability in pre-formed biofilms of *S. aureus* isolate.
- SCEO improved the survival of *G. mellonella* larvae inoculated with *S. aureus*.

ACCEPTED MANUSCRIPT