



Indian Journal of Geo Marine Sciences
Vol. 50 (01), January 2021, pp. 37-43



Evaluation of two freshwater macrophytes, *Ceratophyllum demersum* and *Potamogeton amplifolius* as feed ingredients for Nile tilapia (*Oreochromis niloticus*) fingerlings

A Balkhasher^a, T El-Sayed Ali^{*.b} & A F M El-Sayed^b

^aSaudi Aquaculture Society, Ministry of Environmental Water and Agriculture - Jeddah – 11195, Saudi Arabia

^bOceanography Department, Faculty of Science, Alexandria University, Alexandria – 21568, Egypt

*[E-mail: tameraly@yahoo.com]

Received 15 May 2019; revised 01 October 2020

The present study was carried out to evaluate the potential of two common freshwater macrophytes (*Ceratophyllum demersum* and *Potamogeton amplifolius*) as feed ingredients for Nile tilapia fingerlings, in two consecutive experiments. The first experiment investigated the use of raw, dried *C. demersum* and *P. amplifolius* as sources of energy in the diets of fingerlings. The macrophytes were incorporated in 6 isonitrogenous and isocaloric diets at four levels as a replacement of dietary wheat bran. The test diets were fed to triplicate groups, stocked in 140 L culture aquaria in a recirculating system, three times a day to satiation, for 45 days. The results showed that control, macrophyte-free diet produced significantly improved growth rates and feed utilization efficiency than macrophytes-based diets.

In Experiment 2, fermented *C. demersum* and *P. amplifolius* were incorporated into six isonitrogenous and isocaloric diets. The diets were fed to triplicate groups of fingerlings (5.3 g) for 45 days. Growth rates and feed utilization efficiency of fish fed with fermented *P. amplifolius* at 33 % and 66 % inclusion level were not significantly different from fish fed the control diet. At 100 % inclusion level, fish performance was significantly reduced. On the other hand, fermented *C. demersum* produced extremely poor performance, compared to raw *ceratophyllum*. In conclusion, the present results indicated that fermentation improved the quality of *P. amplifolius*; but not *C. demersum*.

[**Keywords:** Fermentation, Feed utilization, Macrophytes, Nile tilapia]

Introduction

Global tilapia production has witnessed a sharp expansion during the past two decades, and is being cultured in more than 130 countries worldwide. Nowadays, tilapias are considered the second most important farmed finfish group in the world, after carps¹. Global production of farmed tilapia has increased from less than a half million metric tons in the early 1990s to 58 million metric tons in 2016, representing 125 % of freshwater fish production and 107 % of total fish culture, with an average annual growth rate of 135 %¹. Nile tilapia is the dominant farmed species; contributing 71 % to total tilapia production in 2016^(ref. 1).

The global expansion and industrialization of tilapia production has led to stepwise improve in tilapia culture from traditional, low-input, semi-intensive systems to more intensive farming practices, with an increasing dependence on formulated diets. This has created a gap between feed supply and a farmer's demand. Therefore, the major challenge faced by tilapia culture industry is the production of

sufficient quantity of high quality feeds. The sharp increase in feed ingredients in recent years has made the challenge more difficult, and the search for unconventional, locally available ingredients has become inevitable²⁻⁵.

The potential of soft submerged aquatic macrophytes as feed ingredient for herbivorous/omnivorous fishes; such as tilapia, have attracted the attention of many authors, with varying results, depending on cultured species and size and macrophyte species. The most commonly studied macrophytes were hornwort, oxygen weed, water velvet and pondweeds⁶⁻¹⁰. Hornwort (*Ceratophyllum demersum*) is a widely distributed fresh water macrophyte, belonging to family Ceratophyllaceae. The large-leaf pondweed (*Potamogeton amplifolius*) is also another well-known pondweed; which belongs to family Potamogetonaceae. Macrophytes have been used either fresh as a whole or dried as a partial component of a diet. Also, they have been tested as a partial or complete replacement of protein and/or energy sources in pelleted diets. The usage of

macrophytes in fish feeds depends -among other factors- on fish species and size, source, composition and processing of macrophytes, and culture systems¹¹.

A number of studies were carried out on the preference of different wild tilapia for different aquatic weeds. The feed preference of adult blue tilapia (*Oreochromis aureus*) for five aquatic plants was evaluated by Schwartz & Maughan¹². The order of preference was *Najas guadalupensis* and *Chara* sp., filamentous algae (predominantly *Cladophora* sp.), *Potamogeton pectinatus* and *P. nodosus*, respectively. Similarly, when juvenile Nile tilapia were fed on different fresh macrophytes, *Elodea canadensis* was the most preferred, followed by *Potamogeton pectinatus* and *Spirodela polyrhiza* (with equal preference), whereas *Myriophyllum spicatum* showed the lowest preference¹³. The consumption of the water fern (*Azolla pinnata*) by Nile tilapia (*Oreochromis niloticus*)¹⁴ and *Tilapia zillii*¹⁵ deteriorated the growth rates over time. When *Azolla* was incorporated in test diets, they led to a reduction of fish performance beyond 25 % inclusion level¹⁶ for Nile tilapia in South Lake (Guangdong province, China). They found that macrophyte consumption was size-specific, where larger ones consumed mainly macrophytes, while small tilapias were much more dependent on periphyton, seston, or detritus. Moreover, in another study, the authors also found that fresh and dried (pelleted) *Ceratophyllum* improved the growth rate of Nile tilapia reared in hapa-in-pond. Similarly Bag *et al.*¹⁷, evaluated the use of three aquatic weeds namely, lemna (*Lemna minor*), water hyacinth (*Eichhornia crassipes*) and azolla (*Azolla pinnata*) as major feed ingredients in Mozambique tilapia (*O. mossambicus*) feeds. The limna-based diet produced significantly better performance than the other two macrophytes.

Fermentation may improve the nutritive value of feed ingredients, including macrophytes⁸. When molasse-fermented water hyacinth replaced wheat bran at two levels 10 and 20 % substitution levels, significant growth rate response of Nile tilapia fingerling¹⁸ found that fermented water hyacinth was well accepted by catla (*Catla catla*), rohu (*Labeo rohita*), mrigal (*Cirrhinus mrigala*), silver carp (*Hypophthalmichthys molitrix*) and by common carp (*Cyprinus carpio*). However, mrigal and silver carp showed the best growth rates, followed by rohu. Fermentation of water hyacinth may thus be a simple and efficient treatment for utilizing water hyacinths as

a feed or manure in fish culture without the energy-consuming process of palletization¹⁹. Such findings were confirmed in a study realized by Sadique²⁰ who showed the significant effect of molasse-fermented water hyacinth on the growth rate and flesh quality of common carp fingerlings.

Several aquatic macrophytes are widely distributed in irrigation and drainage water bodies in Egypt, as well as in inland and coastal lakes. Water hyacinth (*Eichhornia crassipes*), *Ceratophyllum demersum* and *Potamogeton amplifolius* are the most dominant in these water bodies. However, despite the potentials these macrophytes may have as fish feed ingredients, they have not been evaluated yet. Therefore, the present study was carried out to evaluate the potential of fresh, dry and fermented *C. demersum* and *P. amplifolius* as feed ingredients for Nile tilapia (*Oreochromis niloticus*) fingerlings.

Materials and Methods

Fish and culture facilities

Nile tilapia (*Oreochromis niloticus*) fingerlings (n = 205) were brought from Barsik Fish Farm (Behaira Governorate), Egypt. The fish were transported to the laboratory in Nylon bags (100×30×30 cm) half-filled with water and completed with oxygen gas for fish respiration. After resting and acclimation to water temperature in the lab, for a few minutes, the fish were distributed randomly into the culture aquaria. Fish were fed on a commercial (30 % CP) diet for one week to adapt the laboratory conditions and artificial feeding. The fish were stocked into 140 l aquaria in a closed, self-cleaning recirculating indoor system. The culture system was provided with a biological filter, aeration through an air compressor, and heaters to maintain water temperature at 27 °C. Approximately 10 % of the water volume was replaced by new freshwater daily. Lighting in the culture unit was set at 12:12 L:D cycle. Water quality parameters, including dissolved oxygen (DO), ammonia (NH₄-N), nitrates (NO₃-N) and nitrites (NO₂-N) and pH, were monitored weekly. The average values of these parameters throughout the study were: DO = 6.4±1.3 mg l⁻¹, NH₄ -N = 0.06±0.002 mg l⁻¹, NO₃ -N = 8.4±1.72 mg l⁻¹, NO₂ = 0.00 mg l⁻¹ and pH = 8.0±0.09.

Macrophyte fermentation

The macrophytes were partially dried in an electric oven at 60 °C until about 50 % of their water content

was removed. They were then fermented as described by El-Sayed⁸. Each macrophyte was put in a clean, dry glass aquarium at room temperature (25-30 °C). Five percent sugar cane molasses and 2 ml orthophosphoric acid/kg were added to each aquarium, with continuous mixing. The aquaria were then covered with glass covers. The mixture of macrophytes was remixed every ten days to facilitate decomposition, for two months. After fermentation process, the macrophytes were sun-dried for two days, grinded into powder-like form, using an electric grinder. They were then stored into sealed and labeled plastic bags until used. The proximate composition of fermented macrophytes is given in Table 1.

Test diets

Twelve isonitrogenous (30 - 35 % CP), isocaloric (400 - 450 kcal GE/100 g) test diets were prepared. Fresh and fermented *C. demersum* and *P. amplifolius* were incorporated into the test diets, as an energy source, at 33 %, 66 %, and 100 % as a replacement for wheat bran (Table 2). The chemical analysis of the macrophytes, experimental diets and whole fish was done per the AOAC²¹ guideline. The test diets for each experiment were fed to groups of Nile tilapia fingerlings in triplicates, three times/day (08:00, 12:00 and 16:00 hrs), 7 days a week, for 45 days. Each group of fish was bulk weighed at the start and after every 15 days throughout the experimental period.

Table 1 — Proximate composition of *C. demersum* and *P. amplifolius* before (BF) and after (AF) fermentation and wheat bran (WB) used in the experiment

Component (% DM)	<i>C. demersum</i>			<i>P. amplifolius</i>			(WB)
	BF	AF	Change %	BF	AF	Change %	
Crude protein	9.2	14.0	+ 52	11.1	25.0	+ 125	13.0
Ether extract	1.9	2.2	+ 16	2.3	2.6	+ 13	4.3
Fiber	15.8	14.2	- 10	15.4	10.8	-30	15.0
Nitrogen free extract	63.6	58.5	-8	66.3	54.9	- 17	62.5
Ash	9.5	11.1	+ 17	4.9	6.7	+ 37	5.2
¹ G E (kcal/100 g)	338	351	+ 4	346	397	+ 15	345

¹GE, gross energy, calculated based on 5.65, 9.5 and 4.1 (Kcal/100 g) for protein, lipid, and carbohydrate, respectively

Table 2 — Composition and proximate analyses (percent dry weight) of the test diets

Ingredients (%)	Control	Fermented						Raw (Non-fermented)					
		<i>C. demersum</i>			<i>P. amplifolius</i>			<i>C. demersum</i>			<i>P. amplifolius</i>		
		33 %	66 %	100 %	33 %	66 %	100 %	33 %	66 %	100 %	33 %	66 %	100 %
Fish meal ¹	10	10	10	10	10	8	9	10	10	10	10	10	10
Soy bean meal	54	54	54	54	54	55	55	40	40	40	40	40	40
Wheat bran meal	27	18	9	--	18	10	--	26	12	--	26	12	--
<i>C. demersum</i> meal	---	9	18	27	--	--	--	13	26	38	--	--	--
<i>P. amplifolius</i> meal	--	--	--	--	9	18	27	--	--	--	13	26	38
Soy bean oil	3	3	3	3	3	3	3	4	5	5	4	5	5
Fish oil	2	2	2	2	2	2	2	3	3	3	3	3	3
Vit & min mix ²	2	2	2	2	2	2	2	2	2	2	2	2	2
Binder ³ (CMC)	2	2	2	2	2	2	2	2	2	2	2	2	2
Total	100	100	100	100	100	100	100	100	100	100	100	100	100
Crude protein	35.31	36.18	35.22	35.47	35.78	36.51	36.25	30.50	30.12	31.44	28.86	30.10	31.30
Ether extract	5.81	7.25	5.87	5.13	5.14	6.68	5.27	6.64	7.22	7.10	6.08	6.23	5.53
Crude fiber	3.48	2.60	3.71	4.24	3.46	2.78	2.98	7.95	9.19	9.89	4.46	5.38	6.98
Ash	7.90	12.87	19.13	24.22	10.50	12.56	15.48	12.97	16.65	20.13	11.87	13.12	14.79
NFE ⁴	47.50	41.10	36.07	30.94	45.12	41.47	40.02	40.43	35.99	31.75	44.65	42.87	43.28
GE ⁵ (kcal/100 g)	449	442	403	376	436	440	420	400	387	376	403	401	404

9

¹60 % crude protein (Nile tilapia meal). ²Contains (per g): retinal palmitate, 20000 IU; cholecalciferol, 5000 IU; DL-a-tocopherol, 10 mg; ascorbic acid, 25 mg; Vit K3, (Menadione), 3.5 mg; thiamin hydrochloride, 2 mg; riboflavin, 4.8 mg; pyridoxine hydrochloride, 2.5 mg; cyanocobalamine, 25 mcg; biotin, 10 mcg; nicotinic acid, 25 mg; folic acid, 0.5 mg; Dcalciumpanthothenate, 7.5 mg; sodium sulphate, 50 mg; potassium chloride, 30 mg; manganese sulphate, 15 mg; zinc sulphate, 15 mg; copper sulphate, 2 mg; ferrous sulphate, 15 mg. ³Carboxymethyl cellulose, used as binder. ⁴NFE, nitrogen-free extract, determined by difference. ⁵GE, gross energy, calculated based on 5.65, 9.5 and 4.1 (Kcal/100 g) for protein, lipid, carbohydrate, respectively.

Body composition analysis

At the termination of the study, all fish in each tank were netted, weighed, and frozen for final body composition analyses. Initial body analyses were performed on a sample of fish, which were weighed and frozen prior to the study. Proximate analyses of body water, protein, lipid, and ash were performed according to standard methods²¹.

Statistical analysis

Results of fish growth rates, feed utilization efficiency and body composition of an experiment were subjected to one-way analyses of variance (ANOVA). Orthogonal polynomial procedures were used to compare means at $P = 0.05$. Least significant difference (LSD) was used to test differences among treatment means when F -values from ANOVA were significant.

Results

With respect to fish growth rates, feed utilization efficiency and body composition, the results of the experiment indicated that the inclusion of macrophytes (*Ceratophyllum demersum* and *Potamogeton amplifolius*) in Nile tilapia diets significantly affected their growth rates and feed utilization efficiency (Table 3). The control diet [containing wheat bran (WB) as an energy source] produced significantly better ($P < 0.05$) weight gain, specific growth rate (SGR), feed conversion ratio (FCR) and protein efficiency ratio (PER). The results of inclusion of raw (non-fermented) *C. demersum* and *P. amplifolius* in Nile tilapia diets at all WB-substitution levels significantly reduced growth rates and feed utilization efficiency. This finding suggests that raw *C. demersum* (RC) and raw *P. amplifolius* (RP) macrophytes are not good sources of energy for Nile tilapia fingerlings. However, *C. demersum* was slightly better utilized than *P. amplifolius*. Fermentation improved the quality of *P. amplifolius* (FP), and led to improved performance up to 66 % inclusion level. On the contrary, fermentation of *C. demersum* (FC) had resulted in extremely poor performance of Nile tilapia fingerlings.

Body composition of Nile tilapia fed the test diets in the present study was not significantly affected by dietary treatments (Table 4).

Discussion

The relatively high fiber contents of *C. demersum* and *P. amplifolius* may have also been responsible for

poor digestibility which in turn may have led to poor growth. In support, Edwards²² reported that high fiber levels in the diet are known to retard the growth of fish. Boyd²³ also noted that the coarseness of macrophytes, due to the encrustation by calcium carbonate on their external surfaces, makes them unpalatable.

The present results agreed with those reported on tilapia fed on *C. demersum*²⁴, where the poor growth has been attributed to the poor digestibility of this macrophyte. Poor performance was also recorded when *C. demersum* was fed to Nile tilapia^{6,25,26}.

Similarly, Appler²⁷ demonstrated that most of the aquatic plants including algae contain 40 % or more

Table 3 — Effects of fresh and fermented dietary macrophytes on weight gain and feed utilization efficiency of fingerling Nile tilapia

Treatment	IW ¹	FW ²	Percent gain ³	SGR ⁴	FCR ⁵	PER ⁶
Control	5.7	15.1 ^a	164.9 ^a	2.2 ^a	1.3 ^a	2.6 ^a
RC33	6.1	13.5 ^{ab}	121.3 ^{ab}	1.8 ^{ab}	2.2 ^{bc}	1.5 ^{bc}
RC66	6.9	10.6 ^b	53.6 ^c	1.0 ^c	4.0 ^d	0.8 ^d
RC100	6.3	8.9 ^c	41.3 ^c	0.8	3.9 ^d	0.8 ^d
FC33	3.9	7.7 ^{cd}	97.4 ^d	1.5 ^b	1.6 ^{ab}	2.2 ^{ab}
FC66	3.8	8.0 ^{cd}	110.5 ^c	1.7 ^{ab}	1.9 ^b	1.8 ^b
FC100	3.9	6.5 ^d	66.7 ^{de}	1.1 ^c	3.9 ^d	0.8 ^d
RP33	5.8	9.6 ^c	65.5 ^{de}	1.1 ^c	2.4 ^c	1.5 ^{bc}
RP66	6.1	7.8 ^{cd}	27.9 ^f	0.5 ^d	3.0 ^{cd}	1.1 ^c
RP100	6.1	6.9 ^d	13.1 ^f	0.3 ^d	5.6 ^e	0.6 ^d
FP33	6.4	15.9 ^a	149.0 ^b	2.0 ^a	1.3 ^a	2.7 ^a
FP66	6.4	16.5 ^a	157.8 ^b	2.0 ^a	1.2 ^a	2.8 ^a
FP100	6.4	11.4 ^b	78.1 ^d	1.3 ^b	2.0 ^b	1.8 ^b

Values in the same column with different superscripts are significantly different ($P = 0.05$). ¹Initial weight (IW); ²Final weight (FW); ³Percent gain = 100 (final weight - initial weight)/initial weight; ⁴Specific growth rate (SGR) = 100 (ln final weight - ln initial weight)/time [days]; ⁵Feed conversion ratio (FCR) = dry feed offered/fish weight gain; and ⁶Protein efficiency ratio (PER) = Fish weight gain (g) / protein intake (g).

Table 4 — Effect of fresh and fermented dietary macrophytes on body composition (% dry weight) of Nile tilapia fingerlings

Treatment	Water content	Ether extract	Protein	Ash
Initial	70.9	22.7	65.9	13.7
Control	69.6	16.3	67.8	18.1
RC33	75.0	22.1	61.6	17.6
RC66	74.5	23.2	59.1	18.9
RC100	77.3	23.9	57.4	20.3
FC33	73.2	19.7	61.9	18.7
FC66	74.2	17.2	64.8	17.5
FC100	71.2	18.4	62.1	19.5
RP33	75.8	21.1	62.3	18.6
RP66	80.0	22.7	58.2	18.0
RP100	80.2	21.6	60.1	18.3
FP33	69.8	16.1	62.3	21.5
FP66	68.2	17.1	62.8	18.5
FP100	68.0	17.5	62.3	20.0

carbohydrates, of which only a small fraction consists of mono-saccharides and di-saccharides. Low digestibility of plant materials has been attributed to a preponderance of complex carbohydrates¹⁷. The same findings could be true in the present study, where *C. demersum* and *P. amplifolius* contained more than 50 % of nitrogen free extract, part of which is mono-saccharides and di-saccharides, which may interpret the limited utilization of these macrophytes leading to poor growth. However, more work is needed to verify the mono-saccharides and disaccharides contents of these plants and their digestibility and assimilation by Nile tilapia.

The reduced growth rates and feed efficiency of Nile tilapia fed macrophytes-based diets in the present study may have also been due to the effects of the anti-nutritional factors contained in these macrophytes. The existence of anti-nutritional factors within plant feedstuffs restricts their use in animal feeds²⁸.

On the contrary, some authors recommended the use of macrophytes as feed for fish^{10,29,22}. They noted that *T. zillii* and *T. rendalli* are voracious feeders of submerged macrophytes. Buddington³⁰ reported that *T. zillii* preferred *Najas guadalupensis* as a food source to *Lemna*, *Myriophyllum spicatum* and *Potamogeton pectinatus*. Cassani³¹ noted that grass carp prefer submerged, rather than floating macrophytes when they are supplied in fresh form^{7,32}. Moreover, Hasanuddin *et al.*³³ confirmed the suitability of *Ceratophyllum* sp. for *Oreochromis niloticus*.

It is obvious from the current study that fermented *P. amplifolius* can be considered as a feed ingredient in the diets for Nile tilapia fingerlings up to 66 % level of incorporation. Growth performance indices including SGR, FCR, and PER, of Nile tilapia fingerlings were similar at 33 % and 66 % incorporation of fermented *P. amplifolius* to the reference diet. The utilization efficiency of fermented *P. amplifolius* was significantly better than raw *P. amplifolius*. This means that fermentation improved the quality of *P. amplifolius*.

The better performance of Nile tilapia fed with fermented *P. amplifolius* up to 66 % inclusion level may have been due to the increased level and improved nature of protein of this macrophyte. It has been reported that crude protein was significantly increased in the fermented aquatic macrophytes when compared to the unfermented macrophytes^{19,34}. These

authors noted increase of crude protein content in fermented *Lemna* and *Spirodelamay* through microbial synthesis. The present study found that crude protein content was also affected by fermentation. The effect of fermentation on the protein content was conditional and strongly depends on the plant species.

Fermenting *P. amplifolius* may have also contributed in the removal of the anti-nutrients that might have been present in this macrophyte. For example, Velásquez³⁵ reported that the anti-nutritional substances, including trypsin inhibitor, phytates, tannins (hydrolyzed and condensed), and oxalates in *Azolla* were significantly reduced by the lactic acid fermentation.

Fermentation has also significantly reduced the fiber content of *P. amplifolius* in the present study compared with the terrestrial plants; the fiber structure of aquatic macrophytes is relatively easier to be decomposed by microorganism³⁶. Crude fiber was significantly lower in the fermented aquatic macrophytes when compared to the unfermented samples. The good performance of Nile tilapia fed with fermented *P. amplifolius* in the present study is also in agreement with the other results. El-Sayed⁸ reported that the process of fermentation is necessary when water hyacinth (*Eichhornia crassipes*) is included at levels of 20 % or more in Nile tilapia diets. Fermentation of duckweed had a significant positive effect on the growth performance, weight gain, specific growth ratio and protein efficiency ratio when applied to *Oreochromis niloticus* juveniles fed with low fishmeal diets³⁷.

Fermenting *C. demersum* led to a considerable reduction in fish performance compared to non-fermented format all inclusion levels and therefore, may not be considered as a feed ingredient in the diets for the Nile tilapia fingerlings. The cause of poor growth rates, PER, FCR, PPV and SGR of Nile tilapia fingerlings fed with fermented *C. demersum* is not known, particularly that the non-fermented *C. demersum* has resulted in better performance.

More research on the effects of processing on the quality of *C. demersum* as a feed ingredient for tilapia and other herbivorous fishes is needed. Other processing methods should also be tried in order to verify their effects on the quality of this macrophyte. Body composition of Nile tilapia fed with test diets in the present study was not significantly affected by

dietary treatments. This contrasts the findings of Edwards³⁸ who found that body protein, lipid, and ash were positively correlated with energy contents of dietary macrophytes.

The study concluded that, the inclusion of raw (non-fermented) *C. demersum* and *P. amplifolius* in Nile tilapia diets at all WB-substitution levels significantly reduced fish growth rates and feed utilization efficiency. This finding suggests that raw *C. demersum* and *P. amplifolius* macrophytes are not good sources of energy for Nile tilapia fingerlings. However, *C. demersum* was slightly better than *P. amplifolius*. This is due to the fermentation which improved the quality of *P. amplifolius* and led to improved performance up to 66 % inclusion level. On the contrary, fermentation of *C. demersum* had resulted in extremely poor performance of Nile tilapia fingerlings.

Acknowledgements

The authors are highly appreciated to all colleagues in Oceanography Department, Faculty of Science, Alexandria University who helped them during the experimental period.

Conflict of Interest

The authors declare that there is no conflict of interest.

Author Contributions

The authors have equal participation in preparing the manuscript.

References

- 1 FAO, *The state of the world fisheries and aquaculture*, (Rome, Italy), 2018, pp. 224.
- 2 El-Sayed Ali T, Abdel-Aziz S H, El-Sayed A M & Zeid S, Structural and functional effects of early exposure to 4-nonylphenol on gonadal development of Nile tilapia (*Oreochromis niloticus*): a-histological alterations in ovaries, *Fish Physiol Biochem*, 40 (2014) 1509-1519.
- 3 El-Sayed Ali T, Martinez-Llorens S, Monino A, Jover Cerda M & Tomas-Vidal A, Effects of weekly feeding frequency and previous ration restriction on the compensatory growth and body composition of Nile Tilapia fingerlings, *Egypt J Aquat Res*, 42 (2016) 357-363.
- 4 El-Sayed Ali T, Abdel-Aziz S H, El-Sayed A M & Zeid S, Effects of nonylphenol on plasma steroids, vitellogenin synthesis and sex reversal in Nile tilapia (*Oreochromis niloticus*), *Indian J Geo-Mar Sci*, 46 (3) (2017a) 521-528.
- 5 El-Sayed Ali T, El-Sayed A M, Abdel-Razek Eissa M & Hanafi H, Effects of dietary biogen and sodium butyrate on hematological parameters, immune response and histological characteristics of Nile tilapia (*Oreochromis niloticus*) fingerlings, *Aqua Int*, 26 (2) (2017b) 1-12. <https://doi.org/10.1007/s10499-017-0205-3>.
- 6 Chiayvareesajja S, Sirikul B, Sirimontrapon P, Rakkeaw S & Tansakul R, Comparison between natural feeding alone and supplemental feeding with pellets containing locally available ingredients for cage culture of *Oreochromis niloticus* in Thale Noi, Thailand, In: *The Second International Symposium on Tilapia in Aquaculture*, edited by RSV Pullin, T Bhukaswan, K Tonguthai & J L Maclean, (ICLARM Conference Proceedings No 15, Manila, Philippines), 1988, pp. 323-327.
- 7 Jagdish M, Rana S V S & Agarwal V P, Efficacy of grass carp (*Ctenopharyngo donidella*) in weed control and its growth in Karna Lake (Haryana), *J Inland Fish Soc India*, 27 (1995) 49-55.
- 8 El-Sayed A-F M, Effects of fermentation methods on the nutritive value of water hyacinth for Nile tilapia, *Oreochromis niloticus* (L) fingerlings, *Aquaculture*, 218 (2003) 471-478.
- 9 Henry-Silva G G, Monteiro Camargo A F & Pezzato L E, Apparent digestibility of aquatic macrophytes by Nile tilapia (*Oreochromis niloticus*) and water quality in relation nutrients concentrations, *R Bras de Zootec*, 35 (3) (2006) 641-647.
- 10 Mondal D, The utilization of aquatic weeds in an environmental friendly way of fish feed formulation- A review, *Int Res J Environ Sci*, 7 (2) (2018) 60-66.
- 11 Hasan M & Chakrabarti R, Use of algae and aquatic macrophytes as feed in small-scale aquaculture FAO fisheries and aquaculture technical paper (2009).
- 12 Schwartz D P & Maughan O E, The feeding preferences of *Tilapia aurea* (Steindachner) for five aquatic plants, *Proc Okla Acad Sci*, 64 (1984) 14-16.
- 13 Šetlíková I & Adámek Z, Feeding selectivity and growth of Nile tilapia (*Oreochromis niloticus* L) fed on temperate-zone aquatic macrophytes, *Czech J Anim Sci*, 49 (6) (2004) 271-278.
- 14 El-Sayed A-F M, Effects of substituting fish meal with *Azolla pinnata* in practical diets for fingerling and adult Nile tilapia, *Oreochromis niloticus* (L), *Aquac Res*, 23 (2) (1992) 167-173.
- 15 Abdel-Tawwab M, The Preference of the Omnivorous-Macrophagous, *Tilapia zillii* (Gervais), to Consume a Natural Free-floating Fern, *Azolla pinnata*, 39 (2008) 104-112.
- 16 Rao W, Ning J, Zhong P, Jeppesen E & Liu Z, Size-dependent feeding of omnivorous Nile tilapia in a macrophyte-dominated lake: implications for lake management, *Hydrobiol*, 749 (2015) 125-134.
- 17 Bag M P, Mahapatra S C & Rao P S, Aquatic Weed as Potential Feed for Mozambique tilapia, *Oreochromis mossambicus*, *J Aquac Res Deve*, 3 (8) (2012) 1-4.
- 18 Olah J, Ayyappan S & Purushothaman C S, Processing and utilization of fermented water hyacinth, *Eichhornia crassipes* (Mart) Solms in carp culture, *Aqua Hung*, 6 (1990) 219-234.
- 19 Sayed-Lafi R M, Al-Tameemi R & Gowde A I, Evaluation of raw and fermented water hyacinth (*Eichhornia crassipes*) incorporated diets on growth and feed efficiency of young grass carp (*Ctenopharyngodon idella*) Basrah, *J Agric Sci*, 31 (2018) 31-39.

- 20 Sadique J, Pandey A, Khairnar S O & Naveen Kumar B, Effect of molasses-fermented water hyacinth feed on growth and body composition of common carp, *Cyprinus carpio*, *J Entomol Zool Stud*, 6 (4) (2018) 1161-1165.
- 21 Association of Official Analytical Chemists, AOAC, Official Methods of Analysis AOAC International, Arlington, VA, USA (1995).
- 22 Edwards P, Food Potential of Aquatic Macrophytes Manila, *ICLARM Stud Rev*, 5 (1980) pp. 51.
- 23 Boyd C E, Fresh-water plants: a potential source of protein, *Econ Bot*, 22 (4) (1968) 359-368.
- 24 CIFA (Central Institute of Freshwater Aquaculture), *Annual Report of the Central Inland Fisheries Research Institute*, (Barrackpore), 1981.
- 25 Tantikitti C, Rittibhombhun N, Chaiyakum K & Tansakul R, Economics of tilapia pen culture using various feeds in Thale Noi, Songkhla Lake, Thailand, In: *The Second International Symposium on Tilapia in Aquaculture* Manila, edited by R S V Pullin, T Bhukaswan, K Tonguthai & J L Maclean, (ICLARM Conference Proceedings 15), 1988, pp. 569-574.
- 26 Klinavee S, Tansakul R & Promkuntong W, Growth of Nile tilapia (*Oreochromis niloticus*) fed with aquatic plant mixtures, In: *The Second Asian Fisheries Forum Manila*, edited by R Hirano & I Hanyu, (Asian Fisheries Society), 1990, pp. 283-286.
- 27 Appler H N, Evaluation of *Hydrodictyon reticulatum* as protein source in feeds for *Oreochromis niloticus* and *Tilapia zillii*, *J Fish Biol*, 27 (1985) 327-334.
- 28 Tacon A G J, *Standard Methods for the Nutrition and Feeding of Farmed Fish and Shrimp*, Vol 2, (Nutrient Sources and Composition Argent Laboratories Press, Redmond, WA, USA), 1990, pp. 129.
- 29 Abdel-Tawwab M, Effect of Free-Floating Macrophyte, *Azolla pinnata* on Water Physico-Chemistry, Primary Productivity, and the Production of Nile Tilapia, *Oreochromis niloticus* (L), and Common Carp, *Cyprinus carpio* L, in Fertilized Earthen Ponds, *J Appl Aquac*, 18 (2006) 21-41, DOI: 101300/J028v18n01_02
- 30 Buddington R K, Digestion of an aquatic macrophyte by *Tilapia zillii* (Gervais), *J Fish Biol*, 15 (1979) 449-455.
- 31 Cassani J R, Feeding behaviour of underyearling hybrids of the grass carp, *Ctenopharyngodon idella* (female) and the bighead carp, *Hypophthalmichthysno bilis* (male), on selected species of aquatic plants, *J Fish Biol*, 18 (1981) 127-133.
- 32 Aravindakshan P K, Jena J K, Ayyappan S, Routray P, Muduli H K, *et al.*, Evaluation of production trials with grass carp as a major component in carp polyculture, *J Inland Fish Soc India*, 31 (1999) 64-68.
- 33 Hasanuddin Muslim Putra B S, Nur A & Rimmer M A, Aquatic weed *Ceratophyllum* sp as a low-cost feed for brackish water pond culture of tilapia *Oreochromis niloticus*, *AACL Bioflux*, 9 (2016) 408-413.
- 34 Wee K L, Use of non-conventional feedstuff of plant origin as fish feeds is it practical and economically feasible In: *Fish Nutrition Research in Asia, Proceedings of the Fourth Asian Fish Nutrition Workshop*, Asian Fisheries Society Special Publication, Vol 5, edited by S S De Silva, (Asian Fisheries Society, Manila, Philippines), 1991, pp. 205.
- 35 Cruz Velásquez Y, Kijora C, Wedler E, Danier J & Schulz C, Fermentation properties and nutritional quality of selected aquatic macrophytes as alternative fish feed in rural areas of the Neotropics Animal Breeding in the Tropics and Subtropics, (Humboldt University of Berlin Philippstraße, 13 D-10115, Berlin, Germany), 2011.
- 36 Xiaoa L, Yanga L, Zhanga Y, Gua Y, Jianga L, *et al.*, Solid state fermentation of aquatic macrophytes for crude protein extraction, *Ecol Eng*, 35 (2009) 1668-1676.
- 37 Cruz Velásquez Y, Kijora C, Wuertz S & Schulz C, Effect of fermented aquatic macrophytes supplementation on growth performance, feed efficiency and digestibility of Nile Tilapia (*Oreochromis niloticus*) juveniles fed low fishmeal diets, *Livest Res Rural*, 27 (9) (2015).
- 38 Edwards P, Kamal M & Wee K L, Incorporation of composted and dried water hyacinth in pelleted feed for the tilapia *Oreochromis niloticus* (Peters), *Aquac Fish Management*, 16 (1985) 233-248.