

INTERACTION OF GONADAL STEROIDS AND GROWTH FACTORS IN BRAIN SEX DIFFERENTIATION

Maria C. Fernandez-Galaz¹, Arpad Parducz², Frederick Naftolin³, Ignacio Torres-Aleman¹, and Luis M. Garcia-Segura¹

¹Institute Cajal, CSIC, Madrid, Spain, ²Institute of Biophysics, Biological Research Center, Hungarian Academy of Sciences, Szeged, Hungary, and ³Department of Obstetrics and Gynecology, Yale University School of Medicine, New Haven, Connecticut, USA •

SUMMARY

• *Sex hormones have developmental trophic actions on neurons and glial cells and activational effects in the adult brain. It has been proposed that sex steroids may interact with peptide trophic factors to induce part of their biological effects in the nervous system. The first evidence of such an interaction was provided by Toran-Allerand et al (Brain Res 1980; 184: 517-524), showing that in explant cultures of fetal rodent hypothalamus, estrogen and insulin have synergistic effects on neurite growth, an effect probably mediated by insulin-like growth factor-1 receptors. Recent data indicate that estrogen and insulin-like growth factor-1 signaling pathways interact on hypothalamic neurons to regulate survival and differentiation and that sex steroids interact with a variety of different trophic signals in vivo to regulate neuroendocrine events. These findings suggest that trophic factors may be involved in the genesis of sex differences in the developing brain and in the maintenance of a sexually differentiated brain function in the adult. (BioinedRev 1997; 7: 67-74)*

INTRODUCTION

• Hormonal steroids modulate behavior, brain physiology and neuroendocrine function by exerting a variety of different

Received for publication 5 April 1997 and accepted 3 June 1997.

Correspondence and reprint requests to Dr Luis M. Garcia-Segura, Institute Cajal, CSIC, Av. Dr. Arce 37, E-28002 Madrid, Spain, Tel: 34 (1) 585 4729, Fax: 34 (1) 585 4754

effects on specific neuronal populations. The mechanism of action of hormonal steroids in the nervous system, as in other tissues, involves the activation of specific nuclear receptors that act as transcription factors for specific genes (1,2). In addition, steroid hormones may modulate neuronal excitability by having rapid non-genomic actions (3). Neural activity is also modulated by sex steroid metabolites produced by nerve cells and by local steroids that are synthesized *de novo* in the brain from cholesterol (4). In this review we examine evidence generated in studies on rodents indicating that sex steroids may act as trophic factors for neural cells and as promoters of neural differentiation and plasticity. This trophic and plastic effect of sex steroids may be mediated in part by interaction with other trophic molecules of peptide nature. In particular, complex interactions of one such molecule, insulin-like growth factor-I (IGF-I), and estradiol have been documented recently in hypothalamic neurons.

SEX HORMONES AND CELL COMMUNICATION IN BRAIN

• Due to the complex pattern of cell interactions in the development of the central nervous system (CNS), it is difficult to determine whether the effect of sex steroids on a given neural cell type is directly exerted on that cell or mediated by other neurons or glial cells. Since many neurons express receptors for gonadal steroids and since gonadal hormones and their metabolites may have a variety of direct rapid effects on neuronal membranes, it is presumable that steroids will directly affect many different neurons throughout the brain. However, indi-

reel effects are probably very important as well. Acting on their target neurons, for example, gonadal hormones may induce the release of neurotransmitters and trophic factors that could affect the development and function of other neurons. Furthermore, by promoting neuronal survival and differentiation of their target cells, gonadal hormones could also affect the survival and differentiation of the neurons that contact these cells.

The action of sex hormones in the brain involves not only neuron-to-neuron communication since glial cells are also affected by sex hormones and their metabolites (5). Glial cells are a source of growth factors for neurons, promoting neuronal survival and differentiation. Several laboratories have shown that glial cells in culture express receptors for sex steroids (6, 7), suggesting that at least in early stages of brain development glial cells may be a direct target of these hormones (6, 7). Furthermore, it has been reported that hypothalamic glia *in situ* expresses estrogen receptors, based on immunohistochemical evidence (8). Sex hormones and neurosteroids affect myelination by acting on oligodendrocytes and Schwann cells (9,10) and modulate the response of nerve tissue to injury by acting on microglia and astroglia (5, 11). In addition to being a target for sex steroids, glial cells are also involved in their metabolism (12) and in the synthesis of neurosteroids (10, 13-15) and participate in the organizational and activational effects of sex steroids on synapse formation (16, 17) and synaptic plasticity (18,19).

GONADAL HORMONES ACT AS NEUROTROPHIC FACTORS

- It is classically recognized that sex steroids exert organizational effects during the critical period for brain sexual differentiation and activational effects in the adult brain (20). During the CNS development, gonadal steroids determine the number of neurons in several brain areas. One of the best studied examples is the sexually dimorphic nucleus of the preoptic area in rodents, which is larger in volume in males than in females (21, 22). This difference in nuclear size is at least partially due to the action of gonadal steroids during the critical period, promoting the survival of a specific population of neurons (23). Another example of a sexually dimorphic structure in the CNS of rodents is the spinal nucleus of the bulbocavernosus (24). Androgens prevent normal cell death of motoneurons of this nucleus (25). These studies suggest that sex steroids may act as neurotrophic factors, promoting the survival of specific neurons or regulating apoptosis (26).

Neurotrophic effects of sex steroids have been characterized *in vitro*. Estradiol promotes the survival of cultured neurons from the amygdala, the hypothalamus and the spinal cord (26a-29). The effect of estradiol on the survival of cultured hypothalamic neurons is mimicked by testosterone, but not by the non-aromatizable androgen dihydrotestosterone (28). The effect

appears to be mediated in part through the estrogen receptor since it is saturable and can be blocked by estrogen receptor antagonists, such as tamoxifen and ICI 182,780, and by the inhibition of estrogen receptor synthesis in the cultures by using an antisense oligonucleotide against estrogen receptor mRNA (28, 29).

In addition to the effects on cell number, sex steroids also affect neuronal differentiation in several brain areas, modulating the growth of axons, the number and branching of dendrites and the number of dendritic spines. Estradiol modulates, for instance, the growth of neurites in hypothalamic neurons in explant cultures (30,31) and monolayer cultures (29,32, 33), while in the spinal nucleus of the bulbocavernosus, androgens regulate dendritic outgrowth during the neonatal period and dendritic retraction during the pubertal period (34, 35). Gonadal steroid-dependent sex differences in neuronal processes have been detected in other brain structures, including the preoptic area (36) and hippocampus (37). These effects of gonadal hormones on the differentiation of neuronal processes are mediated, at least in part, by the modulation of cytoskeletal proteins (29, 32, 38).

Hormonal modifications in the growth of neuronal processes may result in changes in the pattern of neuronal connectivity. It is well established that during the critical period gonadal steroids influence the formation of synaptic contacts among neurons, resulting in specific sex differences in neuronal connectivity. Furthermore, the effect of sex hormones on synapses is not restricted to the developmental period. Synapses are plastic structures that may be modified in the adult brain in response to changing physiological conditions, including modifications in hormone levels (18, 39). Gonadal hormones influence synapse formation in areas of the CNS involved in the control of reproductive behavior, such as the ventromedial hypothalamic nucleus, lateral septum and the amygdala and in areas involved in the control of the release of pituitary hormones, such as the hypothalamic arcuate nucleus and the preoptic area. In addition, gonadal hormones may affect neuronal connectivity in cognitive areas, such as the hippocampal formation and the cerebral cortex (18,37, 39-41).

INTERACTION OF SEX STEROIDS AND TROPHIC FACTORS

- The complexity of gonadal steroid action on the brain is not only a consequence of the complexity of cellular interactions in the neural tissue. Interactions between the signaling cascades of membrane receptors for peptidic factors and the steroid hormone receptors should also be taken into consideration. One of the best characterized neurotrophic factors, and the first one discovered, is nerve growth factor (NGF). In some brain areas and during specific developmental stages estradiol and NGF may affect the same target neurons. Genes encoding

NGF and the low-affinity receptor for NGF (p75) are expressed in subpopulations of neurons of hypothalamic neuroendocrine areas (42). Furthermore, it has been shown that estrogen receptors colocalize with p75 in cholinergic neurons of the basal forebrain of developing rodents (43). Estradiol may therefore exert part of its trophic effects modulating the action of NGF. This is further supported by the fact that estradiol is able to modulate the levels of NGF receptors in PC 12 cells, in cerebral cortical cultures and in dorsal root ganglion neurons of adult female rats (44-46). Moreover, putative estrogen responsive elements have been identified in the NGF gene and in the genes for the NGF tyrosine kinase (trkA) and low-affinity receptors (46). By regulating trkA or p75 levels, estradiol may affect programmed neuronal death (26,47).

A putative estrogen receptor element has been identified also in the gene encoding brain-derived neurotrophic factor (48). The receptor trkB for this neurotrophin is expressed in neurons from hypothalamic neuroendocrine areas, as well as trkC, which is the receptor for neurotrophin-3 (42).

In addition to neurotrophins, other trophic factors may also be relevant in regard to the action of sex steroids in the brain. These include basic fibroblast growth factor (bFGF) and transforming growth factors- α (TGF- α) and - β (TGF- β), which appear to be involved in the maturation of the neuroendocrine hypothalamus and/or in the regulation of luteinizing hormone-releasing hormone (LHRH) neurons (49-52). Sex steroids may interact with some of these factors to exert their effects on neurons and glial cells. Also, estrogen modulates the expression of TGF- β in the hypothalamus (50). This factor is produced by hypothalamic glial cells and appears to regulate LHRH neurons (49, 50, 53). Gene expression of TGF- β increases in glial cells at the time of puberty in regions of the hypothalamus involved in LHRH control (50).

Another factor that may be involved in the effects of gonadal hormones on brain is IGF-I. This factor is highly expressed by glia and neurons of the developing brain (54,55) and has prominent trophic actions, stimulating the survival and differentiation of specific neural cell populations, including hypothalamic neurons in culture (28, 56). IGF-I may also act as a hormonal signal and may be involved in the feed-back regulation of growth hormone by affecting the synthesis or release of growth hormone-releasing hormone or somatostatin by hypothalamic neurons (57,58). IGF-I may also affect the reproductive axis by modulating the secretion of LHRH (59-61).

INTERACTION OF ESTRADIOL AND INSULIN-LIKE GROWTH FACTOR I

In several tissues and cell types estrogen upregulates IGF-I gene expression (62-68) and modulates IGF-I action by

affecting the levels of IGF-I receptors (63, 69) and IGF-binding proteins (70,73) which appear to modulate the availability of IGF-I to target cells. Likewise, IGF-I may regulate steroid hormone action by stimulating the synthesis of steroid hormones (74-76) and steroid hormone receptors (77-79). Furthermore, IGF-I, as other growth factors, may activate the estrogen receptor in different cell lines, including neuroblastoma cells (80-85). This activation of the estrogen receptor occurs in the absence of the hormone. It is mediated by IGF-I receptor membrane-associated signaling pathways (82,85) and involves the activation function 2 domain of the estrogen receptor in neuroblastoma cells (85) and the activation function 1 domain in other cell lines (81,82).

The first evidence of an interaction between IGF-I and estrogen on neural cells was obtained in explant cultures of fetal rodent hypothalamus where estrogen and insulin have synergistic effects on neurite growth, an effect that is probably mediated through IGF-I receptors (86). Furthermore, estrogen modulates IGF-I receptors and binding proteins in monolayer hypothalamic cultures (87) and estrogen effects on hypothalamic neuronal survival and neurite growth are mediated by IGF-I (29). Hypothalamic cultures exposed to estradiol or IGF-I show a significant increase in neuronal survival and in the growth of neuronal processes. Inhibition of IGF-I synthesis in the cultures with an antisense oligonucleotide to mRNA results in a significant decrease in the stimulatory effects of 17 β -estradiol on the number of neurons and the extension of neuronal processes (29). This indicates that IGF-I is necessary for the manifestation of the hormonal effect, suggesting that estradiol may induce neuronal survival and differentiation by activation of IGF-I signaling cascades. Also, the effect of IGF-I on the survival and differentiation of hypothalamic neurons depends on the estrogen receptor (29). Both the pure estrogen receptor antagonist ICI 162,780 or an antisense oligodeoxynucleotide to the estrogen receptor mRNA block the effects of IGF-I. This indicates that estrogen receptors are necessary for the action of IGF-I on hypothalamic survival and neurite growth, suggesting that IGF-I may activate, either directly or indirectly, estrogen receptors.

Interactions of sex steroids and IGF-I are also detected in the brain. IGF-I levels increase in the arcuate nucleus and median eminence of male and female rats at the time of puberty, with males reaching higher levels than females (88). This sex difference is abolished by perinatal administration of testosterone to females. Furthermore, in females there is an abrupt increase in IGF-I levels between the morning and the afternoon of the first proestrus. Henceforth, IGF-I levels fluctuate in the arcuate nucleus and median eminence according to the different stages of the estrus cycle. IGF-I levels are high in the afternoon of proestrus, after the peak of estrogen in plasma, remain increased on the morning of the following day and then decrease to basal conditions by the morning of metestrus (88). In addi-

lion, IGF-I levels decrease by ovariectomy and increase in a dose-dependent manner when ovariectomized rats are injected with 17 β -estradiol, an effect that is blocked by the simultaneous administration of progesterone (88). These changes, linked to fluctuations of gonadotrophins, suggest that the modulation by sex steroids of the levels of IGF-I in the mediobasal hypothalamus may be related to LHRH regulation. Moreover, these changes do not reflect modifications in the local synthesis of IGF-T (61,88; see also 89), being probably related to differences in the accumulation of IGF-I from extracerebral sources (61).

CONCLUSION

M Effects of sex hormones on different neuronal populations result in the development of local sex differences in neuronal number and connectivity. Part of these actions of sex hormones may be exerted by the modulation of trophic factor signaling among brain cells. Direct evidence of an interaction of sex hormones and trophic factors on neuronal survival and differentiation has been obtained in the hypothalamus. Furthermore, a variety of trophic factors interact with gonadal hormones in the brain to regulate neuroendocrine events. Recent findings in hypothalamic cultures suggest that the survival and differentiation of neurons expressing hormone receptors and the resulting sexual differentiation of brain circuitry and function may depend on the coordinated play of hormonal and neurotrophic signals acting on neurons and glial cells.

ACKNOWLEDGEMENTS

M This work was supported by Pundacidn Ramdn Areces, DGICYT. NATO (CRG930387), NIH (HD13587), and Pundacidn Endocrinologfa y Nutricidn.

REFERENCES

1. Beato M. Gene regulation by steroid hormones. *Ce//1989*; 56: 335-344
2. Evans RM. The steroid and thyroid hormone receptor superfamily. *J'ci'fMce* 1988; 240: 889-895
3. McEwen BS. Non-genomic and genomic effects of steroids on neural activity. *7/-fMfAv f/zormaco/^c*; 1991; 12: 141-147
4. Baulieu EE, Rohel P. Neurosteroids: a new brain function? *V &<?m/c/ gi'oc/z?m MrV 5m/* 1990; 37: 395-403
5. Garcia-Segura LM, Chowen J A, Naftolin F. Endocrine glia: roles of glial cells in the brain actions of steroid and thyroid hormones and in the regulation of hormone secretion. *FmMfM»Hm(?&)LTm* 1996; 17: 180-211
6. Jung-Testas I, Renoir JM, Gasc JM, Baulieu EE. Estrogen-inducible progesterone receptor in primary cultures of rat glial cells. *Exp Ce;/?ej* 1991; 193: 12-19
7. Santagati S, Melcangi RC, Celotti P, Martini L, Maggi A. Estrogen receptor is expressed in different types of glial cells in culture. *JTVeMroc/iem* 1994; 63: 2058-2064
8. Langub MC, Watson RE. Estrogen receptor-immunoreactive glia, endothelia, and ependyma in guinea pig preoptic area and median eminence: electron microscopy. *E«<Jocn-MoJogy* 1992; 130:364-372
9. Jung-Testas I, Schumacher M, Robel P, Baulieu EE. Actions of steroid hormones and growth factors on glial cells of the central and peripheral nervous system. *V ^(em/^ iKocAem Mo/ gz'o/* 1994; 48: 145-154
10. Koenig HL, Schumacher M, Ferzaz B, Do Thi AN, Ressouches A, Guennoun R *ef a/*. Progesterone synthesis and myelin formation by Schwann cells. *.Science* 1995; 268: 1500-1503
11. Garcia-Estrada J, Del Rio JA, Luquin S, Soriano E, Garcia-Segura LM. Gonadal hormones down-regulate reactive gliosis and astrocyte proliferation after a penetrating brain injury. *gmm 7?gj* 1993; 628: 271-278
12. Melcangi RC, Celotti P, Castano P, Martini L. Differential localization of the 5 α -reductase and the 3 α -hydroxysteroid dehydrogenase in neuronal and glial cultures. *EM&iCRM/o^* 1993; 132:1252-1259
13. Akwa Y, Sananes N, Gouezou M, Robel PI, Baulieu EE, Le Goascogne C. Astrocytes and neurosteroids: Metabolism of pregnenolone and dehydroepiandrosterone. Regulation by cell density. *JCe//gi'o/* 1993; 121: 135-143
14. Hu ZY, Bourreau E, Jung-Testas I, Robel P, Baulieu EE. Neurosteroids. Oligodendrocyte mitochondria convert cholesterol to pregnenolone. *f me M??Z !/«%/ j'c*; *CAM* 1987; 84: 8215-8219
15. Kabbadj K, El-Etr M, Baulieu EE, Robel P. Pregnenolone metabolism in rodent embryonic neurons and astrocytes. *G&z1993*;7:170-175
16. Chowen JA, Busiguina S, Garcia-SeguraLM. Sexual dimorphism and sex steroid modulation of glial fibrillary acidic protein messenger RNA and immunoreactive levels in the rat hypothalamus. */Vewrojc;!g/zce* 1995; 69: 519-532
17. Garcia-Segura LM, Duenas M, Busiguina S, Naftolin F, Chowen JA. Gonadal hormone regulation of neuronal-

- glial interactions in the developing neuroendocrine hypothalamus. *J Steroid Biochem Mol Biol* 1995; 53: 293-298
18. Garcia-Segura LM, Chowen JA, Parducz A, Naftolin F. Gonadal hormones as promoters of structural synaptic plasticity: cellular mechanisms. *Prog Neurobiol* 1994; 44:
 19. Garcia-Segura LM, Luquin S, Parducz A, Naftolin F. Gonadal hormone regulation of glial fibrillary acidic protein immunoreactivity and glial ultrastructure in the rat neuroendocrine hypothalamus. *Glia* 1994; 10: 59-69
 20. Arnold AP, Breedlove SM. Organizational and activational effects of sex steroids on brain and behavior: A reanalysis. *Horm Behav* 1985; 19: 469-498
 21. Gorski RA, Gordon JH, Shryne JE, Southam AM. Evidence for a morphological sex difference within the medial preoptic area of the rat brain. *Brain Res* 1978; 148: 333-346
 22. Gorski RA, Harlan RE, Jacobson CD, Shryne JE, Southam AM. Evidence for the existence of a sexually dimorphic nucleus in the preoptic area of the rat. *J Comp Neurol* 1980; 198: 529-539
 23. Dodson RE, Shryne JE, Gorski RA. Hormonal modification of the number of total and late-arising neurons in culture: sex differences and estrogen effects. *J Neurosci Res* 1988; 33: 266-281
 24. Breedlove SM, Arnold AP. Hormonal control of developing neuromuscular system. I. Sensitive periods for the androgen-induced masculinization of the rat spinal nucleus of the bulbocavernosus. *J Neurosci* 1983; 3: 424-432
 25. Norclee E.T, Nordcn KW, Sengelaub DR, Arnold AP. Androgens prevent normally occurring cell death in a sexually dimorphic spinal nucleus. *Science* 1985; 229: 671-673
 26. Arumae U. Neurotrophins: neural antiapoptotic molecules with neurite growth-promoting properties. *Biomed Rev* 1995;4: 15-28
 - 26a. Arimatsu Y, Halanaka H. Estrogen treatment enhances survival of cultured fetal rat amygdala neurons in a defined medium. *Dev Brain Res* 1986; 26: 151-159
 27. Hauser KF, Toran-Allerand CD. Androgen increases the number of cells in fetal mouse spinal cord cultures: implications for motoneuron survival. *Brain Res* 1989; 485: 157-164
 28. Chowen JA, Torres-Aleman I, Garcia-Segura LM. Tropic effects of estradiol on fetal rat hypothalamic neurons. *Neuroendocrinology* 1992; 56: 895-901
 29. Dueñas M, Torres-Aleman I, Naftolin F, Garcia-Segura LM. Interaction of insulin-like growth factor-I and estradiol signalling pathways on hypothalamic neuronal differentiation. *Neuroscience* 1996; 74: 531-539
 30. Toran-Allerand CD. Sex steroids and the development of the newborn mouse hypothalamus and preoptic area *in vitro*, implications for sexual differentiation. *Brain Res* 1976; 106: 407-412
 31. Toran-Allerand CD, Hashimoto K, Greenough WT, Saltarelli N. Sex steroids and the development of the newborn mouse hypothalamus *in vitro*: III. Effects of estrogen on dendritic differentiation. *Dev Brain Res* 1983; 7: 97-101
 32. Ferreira A, Caceres A. Estrogen-enhanced neurite growth: Evidence for a selective induction of tau and stable microtubules. *J Neurosci* 1991; 11: 392-400
 33. Diaz H, Lorenzo A, Carrer HF, Caceres A. Time lapse study of neurite growth in hypothalamic dissociated neurons in culture: sex differences and estrogen effects. *J Neurosci Res* 1992;33:266-281
 34. Kurz EM, Sengelaub DR, Arnold AP. Androgens regulate the dendritic length of mammalian motoneurons in adulthood. *Science* 1986; 232: 395-398
 35. Goldstein LA, Kurz EM, Sengelaub DR. Androgen regulation of dendritic growth and retraction in the development of a sexually dimorphic spinal nucleus. *J Neurosci* 1990; 10:935-946
 36. Raisman G, Field PM. Sexual dimorphism in the neuropil of the preoptic area of the rat and its dependence on neonatal androgen. *Brain Res* 1973; 54: 1-29
 37. Gould E, Woolley CS, Frankfurt M, McEwen BS. Gonadal steroids regulate dendritic spine density in hippocampal pyramidal cells in adulthood. *J Neurosci* 1990; 10: 1286-1291
 38. Lorenzo A, Diaz H, Carrer H, Caceres A. Amygdala neurons *in vitro*: neurite growth and effects of estradiol. *J Neurosci Res* 1992; 33: 418-435
 39. Matsumoto A. Synaptogenic action of sex steroids in de-

- veloping and adult neuroendocrine brain. *Psychoneuroendocrinology* 1991; 16:25-40
40. Olmos G, Naftolin F, Percz J, TranquePA, Garcia-Segura LM. Synaptic remodelling in the rat arcuate nucleus during the estrous cycle. *Neuroscience* 1989; 32: 663-667
 41. Parducz A, Percz J, Garcia-Segura LM. Estradiol induces plasticity of GABAergic synapses in the hypothalamus. *Neuroscience* 1993; 53:395-401
 42. Berg von der Emtle K, Decs WL, Hiney JK, Hill DF, Dissen GA, Costa ME *et al.* Neurotrophins and the neuroendocrine brain: different neurotrophins sustain anatomically and functionally segregated subsets of hypothalamic dopaminergic neurons. *J Neurosci* 1995; 15: 4223-4237
 43. Toran-Allerand CD, Miranda RC, Bentham W, Sohrabji F, Brown EJ, Hochberg RB *et al.* Estrogen receptors co-localize with low-affinity NGF receptors in cholinergic neurons of the basal forebrain. *Proc Natl Acad Sci USA* 1992;89:4668-4672
 44. Sohrabji F, Greene LA, Miranda RC, Toran-Allerand D. Reciprocal regulation of estrogen and NGF receptors by their ligands in PC 12 cells. *J Neurobiol* 1994; 25: 974-988
 45. Sohrabji F, Miranda C, Toran-Allerand CD. Estrogen differentially regulates estrogen and nerve growth factor receptor mRNA in adult sensory neurons. *J Neurosci* 1994 14: 459-471
 46. Toran-Allerand CD. Mechanisms of estrogen action during neural development: mediation by interactions with the neurotrophins and their receptors? *J Steroid Biochem Mol Biol* 1996; 56: 169-178
 47. Frade JM, Rodrigucz-Tebar A. Barde YA. Induction of cell death by endogenous nerve growth factor through its p75 receptor. *Nature* 1996; 383: 166-168
 48. Sohrabji F, Miranda RC, Toran-Allerand CD. Identification of a putative estrogen receptor element in the gene encoding brain-derived neurotrophic factor. *Proc Natl Acad Sci USA* 1995; 92: 11110-11114
 49. Ojeda SR, Dissen G A. Junior MP. Neurotrophic factors and female sexual development. *Front Neuroendocrinol* 1992 13:120-162.
 50. Ma YJ, Junier MP, Costa ME, Ojeda SR. Transforming growth factor- β gene expression in the hypothalamus is developmentally regulated and linked to sexual maturation. *Neuron* 1992;9:657-670
 51. Melcangi RC, Galbiati M, Messi E, PivaF, Martini L, Motta M. Type 1 astrocytes influence luteinizing hormone-releasing hormone release from the hypothalamic cell line GT1 -1: is transforming growth factor- β the principle involved? *Endocrinology* 1995; 136:679-686
 52. Wetsel WC, Hill DF, Ojeda SR. Basic fibroblast growth factor regulates the conversion of pro-luteinizing hormone-releasing hormone (Pro-LHRH) to LHRH in immortalized hypothalamic neurons. *Endocrinology* 1996; 137: 2606-2616
 53. Voigt P, Ma YJ, Gonzalez D, Fahrenbach WH, Wetsel WC, Berg von der Emde K *et al.* Neural and glial-mediated effects of growth factors acting via tyrosine kinase receptors on luteinizing hormone-releasing hormone neurons. *Endocrinology* 1996; 137: 2593-2605
 54. Bondy C, Werner H, Roberts CT, LeRoith D. Cellular pattern of type I insulin-like growth factor receptor gene expression during maturation of the rat brain: Comparison with insulin-like growth factors I and II. *Neuroscience* 1992; 46: 909-923
 55. Garcia-Segura LM, Perez J, Pons S, Rejas MT, Torres-Aleman I. Localization of insulin-like growth factor I (IGF-I)-like immunoreactivity in the developing and adult rat brain. *Brain Res* 1991; 560: 167-174
 56. Torres-Aleman I, Naftolin F, Robbins RJ. Trophic effect of insulin-like growth factor-I on fetal rat hypothalamic cells in culture. *Neuroscience* 1990; 35: 601-608
 57. Berelowitz M, Szabo M, Frohman LA, Firestone S, Chu L, HintzRL. Somatomedin C mediates growth hormone negative feedback by effects on both the hypothalamus and the pituitary. *Science* 1981; 212: 1279-1281
 58. Tannenbaum GS, Guyda HJ, Posner BI. Insulin-like growth factors: A role in growth hormone negative feedback and body weight regulation via brain. *Science* 1983; 220: 77-79
 59. Bourguignon JP, Gerard A, Alvarez Gonzalez ML, Franchimont P. Acute suppression of gonadotropin-releasing hormone secretion by insulin-like growth factor-I and subproducts: an age-dependent endocrine effect. *Neuroendocrinology* 1993; 58: 525-530
 60. Hiney JK, Ojeda SR, Les Decs W. Insulin-like growth factor-I: a possible metabolic signal involved in the regulation of female puberty. *Neuroendocrinology* 1991; 54:

- 420-423
61. Hiney **JK**, Sri vastava V, Nyherg CL, OjedaSR, Les Dees W. Insulin-like growth factor-I of peripheral origin acts centrally to accelerate the initiation of female puberty. *Endocrinology* 1996; 137: 3717-3728
 62. Gahary A, Chakrabarti S, Murphy LJ. Localization of the sites of synthesis and action of insulin-like growth factor-I in the rat uterus. *Mol Endocrinol* 1990; 4: 191-195
 63. Hemandcz ER. Regulation of the genes for insulin-like growth (actor (IGF) I and II and their receptors by steroids and gonadotropins in the ovary. *J Steroid Biochem Mol Biol* 1995; 53:219-221.
 64. Kapur S, Tamada H, Dey SK, Andrews GK. Expression of insulin-like growth factor-I (IGF-I) and its receptor in the peri-implantation mouse uterus, and cell-specific regulation of IGF-I gene expression by estradiol and progesterone. *Biol Reprod* 1992; 46: 208-219
 65. Miehels KM, Ee WH, Seltzer A, Saavedra JM, Bondy CA. Up-regulation of pituitary [¹²⁵I] insulin-like growth factor-I (IGF-I) binding and IGF-I gene expression by *estrogen*. *Endocrinology* 1993; 132:23-29
 66. Murphy E.T, Murphy LC, Friesen HG. Estrogen induces insulin-like growth factor-I expression in the rat uterus. *Mol Endocrinol* 1987; 1:445-450
 67. Simmen RCM, Simmen FA, Hoi'ig A, Farmer SJ, Bazer FW. Hormonal regulation of insulin-like growth factor gene expression in pig uterus. *Endocrinology* 1990; 127: 2166-2174
 68. Umayahara Y, Kawamori R, Watada H, Imano E, Iwama N, Morishima T *et al*. Estrogen regulation of the insulin-like growth factor-I gene transcription involves an AP-1 *enhimccr*. *JBiolChein* 1994; 269: 16433-16442
 69. Wimalascna J, Median D, Dostal R, Foster JS, Cameron M, Smith M. Growth factors interact with estradiol and gonadotrophins in the regulation of ovarian cancer cell growth and growth factor receptors. *Oncol Res* 1993; 5: 325-337
 70. Krywicki RF, Figueroa JA, Jackson JG, Kozelsky TW, Shimasaki S, von Hoff *DDetal*. Regulation of insulin-like growth factor-binding proteins in ovarian cancer cells by oestrogen. *EurJ Cancer* 1993; 29A: 2015-2019
 71. Molnar P, Murphy LJ. Effects of oestrogen on rat uterine expression of insulin-like growth factor-binding proteins. *J Mol Endocrinol* 1994; 13: 59-67
 72. Owens PC, Gill PG, De Young NJ, Weger MA, Knowels SE, Moyse KJ. Estrogen and progesterone regulate secretion of insulin-like growth factor-binding proteins by human breast cancer cells. *Biochem Biophys Res Commun* 1993; 193:467-473
 73. Yallampalli C, Rajaraman S, Nagamani M. Insulin-like growth factor binding proteins in the rat uterus and their regulation by oestradiol and growth hormone. *J Reprod Fertil* 1993; 97: 501-505
 74. Constantino CX, Keyes PL, Kostyo JL. Insulin-like growth factor-I stimulates steroidogenesis in rabbit luteal cells. *Endocrinology* 1991; 128: 1702-1708
 75. Erickson GF, Garzo VG, Magoffin DA. Insulin-like growth factor-I (IGF-I) regulates aromatase activity in human granulosa luteal cells. *J Clin Endocrinol Metab* 1989; 69: 716-724
 76. Hernandez ER, Resnick CE, Svoboda ME, Van Wyk J, Payne DW, Adashi EY. Somatomedin-C/insulin-like growth factor-I as an enhancer of androgen biosynthesis by cultured rat ovarian cells. *Endocrinology* 1988; 122: 1603-1612
 77. Aronica SM, Katzenellenbogen BS. Progesterone receptor regulation in uterine cells: stimulation by estrogen, cyclic adenosine 3',5'-monophosphate, and insulin-like growth factor-I and suppression by antiestrogens and protein kinase *inhibitors*. *Endocrinology* 1991; 128: 2045-2052
 78. Cho H, Aronica SM, Katzenellenbogen BS. Regulation of progesterone receptor gene expression in MCF-7 breast cancer cells: a comparison of the effects of cyclic adenosine 3',5'-monophosphate, estradiol, insulin-like growth factor-I, and serum factors. *Endocrinology* 1994; 134:658-664
 79. Katzenellenbogen BS, Norman MJ. Multihormonal regulation of the progesterone receptor in MCF-7 human breast cancer cells: interrelationships among insulin/insulin-like growth factor-I, serum, and *estrogen*. *Endocrinology* 1990; 126:891-898
 80. Aronica SM, Katzenellenbogen BS. Stimulation of estrogen receptor-mediated transcription and alteration in the phosphorylation state of the rat uterine estrogen receptor by estrogen, cyclic adenosine monophosphate, and *insulin-like growth factor-I*. *Mol Endocrinol* 1993; 7:743-752

-
- 8). Ignar-Trowbridge DM, Pimentel M, Parker MG, McLachlan JA, Korach KS. Peptide growth factor cross-talk with the estrogen receptor requires the A/B domain and occurs independently of protein kinase C. *Endocrinology* 1996; 137: 1735-1744
82. Kato S, Enclou H, Masuhiro Y, Kitamoto T, Uchiyama S, Sasaki H *et al.* Activation of the estrogen receptor through phosphorylation by mitogen-activated protein kinase. *Science* 1995; 270: 1491-1494
83. Ma ZQ, Santagali S, Patrone C, Pollio G, Vegeto E, Maggi A. Insulin-like growth factor activates estrogen receptor to control the growth and differentiation of the human neuroblastoma cell line SK-N-BE. *Mol Endocrinol* 1994; 8: 910-918
84. Newton CJ, Buric R, Trapp T, Brockmeier S, Fagotto U, Stalla GK. The insulin-like growth factor (IGF) transduces growth factor signals. *J Steroid Biochem Mol Biol* 1994; 48:481-486
85. Patrone C, Ma ZQ, Pollio G, Agrati P, Parker MG, Maggi A. Cross-coupling between insulin and estrogen receptor in human neuroblastoma cells. *Mol Endocrinol* 1996; 10:499-507
86. Toran-Allerand CD, Ellis E, Pfenninger KH. Estrogen and insulin synergism in neurite growth enhancement *in vitro*: Mediation of steroid effects by interactions with growth factors? *Dev Brain Res* 1988; 41: 87-100
87. Pons S, Torres-Aleman I. Estradiol modulates insulin-like growth factor-I receptors and binding proteins in neurons from the hypothalamus. *J Neuroendocrinol* 1993; 5: 267-
88. Dueñas M, Euquin S, Chowen JA, Torres-Aleman I, Naftolin F, Garcia-Segura EM. Gonadal hormone regulation of insulin-like growth factor-I-like immunoreactivity in hypothalamic astroglia of developing and adult rats. *Neuroendocrinology* 1994; 59: 528-538
89. De Cupis A, Favoni RE. Oestrogen/growth factor cross-talk in breast carcinoma: a specific target for novel antiestrogens. *Trends Pharmacol Sci* 1997; 18: 245-251