

## INTERACTION OF HEAT SHOCK PROTEIN (hsp90) WITH THE CYTOSKELETON: POTENTIAL IMPLICATIONS IN INTRACELLULAR TRANSPORT

Christos Slournaras, Andrew N. Margioris, and Aehille Gravanis

*Department of Biochemistry, Clinical Chemistry, and Pharmacology, School of Medicine, University of Crete, Heraklion, Greece*

### SUMMARY

*In this article we will summarize the details concerning the association of 90 kDa heat shock protein (hsp90) with cytoskeletal structures and we will discuss the potential involvement of these interactions in the translocation of steroid hormone receptors to the nucleus.*

*In cultured mammalian cells hsp90 has been found to be colocalized with both microtubules and cytokeratin intermediate filaments, whereas no association with actin filaments and vimentin intermediate filaments has been established.*

*The colocalization of hsp90 with microtubules and cytokeratin in intact cells raises the possibility that cytoskeletal structures could be used as "rails" for the direct movement of the steroid hormone receptor via association-dissociation with hsp90 molecules from the cytoplasmic site of translocation to the nuclear site of action.*

### INTRODUCTION

The cytoskeleton, the integral structure network of the cell, has been previously compared to the bones and muscles of the body. However, its function is much more complex in many ways allowing participa-

tion in the regulation of cell shape, cell movement, cell division, endocytosis and adhesion. The cytoskeleton is composed of three well characterized classes of filamentous structures: (i) microtubules which are hollow cylinders of approximately 25 nm outer diameter and indefinite length; they are constructed from identical subunits of tubulin, a globular protein of approximately 55 kDa. (ii) actin filaments which are 6-8 nm in diameter; they have the basic structure of double-stranded helix and are made from assembled actin molecules and the associated tropomyosin and troponin. and (iii) intermediate filaments which are composed of different types of proteins assembled in structures of 8-11 nm in diameter; there are five major types of intermediate filaments described according to their protein subunits: vimentin, desmin, cytokeratins, neurofilament proteins (NF-L, NF-M, NF-H), and glial fibrillary acidic protein (GFAP), which are specifically expressed in particular cell types.

The research on the cellular organization and distribution of the filamentous structures was made possible through the application of labelled antibody procedures. The first of these studies some 20 years ago used antibodies against actin to visualize actin filaments (1) similar studies with antibodies against tubulin (2) and intermediate filament proteins (5,6) followed soon. Through the following 2 years the immunofluorescence mi-

scopy techniques were fully established providing a convenient and useful tool for determining the organization and distribution of cytoskeletal proteins and structures, as well as their association with each other or with other cellular components.

The analysis of cytoskeletal preparations supports the understanding that the three major cytoskeletal structures are interconnected and their functions are coordinated.

Compared to the remarkable progress at the structural level, our knowledge of the biochemical basis of these interactions as well as of the physiological role of the cytoskeletal elements are very poor. One of the most striking features of the concept of interaction of cytoskeletal structures, is their involvement in intracellular transport processes. It is a challenging aim to search for lines of evidence regarding the use of cytoskeletal structures in the intracellular protein transport. Studies of the colocalization of hsp90 with the cytoskeletal proteins provide an interesting area to approach such an aim.

The cellular function of hsp90 is still unknown (7), despite of its ubiquitous presence among eukaryotic cells. Hsp90 associates with the oncoprotein pp60<sup>SIL</sup> as well as with some other sarcoma virus transforming proteins (8-10). Hsp90 is also associated with steroid receptors in their untransformed, non-DNA binding state in cytosol preparation (11-13). Two functions have been described to receptor-bound hsp90: (i) masking the receptor DNA binding domain, and (ii) maintenance of the ligand binding domain in a functional hormone-binding conformation (14). Thus, modulation of important functions of the steroid receptors has been attributed to hsp90, including prevention of DNA binding and optimization of transcriptional activity.

Searching to attribute a biological role and function for hsp90, efforts were undertaken recently to study the interaction of this protein with the cytoskeleton. Isolated hsp90 showed the ability to bind to F-actin (15,16), while no association of the protein with actin filaments could be demonstrated in intact cells. Using immunofluorescence microscopy, the colocalization of hsp90 with microtubules has been

shown in cytosolic preparations of several mammalian cell lines (17,18). Moreover, the interaction of hsp90 with cytokeratin intermediate filaments has been also postulated in a human endometrial adenocarcinoma cell line (19).

In the present paper we will summarize the details of our studies concerning the association of hsp90 with cytoskeletal proteins and structures and we will discuss the potential involvement of this interaction in the intracellular translocation of steroid hormone receptors.

### **hsp90 EXPRESSION IN VARIOUS CELL TYPES**

- A wide range of cells, when exposed to elevated temperatures or other environmental stresses, synthesize specific proteins called heat shock proteins (20,21). They play important roles in cellular functions, suggesting that they are essential for living cells (22-24). Using immunofluorescence microscopy techniques, hsp90 has been found almost exclusively in the cytoplasm of chicken embryo fibroblasts (25), HeLa cells (26), and *Drosophila* cells (27). Recent studies demonstrated the presence of hsp90 also in the nucleus (25,28,29).

The determination of the cellular localization of hsp90 and particularly of its colocalization with filamentous structures could provide us with important clues for its function. Collier and Schlesinger (25) reported a fibrous distribution of hsp90 in less dense edges of chicken fibroblasts. Sanchez et al (17) showed that monoclonal antibodies against tubulin and hsp90 could be used to immunoadsorb hsp90 and tubulin from the cytoplasm of L929 fibroblasts. Moreover, they reported a colocalization of hsp90 with cytoplasmic microtubules.

However, other groups using two distinct monoclonal antibodies, which immunoprecipitate both free hsp90 and hsp90 complexed with other proteins, reported no immune-specific coadsorption of either  $\alpha$ - or  $\beta$ -tubulin (30).

Conclusive data concerning an association of hsp90 with filamentous structures were provided only recently (18,19). If hsp90 is colocalized with cytoskeletal

elements in their filamentous form, it should also be present in the Triton-insoluble cytoskeleton. Based on such an assumption, Fostinis et al (19) prepared this fraction from Ishikawa human endometrial cells. Indeed, when these preparations were stained with anti-hsp90 antibodies, filamentous structures were clearly observed, indicating that hsp90 is present in the cytosol- and membrane-free cellular fragment of cytoskeleton (Fig. 1).

#### hsp90 DOES NOT SEEM TO ASSOCIATE WITH F-ACTIN

- Koyasu et al first reported that hsp90 purified from a mouse lymphoma cell line showed the ability to bind to purified F-actin (15). This binding which was saturable in a molar<sup>1</sup> ratio of about 1 hsp90 dimer to 10 actin molecules was proposed to be Ca<sup>2+</sup>-dependent, since it was regulated by calmodulin (16). It was concluded that Ca<sup>2+</sup>-calmodulin inactivates the ability of hsp90 to bind to F-actin by forming a Ca<sup>2+</sup>-calmodulin-hsp90 complex. The K<sub>D</sub>




Figure 1. Immunofluorescence microscopy in preparations of the Triton-insoluble cytoskeleton of Ishikawa cells: Immunostaining of hsp90 with anti-hsp90 antibodies, clearly showing filamentous structures.

values for the binding of hsp90 to F-actin were found to be relatively high, indicating a weak interaction (16). Moreover, hsp90 was dissociated from F-actin by the binding of tropomyosin to F-actin. However, this association was demonstrated only with purified hsp90 and F-actin, when both components were present in solution in high concentrations'. No localization of hsp90 with stress fibers could be demonstrated in cytosol preparations or in intact cells (-15). Thus, the question remained open if hsp90 really associates with actin filaments in living cells.

Using double immunofluorescence staining, a much more detailed study on the cellular colocalization of hsp90 with actin filaments has been performed by Fostinis et al (19). In this work, Ishikawa human endometrial cells were treated with 10<sup>-5</sup>M cytochalasin B in order to destroy actin filaments; this was followed by double staining with rhodamine phalloidin and anti-hsp90 antibodies. Under these experimental conditions the actin filament staining of the cells totally disappeared, while residual filamentous structures were labelled with anti-hsp90 antibodies (19). From these results it was concluded that hsp90 is not associated with actin filaments in cultured Ishikawa cells. These observations supported the suggestion that the *in vitro* reported binding of hsp90 to F-actin does not reflect the real situation in the living cell. Thus, the hypothesis regarding the involvement of hsp90 in the transport of biologically important proteins, such as steroid hormone receptors, through their interaction with actin (16), seems to be unlikely,


#### THE COLOCALIZATION OF hsp90 WITH MICROTUBULES

- Recent studies on the coadsorption of hsp90 and tubulin in cytosolic preparations with several monoclonal anti-hsp90 and anti-tubulin antibodies provide inconclusive data (17,30). It was previously reported that hsp90 has a rather diffused cytoplasmic distribution without relation to filamentous structure<sup>^</sup> (25,27,31). However, some of these studies were performed with round-shaped cells having a relatively small cytoplasmic volume, such as *Drosophila* or HeLa cells (27,31), where microtubules are much more difficult to be observed. Moreover, the mitotic spindle could not be observed because these cells were not dividing.

distribution of hsp90 in intact cultured PtK cells using indirect immunofluorescence microscopy with a monoclonal anti-hsp90 antibody (AC88). They showed that in interphase cells the pattern of the fluorescence staining was distributed in a filamentous way throughout the cytoplasm and in metaphase cells it was located on the mitotic spindle. In addition these studies have further provided biochemical evidence for an association of hsp90 with tubulin. Indeed, anti-tubulin and AC88 antibodies were able to immunoadsorb hsp90 and tubulin, respectively, from the cytosol of PtK cells. These results suggested that hsp90 seems to associate with tubulin-containing complexes in intact PtK cells. Recently, the same group presented a more detailed study on the colocalization of hsp90 with microtubules in mammalian cells (18). In this report, two antibodies against hsp90, a monoclonal (AC88) and a polyclonal, were used to study the localization of hsp90 in rat endothelial and PtK, epithelial cells. Both

antibodies produced an immunofluorescence pattern: identical to microtubular networks, indicating that hsp90 binds to microtubules in all stages of the cell cycle. In all cells studied, AC88 is found on the microtubules of interphase cells. Moreover, the mitotic apparatus was also labelled by AC88 and by a polyclonal antibody in all stages of mitosis. The hsp90-microtubules association was further supported by the finding that labelling of these cytoplasmic structures was eliminated when anti-hsp90 antibodies were preincubated with purified hsp90.

The colocalization of hsp90 with microtubules was also recently shown in Ishikawa cells (19). Using a polyclonal anti-hsp90 antibody, a filamentous morphology of hsp90 staining and a coloration of the mitotic spindle was clearly observable in these cells (Fig. 2), thus confirming the previously published data (17,18).



**Figure 2.** Immunofluorescence microscopy of Ishikawa cells stained with anti-hsp90 antibodies. Demonstration of cytoplasmic filamentous structures (a) and the mitotic spindle (b).

---

### IMMUNOSTAINING OF hsp90 IN THE PRESENCE OF CYTOSKELETAL INHIBITORS

M Redmond *et al* (18) reported that treatment of rat endothelial cells with the antitubulin agent colcemid caused the disappearance of microtubules, while an identical weak labeling was also observed when cells were stained with anti-hsp90 antibodies, suggesting that most probably hsp90 colocalizes only with microtubules. However, Fostinisei *et al* (19) showed that a clear residual filamentous staining with anti hsp90 antibodies was still present, when the cells were incubated with colchicine or triethyllead, two different and well established antitubulin agents (33,34). Furthermore, under exactly the same experimental conditions a total breakdown of the microtubules was demonstrated, when cells were stained for tubulin. From these results it was concluded that hsp90 most probably associates not only with microtubules but also with structures resistant to antitubulin agents.

What could be the interpretation of these contradictory reports?

Redmond *et al* (18) used an overnight cell fixation procedure in formaldehyde, followed by the immunostaining of the proteins. This fixation procedure, however, seems to be very drastic, it has been shown to favour the rearrangement of some proteins or even the falling off in the staining of cytoskeletal structures (32). Moreover, Fostinisei *et al* (19) reported that using the formaldehyde method for longer than 10 minutes to fix Ishikawa cells, an increasing falling off in the immunostaining of microtubules and vimentin intermediate filaments became evident. Thus, the residual hsp90-labelled structures still present in Ishikawa cells treated by antitubulins are most probably visualized because of using quick and mild fixation procedures. In consequence, the use of a mild cell fixation technique seems to be very important to preserve the labelled filaments.

---

### STUDYING THE INTERACTION OF hsp90 WITH INTERMEDIATE FILAMENT PROTEINS

- It seems that hsp90 associates not only with microtubules but most probably also with other

filamentous structures. Since all reported data point out that hsp90 does not colocalize with actin filaments, the interaction of this protein with intermediate filament structures should be possible. Owing to the failure of specific inhibitors for intermediate filament proteins, this hypothesis is difficult to be tested. Actin filament and microtubule inhibitors, inducing structural rearrangements in the organization of this cytoskeletal protein family, can be used (35,36).

This was recently attempted by Fostinisei *et al* (19), using triethyllead, colchicine or cytochalasin B, and mixtures of these inhibitors to study the interaction of hsp90 with vimentin and cytokeratin intermediate filaments. The concentration of the cytoskeletal inhibitors used were in all cases, higher of those causing the full breakdown of microtubules and actin filaments. This was essential to exclude any remaining hsp90 labelling with microtubules still intact.

---

### hsp90 SEEMS TO NOT ASSOCIATE WITH VIMENTIN INTERMEDIATE FILAMENTS

---

- When cells treated by triethyllead were stained for vimentin, a clear rearrangement of the filaments became evident, presenting all morphological characteristics described for this interaction (35). However, under the same experimental conditions the immunostaining of hsp90 was noticeably different. These data indicate that hsp90 does not associate with vimentin intermediate filaments. Double staining immunofluorescence experiments exposing both staining patterns on the same film plaque further supported this hypothesis (19). This technique uses second antibodies conjugated with different chromophores and is now widely used for the study of the cellular colocalization of proteins with structural elements.

When cells were double stained with FITC-staining for vimentin (green) and with Texas red staining for hsp90 (red), followed by exposure of both staining patterns on the same film plaque, separate colored green and red filaments were shown (19). This was a clear indication that hsp90 does not colocalize with vimentin. This finding was further supported by parallel experiments with cells treated with

triethyllead. A reorganization of vimentin filaments (green) became evident, without affecting the red hsp90 staining (19).


#### **hsp90 CO LOCALIZATION WITH CYTOKERATINS**

- Using the above procedures of single and double staining immunofluorescence in Ishikawa cells, it was demonstrated that hsp90 associates with cytokeratin intermediate filaments (19).

This became evident in cells treated with a mixture of cytochalasin B and colchicine. Staining of the cells with anti-vimentin and anti-hsp90 antibodies offered figures with similarly reorganized filaments, showing an open lattice form including filament focal centers. This reorganization characteristics have been described for keratin cytoskeleton after combined treatment with these anticytoskeletal agents (36).

Double staining immunofluorescence microscopy in Triton-in soluble cytoskeletal preparations further corroborated this observation. By exposing both staining patterns on the same picture exclusively yellow filament structures were observed, when the cells were double stained for cytokeratins with FITC-staining (green) and for hsp90 with Texas-red staining (red). This is shown in Fig. 3a, where the absence of any selectively green or red fluorescence is noticed.

Even more convenient, were the immunofluorescence staining patterns of the Triton-insoluble preparations of cells treated with the cytochalasin B/colchicine mixture. Double stained and double exposed cells showed exclusively yellow colored filamentous structures which clearly demonstrated the well known reorganization characteristics of cytokeratin filaments (Fig. 3b).



*Figure 3. Double staining immunofluorescence microscopy exposing both staining patterns on the same film plaque, (a) Triton-Insoluble cytoskeleton of Ishikawa cells double stained for cytokeratins (green) and hsp90 (red). (b) Same as (a) but after confined treatment with cytochalasin B and colchicine, showing the reorganization characteristics described for keratin cytoskeleton. Note the exclusively yellow colored filamentous structures and absence of any separate red or green fluorescence. (Experimental details in Kef 19).*

**SPECIFICITY OF THE ANTI-hsp90 AND ANTICYTOSKELETAL PROTEIN ANTIBODIES**

• Since all data summarized in this review deal with immunoreactions of different anti-hsp90 and anti-cytoskeletal protein antibodies it was very important to exclude any kind of cross reactions with

investigated cytoskeletal proteins and cytosolic preparations, respectively. For this purpose, Western blotting analyses of cytosolic preparations were performed, using anti-hsp90, anti-tubulin, anti-vimentin and anti-cytokeratin antibodies. No cross reaction of anti-hsp90 antibodies with purified cytoskeletal proteins were observed. Moreover, all




Figure 4. Western blotting analysis of cytosolic preparations of Ishikawa cells using the following antibodies: (a) anti-hsp90, (b) anti-tubulin, (c) anti-vimentin, (d) anti-cytokeratin (8. S3). No cross reaction of anti-hsp90 antibodies could be visualized with cytosolic preparations of tubulin, vimentin, or cyokeratins. (e) Coumassie blue staining of (1) cytosolic preparation of Ishikawa cells, (2) purified tubulin and (3) purified vimentin.

anti-cytoskeletal antibodies did not cross-react with proteins in the range of 90 kD of the cytosolic preparations (Fig. 4). Thus the observed association of hsp90 with microtubules and cytokeratins is not due to unspecific reaction of the anti-hsp90 antibody with these cytoskeletal structures. Similar conclusions concerning the specificity of the anti-hsp90 antibodies were made also by Redmond et al (18).

#### BIOLOGICAL SIGNIFICANCE OF THE INTERACTION OF hsp90 WITH CYTOSKELETAL STRUCTURES

Hsp90 is an ubiquitous, abundant and highly conserved protein performing essential cellular functions. Its involvement in transport processes in the cell has been speculated in the past (8). In the case of newly synthesized pp60<sup>src</sup> and some other avian transforming virus protein kinases, it was postulated that hsp90 is implicated in their translocation from the site of synthesis to the plasma membrane (8,37,38). Moreover, in an earlier review dealing with the physiological role of heat shock proteins, the implication of hsp90 in the transport of steroid receptors to the nucleus has been also discussed (7).

This hypothesis was further encouraged by the findings of the hsp90 colocalization with microtubules (17-19) and cytokeratins (19) in intact cells, as well as by the association of cytosolic glucocorticoid receptor with cytoplasmic microtubules (39, 40). Recent lines of evidence suggest that cytoskeletal structures may also be involved in the intracellular protein transport (41). Thus, the colocalization of hsp90 with microtubules and cytokeratins and its association with the steroid hormone receptors raises the possibility that cytoskeletal structures could be used as "rails" for the intracellular transport of steroid hormone receptors to the nucleus. This assumption is promoted by the colocalization of hsp90 with cytokeratins, since microtubules do not have direct links with the nucleus, while the intermediate filament protein lamin creates the nuclear lamina (42). Thus, we think that the colocalization of hsp90 with both cytoskeletal structures, the microtubules and the cytokeratin intermediate filaments, offer a well organized intracellular "highway" for the efficient movement of the steroid hormone receptor via association-

dissociation with hsp90 molecules, from the cytoplasmic site of synthesis to the nuclear site of action. Once the "highway" is ready, the motors (43) are needed to ensure the transport on it. That may also be tested in the case of hsp90-steroid hormone receptor translocation into the cell nucleus.

#### ACKNOWLEDGMENTS

This work was supported by grants from the Greek Secretariat of Research and Technology (TET, ENEA 91) and the University of Crete Research Committee to Christos Stournaras.

#### REFERENCES

1. Lazarides E, Weber K. Actin antibody: The specific visualization of actin filaments in non-muscle cells. *Proc Natl Acad Sci USA* 1974; 71:2268-2272
2. Weber K, Pollack R, Bibring T. Antibody against tubulin: The specific visualization of cytoplasmic microtubules in tissue cultured cells. *Proc Natl Acad Sci USA* 1975; 72:459-463
3. Weber K, Bibring T, Osborn M. Specific visualization of tubulin containing structures in tissue-culture cells by immunofluorescence. Cytoplasmic microtubules, vinblastine-induced paracrystals, and mitotic figures. *Exp Cell Res* 1975; 95: 111-120
4. Brinkley BR, Hiller GM, Highfield DP. Cytoplasmic microtubules in normal and transformed cells in cultures: analysis by tubulin antibody and immunofluorescence. *Proc Natl Acad Sci USA* 1975; 72: 4981-4985
5. Osborn M, Franke WW, Weber K. Visualization of a system of filaments 7-10 nm thick in cultured cells of an epithelial line (PtK2) by immunofluorescence microscopy. *Proc Natl Acad Sci USA* 1977; 74: 2490-2494
6. Hynes RO, Deslree AT, 10 nm filaments in normal



- and transformed cell. *Cell* 1978; 13: 151-163
7. Schlesinger MJ. Heat shock proteins: the search for functions. *J Biol Chem* 1986; 103: 321-325
  8. Brugge JS, Erikson E, Erikson RL. The specific interaction of the Rous sarcoma virus transforming protein with two cellular proteins. *Cell* 1981; 25: 363-372
  9. Opperman H, Levinson W, Bishop JM. A cellular protein that associates with the transforming protein of Rous sarcoma virus is also a heat shock protein. *Proc Natl Acad Sci USA* 1981; 78: 1067-1071
  10. Lipsich LA, Cutt JR, Brugge JS. Association of the transforming proteins of Rous, Fujinami and Y73 avian sarcoma viruses with the same two cellular proteins. *Mol Cell Biol* 1982; 2: 875-880
  11. Catelli MG, Binart N, Jung-Testas I, Renoir JM, Baulieu EE, Feramisco JR, Welch WJ. The common 90-kd protein component of non-transformed steroid receptors is a heat shock protein. *EMBO J* 1985; 4: 3131-3135
  12. Sanchez ER, Toft DO, Schlesinger MJ, Pratt WB. Evidence that the 90-kDa phosphoprotein associated with the untransformed L-cell glucocorticoid receptor is a murine heat shock protein. *J Biol Chem* 1985; 260: 12398-12401
  13. Schuh S, Yonemoto W, Brugge J, Bauer VJ, Riehl RM, Sullivan WP, Toft DO. A 90000-dalton binding protein common to both steroid receptors and the Rous sarcoma virus transforming protein. *J Biol Chem* 1985; 260: 14292-14296
  14. Baulieu EE, Binart N, Cadepond F, Catelli MG, Chambrat B, Gamier JM, Groyer-Schweizer G, Oblin ME, Radanyi C, Redeuilh G, Renoir JM, Sabbah M. In: Gustafsson JA, Eriksson H, Carlstedt-Duke J, editors. *The Steroid/Thyroid Hormone Receptor Family and Gene Regulation*. Birkhauser Verlag, Basel 1989; 301-318
  15. Koyasu S, Nishida E, Kadowaki T, Matsuzawa M, Iida K, Harada F, Kasuga M, Sakai H, Yahara I. The mammalian heat shock proteins hsp90 and hsp 100 are actin binding proteins. *Proc Natl Acad Sci USA* 1986; 83: 8054-8058
  16. Nishida E, Koyasu S, Sakai H, Yahara I. Calmodulin regulated binding of the 90-kDa heat shock protein to actin filaments. *J Biol Chem* 1986; 261: 16033-16036
  17. Sanchez ER, Redmond T, Scherrer LC, Bresnick EH, Welsh MJ, Pratt WB. Evidence that the 90 kilodalton heat shock protein is associated with tubulin-containing complexes in L cell cytosol and in intact PtK cells. *Mol Endocrinol* 1988; 2: 756-760
  18. Redmond T, Sanchez ER, Bresnick EH, Schlesinger MJ, Toft DO, Pratt WB, Welsh MJ. Immunofluorescence localization of the 90-kDa heat shock protein and microtubules in interphase and mitotic mammalian cells. *Eur J Cell Biol* 1989; 50: 66-75
  19. Fostinis Y, Theodoropoulos PA, Gravanis A, Stourmaras C. Heat shock protein HSP90 and its association with the cytoskeleton: a morphological study. *Biochem Cell Biol* 1992; 70: 779-786
  20. Ashburner M, Bonner JJ. Induction of gene activity in *Drosophila* by heat shock. *Cell* 1979; 17: 241-254
  21. Schlesinger MJ, Ashburner M, Tissieres A, editors. *Heat Shock from Bacteria to Man*. Cold Spring Harbor Laboratory, Cold Spring Harbor New York 1982
  22. Bond U, Schlesinger MJ. Ubiquitin is a heat shock protein in chicken embryo fibroblasts. *Mol Cell Biol* 1985; 5: 949-956
  23. Iida H, Yahara I. Yeast heat shock protein of Mr 48000 is an isoprotein of enolase. *Nature* 1985; 315: 688-690
  24. Ungewickell E. The 70 kD mammalian heat shock proteins are structurally and functionally

- related to the uncoating protein that releases clathrin triskelia from coated vesicles. *EMBO J* 1985; 4: 3385-3391
25. Collier NC, Schlesinger MJ. The dynamic state of heat shock proteins in chicken embryo fibroblasts. *J Cell Biol* 1986; 103: 1495-1507
  26. Lai BT, Chin NW, Stanek AE, Keh W, Lanks KW. Quantitation and intracellular localization of the 85 kD heat shock protein by using monoclonal and polyclonal antibodies. *Mol Cell Biol* 1984; 4: 2802-2810
  27. Carbajal ME, Duband JL, Lettre F, Valet JP, Tanguay RM. Cellular localization of *Drosophila* 83-kilodalton heat shock protein in normal, heat-shocked and recovering cultured cell with a specific antibody. *Biochem Cell Biol* 1986; 64: 816-825
  28. Kwang IK, Devin J, Cadepond F, Jibard N, Guichon-Mantel A, Baulien EE, Catelli MG. In vivo functional protein-protein interaction: nuclear targeted HSP90 shifts cytoplasmic steroid receptor mutants into the nucleus. *Proc Natl Acad Sci USA* 1994; In press
  29. Welch WJ, Feramisco JR. Purification of the major mammalian heat shock proteins. *J Biol Chem* 1982; 257: 14949-14959
  30. Perdew GH, Whitlaw ML. Evidence that the 90-kDa heat shock protein (HSP90) exists in cytosol in heteromeric complexes containing HSP70 and three other proteins with MW of 63000, 56000 and 50000. *J Biol Chem* 1991; 266:6708-6713
  31. Van Bergen en Henegouwen PMP, Berbers G, Linnemans WAM, van Wijk R. Subcellular localization of the 84000 dalton heat shock protein in mouse neuroblastoma cells: evidence for a cytoplasmic and nuclear location. *Eur J Cell Biol* 1987; 43: 469-478
  32. Leong AS, Gilham PN. The effect of progressive formaldehyde fixation on the preservation of tissue antigens. *Pathol* 1989; 21: 266-268
  33. Faulstich H, Stournaras C. Potentially toxic concentrations of triethyllead in Black Forest rainwater samples. *Nature* 1985; 317: 714-715
  34. Zimmermann H-P, Faulstich H, Hansch GM, Doendes KH, Stournaras C. The interaction of triethyllead with tubulin and microtubules. *Mutat Res* 1988; 201: 293-302
  35. Zimmermann H-P, Plagens U, Vorgias CE, Traub P. Changes in the organization of non-epithelial intermediate filaments induced by triethyllead chloride. *Exp Cell Res* 1986; 167: 360-368
  36. Knapp LW, O'Guin WM, Sawyer RH. Rearrangement of the keratin cytoskeleton after combined treatment with microtubule and microfilament inhibitors. *J Cell Biol* 1983; 97:1788-1794
  37. Ben-Ze'ev A. The cytoskeleton in cancer cells. *Biochem Biophys Acta* 1985; 780: 197-212
  38. Brugge JS. Interaction of Rous sarcoma virus protein ppeO<sup>TM</sup> with cellular proteins pp50 and pp90. *Curr Top Microbiol Immunol* 1986; 123:1-22
  39. Akner G, Sundqvist K-G, Denis M, Wikstrom A-C, Gustafsson J-A. Immunocytochemical localization of glucocorticoid receptor in human gingival fibroblasts and evidence for a colocalization of glucocorticoid receptor with cytoplasmic microtubules. *Eur J Cell Biol* 1990; 53: 390-401
  40. Akner G, Mossberg K, Wikstrom A-C, Sundqvist K-G, Gustafsson J-A. Evidence for colocalization of glucocorticoid receptor with cytoplasmic microtubules in human gingival fibroblasts, using two different monoclonal anti-GR antibodies, confocal laser scanning microscopy and image analysis. *J Steroid Biochem* 1991; 39: 419-432
  41. Amos LA, Amos WM. Cytoplasmic transport in axons. *J Cell Sci* 1987; 87: 1-2

42. Georgatos SD, Blobel G. Lamin B constitutes an intermediate filament attachment site at the nuclear envelope. *J Cell Biol* 1987; 105: 117-125
43. Vale RD. Microtubule motors: many new models off the assembly line. *Trends Biochem Sci* 1992; 17: 300-304

Received for publication 29 August 1994

*Address for correspondence:*  
*Dr Christos Stournaras*  
*Department of Biochemistry*  
*School of Medicine*  
*University of Crete*  
*GR-71110 Heraklion*  
*GREECE*

*Tel: 003081 547070*  
*Fax: 003081542112*