

ADJUNCTIVE ANTIBIOTIC THERAPY IN CASES OF PERIODONTITIS AND PERI-IMPLANTITIS

Elitsa Sabeva

*Department of Periodontology and Dental Implantology, Faculty of Dental Medicine,
Medical University of Varna*

ABSTRACT

INTRODUCTION: Periodontitis and peri-implantitis are plaque-induced diseases, which are associated with certain bacteria. The conventional treatment aims primarily at tooth or implant surface debridement. In certain cases, surgical therapy is necessary. To improve the outcome of the treatment, systemic or topical antibiotics could be considered.

AIM: The aim of this review is to investigate the effect of the adjunctive systemically or topically delivered antibiotics on the outcome of the periodontitis and peri-implantitis treatment.

MATERIALS AND METHODS: Articles related to the topic were searched in the PubMed database. Articles published from 2000 to 2019, only in English language, were included in the review. The search was conducted with a variety of keywords in different combinations being used. These were: “periodontitis”, “peri-implantitis”, “antibiotic”, “local”, “metronidazole”, “amoxicillin”, “azithromycin”, “tetracycline”.

RESULTS: Forty-seven articles were included in the review. The selected articles described the adjunctive systemic and/or local delivery of metronidazole, azithromycin, the combination of metronidazole and amoxicillin and some representatives of the tetracycline group, such as doxycycline and minocycline in cases of periodontitis and peri-implantitis.

CONCLUSION: The adjunctive systemic and topical antibiotic therapy could benefit the outcome of periodontitis and peri-implantitis treatment.

Keywords: *antibiotic, peri-implantitis, periodontitis, treatment*

Address for correspondence:

Elitsa Sabeva
Faculty of Dental Medicine
Medical University of Varna
84 Tzar Osvoboditel Blvd
9002 Varna
e-mail: Elitsa.Sabeva@mu-varna.bg

Received: November 18, 2019

Accepted: December 7, 2019

INTRODUCTION

Periodontitis and peri-implantitis are plaque-induced diseases, which are associated with certain bacteria. The conventional treatment aims primarily at tooth or implant surface debridement. In certain cases, surgical therapy is necessary. To improve the outcome of the treatment, systemic or topical antibiotics could be considered.

AIM

The aim of this review is to investigate the effect of the adjunctive systemically or topically delivered antibiotics on the outcome of the periodontitis and peri-implantitis treatment.

MATERIALS AND METHODS

Articles related to the topic were searched in the PubMed database. Articles published from 2000 to 2019, only in English language, were included in the review. The search was conducted with a variety of keywords in different combinations being used. These were: “periodontitis”, “peri-implantitis”, “antibiotic”, “local”, “metronidazole”, “amoxicillin”, “azithromycin”, “tetracycline”.

Metronidazole

Systemic metronidazole, in addition to the non-surgical therapy of peri-implantitis, is an effective treatment in terms of radiographic parameters and probing depth (PD) (1). According to a study, bacterial genes encoding for tetracycline resistance are typical for healthy sulci around implants and for cases of peri-implantitis. To a lesser extent, resistance to beta-lactams is observed. The authors did not detect any resistance to metronidazole and support its use (2).

Lan et al. (3) introduced personalized poly-ε-caprolactone/alginate rings loaded with metronidazole, whose purpose was to ensure controlled release of the agent. The authors concluded that this system has the potential to reduce the bacterial biofilm development and thereby could be used for peri-implantitis prevention.

Tang et al. (4) compared 25% metronidazole gel and ultrasonic scaling with carbon fiber tip and concluded that both treatment modalities are reliable for peri-implantitis therapy.

The local application of metronidazole could lead to a destabilization of the supragingival biofilm (5). The combination of topically delivered metronidazole and zinc hyaluronate has a comparable effect to that of amoxicillin on the oral pathogens (6). The topical delivery of 5% metronidazole along with the debridement could also have a positive effect on the periodontitis therapy (7).

Systemic administration of metronidazole combined with full-mouth disinfection is an efficient

strategy in cases where *P. gingivalis* and *T. forsythia* are detected (8).

Amoxicillin + Metronidazole

The systemic administration of metronidazole and amoxicillin together is used in periodontitis (9,10,11,12,13) and in peri-implantitis treatment (14).

In vitro evaluation of the antibiotic resistance of the bacterial pathogens, typical for the submucosal microbiota in peri-implantitis patients demonstrated that these bacteria are frequently resistant to doxycycline, clindamycin, metronidazole and amoxicillin. The authors also found that the bacteria are only infrequently resistant to both metronidazole and amoxicillin (15).

The combination of metronidazole and amoxicillin, in addition to scaling and root planing (SRP), has led to improvement of the results of the periodontal treatment of patients with type 2 diabetes mellitus (16). El-Makaky and Shalaby (17) also studied the effect of periodontal therapy on the clinical parameters and metabolic response in patient with uncontrolled type 2 diabetes suffering from periodontitis and concluded that the non-surgical therapy of periodontal disease combined with metronidazole and amoxicillin resulted in better metabolic outcome and improved clinical parameters in regard to the periodontitis. The authors observed bleeding on probing (BOP), pocket depth, plaque, clinical attachment level (CAL), and glycated hemoglobin. Other authors also came to a similar conclusion: the adjunctive amoxicillin and metronidazole administration increased the positive effects of the periodontal therapy, considering clinical parameters, such as PD, CAL and BOP (18).

McGowan et al. (19) recommended a 7-day course of metronidazole and amoxicillin. A study supported the regimen of 500 mg metronidazole and 500 mg amoxicillin, three times a day for 7 days in combination with SRP. The authors compared a 3-day and a 7-day course and reported better results with the longer antibiotic course, considering improved clinical parameters, decreased detection of some periopathogens and oxidative stress (20).

As an alternative of the combination of metronidazole and amoxicillin, Araujo et al. (21) proposed clarithromycin, although the authors suggested that this antibiotic was not superior to the aforementioned

combination. Both antibiotic therapies led to reduced bacterial count of *A. actinomycetemcomitans* and *P. gingivalis*. The duration of both treatments was 7 days with the dose of metronidazole and amoxicillin being 400 mg three times a day and 500 mg three times a day, respectively. The dosage of clarithromycin was 500 mg twice a day.

The systemic use of amoxicillin and metronidazole in addition to surgical protocol for peri-implantitis treatment including also surgical access debridement and decontamination of implant surface resulted in a positive effect, which was maintained for 12 months (14).

During a peri-implantitis treatment, Stein et al. (22) administered amoxicillin and metronidazole for a period of 7 days to the patients with severe periodontitis. Other authors did not observe any improvement of the results considering the clinical and microbiological parameters, adding metronidazole and amoxicillin to the non-surgical peri-implantitis treatment (23).

Tetracyclines

Local antimicrobial agents may improve the outcomes of periodontal surgical treatment and may reduce the peri-implantitis manifestations (24). Treatment of peri-implantitis by topical application of tetracyclines led to improvement of the microbiological and clinical parameters (25).

Locally delivered tetracyclines are available in different forms, such as ointment (26), fibers (25, 27, 28), and nanoparticles (29). Topical application of tetracycline is associated with significant positive effect on the clinical parameters, such as PD, sulcular bleeding index, and CAL (30). According to a study, the use of systemic and local antibiotics could decrease the inflammation at the surgical site after implant placement (31). Minocycline hydrochloride ointment and erythromycin eye ointment were applied locally at the implant surgery site. The effect on the healing was assessed and certain parameters were measured. The authors concluded that the effect of both ointments was comparable in regard to the early stage of the healing process (31). Local minocycline demonstrated considerable anti-inflammatory effects (29). The combination of tetracycline fibers and SRP is a reliable treatment, resulting in an improvement of the periodontal health (32). Tetracy-

cline fibers combined with SRP are superior to SRP alone considering the clinical parameters, such as pocket depth and CAL (27).

Topical application of minocycline in combination with surgical treatment of peri-implantitis improves the outcome of the therapy regarding the clinical and radiological results (33). Radiographic and clinical parameters are also improved using nonsurgical treatment of peri-implantitis combined with irrigation with chlorhexidine and local application of minocycline hydrochloride (34).

Topical application of controlled-release doxycycline was included in an anti-infective non-surgical peri-implantitis treatment protocol, which demonstrated improvement of peri-implant health, considering the clinical parameters (35). A new gel was proposed, which included metronidazole and doxycycline in its formulation. The gel was efficient in vitro against planktonic species and against bacterial biofilm. The effect continued 13 days. The authors concluded that its use in cases of peri-implantitis could be discussed (36). Nano-doxycycline gel in addition to SRP also demonstrated anti-inflammatory effect (37).

According to a study, systemic doxycycline treatment did not influence *A. actinomycetemcomitans* infection associated with peri-implantitis. The authors also concluded that periodontitis of the remaining teeth could contribute to the development of peri-implantitis (38). Residues of a minocycline hydrochloride agent could be found on the implant surface. Because it was hydrophobic it could not be removed well using water.

Residual local agents on the implant surface in cases of peri-implantitis might compromise the reosseointegration and lead to bacterial colonization (39). In a study in dogs, it was reported that the application of topical tetracycline hydrochloride solution was an effective method for decontamination of implant surfaces, which were infected (40).

Azithromycin

Systemic delivery of azithromycin in combination with mechanical debridement could contribute to the management of peri-implant mucositis in the therapy of peri-implantitis (41). Some authors concluded that the systemic administration of azithromycin in addition to the surgical peri-implantitis

treatment was not superior to the surgical therapy alone (42). According to Tada et al. (43) the reduction of the bacterial number, caused by the systemic azithromycin, was followed by new increase thereafter. Gomi et al. (44) concluded that the increase of the periodontal bacteria was observed 6 months after the therapy, which included combination of azithromycin and full-mouth SRP. According to O'Rourke (45), systemically delivered azithromycin could contribute to the positive effect of the non-surgical periodontal treatment, especially when the pocket depth was ≥ 6 mm. Yashima et al. (46) reported comparable results from full-mouth SRP, with and without systemic administration of azithromycin, considering the clinical parameters. Renatus et al. (47) supported the opinion about the positive effect of the systemic delivery of azithromycin during periodontitis treatment.

RESULTS

Forty-seven articles were included in the review. The selected articles described the adjunctive systemic and/or local delivery of metronidazole, azithromycin, the combination of metronidazole and amoxicillin, and some representatives of the tetracycline group, such as doxycycline and minocycline in cases of periodontitis and peri-implantitis.

DISCUSSION

The majority of the studies supported the topical application of tetracyclines in adjunct to the periodontitis and peri-implantitis treatment (24,25,27,29,30,32-35,37). One study reported debatable results because of the difficulty of removing completely one of the minocycline hydrochloride agents of the implant surface (39).

Many authors observed positive effect of the systemic administration of the metronidazole and amoxicillin combination in adjunct to the periodontal treatment in terms of the outcomes of the therapy (16-18,20).

Topical metronidazole was found efficient in cases of periodontitis and peri-implantitis (4-7).

The reports about the effect of systemically administered azithromycin on the peri-implantitis and periodontitis treatment are debatable. According to some of the authors the adjunctive systemic delivery of azithromycin benefits the periodontitis (45)

and peri-implantitis treatment (41), while according to others, the results of the treatment with and without azithromycin administration were comparable (42,46).

CONCLUSION

The adjunctive systemic and topical antibiotic therapy could benefit the outcome of periodontitis and peri-implantitis treatment.

REFERENCES

1. Liñares A, Pico A, Blanco C, Blanco J. Adjunctive systemic metronidazole to nonsurgical therapy of peri-implantitis with intrabony defects: a retrospective case series study. *Int J Oral Maxillofac Implants.* 2019;34(5):1237-45. doi: 10.11607/jomi.7343.
2. Koukos G, Papadopoulos C, Tsalikis L, Sakellari D, Arsenakis M, Konstantinidis A. Prevalence of antibiotic resistance genes in subjects with successful and failing dental implants. A pilot study. *Open Dent J.* 2015;8:257-63. doi: 10.2174/1874210601408010257.
3. Lan SF, Kehinde T, Zhang X, Khajotia S, Schmidtke DW, Starly B. Controlled release of metronidazole from composite poly- ϵ -caprolactone/alginate (PCL/alginate) rings for dental implants. *Dent Mater.* 2013;29(6):656-65. doi: 10.1016/j.dental.2013.03.014.
4. Tang Z, Cao C, Sha Y, Lin Y, Wang X. Effects of non-surgical treatment modalities on peri-implantitis. *Zhonghua Kou Qiang Yi Xue Za Zhi.* 2002;37(3):173-5.
5. Ré AC, Ferreira MP, Freitas O, Aires CP. Local antibiotic delivery in periodontitis: drug release and its effect on supragingival biofilms. *Biofouling.* 2016;32(9):1061-6. doi: 10.1080/08927014.2016.1230735.
6. Léber A, Budai-Szűcs M, Urbán E, Vályi P, Gácsi A, Berkó S, et al. Combination of zinc hyaluronate and metronidazole in a lipid-based drug delivery system for the treatment of periodontitis. *Pharmaceutics.* 2019;11(3). pii: E142. doi: 10.3390/pharmaceutics11030142.
7. Paul TP, Emmatty R, Pulikkottil JJ, Rahman AA, Kumar SA, George N. Comparative evaluation of sustained release collagen device containing 5% metronidazole (metrogene) along with and without scaling and root planing at regular intervals with treatment of chronic periodontitis: a case control study. *J Int Oral Health.* 2015;7(6):18-22.

8. Preus HR, Gjermo P, Scheie AA, Baelum V. The effect of metronidazole on the presence of *P. gingivalis* and *T. forsythia* at 3 and 12 months after different periodontal treatment strategies evaluated in a randomized, clinical trial. *Acta Odontol Scand.* 2015;73(4):258-66. doi: 10.3109/00016357.2014.920106.
9. Martins ES, César-Neto JB, Albuquerque-Souza E, Rebeis ES, Holzhausen M, Pannuti CM, et al. One-year follow-up of the immune profile in serum and selected sites of generalized and localized aggressive periodontitis. *Cytokine.* 2019;116:27-37. doi: 10.1016/j.cyto.2018.12.019.
10. Rajendran M, Looney S, Singh N, Elashiry M, Meghil MM, El-Awady AR, et al. Systemic antibiotic therapy reduces circulating inflammatory dendritic cells and Treg-Th17 plasticity in periodontitis. *J Immunol.* 2019;202(9):2690-9. doi: 10.4049/jimmunol.1900046.
11. Teughels W, Dhondt R, Dekeyser C, Quirynen M. Treatment of aggressive periodontitis. *Periodontol.* 2000. 2014;65(1):107-33. doi: 10.1111/prd.12020.
12. Ikram S, Hassan N, Baig S, Borges KJJ, Raffat MA, Akram Z. Effect of local probiotic (*Lactobacillus reuteri*) vs systemic antibiotic therapy as an adjunct to non-surgical periodontal treatment in chronic periodontitis. *J Investig Clin Dent.* 2019;10(2):e12393. doi: 10.1111/jicd.12393.
13. Bizzarro S, Van der Velden U, Loos BG. Local disinfection with sodium hypochlorite as adjunct to basic periodontal therapy: a randomized controlled trial. *J Clin Periodontol.* 2016;43(9):778-88. doi: 10.1111/jcpe.12578.
14. Heitz-Mayfield LJA, Salvi GE, Mombelli A, Faddy M, Lang NP. Anti-infective surgical therapy of peri-implantitis. A 12-month prospective clinical study. *Clin Oral Implants Res.* 2012;23(2):205-10. doi: 10.1111/j.1600-0501.2011.02276.x.
15. Rams TE, Degener JE, van Winkelhoff AJ. Antibiotic resistance in human peri-implantitis microbiota. *Clin Oral Implants Res.* 2014;25(1):82-90. doi: 10.1111/clr.12160.
16. Tamashiro NS, Duarte PM, Miranda TS, Maciel SS, Figueiredo LC, Faveri M, et al. Amoxicillin plus metronidazole therapy for patients with periodontitis and type 2 diabetes: a 2-year randomized controlled trial. *J Dent Res.* 2016;95(7):829-36. doi: 10.1177/0022034516639274.
17. El-Makaky Y, K Shalaby H. The effects of non-surgical periodontal therapy on glycemic control in diabetic patients (randomized controlled trial). *Oral Dis.* 2019. doi: 10.1111/odi.13256.
18. Zandbergen D, Slot DE, Niederman R, Van der Weijden FA. The concomitant administration of systemic amoxicillin and metronidazole compared to scaling and root planing alone in treating periodontitis: =a systematic review=. *BMC Oral Health.* 2016;16:27. doi: 10.1186/s12903-015-0123-6.
19. McGowan K, McGowan T, Ivanovski S. Optimal dose and duration of amoxicillin-plus-metronidazole as an adjunct to non-surgical periodontal therapy: A systematic review and meta-analysis of randomized, placebo-controlled trials. *J Clin Periodontol.* 2018;45(1):56-67. doi: 10.1111/jcpe.12830.
20. Boia S, Boariu M, Baderca F, Rusu D, Muntean D, Horhat F, et al. Clinical, microbiological and oxidative stress evaluation of periodontitis patients treated with two regimens of systemic antibiotics, adjunctive to non-surgical therapy: A placebo-controlled randomized clinical trial. *Exp Ther Med.* 2019;18(6):5001-15. doi: 10.3892/etm.2019.7856.
21. Araujo CF, Andere NMRB, Castro Dos Santos NC, Mathias-Santamaria IF, Reis AA, de Oliveira LD, et al. Two different antibiotic protocols as adjuncts to one-stage full-mouth ultrasonic debridement to treat generalized aggressive periodontitis: A pilot randomized controlled clinical trial. *J Periodontol.* 2019;90(12):1431-40. doi: 10.1002/JPER.18-0399.
22. Stein JM, Hammächer C, Michael SS. Combination of ultrasonic decontamination, soft tissue curettage, and submucosal air polishing with povidone-iodine application for non-surgical therapy of peri-implantitis: 12 Month clinical outcomes. *J Periodontol.* 2017. doi: 10.1902/jop.2017.170362.
23. Shibli JA, Ferrari DS, Siroma RS, Figueiredo LC, Faveri M, Feres M. Microbiological and clinical effects of adjunctive systemic metronidazole and amoxicillin in the non-surgical treatment of peri-implantitis: 1 year follow-up. *Braz Oral Res.* 2019 Sep 30;33(suppl 1):e080. doi:10.1590/1807-3107bor-2019.vol33.0080.
24. Paquette DW, Ryan ME, Wilder RS. Locally delivered antimicrobials: clinical evidence and relevance. *J Dent Hyg.* 2008;82 Suppl 3:10-5.

25. Mombelli A, Feloutzis A, Brägger U, Lang NP. Treatment of peri-implantitis by local delivery of tetracycline. Clinical, microbiological and radiological results. *Clin Oral Implants Res.* 2001;12(4):287-94. doi: 10.1034/j.1600-0501.2001.012004287.x.
26. Da Rocha HA, Silva CF, Santiago FL, Martins LG, Dias PC, De Magalhães D. Local drug delivery systems in the treatment of periodontitis: a literature review. *J Int Acad Periodontol.* 2015;17(3):82-90.
27. Sinha S, Kumar S, Dagli N, Dagli RJ. Effect of tetracycline HCl in the treatment of chronic periodontitis - A clinical study. *J Int Soc Prev Community Dent.* 2014;4(3):149-53. doi: 10.4103/2231-0762.142011.
28. Boyeena L, Koduganti RR, Panthula VR, Jammula SP. Comparison of efficacy of probiotics versus tetracycline fibers as adjuvants to scaling and root planing. *J Indian Soc Periodontol.* 2019;23(6):539-544. doi: 10.4103/jisp.jisp_590_18.
29. Martin V, Ribeiro IAC, Alves MM, Gonçalves L, Almeida AJ, Grenho L, et al. Understanding intracellular trafficking and anti-inflammatory effects of minocycline chitosan-nanoparticles in human gingival fibroblasts for periodontal disease treatment. *Int J Pharm.* 2019;572:118821. doi: 10.1016/j.ijpharm.2019.118821.
30. Nadig PS, Shah MA. Tetracycline as local drug delivery in treatment of chronic periodontitis: A systematic review and meta-analysis. *J Indian Soc Periodontol.* 2016;20(6):576-83. doi: 10.4103/jisp.jisp_97_17.
31. Xu L, Wang Y, Nguyen VT, Chen J. Effects of topical antibiotic prophylaxis on wound healing after flapless implant surgery: a pilot study. *J Periodontol.* 2016;87(3):275-80. doi: 10.1902/jop.2015.150464.
32. Dang AB, Chaubey KK, Thakur RK, Mohan R, Chowdhary Z, Tripathi R. Comparative evaluation of efficacy of three treatment modalities - tetracycline fibers, scaling and root planing, and combination therapy: A clinical study. *J Indian Soc Periodontol.* 2016;20(6):608-13. doi: 10.4103/jisp.jisp_52_17.
33. Cha JK, Lee JS, Kim CS. Surgical therapy of peri-implantitis with local minocycline: a 6-month randomized controlled clinical trial. *J Dent Res.* 2019;98(3):288-95.
34. Heo S, Kim HJ, Joo JY, Lee J, Kim SJ, Choi J. Simplified nonsurgical treatment of peri-implantitis using chlorhexidine and minocycline hydrochloride. *J Periodontal Implant Sci.* 2018;48(5):326-33. doi: 10.5051/jpis.2018.48.5.326.
35. Mensi M, Scotti E, Calza S, Pilloni A, Grusovin MG, Mongardini C. A new multiple anti-infective non-surgical therapy in the treatment of peri-implantitis: a case series. *Minerva Stomatol.* 2017;66(6):255-66. doi: 10.23736/S0026-4970.17.04054-7.
36. Nastri L, De Rosa A, De Gregorio V, Grassia V, Donnarumma G. A new controlled-release material containing metronidazole and doxycycline for the treatment of periodontal and peri-implant diseases: formulation and in vitro testing. *Int J Dent.* 2019;2019:9374607. doi: 10.1155/2019/9374607.
37. Madi M, Pavlic V, Samy W, Alagl A. The anti-inflammatory effect of locally delivered nano-doxycycline gel in therapy of chronic periodontitis. *Acta Odontol Scand.* 2018;76(1):71-6. doi: 10.1080/00016357.2017.1385096.
38. van Winkelhoff AJ, Wolf JW. Actinobacillus actinomycetemcomitans-associated peri-implantitis in an edentulous patient. A case report. *J Clin Periodontol.* 2000;27(7):531-5. doi: 10.1034/j.1600-051x.2000.027007531.x.
39. Lee JB, Kweon HH, Cho HJ, Kim CS, Kim YT. Characteristics of local delivery agents for treating peri-implantitis on dental implant surfaces: a preclinical study. *J Oral Implantol.* 2019;45(2):116-26. doi: 10.1563/aaid-joi-D-17-00261.
40. Ramos UD, Suaid F, Wikesjö UME, Susin C, Vital PC, de Souza SLS, et al. Microbiologic effect of two topical anti-infective treatments on ligature-induced peri-implantitis: A pilot study in dogs. *J Periodontol.* 2018;89(8):995-1002. doi: 10.1002/JPER.17-0630.
41. Gershenfeld L, Kalos A, Whittle T, Yeung S. Randomized clinical trial of the effects of azithromycin use in the treatment of peri-implantitis. *Aust Dent J.* 2018. doi: 10.1111/adj.12614.
42. Hallström H, Persson GR, Lindgren S, Renvert S. Open flap debridement of peri-implantitis with or without adjunctive systemic antibiotics: A randomized clinical trial. *J Clin Periodontol.* 2017;44(12):1285-93. doi: 10.1111/jcpe.12805.

43. Tada H, Masaki C, Tsuka S, Mukaibo T, Kondo Y, Hosokawa R. The effects of *Lactobacillus reuteri* probiotics combined with azithromycin on peri-implantitis: A randomized placebo-controlled study. *J Prosthodont Res.* 2018;62(1):89-96. doi: 10.1016/j.jpor.2017.06.006.
44. Gomi K, Matsushima Y, Ujiie Y, Shirakawa S, Nagano T, Kanazashi M, et al. Full-mouth scaling and root planing combined with azithromycin to treat peri-implantitis. *Aust Dent J.* 2015;60(4):503-10. doi: 10.1111/adj.12257.
45. O'Rourke VJ. Azithromycin as an adjunct to non-surgical periodontal therapy: a systematic review. *Aust Dent J.* 2017;62(1):14-22. doi: 10.1111/adj.12448.
46. Yashima A, Morozumi T, Yoshie H, Hokari T, Izumi Y, Akizuki T, et al. Biological responses following one-stage full-mouth scaling and root planing with and without azithromycin: Multicenter randomized trial. *J Periodontol Res.* 2019;54(6):709-19. doi: 10.1111/jre.12680.
47. Renatus A, Herrmann J, Schönfelder A, Schwarzenberger F, Jentsch H. Clinical efficacy of azithromycin as an adjunctive therapy to non-surgical periodontal treatment of periodontitis: a systematic review and meta-analysis. *J Clin Diagn Res.* 2016;10(7):ZE01-7. doi: 10.7860/JCDR/2016/20176.8115.