

---

Reports

---

2021

**Estimation of Juvenile Striped Bass Relative Abundance in the Virginia Portion of Chesapeake Bay Annual Progress Report: 2020-2021**

Jack R. Buchanan  
*Virginia Institute of Marine Science*

Mary C. Fabrizio  
*Virginia Institute of Marine Science*

Troy D. Tuckey  
*Virginia Institute of Marine Science*

Troy D. Tuckey  
*Virginia Institute of Marine Science*

Follow this and additional works at: <https://scholarworks.wm.edu/reports>



Part of the [Aquaculture and Fisheries Commons](#), and the [Marine Biology Commons](#)

---

**Recommended Citation**

Buchanan, J. R., Fabrizio, M. C., Tuckey, T. D., & Tuckey, T. D. (2021) Estimation of Juvenile Striped Bass Relative Abundance in the Virginia Portion of Chesapeake Bay Annual Progress Report: 2020-2021. Virginia Institute of Marine Science, William & Mary. doi: 10.25773/fham-5v16

This Report is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in Reports by an authorized administrator of W&M ScholarWorks. For more information, please contact [scholarworks@wm.edu](mailto:scholarworks@wm.edu).



VIRGINIA INSTITUTE OF MARINE SCIENCE

## **Estimation of Juvenile Striped Bass Relative Abundance in the Virginia Portion of Chesapeake Bay**

**ANNUAL PROGRESS REPORT: 2020 - 2021**

**Jack R. Buchanan  
Mary C. Fabrizio  
Troy D. Tuckey**

Virginia Institute of Marine  
Science  
William & Mary  
Gloucester Point, Virginia



U.S. Fish and Wildlife Service Sport Fish Restoration Project F87R32  
Submitted to Virginia Marine Resources Commission May 2021



doi: 10.25773/fham-5v16



Cover image

Field Assistants Connor Morang and Jordan Wheatley seine for juvenile striped bass at  
on the Rappahannock River. © Jack Buchanan/VIMS.

## TABLE OF CONTENTS

EXECUTIVE SUMMARY .....	iii
PREFACE .....	v
INTRODUCTION .....	1
METHODS .....	2
RESULTS AND DISCUSSION .....	6
CONCLUSION .....	18
ACKNOWLEDGMENTS .....	19
LITERATURE CITED .....	20
TABLES .....	23
FIGURES .....	39
APPENDICES.....	58

## EXECUTIVE SUMMARY

The 2020 Striped Bass juvenile abundance index was 13.89 and was greater than the reference mean of 7.77 from 1980-2009. Abundance indices were above average in the James and Rappahannock rivers and average in the York River in 2020 compared with the river-specific reference means (1980-2009). The 2020 means, however, may not be directly comparable to the reference means because 19% of index hauls in 2020 were not completed due to Tropical Storm Isaias and precautions taken to protect researchers against COVID-19. When the missed index hauls are accounted for in the reference mean calculations, the overall 2020 mean (13.89) is not significantly different from the reference mean (8.68), suggesting an average year of juvenile Striped Bass recruitment in Virginia waters. Relatively low catches of young-of-the-year Striped Bass from sites upriver and downriver of core nursery areas suggest Striped Bass largely remained within core nursery areas in 2020.

Juvenile abundance indices for American Shad, Alewife, Atlantic Silverside, Blueback Herring, Spot, and Spottail Shiner were generally below their historic averages in 2020. In contrast, juvenile abundance indices for Atlantic Croaker, Banded Killifish, and Inland Silverside were generally greater than historic averages in Virginia waters in 2020. In addition, juvenile White Perch abundance indices in 2020 were above historic averages in the James, York and Rappahannock river systems. Together, these results suggest modest production of forage fish prey for piscivores in Virginia.

Sampling at several auxiliary sites was not possible in 2020. On the James River, new restrictions involving access to the Fort Eustis Army Base required finding a permanent replacement site for J22 which had been sampled from the Army Base. Site J26, approximately four miles upriver of site J22, exhibits a similar depth profile and substratum to J22 and was deemed an acceptable permanent replacement. On the Rappahannock River, site R75 was inaccessible due to downed trees. Site R76, approximately 0.25 miles upriver of site R75, was considered an acceptable replacement for the 2020 survey. On the Pamunkey River, site P56 permanently replaced site P55 in

2020 due to the presence of dense stands of *Hydrilla* at site P55 that impeded proper sampling.

Finally, we note the presence of construction at index site J29, located about 0.10 miles upriver from the mouth of College Creek, an area with exceedingly strong currents. Construction efforts are underway to reduce currents, but such modification may alter the hydrology of site J29 once construction is completed. An altered hydrology at J29 may affect striped bass abundance and availability at the site and would compromise our ability to compare catches with those from previous sampling years. During the 2020 survey, researchers assessed multiple locations within 0.25 miles of J29 for consideration as a potential replacement site, but were unsuccessful in identifying a replacement site. Assuming construction continues in 2021, a broader search to find a replacement site for J29 will be conducted.

## **PREFACE**

The primary objective of the Virginia Institute of Marine Science juvenile Striped Bass seine survey is to monitor the relative annual recruitment of juvenile Striped Bass in the principal Virginia nursery areas of Chesapeake Bay. The U.S. Fish and Wildlife Service initially funded the survey from 1967 to 1973 with funds from the Commercial Fisheries Development Act of 1965 (PL88-309). Beginning in 1980, funds were provided by the National Marine Fisheries Service under the Emergency Striped Bass Study program (PL96-118, 16 U.S.C. 767g, the “Chafee Amendment.” Commencing with the 1989 annual survey, the work was jointly supported by Wallop-Breaux funds (Sport Fish Restoration and Enhancement Act of 1988 PL100-488, the “Dingell-Johnson Act”), administered through the U.S. Fish and Wildlife Service, and the Virginia Marine Resources Commission. This report summarizes the results of the 2020 sampling period and compares these results with previous years.

## INTRODUCTION

Striped Bass (*Morone saxatilis*) is one of the most recreationally sought-after fish species on the east coast of the United States. Decreases in the harvest of Striped Bass in the 1970s paralleled the steady decline in abundance of Striped Bass along the east coast; Chesapeake Bay stock abundances were particularly depressed. Declines in harvests mirrored declines in juvenile recruitment (Goodyear 1985). Because the tributaries of Chesapeake Bay were identified as primary spawning and nursery areas, fishery managers enacted regulations intended to halt and reverse the decline of Striped Bass in Chesapeake Bay and elsewhere within its native range (ASMFC 2003).

In 1981, the Atlantic States Marine Fisheries Commission (ASMFC) developed the Atlantic Coast Striped Bass Interstate Fisheries Management Plan (FMP), which included recommendations aimed to improve the stock status. The Virginia Marine Resources Commission (VMRC) adopted the plan in March 1982 (Regulation 450-01-0034). As Striped Bass populations continued to decline, Congress passed the Atlantic Striped Bass Conservation Act (PL 98-613) in 1984, which required states to follow and enforce management measures in the FMP or face a moratorium on Striped Bass harvests. Since 1981 the FMP has been amended six times to address changes in the management of the stocks. Amendment 6 to the plan, adopted in February 2003, requires "producing states" (i.e., Virginia, Maryland, Delaware and New York) to develop and support programs that monitor Striped Bass recruitment. More recently, Addendum VI to Amendment 6 called for an 18% reduction in removals relative to 2017 removals to reduce fishing mortality rates on the stock, because the 2018 benchmark assessment found that the Striped Bass stock was overfished and experiencing overfishing. In 2020, Amendment 7 was drafted to address multiple management issues, including fishery goals and objectives, biological reference points, management triggers, stock rebuilding, regional management, conservation equivalency, recreational release mortality, recreational accountability, and allocation of the coastal commercial quota.



Initially, the Virginia program used a 6 ft. x 100 ft. x 0.25 in. mesh (2 m x 30.5 m x 6.4 mm) bag seine, but comparison hauls with Maryland gear (4 ft. x 100 ft. x 0.25 in. mesh; 1.2m x 30.5m x 6.4mm mesh) showed virtually no statistical differences in catch, and Virginia adopted the "Maryland seine" after 1987 (Colvocoresses 1987). The gear comparison study aimed to standardize methods and promote a bay-wide recruitment estimate (Colvocoresses and Austin 1987). This was never realized due to remaining differences in the methods of estimation of means (MD: arithmetic index; VA: geometric index). A bay-wide index using a geometric mean weighted by spawning area in each river was proposed in 1993 (Austin et al. 1993) but has not been implemented. In 2009, computations of a bay-wide geometric mean juvenile abundance index (JAI) were found to be correlated with abundance estimates of adult fish from fishery-independent monitoring (Woodward 2009).

Primary objectives for the 2020 program were to:

1. estimate the relative abundance of the 2020 year class of Striped Bass in the James, York and Rappahannock river systems,
2. quantify environmental conditions at the time of collection, and
3. examine relationships between juvenile Striped Bass abundance and environmental and biological data.

## **METHODS**

Field sampling was conducted during five biweekly periods (rounds) from 6 July to 9 September 2020. Pilot sampling at two sites in the James River during early-June revealed zero Stiped Bass, which suggested that sampling did not need to be initiated in mid-June, as was done in 2012, 2016, 2017, and 2019 (Buchanan et al. 2020). During each round, seine hauls were conducted at 18 index stations and 21 auxiliary stations in the James, York and Rappahannock river systems (Figure 1). Auxiliary sites were added to the survey in 1989 to provide better geographic coverage and increase sample sizes within each river system. Such monitoring was desirable in light of increases in Striped Bass stock size during the 1980s and hypothesized expansion of the nursery grounds in

years of high juvenile abundance. However, due to Tropical Storm Isaias and precautions taken to protect researchers from COVID-19, 19% of index sites and 8% of auxiliary sites were not sampled in 2020 (Table 1).

Three auxiliary sites were relocated for all sampling rounds in 2020. On the Rappahannock River, site R75 was inaccessible due to downed trees. A suitable replacement site (R76) was found less than 0.25 miles upriver of site R75. On the Pamunkey River, site P56 permanently replaced P55 in 2020 due to the presence of dense stands of *Hydrilla*, which reduce seine net sampling efficiency (Machut and Fabrizio 2010). Site P56 was identified as a suitable replacement for site P55 during the 2019 survey and is located less than 0.25 miles upriver of site P55 (Buchanan et al. 2020). On the James River, increasing difficulties and restrictions involving access to Fort Eustis Army Base required finding a replacement site for J22. Site J26 permanently replaced site J22 in 2020. Site J26 is approximately four miles upriver of site J22 and exhibits a similar depth profile and substratum as site J22.

As in previous years, collections were made by deploying a 100 ft. (30.5 m) long, 4 ft. (1.2 m) deep, and 0.25 in (6.4 mm) mesh minnow seine perpendicular to the shoreline until either the net was fully extended or a depth of approximately 4 ft. (1.2 m) was encountered and then pulling the offshore end down-current and back to the shore. During each round, a single haul was made at each auxiliary station while duplicate hauls, with an interlude of at least 30 minutes, were made at each index station. Every fish collected during a haul was removed from the net and placed into water-filled buckets. All Striped Bass were measured to the nearest mm fork length (FL), and for all other species, a subsample of up to 25 individuals was measured to the nearest mm FL (or total length if appropriate). At index stations, fish collected during the first haul were held in a water-filled bucket until the second haul was completed. All captured fish, except those preserved for life history studies, were returned to the water at the conclusion of sampling. Sampling time, tidal stage, and weather conditions were recorded at each sampling location. Salinity, water temperature, and dissolved

oxygen concentration were measured after the first haul using a YSI water quality sampler.

From 1999 to 2015, the VIMS seine survey used a net comprised of 0.25 inch knotless oval mesh. However, this netting was no longer available from the manufacturer in 2015, so a new net was constructed from 0.25 inch knotless rhomboid mesh material. To test whether the change in mesh material influenced the relative catch efficiency of the net, paired hauls of old and new nets were conducted during the 2015 sampling season, and these data were used to estimate species-specific calibration factors for juvenile Striped Bass and White Perch (Fabrizio et al. 2017). The estimated calibration factor was 0.5175 for Striped Bass and 0.6537 for White Perch, implying that the new net captured more Striped Bass and White Perch than the old net (i.e., catches in the new net were adjusted by multiplying by the calibration factor; Fabrizio et al. 2017). However, due to low sample sizes ( $n < 30$ ), these calibration factors were viewed as preliminary (Gallagher et al. 2017) and additional paired hauls were conducted during the 2017 sampling season. The addition of 2017 data markedly increased sample sizes ( $n > 70$ ), and resulted in calibration factors that were not significantly different from 1 for either species (Appendix Table 1). Therefore, catch data for Striped Bass and White Perch were not adjusted when estimating indices of abundance from catches observed in the new net.

In this report, comparisons of Striped Bass recruitment indices with prior years are made for the “primary nursery” area only (Colvocoresses 1984), using data collected from months and areas sampled during all years (i.e., index stations). Catch data from auxiliary stations are not included in the calculation of the annual indices. The index of relative abundance for young-of-the-year Striped Bass is calculated as the adjusted overall mean catch per seine haul such that

$$\text{Index} = (\exp(\ln[(\text{totnum})+1]) - 1) \times 2.28$$

where *totnum* is the total number of Striped Bass per seine haul; catches from the first and second seine haul at each index station are considered in this calculation. Because the frequency distribution of the catch is skewed (Colvocoresses 1984), a logarithmic

transformation ( $\ln((totnum)+1)$ ) was applied to the data prior to analysis (Sokal and Rohlf 1981). Mean values are back-transformed and scaled arithmetically ( $\times 2.28$ ) to allow comparisons with Maryland indices. Thus, a “scaled” index refers to an index that is directly comparable with the Maryland index.

Even with a 30-minute interlude between hauls at index stations, second hauls cannot be considered independent samples and their use violates a key assumption necessary for making inferences from a sample mean (Rago et al. 1995). Previous reports consistently documented lower catches on average in the second haul (e.g., Hewitt et al. 2007, 2008), a result which artificially lowers the geometric mean when data from both hauls are included in the index computation. In accordance with suggestions made by Rago et al. (1995), the Virginia juvenile Striped Bass index was also recomputed using only the first haul at each index station. Additionally, the rehabilitation of Chesapeake Bay Striped Bass stocks and subsequent relaxation of fisheries regulations in Chesapeake Bay in 1990 (ASMFC 2003) allow examination of the recruitment of Striped Bass during three periods:

- 1967 - 1973: an early period of monitoring;
- 1980 - 1989: a decade reflecting severe population depression during which temporary fishing moratoria were in place; and,
- 1990 - 2012: a period of post-recovery and regulation targeting the development of a sustainable fishery. Note that the estimated spawning stock biomass (SSB) has been below the SSB threshold since 2013 and hence, the stock has been overfished according to the 2018 benchmark stock assessment.

The 2020 annual index calculated from both hauls was compared with the average index from 1980-2009 (hereafter referred to as the reference period) to reflect the fixed time period used in the definition of recruitment failure in Virginia, as stipulated by Addendum II to Amendment 6 of the Striped Bass fishery management plan (ASMFC 2010). In addition, an average index value for 1990-2012 was calculated using only the first haul at each index site to provide a benchmark for interpreting recruitment

strength during the post-recovery period and was compared with the 2020 annual index.

Throughout this report, mean catch rates are compared using 95% confidence intervals. Reference to “significant” differences between geometric means in this context will be restricted to cases of non-overlapping confidence intervals. Because standard errors are calculated from transformed (logarithmic) values, confidence intervals for the back-transformed and scaled indices are non-symmetrical.

## **RESULTS AND DISCUSSION**

### ***Juvenile Index of Abundance for Virginia***

We collected 1,836 young-of-the-year Striped Bass in 2020 from 145 seine hauls at index stations and 345 individuals from 97 hauls at auxiliary stations (Table 1). Using index-station catches from both hauls, the estimated Striped Bass recruitment index in 2020 was 13.89 (lower confidence interval [LCI] = 11.08, upper confidence interval [UCI] = 17.29; Table 2), which is significantly greater than the average of 7.77 during the reference period (LCI = 6.01, UCI = 9.89; Figure 2). Using index station catches from only the first haul in 2020, 1,256 young-of-the-year Striped Bass were collected, resulting in an index of 15.29 (LCI = 11.43, UCI = 20.24, Table 3), which was not significantly different than the first-haul reference period index of 9.57 (LCI = 7.43, UCI = 12.17). The first haul index was also not significantly different from the mean index estimated for the post-recovery period from 1990-2012 (post-recovery index = 11.91; LCI = 9.25, UCI = 15.17).

Prior to 2011, annual recruitment indices were calculated from all collections made during a sampling year including samples taken before July and after mid-September. In particular from 1967 to 1973, seine sampling extended into October and occasionally into December (1973). Current protocols conclude sampling in late-August or mid-September because after this time, sampling efficiency decreases due to increased avoidance of the sampling gear and movement of juveniles into deeper waters. Indices calculated from data that include catches after this period are therefore biased low. Starting in 2011, recruitment calculations were made using catch data from

the currently established sampling season (July through mid-September, or late-June through August) to permit uniform comparisons of annual recruitment (Tables 2-4).

Striped Bass recruitment success in the Virginia portion of Chesapeake Bay is variable among years and among nursery areas within years. Since the termination of the Striped Bass fishing moratorium in 1990, strong year classes have been observed approximately every decade (1993, 2003, and 2011). The highest recruitment index observed by the Virginia seine survey occurred in 2011. Average to above-average recruitment years occurred between 2003 and 2011, and more recently from 2013 to 2020 (Figure 2). Below-average year classes were observed in 1991, 1999, 2002, and 2012 (Figure 2). In the past decade, recruitment has been average or above average in all but one year (2012), indicating production has been relatively consistent in Virginia nurseries during this time. Under current ASMFC regulations (ASMFC 2010), management action is triggered after three consecutive years of low recruitment in producing states (i.e., the index value is below the first quartile in the time series; Figure 1). Such periods of persistently low recruitment have previously occurred in Virginia from 1971-1973 and 1980-1983 (Figure 2).

Continued monitoring of regional recruitment success will be important in identifying management strategies to protect the spawning stock of Chesapeake Bay Striped Bass, particularly now that the stock is experiencing overfishing and that spawning stock biomass is below the threshold. Research suggests that a Chesapeake Bay-wide index, computed from Virginia and Maryland data combined, will provide a better estimate of recruitment strength and serve as a better predictor of subsequent adult Striped Bass abundance within the Bay (Woodward 2009). This may be particularly appropriate in years when indices from Virginia and Maryland provide divergent estimates of year-class strength (such as 2019, when Virginia reported average recruitment and Maryland reported below-average recruitment for Striped Bass); such differences may arise due to annual changes in the relative contribution of nursery areas throughout Chesapeake Bay.

### ***Juvenile Index of Abundance for Individual Watersheds***

Using index-station catches from both hauls, the estimated Striped Bass recruitment indices in the three Virginia watersheds during 2020 showed variation relative to their individual means from the reference period (1980-2009; Table 4; Figure 3). The 2020 JAI for the James River drainage was 20.07 (LCI = 15.01, UCI = 26.60), which was significantly greater than the reference period index of 10.41 (LCI = 7.83, UCI = 13.64; Table 4). The 2020 JAI for the York River drainage was 7.68 (LCI = 5.57, UCI = 10.37), which was not significantly different from the reference period index of 5.85 (LCI = 4.50, UCI = 7.48; Table 4). Similar to the James River, the 2020 JAI for the Rappahannock River drainage was 19.20 (LCI = 11.43, UCI = 31.39), which was significantly greater than the reference period index of 7.90 (LCI = 5.63, UCI = 10.82, Table 4).

The James River drainage includes the James River proper and the Chickahominy River. Although the 2020 index for the James River drainage was significantly greater than the long-term average, examination of the 2020 JAIs for the James River proper and Chickahominy River revealed no differences from the average indices for the reference period. The 2020 JAI for the James River main stem (excluding the Chickahominy River) was 19.12 (LCI = 12.61, UCI = 28.47), which was not significantly different from the reference period index of 9.72 (LCI = 7.06, UCI = 13.12; Table 4). The 2020 JAI for the Chickahominy River was 21.96 (LCI = 15.74, UCI = 30.32), which was not significantly different than the reference period index of 11.95 (LCI = 8.70, UCI = 16.15; Table 4).

The core nursery area within the James River drainage consists of six mid-river stations: four in the James River (J36, J42, J46, J51) and two in the Chickahominy River (C1, C3). Historically, these six stations tended to have relatively high and stable abundance. In 2020, 50% of all young-of-the-year Striped Bass collected from the James River drainage were captured from this core nursery area (Table 1). The remaining Striped Bass were captured at upriver (39%) or downriver sites (11%; Table 1).

No index sites are located along the main stem of the York River; thus, the watershed JAI is estimated from catches at sites located within the two principle York River tributaries, the Mattaponi and Pamunkey rivers. The 2020 Pamunkey River JAI of 7.30 (LCI = 4.59, UCI = 11.08) was not significantly different than the reference period index of 6.90 (LCI = 4.90, UCI = 9.44; Table 4), and the 2020 Mattaponi River index of 7.98 (LCI = 5.03, UCI = 12.12) was also not significantly different from the reference period average of 5.16 (LCI = 4.06, UCI = 6.45; Table 4). There are distinct core nursery areas within the Pamunkey (P45, P50) and Mattaponi rivers (M33, M37, M41, M44), which generally exhibit high and stable catches compared with other sites in these rivers. This pattern held true in 2020, as the majority of Striped Bass were captured within the core nursery area in the Pamunkey (70%) and Mattaponi (85%) rivers. Overall, approximately 32% of Striped Bass in the York River drainage were collected from the Pamunkey River and 58% from the Mattaponi River in 2020; the remainder (10%) were from the York River auxiliary stations (Table 1).

The 2020 JAI for the Rappahannock River drainage was 19.20 (LCI = 11.43, UCI = 31.39), which was significantly greater than the reference period index of 7.90 (LCI = 5.63, UCI = 10.82, Table 4). The core nursery area within the Rappahannock River consists of the three uppermost index sites (R44, R50, R55) that have consistently dominated the catches in this drainage for more than two decades. In 2020, 95% of the total Rappahannock River catch was taken within the core nursery area (Table 1). The remaining Striped Bass were captured at upriver (4%) or downriver sites (2%; Table 1).

### ***Striped Bass Collections from Auxiliary Stations***

Figures 4-6 illustrate the spatial distribution of the 2020 year class of Striped Bass throughout the areas sampled by this survey. Note that the scaling of CPUE is not constant across the figures. The 1989 addition of auxiliary stations provided increased spatial coverage in the James, York and Rappahannock drainages, and the upriver and downriver auxiliary sites allowed delineation of the upper and lower limits of the nursery. These auxiliary stations help reveal spatial changes in the nursery areas that



may occur due to annual changes in river flow and other factors. Additionally, in years of low or high juvenile abundance, the nursery area may contract or expand spatially. We observed relatively low catches of young-of-the-year Striped Bass at upriver and downriver auxiliary sites in 2020, which suggests that fish mostly remained within the core nursery area.

During 2020, juvenile Striped Bass were captured at all auxiliary sites in the James River except site J12. Catches of juvenile Striped Bass in the James River were relatively low at the upper- and lower-most sites, with the exception of site J56, which produced the highest average index of all sites (Tables 1 and 5; Figure 4). Striped Bass were collected from all auxiliary sites in the Pamunkey and Mattaponi rivers in 2020, although only one individual was captured at the uppermost Pamunkey river site, P56 (Tables 1 and 5; Figure 5). In the York River main stem, relatively few Striped Bass were collected from the three auxiliary stations, and for the second consecutive year, no fish were collected at the lowermost York River site (Y15) in 2020 (Table 5).

We previously suggested that the lack of juvenile Striped Bass at auxiliary stations in the upper reaches of the York River watershed may have been due to the inability to accurately sample in the dense *Hydrilla* vegetation that typically occurs at these sites (Machut and Fabrizio 2010). In 2020, we detected few juvenile Striped Bass at the uppermost auxiliary sites in the Pamunkey (P55) and Mattaponi (M52) rivers (Table 1), but not all fish may have been detected in the area due to low capture efficiencies associated with hauling a seine net through dense aquatic vegetation. Catches in recent years at these two sites may have been affected by the altered state of the nearshore area of these sites. For example, the presence of dense *Hydrilla* stands at sites P55 and M52 may be forcing Striped Bass into deeper waters; alternatively, Striped Bass may utilize *Hydrilla* habitats but remain unavailable to the sampling gear. The continued sampling difficulties at these stations suggested a need to examine alternative collection methods within this region to determine the abundance of juvenile Striped Bass in nearshore areas where *Hydrilla* is present.

Relatively low numbers of Striped Bass were collected at upriver Rappahannock River auxiliary stations during 2020. In recent years, few fish have been collected at downriver sites in the Rappahannock River (R12, R21, R28) even though these sites have favorable substrate and no obstructions to compromise seining. A similar pattern was observed in 2020 with zero individuals collected at downriver sites R12 and R21, and seven individuals collected at downriver site R28 (Table 1; Figure 6).

### ***Comparison among Sampling Rounds***

Indices of juvenile abundance calculated by sampling round in 2020 were not significantly different from the averages calculated during the 1980-2009 reference period, with the exception of Round 3 (21.07, LCI = 12.23, UCI = 35.29), which was significantly greater than its historical average (7.26, LCI = 5.44, UCI = 9.50) (Table 6). The largest number of young-of-the-year Striped Bass was collected during rounds 1 and 2 in 2020, with fewer observed in subsequent rounds (Table 6). This follows patterns observed during the reference period, such that 55% of the Striped Bass captured within the primary nursery areas of Virginia were captured during the first two rounds of sampling. In 2020, we captured 28% of all juvenile Striped Bass in round 1; this was followed by a slight decline (-0.2%) in the number of Striped Bass captured in round 2, a pattern that was similar to average declines observed between rounds 1 and 2 during the reference period (-22%). There were also modest declines in 2020 catches during the third (-40%), fourth (-8%), and fifth (-31%) rounds, which were broadly similar to reference period averages.

### ***Environmental Conditions and Potential Relationships to Striped Bass Abundance***

The juvenile Striped Bass seine survey routinely records temperature, salinity and dissolved oxygen at each site during each round of sampling (see Methods). Environmental conditions during each round in 2020 were compared graphically with long-term average conditions to assess changes in habitat condition for juvenile Striped Bass (Figures 7-9). For temperature and salinity, the long-term average was calculated

using observations from 1989 to 2019; this allowed us to include all years when auxiliary stations were sampled, thereby maximizing and standardizing the spatial extent of sampling (Figure 1). Dissolved oxygen has been measured since 1992, so the long-term average was calculated using observations from 1992 to 2019. In all cases, conditions in 2020 were compared with those in the period 1989 to 2019 (temperature, salinity) or 1992 to 2019 (dissolved oxygen).

Water temperatures tend to follow a well-defined pattern of high temperatures in rounds 1 and 2, followed by declining temperatures as the sampling season progresses (rounds 3, 4, and 5; Figure 7). This pattern was altered slightly in 2020: mean water temperatures were above historic averages during rounds 1, 2, 3, and 5 ranging from 26-34°C (Figure 7). During round 4, mean water temperatures generally declined to below historic averages. These high water temperatures were largely consistent with statewide average air temperatures from July to September of 2020, which were “above average” in Virginia (NCDC 2020). Relatively high water temperatures in Striped Bass nursery areas have now occurred in eight consecutive years, with a similarly high range of temperatures observed since 2013 (Buchanan et al. 2020). This temperature pattern did not seem to affect catches in previous years, however. Similarly, catch rates in 2020 followed the historic pattern with respect to water temperature: 100% of juvenile Striped Bass were captured at temperatures exceeding 25°C (Table 7). Water temperatures in tidal tributaries reflect not only long-term, regional climate patterns, but also significant day-to-day and local variation. Shallow shoreline areas are easily affected by local events such as thunderstorms and small-scale spatial and temporal variations associated with time of sampling (e.g., morning versus afternoon, riparian shading, tidal stage). As noted in previous reports, the relationship between declining Striped Bass catches and decreasing temperatures during rounds 3, 4, and 5 that was typically observed prior to 2020 is considered to be largely the result of a coincident downward decline in catch rates and water temperatures as the season progresses (after early-August) rather than any direct effects of water temperature on juvenile fish distribution.

Across years, mean salinity tends to increase steadily from rounds 1 to 3, then levels off during rounds 4 and 5 (Figure 8). In 2020, average salinities generally overlapped with or were above historical averages on all rivers, except for the James and Chickahominy Rivers, during rounds 1 and 2 (Figure 8). In round 3, average salinities generally dropped below historical averages, most likely due to Tropical Storm Isaias. In round 4, average salinities continued to drop. In round 5, however, an increase in average salinities was observed. As observed in the past, greater catches of young-of-the-year Striped Bass in 2020 were obtained at salinities less than 5 ppt on average (Table 5). In 2020, salinities less than 5 ppt were observed further downriver than usual. No index stations had salinities exceeding 10.0 ppt on average in 2020, whereas the highest mean salinity of 14.2 ppt (observed at Y15 in 2020) was lower than the long-term average at that site (Table 5).

Mean dissolved oxygen (DO) concentrations in 2020 generally overlapped with or were above long-term averages during rounds 1 and 2 within all rivers (Figure 9). During round 3, DO concentrations were above historical averages in the James, Chickahominy, and Rappahannock Rivers, but were below in the Mattaponi, Pamunkey, and York Rivers. In rounds 4 and 5, DO concentrations were generally lower than historical averages. A notable exception to this pattern was in the York River, where DO concentrations were below historical averages during all rounds except round 4. Relationships between DO and juvenile Striped Bass catches are difficult to ascertain, as lower-than-average DO conditions occur inconsistently through time and across sampling sites. In previous years, high seasonal catches at index stations occurred during periods when DO concentrations were more than one standard error (SE) below the historic average, as well as when DO concentrations were within one SE of the historic average. Thus, DO concentrations do not appear to be a primary driver of abundance of juvenile Striped Bass in nursery areas that are sampled by the seine.

Striped Bass recruitment variability may be partially explained by regional climate patterns during winter and spring (Wood 2000). For example, abundance of young Striped Bass in the Patuxent River is positively associated with high freshwater

flow during the preceding winter (Wingate and Secor 2008). One of the strongest Striped Bass year classes in Virginia was produced in 2011, which was characterized by relatively high freshwater flow in winter and spring (Machut and Fabrizio 2012). Freshwater flow in Virginia tidal tributaries varies seasonally, with monthly averages since 1967 showing relatively high flow during the winter, peaks in early-spring (March-April), followed by steady declines through the late-spring and summer (Figure 10). In 2020, freshwater flow was below average to average from January to July in most rivers, except the James River, which exhibited above average discharge in February, April, May, and June (Figure 10). Freshwater flow peaked in August, most likely due to Tropical Storm Isaias, and remained above average in September in most rivers (Figure 10). Statewide precipitation during the winter and spring of 2020 (December 2019-May 2020) was “much above average” in Virginia relative to historical conditions since 1895 (NCDC 2020). The high precipitation and freshwater flow during several months in 2020 may have contributed to the 2020 Striped Bass year-class success in the Chesapeake Bay. However, it remains unclear if regional climate patterns and/or other factors exert effects on variations in recruitment of juvenile Striped Bass.

#### ***Additional Abundance Indices Calculated from the Seine Survey***

A variety of fish species are collected annually by the juvenile Striped Bass seine survey due to a sampling regime that spans the euryhaline to freshwater zone. The five most common species encountered in 2020 were Threadfin Shad (*Dorosoma petenense*), White Perch (*Morone americana*), Spottail Shiner (*Notropis hudsonius*), Blueback Herring (*Alosa aestivalis*), and Bay Anchovy (*Anchoa mitchilli*). In 2020, more than 49,000 individuals comprising 74 species were collected (Table 8). Indices of abundance were estimated for 10 of these species (in addition to Striped Bass) based on catches from only the first haul at a subset of index and auxiliary stations. A different subset of stations was used for each species, based on the range of sites where the species was commonly encountered within each tributary from 1967-2010.

One of the most common species captured annually by the seine survey, White Perch, supports important recreational fisheries in Chesapeake Bay (Murdy et al. 1997, NMFS 2017). The general overlap in spawning time and use of nursery grounds by White Perch and Striped Bass suggest that the seine survey may adequately sample juvenile White Perch and that calculation of a recruitment index for this species is appropriate. Colvocoresses (1988) found a strong correlation between a young-of-the-year White Perch index (geometric mean) calculated from seine survey data and an index obtained for harvest-sized White Perch from a trawl survey. In years of low abundance (e.g., 1985) the proportion of seine hauls containing White Perch may be as low as 40%; whereas in years of high abundance (e.g., 2011), White Perch may be found in 95% of seine hauls. A delta-lognormal index was developed to address this inter-annual variation and to accommodate data with a high proportion of zero hauls. We used Cox's method (Fletcher 2008) to estimate the mean abundance based on the delta-lognormal distribution, and calculated 95% confidence intervals from 1,000 bootstrap samples as described by Fletcher (2008). This approach remains under development, so we report only the means here.

Throughout the 2020 sampling period, 6,305 young-of-the-year White Perch were collected from 133 seine hauls at 30 sites (11 sites in the James, 10 in the York and 9 in the Rappahannock). Because White Perch movement among Virginia tributaries is unlikely (Mulligan and Chapman 1989), we presume each tributary supports a distinct stock and report juvenile abundance for each river system separately (Table 9; Figures 11-14). Generally, river-specific JAIs for White Perch suggest above-average recruitment in the James, York, and Rappahannock rivers in 2020 (Figures 12-14). Although we feel confident in the estimation of annual mean relative abundance of White Perch, alternative approaches for estimating confidence intervals need to be examined. The White Perch JAI developed by the seine survey compliments the juvenile White Perch index currently reported by the VIMS Juvenile Fish Trawl Survey (Tuckey and Fabrizio 2020); however, unlike the index reported by the trawl survey, the seine survey index is based on catches from tidal brackish and freshwater zones.

Atlantic Croaker (*Micropogonias undulatus*) is another economically and recreationally important fish (Murphy et al. 1997, NMFS 2017) regularly collected by the seine survey. Young-of-the-year Atlantic Croaker are collected at predominantly mesohaline regions during rounds 1 to 3, before fish are able to avoid capture by the net (Williams and Fabrizio 2011). Murphy et al. (1997) report peak spawning of Atlantic Croaker from August to October; thus, young-of-the-year fish collected during 2020 were spawned during fall 2019. Similar to White Perch, Atlantic Croaker raw catches exhibit high annual variability in the proportion of nonzero hauls. To address this variation and accommodate data with a high proportion of zero hauls we developed a delta-lognormal index for Atlantic Croaker (as described above). Atlantic Croaker are coastal shelf spawners, and their larvae migrate into Chesapeake Bay and enter nursery areas in the tributaries. Therefore, we report a Virginia-wide estimate of juvenile abundance (Table 10; Figure 15). Based on 2020 catches from 21 stations during rounds 1 to 3, we encountered 749 young-of-the-year Atlantic Croaker and these fish were captured in 40 seine hauls (Table 10; Figure 15). Periods of strong recruitment from 1992-1995, 1997-1998, and 2007-2009 correspond with patterns observed by the VIMS Juvenile Fish Trawl Survey (Tuckey and Fabrizio 2020). In 2020, an above-average year class for Atlantic Croaker appears to have occurred.

Spot (*Leiostomus xanthurus*), like Atlantic Croaker, is another economically and recreationally important species that is collected by the seine survey and reported as a Virginia-wide estimate of juvenile abundance (Table 11; Figure 16). Based on catches from 21 stations during 5 rounds in 2020, 305 young-of-the-year Spot were collected in 50 seine hauls. Using the delta-lognormal approach, we observed a below-average year class for Spot in 2020, similar to estimates from the previous five years (Table 11; Figure 16).

Indices of abundance for common forage species within the tidal nearshore zone were computed for Spottail Shiner (32 stations; Table 12), Atlantic Silverside (24 stations; Table 13), Inland Silverside (*Menidia beryllina*; 36 stations; Table 14), and Banded Killifish (*Fundulus diaphanus*; 32 stations; Table 15). Catches from 5 rounds

were used to estimate abundance indices for these species. The 2020 Spottail Shiner delta-lognormal mean of 26.3 was similar to the historic average of 27.8 (Table 12). The 2020 Atlantic Silverside delta-lognormal mean of 30.4 was lower than the historic average of 49.7 (Table 13). The 2020 Inland Silverside abundance index of 10.2 was higher than the historic average of 5.2 (Table 14). The 2020 Banded Killifish delta-lognormal mean of 9.1 was higher than the historic average of 5.4 (Table 15). Together, these results suggest modest production of forage fish prey was available for piscivores in Virginia waters in 2020. In addition, we note that abundance indices for the three freshwater forage species (Spottail Shiner, Inland Silverside and Banded Killifish) have been increasing since 1989, with each species displaying a statistically significant temporal trend.

Indices of abundance derived from seine survey collections are reported for species of management importance to fulfill Commonwealth compliance requirements to the ASMFC; these species include American Shad (Watkins et al. 2011), Alewife, Blueback Herring, and Atlantic Menhaden (VMRC 2010). Abundance estimates for juvenile American Shad from the seine survey were highly correlated with those from push-net sampling (Wilhite et al. 2003), providing support for the seine survey-based index. These indices are provided to VMRC when requested and are also reported here. Alosines greatly contribute to the dynamics of freshwater, estuarine, and marine habitats serving as prey for many large, predatory fishes and consuming large amounts of plankton. Many stocks of alosine species are currently at record lows or of unknown status because of a lack of data to assess populations accurately, especially within riverine environments. Data collected on American Shad, Alewife, and Blueback Herring from the seine survey are critical for assessing populations in the James, York, and Rappahannock rivers. The 2020 geometric mean abundance index for American shad was relatively high in the Rappahannock River, but below average in the James and York rivers (Figure 17). The 2020 geometric mean abundance index for Alewife was average in the Rappahannock River, but below average in the James and York (Figure 18). The



2020 geometric mean abundance indices for Blueback Herring were average in the three river systems (Figure 19).

## **CONCLUSION**

The 2020 juvenile abundance index (JAI) for Striped Bass (13.89) appears to be above average when compared with the reference period (7.77) for Virginia waters. The relatively high estimate for 2020 appears to have resulted from the inability of researchers to sample 19% of index sites in 2020. When the missed index sites are accounted for in the reference period index calculation, the average for the reference period is 8.68 (LCI = 6.69, UCI = 11.11). When compared to the recalculated reference period index, the 2020 index suggests an average year (Appendix Table 2). Reference period indices were also recalculated for each of the rivers and survey rounds (Appendix Table 3). When the 2020 JAIs for each river and round are compared to their respective recalculated reference period indices, all suggest average recruitment. Continued monitoring of juvenile Striped Bass abundance is important in predicting recruitment to the Striped Bass fisheries in the Chesapeake Bay and along the Atlantic coast. A critical characteristic of the long-term annual seine survey conducted in the Chesapeake Bay is the ability to identify years of below-average recruitment which, if persistent, serve as an early warning to managers of potential declines in Striped Bass spawning stock biomass or deteriorating conditions in nursery areas. Juvenile White Perch and Atlantic Croaker abundance indices in 2020 were higher than the historic averages for these species. The Spot abundance index was below average in 2020. Abundance indices for three Alosine species were average or below average in the James and York rivers and average or above average in the Rappahannock River in 2020, relative to index values in previous years.

## **ACKNOWLEDGMENTS**

We are indebted to the many landowners who graciously allowed us access to their waterfront properties. We thank the Mariners' Museum, Jamestown 4-H Camp, June Parker Marina, and Chickahominy Riverfront Park for their permission to sample. Additional thanks go to Jordan Point Marina, June Parker Marina and Chickahominy Riverfront Park for permission to use their boat ramps. Summer technicians in 2020 were Connor Morang and Jordan Wheatley. We also thank VIMS students/staff who assisted in the field, particularly Daniel Royster and Matthew Oliver. Funding was provided by a grant from the United States Fish and Wildlife Service Sport Fish Restoration Project (F87R32) through the Virginia Marine Resources Commission to the Virginia Institute of Marine Science.

## LITERATURE CITED

- ASMFC (Atlantic States Marine Fisheries Commission). 2003. Amendment #6 to the Interstate Fishery Management Plan for Atlantic Striped Bass. Fisheries Management Report 41. Atlantic States Marine Fisheries Commission, Washington, D.C. 63 p.
- ASMFC (Atlantic States Marine Fisheries Commission). 2010. Addendum II to Amendment #6 to the Interstate Fishery Management Plan: definition of recruitment failure. Atlantic States Marine Fisheries Commission, Washington, D.C. 18 p.
- Austin, H.M., J.A. Colvocoresses and T.A. Mosca III. 1993. Develop a Chesapeake Baywide young-of-the-year Striped Bass index: final report to the Chesapeake Bay Stock Assessment Committee (CBSAC). Cooperative Agreement NA16FUO393-01. Virginia Institute of Marine Science, Gloucester Point, Virginia. 59 p. + 2 app.
- Buchanan, J.R., M.C. Fabrizio and T.D. Tuckey. 2020. Estimation of Juvenile Striped Bass relative abundance in the Virginia portion of Chesapeake Bay. Annual Report 2018. Virginia Institute of Marine Science, Gloucester Point, VA. 62 p.
- Colvocoresses, J.A. 1984. Striped Bass research, Virginia. Part I: Juvenile Striped Bass seining program. Annual Report 1987-88. Virginia Institute of Marine Science, Gloucester Point, Virginia. 64 p.
- Colvocoresses, J.A. 1987. Intercalibration and refinement of estimates of abundance of Chesapeake Bay juvenile Striped Bass. NOAA Tech. Rept. TRS-SAC-91-010, 28 p.
- Colvocoresses, J.A. 1988. Comparisons among York River White Perch stock abundance measures. NOAA Tech. Rept. TRS-SAC-91-021, 18 p.
- Colvocoresses, J.A. and H.M. Austin. 1987. Development of an index of juvenile Striped Bass abundance for the Chesapeake Bay System: I. An evaluation of present measures and recommendations for future studies. Special Science Report 120. Virginia Institute of Marine Science, Gloucester Point, VA. 108 p.
- Fabrizio, M.C., T.D. Tuckey, O.M. Phillips and B.K. Gallagher. 2017. Tracking decadal changes in Striped Bass recruitment: A calibration study of seine surveys in Chesapeake Bay. Virginia Institute of Marine Science, Gloucester Point, VA. 71 p.
- Fletcher, D. 2008. Confidence intervals for the mean of the delta-lognormal distribution. *Environmental and Ecological Statistics* 15: 175 - 189.
- Gallagher, B.K., M.C. Fabrizio and T.D. Tuckey. 2017. Estimation of juvenile Striped Bass relative abundance in the Virginia portion of Chesapeake Bay. Annual Report 2016. Virginia Institute of Marine Science, Gloucester Point, VA. 59 p.
- Gallagher, B.K., M.C. Fabrizio and T.D. Tuckey. 2018. Estimation of juvenile Striped Bass relative abundance in the Virginia portion of Chesapeake Bay. Annual Report 2017. Virginia Institute of Marine Science, Gloucester Point, VA. 60 p.

- Goodyear, C.P. 1985. Relationship between reported commercial landings and abundance of young Striped Bass in Chesapeake Bay, Maryland. *Transactions of the American Fisheries Society* 114: 92 – 96.
- Hewitt, A.H., J.K. Ellis and M.C. Fabrizio. 2007. Estimation of juvenile Striped Bass relative abundance in the Virginia portion of Chesapeake Bay. Annual Report 2006. Virginia Institute of Marine Science, Gloucester Point, VA. 31 p.
- Hewitt, A.H., L.S. Machut and M.C. Fabrizio. 2008. Estimation of juvenile Striped Bass relative abundance in the Virginia portion of Chesapeake Bay. Annual Report 2007. Virginia Institute of Marine Science, Gloucester Point, VA. 28 p.
- Machut, L.S., and M.C. Fabrizio. 2010. Estimation of juvenile Striped Bass relative abundance in the Virginia portion of Chesapeake Bay. Annual Report 2009. Virginia Institute of Marine Science, Gloucester Point, VA. 45 p.
- Machut, L.S., and M.C. Fabrizio. 2012. Estimation of juvenile Striped Bass relative abundance in the Virginia portion of Chesapeake Bay. Annual Report 2011. Virginia Institute of Marine Science, Gloucester Point, VA. 55 p.
- Mulligan, T.J., and R. W. Chapman. 1989. Mitochondrial DNA analysis of Chesapeake Bay White Perch, *Morone americana*. *Copeia* 3: 679 – 688.
- Murdy, E.O., R.S. Birdsong, and J.A. Musick. 1997. *Fishes of Chesapeake Bay*. Smithsonian Institution Press, Washington, D. C. 324 p.
- NCDC (National Climate Data Center). 2020. National temperature and precipitation maps. Site accessed October 2020. <https://www.ncdc.noaa.gov/temp-and-precip/us-maps/>.
- NMFS (National Marine Fisheries Service). 2017. Annual commercial landing statistics. Site accessed October 2017. [https://www.st.nmfs.noaa.gov/st1/commercial/landings/annual\\_landings.html](https://www.st.nmfs.noaa.gov/st1/commercial/landings/annual_landings.html).
- Peer, A.C., and T.J. Miller. 2014. Climate change, migration phenology, and fisheries management interact with unanticipated consequences. *North American Journal of Fisheries Management* 34(1): 94-110.
- Phillips, O. 2019. Recruitment characteristics of juvenile Striped Bass (*Morone saxatilis*) across recovery periods, year classes, and subestuaries in the Chesapeake Bay. M.S. thesis, William & Mary.
- Rago, P., D. Stephan, and H. Austin. 1995. ASMFC Special Report 48. Report of the juvenile indices abundance workshop, January 1992, Kent Island, MD. 83 p.
- Sokal, R.R. and F.J. Rohlf. 1981. *Biometry*. W.H. Freeman and Co., San Francisco, CA. 851 p.
- Tuckey, T.D., and M.C. Fabrizio. 2020. Estimating relative juvenile abundance of ecologically important finfish in the Virginia portion of Chesapeake Bay. Final Report to the Virginia Marine Resources Commission.

- United States Geological Survey (USGS). 2020. Current conditions for Virginia: streamflow (USGS Water Data for the Nation). Site accessed October 2020. <https://waterdata.usgs.gov/va/nwis/current/?type=flow>.
- Watkins, B.J. Olney, and R. O'Reilly. 2011. A summary of Virginia's American Shad fisheries in 2010 and results of monitoring and restoration programs: annual compliance report to the Atlantic States Marine Fisheries Commission American Shad Technical Committee, Virginia Institute of Marine Sciences, Gloucester Point, VA. 43 pp.
- Wilhite, M.L., K.L. Maki, J.M. Hoenig, and J.E. Olney. 2003. Towards validation of a juvenile index of abundance for American Shad in the York River, Virginia. Pages 285 - 294 *in* K. E. Limburg and J. A. Waldman (eds.) Biodiversity Status and Conservation of the World's Shads. American Fisheries Society Symposium 35, Bethesda, MD.
- Williams, B.D. and M. C. Fabrizio. 2011. Detectability of estuarine fishes in a beach seine survey of tidal tributaries of lower Chesapeake Bay. *Transactions of the American Fisheries Society* 140: 1340-1350.
- Wingate, R.L., and D. H. Secor. 2008. Effects of winter temperature and flow on a summer-fall nursery fish assemblage in the Chesapeake Bay, Maryland. *Transactions of the American Fisheries Society* 137: 1147 - 1156.
- Wood, R.J. 2000. Synoptic scale climatic forcing of multispecies fish recruitment patterns in Chesapeake Bay. Ph.D. Dissertation. William & Mary, Williamsburg, VA.
- Woodward, J.R. 2009. Investigating the relationships between recruitment indices and estimates of adult abundance for Striped Bass, Weakfish, and Atlantic Croaker. Master's thesis. William & Mary, Williamsburg, VA.
- VMRC (Virginia Marine Resources Commission). 2010. Atlantic Menhaden compliance report for Virginia: Report to the Atlantic States Marine Fisheries Commission. Fisheries Management Division, Newport News, VA. 16 pp.



**Table 2.** Catch of young-of-the-year Striped Bass in the primary nursery areas of Virginia (index stations) summarized by year, where  $x$  = total fish,  $\text{Index} = (\exp(\ln(x + 1)) - 1) \times 2.28$ , SD = Standard Deviation, and SE = Standard Error.

Year	Total Fish (x)	Mean $\ln(x+1)$	SD	Index	CI ( $\pm 2$ SE)	N (Hauls)
1967	191	1.18	1.00	5.17	3.20-7.86	42
1968	184	1.04	0.92	4.15	2.68-6.06	50
1969	193	0.97	0.94	3.73	2.39-5.46	55
1970	345	1.39	1.11	6.88	4.52-10.06	56
1971	165	0.90	0.90	3.34	2.17-4.81	60
1972	84	0.45	0.59	1.28	0.87-1.75	90
1973	133	0.60	0.82	1.86	1.12-2.76	70
1980	228	0.74	0.90	2.52	1.68-3.53	89
1981	165	0.52	0.69	1.56	1.10-2.09	116
1982	323	0.78	0.97	2.71	1.85-3.74	106
1983	296	0.91	0.83	3.40	2.53-4.42	102
1984	597	1.09	1.06	4.47	3.22-6.02	106
1985	322	0.72	0.86	2.41	1.78-3.14	142
1986	669	1.12	1.04	4.74	3.62-6.06	144
1987	2,191	2.07	1.23	15.74	12.40-19.83	144
1988	1,348	1.47	1.13	7.64	6.10-9.45	180
1989	1,978	1.78	1.12	11.23	9.15-13.68	180
1990	1,249	1.44	1.10	7.34	5.89-9.05	180
1991	667	0.97	0.95	3.76	2.96-4.68	180
1992	1,769	1.44	1.24	7.35	5.72-9.31	180
1993	2,323	2.19	0.98	18.11	15.35-21.30	180
1994	1,510	1.72	1.03	10.48	8.66-12.60	180
1995	926	1.22	1.05	5.45	4.33-6.75	180
1996	3,759	2.41	1.23	23.00	18.77-28.07	180
1997	1,484	1.63	1.10	9.35	7.59-11.41	180
1998	2,084	1.92	1.14	13.25	10.82-16.12	180
1999	442	0.80	0.86	2.80	2.19-3.50	180
2000	2,741	2.09	1.24	16.18	13.06-19.92	180
2001	2,624	1.98	1.27	14.17	11.33-17.60	180
2002	813	1.01	1.09	3.98	3.05-5.08	180
2003	3,406	2.40	1.18	22.89	18.84-27.71	180
2004	1,928	1.88	1.04	12.70	10.54-15.22	180
2005	1,352	1.61	1.05	9.09	7.45-11.02	180
2006	1,408	1.69	1.04	10.10	8.31-12.18	180
2007	1,999	1.83	1.18	11.96	9.66-14.70	180
2008	1,518	1.50	1.17	7.97	6.33-9.93	180
2009	1,408	1.55	1.10	8.42	6.80-10.32	180
2010	1,721	1.61	1.25	9.07	7.14-11.40	180
2011	4,189	2.56	1.19	27.09	22.30-32.80	178
2012	408	0.78	0.83	2.68	2.10-3.33	179
2013	1,620	1.76	1.08	10.94	8.97-13.25	180
2014	2,293	1.78	1.26	11.30	8.98-14.09	181
2015	1,879	1.84	1.13	12.00	9.78-14.64	179
2016	1,557	1.58	1.17	8.74	6.98-10.84	180
2017	2,060	1.61	1.28	9.17	7.18-11.57	180
2018	1,875	1.74	1.19	10.72	8.61-13.24	180
2019	1,624	1.65	1.14	9.54	7.69-11.74	180
2020	1,836	1.96	1.15	13.89	11.08-17.29	145
Reference (1980-2009)	43,527	1.48	0.53	7.77	6.01-9.89	30 (years)

**Table 3.** Catch of young-of-the-year Striped Bass in the primary nursery areas of Virginia using only the 1st haul (Rago et al. 1995), where  $x$  = total fish, Index =  $(\exp(\ln(x + 1)) - 1) \times 2.28$ , SD = Standard Deviation, and SE = Standard Error.

Year	Total Fish (x)	Mean $\ln(x+1)$	SD	Index	CI ( $\pm 2$ SE)	N (Hauls)
1967	191	1.18	1.00	5.17	3.20-7.86	42
1968	184	1.04	0.92	4.15	2.68-6.06	50
1969	193	0.97	0.94	3.73	2.39-5.46	55
1970	345	1.39	1.11	6.88	4.52-10.06	56
1971	165	0.90	0.90	3.34	2.17-4.81	60
1972	84	0.45	0.59	1.28	0.87-1.75	90
1973	133	0.60	0.82	1.86	1.12-2.76	70
1980	216	0.82	0.96	2.90	1.85-4.21	72
1981	112	0.64	0.74	2.05	1.28-2.99	58
1982	172	0.86	0.96	3.10	1.86-4.71	54
1983	185	0.97	0.94	3.74	2.36-5.54	52
1984	377	1.27	1.09	5.81	3.72-8.63	53
1985	216	0.94	0.92	3.54	2.40-4.97	71
1986	449	1.35	1.07	6.53	4.56-9.06	72
1987	1,314	2.27	1.22	19.77	14.25-27.13	72
1988	820	1.57	1.21	8.66	6.20-11.85	90
1989	1,427	2.06	1.18	15.68	11.71-20.77	90
1990	720	1.58	1.12	8.76	6.44-11.70	90
1991	462	1.17	1.05	5.04	3.59-6.85	90
1992	1,143	1.65	1.31	9.63	6.76-13.41	90
1993	1,241	2.34	0.89	21.36	17.31-26.25	90
1994	969	1.93	1.09	13.37	10.17-17.40	90
1995	559	1.37	1.07	6.71	4.89-8.99	90
1996	2,326	2.60	1.27	28.29	21.11-37.69	90
1997	931	1.83	1.14	11.92	8.90-15.76	90
1998	1,365	2.12	1.22	16.66	12.35-22.23	90
1999	274	0.92	0.91	3.43	2.43-4.64	90
2000	1,528	2.22	1.23	18.70	13.91-24.90	90
2001	1,671	2.16	1.32	17.52	12.70-23.89	90
2002	486	1.17	1.13	5.03	3.48-7.01	90
2003	2,042	2.50	1.26	25.61	19.09-34.13	90
2004	1,129	2.07	1.04	15.75	12.19-20.19	90
2005	835	1.79	1.07	11.42	8.64-14.90	90
2006	767	1.76	1.06	11.02	8.34-14.36	90
2007	1,271	2.09	1.21	16.07	11.95-21.39	90
2008	867	1.70	1.11	10.15	7.56-13.42	90
2009	861	1.72	1.11	10.47	7.81-13.83	90
2010	994	1.75	1.26	10.83	7.78-14.82	90
2011	2,397	2.70	1.17	31.69	24.29-41.16	90
2012	265	0.92	0.87	3.47	2.50-4.63	90
2013	900	1.83	1.11	11.99	9-15.76	90
2014	1,401	2.01	1.24	14.81	10.87-19.93	90
2015	978	1.92	1.09	13.21	10.02-17.22	90
2016	783	1.60	1.16	9.06	6.60-12.21	90
2017	1,200	1.69	1.29	10.09	7.13-13.96	90
2018	1,072	1.80	1.24	11.54	8.37-15.66	90
2019	880	1.70	1.18	10.24	7.49-13.77	90
2020	1,256	2.04	1.17	15.29	11.43-20.24	89
1980-2009	26,735	1.65	0.54	9.57	7.43-12.17	30 (years)
1990-2012	25,103	1.83	0.50	11.91	9.25-15.17	23 (years)



**Table 4.** Catch of young-of-the-year Striped Bass per seine haul at index stations in 2020 summarized by drainage and river.

Drainage River	<u>2020</u>				<u>Reference Period</u> (1980-2009)				
	Total Fish	Index	C.I. (± 2 SE)	N (hauls)	Total Fish	Index	C.I. (± 2 SE)	N (years)	
JAMES	620	20.07	15.01-26.60	46	17,650	10.41	7.83-13.64	30	
James	436	19.12	12.61-28.47	30	10,727	9.72	7.06-13.12	30	
Chickahominy	184	21.96	15.74-30.32	16	6,923	11.95	8.70-16.15	30	
YORK	319	7.68	5.57-10.37	56	12,470	5.85	4.50-7.48	30	
Pamunkey	121	7.30	4.59-11.08	24	6,442	6.90	4.90-9.44	30	
Mattaponi	198	7.98	5.03-12.12	32	6,028	5.16	4.06-6.45	30	
RAPPAHANNOCK	897	19.20	11.43-31.39	43	13,407	7.90	5.63-10.82	30	
Overall	1,836	13.89	11.08-17.29	145	43,527	7.77	6.01-9.89	30	

**Table 5.** Striped Bass indices and average site salinity during 2020 compared to average index values during the auxiliary monitoring period (1989-2019), with corresponding average salinities (Avg. Sal., ppt). The York drainage includes Pamunkey and Mattaponi rivers. Index stations are indicated by bold font. Indices are calculated using only the 1st haul (Rago et al. 1995). Sites J26, P56, and R76 were established in 2020, thus, auxiliary monitoring period data are unavailable.

Drainage														
<b>JAMES</b>														
	Station	J12	J26	<b>J29</b>	<b>J36</b>	J42	<b>C1</b>	<b>C3</b>	<b>J46</b>	J51	<b>J56</b>	J62	J68	J77
1989-2019	Avg. Sal.	14.4	-	4.8	2.4	1.3	1.4	1.2	0.5	0.2	0.2	0.2	0.1	0.1
	Index	1.6	-	11.1	17.2	11.2	22.7	12.0	25.5	17.5	9.9	11.7	6.0	2.9
2020	Avg. Sal.	12.8	3.6	1.8	0.5	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1
	Index	0.0	20.4	11.0	5.7	5.7	36.1	23.1	26.8	23.1	100.8	3.7	7.3	1.8
<b>YORK</b>														
	Station	Y15	Y21	Y28	P36	<b>P42</b>	<b>P45</b>	<b>P50</b>	P56					
1989-2019	Avg. Sal.	16.5	13.6	10.6	4.1	1.7	0.7	0.4	-					
	Index	1.4	2.6	7.1	11.3	5.2	13.2	19.2	-					
2020	Avg. Sal.	14.2	12.8	7.0	2.3	0.8	0.2	0.1	0.1					
	Index	0.0	9.8	6.7	9.2	4.3	8.0	16.1	0.3					
	Station				<b>M33</b>	M37	<b>M41</b>	<b>M44</b>	<b>M47</b>	M52				
1989-2019	Avg. Sal.				4.5	2.2	1.1	0.4	0.2	0.1				
	Index				9.3	10.9	7.6	9.1	5.9	1.3				
2020	Avg. Sal.				2.4	1.0	0.2	0.1	0.0	0.0				
	Index				15.6	18.8	4.4	19.5	7.7	3.7				
<b>RAPPAHANNOCK</b>														
	Station	R12	R21	<b>R28</b>	<b>R37</b>	R41	<b>R44</b>	<b>R50</b>	<b>R55</b>	R60	R65	R69	R76	
1989-2019	Avg. Sal.	14.0	12.6	9.9	5.1	2.9	1.7	0.7	0.5	0.2	0.2	0.1	-	
	Index	0.7	0.7	4.6	3.3	6.0	12.1	21.8	46.9	6.1	4.7	2.5	-	
2020	Avg. Sal.	13.9	12.3	8.8	3.2	1.3	0.6	0.2	0.1	0.0	0.0	0.0	0.0	
	Index	0.0	0.0	1.0	1.5	1.0	31.5	58.3	94.8	4.5	6.7	4.3	1.0	

**Table 6.** Catch of young-of-the-year Striped Bass at index stations in 2020 summarized by sampling round.

Month (Round)	N (hauls)	<u>2020</u>			Change From Previous Round	<u>Reference Period (1980-2009)</u>				Change From Previous Round
		Total Fish	Index	C.I. ( $\pm$ 2 SE)		N (years)	Total Fish	Index	C.I. ( $\pm$ 2 SE)	
July (1 <sup>st</sup> )	34	520	12.8	7.29-21.5		30	13,467	11.97	9.15-15.48	
(2 <sup>nd</sup> )	35	519	17.22	10.72-26.98	-0.2%	30	10,535	9.11	6.84-11.95	-21.8%
Aug. (3 <sup>rd</sup> )	20	313	21.07	12.23-35.29	-39.7%	30	7,838	7.26	5.44-9.50	-25.6%
(4 <sup>th</sup> )	35	287	10.02	6.46-15.02	-8.3%	26	6,907	6.88	5.12-9.04	-11.9%
Sept. (5 <sup>th</sup> )	21	197	12.44	7.2-20.57	-31.4%	23	4,780	6.04	4.73-7.61	-30.8%

**Table 7.** Catch of young-of-the-year Striped Bass per seine haul in the primary nursery areas of Virginia in 2020 summarized by water temperature.

Temp (°C)	<u>2020</u>				<u>Reference Period</u> (1980-2009)			
	Total Fish	Index	C.I. (± 2 SE)	N (sites)	Total Fish	Index	C.I. (± 2 SE)	N (sites)
15.0-19.9	-	-	-	-	47	1.98	0.46-4.34	19
20.0-24.9	-	-	-	-	2,430	4.13	3.61-4.7	568
25.0-29.9	641	9.81	7.4-12.83	79	33,808	9.11	8.66-9.57	3,588
> 30.0	1,195	20.60	14.6-28.74	66	6,871	9.66	8.6-10.82	679

**Table 8.** Fish species collected during the 2020 seine survey (index and auxiliary stations).

<b>Scientific Name</b>	<b>Common Name</b>	<b>Total Caught</b>
<i>Dorosoma petenense</i>	Threadfin Shad	9556
<i>Morone americana</i>	White Perch	8905
<i>Notropis hudsonius</i>	Spottail Shiner	3934
<i>Alosa aestivalis</i>	Blueback Herring	3478
<i>Anchoa mitchilli</i>	Bay Anchovy	3187
<i>Menidia menidia</i>	Atlantic Silverside	2600
<i>Brevoortia tyrannus</i>	Atlantic Menhaden	2346
<i>Morone saxatilis</i>	Striped Bass	2181
<i>Menidia beryllina</i>	Inland Silverside	2119
<i>Dorosoma cepedianum</i>	Gizzard Shad	1823
<i>Fundulus heteroclitus</i>	Mummichog	1592
<i>Fundulus diaphanus</i>	Banded Killifish	1553
<i>Micropogonias undulatus</i>	Atlantic Croaker	1086
<i>Trinectes maculatus</i>	Hogchoker	841
<i>Alosa sapidissima</i>	American Shad	771
<i>Hybognathus regius</i>	Eastern Silvery Minnow	665
<i>Notropis analostanus</i>	Satinfin Shiner	481
<i>Lepomis macrochirus</i>	Bluegill	403
<i>Leiostomus xanthurus</i>	Spot	334
<i>Fundulus majalis</i>	Striped Killifish	203
<i>Anchoa hepsetus</i>	Striped Anchovy	146
<i>Etheostoma olmstedii</i>	Tessellated Darter	123
<i>Ictalurus furcatus</i>	Blue Catfish	123
<i>Membras martinica</i>	Rough Silverside	107
<i>Menticirrhus americanus</i>	Southern Kingfish	102
<i>Lepomis auritus</i>	Redbreast Sunfish	89
<i>Notemigonus crysoleucas</i>	Golden Shiner	71
<i>Micropterus punctulatus</i>	Spotted Bass	69
<i>Lepomis gibbosus</i>	Pumpkinseed	62
<i>Mugil cephalus</i>	Striped Mullet	62
<i>Strongylura marina</i>	Atlantic Needlefish	52
<i>Enneacanthus gloriosus</i>	Bluespotted Sunfish	47
<i>Bairdiella chrysoura</i>	Silver Perch	37
<i>Ictalurus punctatus</i>	Channel Catfish	36
<i>Gambusia affinis</i>	Mosquitofish	31
<i>Anguilla rostrata</i>	American Eel	25
<i>Ictalurus catus</i>	White Catfish	22
<i>Cynoscion regalis</i>	Weakfish	21
<i>Carpoides cyprinus</i>	Quillback	21
<i>Morone saxatilis</i> age 1+	Striped Bass Age 1+	18

**Table 8.** (continued)

<b>Scientific Name</b>	<b>Common Name</b>	<b>Total Caught</b>
<i>Micropterus salmoides</i>	Largemouth Bass	16
<i>Alosa mediocris</i>	Hickory Shad	15
<i>Pomatomus saltatrix</i>	Bluefish	13
<i>Perca flavescens</i>	Yellow Perch	13
<i>Synodus foetens</i>	Inshore Lizardfish	12
<i>Opisthonema oglinum</i>	Atlantic Thread Herring	11
<i>Lepisosteus osseus</i>	Longnose Gar	11
<i>Symphurus plagiusa</i>	Blackcheek Tonguefish	10
<i>Mugil curema</i>	White Mullet	9
<i>Elops saurus</i>	Ladyfish	9
<i>Pomoxis nigromaculatus</i>	Black Crappie	8
<i>Alosa pseudoharengus</i>	Alewife	7
<i>Scomberomorus maculatus</i>	Spanish Mackerel	7
<i>Sciaenops ocellatus</i>	Red Drum	6
<i>Pogonius cromis</i>	Black Drum	4
<i>Cyprinus carpio</i>	Common Carp	4
<i>Cynoscion nebulosus</i>	Spotted Seatrout	4
<i>Ictalurus nebulosus</i>	Brown Bullhead	3
<i>Syngnathus fuscus</i>	Northern Pipefish	3
<i>Alosa spp</i>	Herring Spp	3
<i>Paralichthys dentatus</i>	Summer Flounder	2
<i>Menticirrhus saxatilis</i>	Northern Kingfish	2
<i>Rhinoptera bonasus</i>	Cownose Ray	2
<i>Peprilus alepidotus</i>	Harvestfish	1
<i>Trachinotus carolinus</i>	Florida Pompano	1
<i>Chaetodipterus faber</i>	Atlantic Spadefish	1
<i>Moxostoma macrolepidotum</i>	Shorthead Redhorse	1
<i>Cyprinodon variegatus</i>	Sheepshead Minnow	1
<i>Astroscopus guttatus</i>	Northern Stargazer	1
<i>Micropterus dolomieu</i>	Smallmouth Bass	1
<i>Lepomis microlophus</i>	Redear Sunfish	1
<i>Syngnathus louisianae</i>	Chain Pipefish	1
<i>Sander vitreus</i>	Walleye	1
<i>Gobionellus boleosoma</i>	Darter Goby	1
Total		49,507

**Table 9.** Delta-lognormal mean of young-of-the-year White Perch from select seine survey stations by river system and year.

Year	James River		York River		Rappahannock River		N (hauls)
	# of Fish	Delta Mean	# of Fish	Delta Mean	# of Fish	Delta Mean	
1967	341	26.3	6	0.7	256	34.0	26
1968	48	2.4	10	0.7	125	6.9	19
1969	446	21.6	106	7.4	242	14.0	39
1970	1,582	78.2	7	0.5	267	23.5	48
1971	334	16.6	17	1.5	311	23.2	44
1972	38	1.4	247	7.1	392	42.5	57
1973	34	1.4	71	4.1	296	15.9	53
1980	62	2.3	211	15.6	145	9.3	34
1981	97	3.2	18	0.6	133	8.8	41
1982	18	1.3	292	20.2	126	16.5	28
1983	162	10.5	175	9.9	128	13.7	40
1984	94	5.6	100	5.4	156	24.7	44
1985	23	1.0	88	3.2	31	2.3	25
1986	421	18.8	79	2.9	336	39.1	49
1987	712	39.3	880	63.2	1,177	60.5	63
1988	457	22.1	69	2.2	287	13.7	61
1989	424	13.0	807	28.2	1,349	49.6	104
1990	235	5.9	70	1.7	487	11.7	84
1991	296	6.4	169	4.2	387	13.5	91
1992	338	7.7	4	0.1	395	11.9	67
1993	3,812	107.8	344	7.6	1,177	46.5	113
1994	608	17.8	420	9.4	655	19.1	125
1995	741	18.8	17	0.3	418	12.2	93
1996	4,784	166.9	1,654	66.5	2,294	78.9	126
1997	1,703	59.0	305	8.3	248	6.3	102
1998	1,432	35.5	195	4.7	457	18.5	108
1999	159	3.4	1	0.0	486	13.2	67
2000	1,540	38.5	1,363	40.0	1,184	34.2	121
2001	948	20.8	799	21.1	1,126	32.3	123
2002	790	19.1	129	2.7	275	7.0	83
2003	1,364	35.7	1,132	27.8	1,849	70.4	120
2004	1,030	23.8	799	22.0	670	17.9	130
2005	1,871	54.9	579	15.3	834	28.1	122
2006	2,064	44.9	95	2.8	388	10.0	99
2007	2,896	69.2	417	22.7	830	24.5	113
2008	1,627	40.5	184	4.1	1,512	69.6	107
2009	3,825	125.2	10	0.2	1,813	77.7	90
2010	3,085	100.1	1,632	43.6	728	19.1	130
2011	15,805	709.0	4,112	132.6	4,169	164.6	140
2012	1,233	25.1	47	1.0	338	8.8	99
2013	1,640	43.3	433	10.4	623	17.5	119
2014	2,198	71.4	2,373	62.0	841	22.0	120
2015	1,518	32.6	1,621	53.5	1,017	25.3	139
2016	1,474	32.0	980	30.8	1,286	41.2	121
2017	3,804	113.9	460	10.6	2,576	101.6	126
2018	4,757	111.1	1,025	30.7	1,976	56.6	136
2019	2,961	63.7	1,746	42.2	2,529	70.6	141
2020	3,658	87.4	867	20.6	1,780	64.0	133

**Table 10.** Delta-lognormal mean of young-of-the-year Atlantic Croaker from select seine survey stations in Virginia tributaries of Chesapeake Bay by year.

<b>Year</b>	<b>Total Fish</b>	<b>Delta Mean</b>	<b>N (hauls)</b>
1980	167	5.3	20
1981	0	0	0
1982	52	1.1	5
1983	114	5.4	10
1984	17	0.5	4
1985	129	4.1	14
1986	9	0.7	4
1987	46	1.8	9
1988	10	0.6	4
1989	112	1.4	16
1990	20	0.3	2
1991	636	10.0	48
1992	717	11.6	41
1993	1,115	30.1	47
1994	862	16.9	39
1995	598	13.8	36
1996	18	0.4	3
1997	955	27.1	48
1998	840	14.6	43
1999	519	9.4	38
2000	21	0.3	10
2001	35	0.9	11
2002	146	2.2	29
2003	8	0.1	4
2004	185	4.7	20
2005	177	6.5	24
2006	399	6.7	37
2007	329	16.3	21
2008	1,306	71.4	52
2009	1,724	50.1	46
2010	76	2.0	13
2011	36	0.5	10
2012	953	22.8	49
2013	771	16.4	36
2014	9	0.2	2
2015	7	0.1	2
2016	483	12.8	23
2017	230	6.4	24
2018	65	0.6	13
2019	68	1.7	13
2020	749	11.6	40
Overall (1980-2019)	13,964	10.0	40 (years)



**Table 11.** Delta-lognormal mean of young-of-the-year Spot from select seine survey stations in Virginia tributaries of Chesapeake Bay by year.

Year	Total Fish	Delta Mean	N (hauls)
1967	73	2.3	14
1968	655	11.6	38
1969	528	9.6	50
1970	57	0.6	25
1971	704	11.8	58
1972	443	2.6	54
1973	2,306	49.0	72
1980	2,174	25.0	72
1981	829	14.5	43
1982	631	91.7	18
1983	130	5.6	17
1984	899	30.6	19
1985	406	12.2	26
1986	1,338	60.1	33
1987	161	5.1	15
1988	943	20.9	37
1989	1,319	21.1	52
1990	1,050	11.1	62
1991	1,069	12.7	74
1992	525	5.9	65
1993	961	10.9	74
1994	990	9.9	60
1995	237	2.3	40
1996	728	11.6	44
1997	1,900	25.3	78
1998	881	15.6	55
1999	888	11.0	78
2000	465	6.1	46
2001	484	6.5	53
2002	185	1.7	44
2003	470	5.9	27
2004	581	6.1	51
2005	2,711	27.6	87
2006	471	5.1	66
2007	977	17.0	77
2008	906	9.7	84
2009	1,208	13.9	73
2010	2,801	30.4	87
2011	669	12.4	60
2012	581	6.6	66
2013	635	12.1	58
2014	566	13.0	45
2015	44	0.5	11
2016	113	1.3	27
2017	221	2.6	42
2018	294	3.1	34
2019	316	3.4	43
2020	305	3.0	50
Overall (1967-2019)	37,523	12.8	47 (years)

**Table 12.** Delta-lognormal mean of young-of-the-year Spottail Shiner from select seine survey stations in Virginia tributaries of Chesapeake Bay by year.

<b>Year</b>	<b>Total Fish</b>	<b>Delta Mean</b>	<b>N (hauls)</b>
1989	2,843	22.3	115
1990	2,019	15.3	104
1991	1,394	10.8	94
1992	2,313	17.5	99
1993	1,708	12.8	99
1994	2,286	18.6	110
1995	2,212	18.0	105
1996	2,182	18.4	109
1997	3,568	25.9	105
1998	2,100	16.3	101
1999	1,149	8.3	81
2000	4,857	40.2	113
2001	2,721	21.7	113
2002	1,381	9.9	71
2003	3,070	23.4	126
2004	5,133	42.0	127
2005	3,597	30.6	112
2006	3,464	29.2	107
2007	3,837	33.7	111
2008	2,147	17.9	95
2009	3,035	24.1	101
2010	3,989	27.0	105
2011	6,284	58.5	122
2012	4,022	30.8	103
2013	4,325	33.7	109
2014	3,401	24.8	125
2015	4,463	33.8	131
2016	3,397	25.1	122
2017	5,436	43.6	112
2018	6,528	60.3	125
2019	8,169	70.8	124
2020	3,436	26.3	121
Overall (1989-2019)	107,030	27.8	31 (years)

**Table 13.** Delta-lognormal mean of young-of-the-year Atlantic Silverside from select seine survey stations in Virginia tributaries of Chesapeake Bay by year.

<b>Year</b>	<b>Total Fish</b>	<b>Delta Mean</b>	<b>N (Hauls)</b>
1989	1,089	10.8	41
1990	2,917	46.6	51
1991	2,855	42.2	68
1992	6,087	122.8	58
1993	2,364	32.0	59
1994	2,305	32.4	52
1995	3,079	41.3	59
1996	4,871	93.4	52
1997	1,160	13.3	55
1998	2,434	26.4	66
1999	6,822	68.6	88
2000	3,778	43.9	65
2001	4,015	53.4	73
2002	5,387	67.0	96
2003	3,351	55.4	35
2004	1,503	21.8	39
2005	1,979	22.1	69
2006	2,847	31.7	67
2007	2,067	29.5	68
2008	3,454	34.7	58
2009	2,916	37.4	72
2010	1,723	18.4	86
2011	3,585	47.2	75
2012	1,381	13.9	68
2013	6,814	95.1	59
2014	4,891	69.6	67
2015	7,542	103.1	74
2016	2,397	27.1	56
2017	5,259	80.5	73
2018	8,071	136.9	46
2019	2,561	32.5	54
2020	2,500	30.4	53
Overall (1989-2019)	111,504	49.7	31 (years)

**Table 14.** Delta-lognormal mean of young-of-the-year Inland Silverside from select seine survey stations in Virginia tributaries of Chesapeake Bay by year.

<b>Year</b>	<b>Total Fish</b>	<b>Delta Mean</b>	<b>N (Hauls)</b>
1989	495	3.0	86
1990	591	3.8	76
1991	286	1.8	66
1992	339	1.8	60
1993	385	2.3	59
1994	171	1.0	49
1995	109	0.7	48
1996	807	5.4	60
1997	201	1.2	57
1998	213	1.4	61
1999	307	1.9	58
2000	729	5.1	77
2001	660	4.1	66
2002	498	3.0	67
2003	574	3.4	98
2004	1,125	6.6	84
2005	419	2.5	78
2006	1,184	7.5	88
2007	861	5.4	78
2008	704	3.9	92
2009	1,751	9.8	113
2010	1,507	8.8	78
2011	1,476	7.6	89
2012	962	5.2	111
2013	1,658	10.3	109
2014	1,849	10.7	107
2015	1,618	9.9	108
2016	2,160	10.9	119
2017	1,627	9.2	117
2018	1,095	6.3	105
2019	1,277	8.1	105
2020	1,686	10.2	105
Overall (1989-2019)	27,638	5.2	31 (years)

**Table 15.** Delta-lognormal mean of young-of-the-year Banded Killifish from select seine survey stations in Virginia tributaries of Chesapeake Bay by year.

<b>Year</b>	<b>Total Fish</b>	<b>Delta Mean</b>	<b>N (Hauls)</b>
1989	236	1.5	47
1990	238	1.6	50
1991	263	2.0	42
1992	153	1.1	35
1993	264	2.0	41
1994	203	1.4	43
1995	287	2.1	38
1996	654	5.0	64
1997	365	2.6	60
1998	311	2.2	61
1999	297	2.1	49
2000	252	1.7	54
2001	355	2.3	70
2002	364	2.6	49
2003	802	5.7	68
2004	1,383	9.7	89
2005	715	5.6	68
2006	498	4.0	48
2007	692	5.1	75
2008	1,025	6.7	87
2009	1,208	9.0	85
2010	1,965	14.8	97
2011	1,958	14.1	88
2012	1,865	13.6	97
2013	638	4.5	70
2014	715	4.6	87
2015	879	5.4	93
2016	1,834	13.2	108
2017	697	4.5	105
2018	849	5.7	94
2019	1,714	11.4	108
2020	1,232	9.1	91
Overall (1989-2019)	23,679	5.4	31 (years)

## FIGURES




Figure 1. Juvenile Striped Bass seine survey stations. Station numbers denote the approximate river mile from the mouth.




Figure 2. Scaled geometric mean of young-of-the-year Striped Bass in the primary nursery areas of Virginia (index stations) by year. Vertical bars are 95% confidence intervals as estimated by  $\pm 2$  standard errors of the mean. Horizontal lines indicate the arithmetic mean (thin solid), confidence intervals (dashed) and 1<sup>st</sup> quartile (thick solid) during the reference period from 1980-2009 (ASMFC 2010).




Figure 3. Scaled geometric mean of young-of-the-year Striped Bass in the primary nursery areas of Virginia (index stations) by drainage and river.






Figure 4. Catch per unit effort of juvenile Striped Bass by station in the James River drainage during each round in 2020. Data are shown for index (black) and auxiliary (red) stations, using the first haul only. The core nursery area is delineated by thick black lines. Hauls were not completed at all index and auxiliary stations during all rounds in 2020 (see Methods).




Figure 5. Catch per unit effort of juvenile Striped Bass by station in the York River drainage during each round in 2020. Data are shown for index (black) and auxiliary (red) stations, using the first haul only. Core nursery areas in the Pamunkey and Mattaponi rivers are delineated by thick black lines. Hauls were not completed at all index and auxiliary stations during all rounds in 2020 (see Methods).




Figure 6. Catch per unit effort of juvenile Striped Bass by station in the Rappahannock River drainage during each round in 2020. Data are shown index (black) and auxiliary (red) stations, using the first haul only. The core nursery area is delineated by thick black lines. Hauls were not completed at all index and auxiliary stations during all rounds in 2020 (see Methods).




Figure 7. Mean water temperature and 95% confidence intervals during each round (x-axis) in each river during 2020 (thin line and error bars) and the auxiliary monitoring period from 1989-2019 (thick line and shaded region).






Figure 8. Mean salinity and 95% confidence intervals during each round (x-axis) in each river during 2020 (thin line and error bars) and the auxiliary monitoring period from 1989-2019 (thick line and shaded region). Note that the scale of the y-axis varies by river.




Figure 9. Mean dissolved oxygen and 95% confidence intervals during each round (x-axis) in each river during 2020 (thin line and error bars) and the monitoring period from 1992-2019 (thick line and shaded region). Note that dissolved oxygen was not measured on the seine survey before 1992.




Figure 10. Mean freshwater flow and 95% confidence intervals during each month from January to September (x-axis) in each river during 2020 (thin line and error bars) and the historical monitoring period from 1967-2019 (thick line and shaded region). Data are from USGS (2020).




Figure 11. Delta-lognormal mean of young-of-the-year White Perch from select seine survey stations by drainage and year.






Figure 12. Delta-lognormal mean of young-of-the-year White Perch from the James River nursery area from 1967-2020. The time series average is shown by the horizontal line.




Figure 13. Delta-lognormal mean of young-of-the-year White Perch from the York River nursery area from 1967-2020. The time series average is shown by the horizontal line.




Figure 14. Delta-lognormal mean of young-of-the-year White Perch from the Rappahannock River nursery area from 1967-2020. The time series average is shown by the horizontal line.




Figure 15. Delta-lognormal mean of young-of-the-year Atlantic Croaker from select seine survey stations in Virginia tributaries of Chesapeake Bay from 1980-2020. The time series average is shown by the horizontal line.




Figure 16. Delta-lognormal mean of young-of-the-year Spot from select seine survey stations in Virginia tributaries of Chesapeake Bay from 1967-2020. The time series average is shown by the horizontal line.




Figure 17. Scaled geometric mean of American Shad in the primary nursery areas of Virginia by drainage and river, using the first haul only.




Figure 18. Scaled geometric mean of Alewife in the primary nursery areas of Virginia by drainage, using the first haul only.




Figure 19. Scaled geometric mean of Blueback Herring in the primary nursery areas of Virginia by drainage, using the first haul only.



## APPENDICES

**Appendix Table 1.** Calibration factors, 95% confidence intervals and sample sizes (N = number of paired hauls) for Striped Bass and White Perch based on paired hauls of the old and new seine nets in 2015 and 2017. Calibration factors are used to adjust catches from the new net and result in old net equivalent catches (see Fabrizio et al. 2017 for details). In the table below, calibration factors were estimated with (2015 and 2017) and without (2015) the addition of observations from 2017. Note that the 95% confidence intervals for these species overlap with 1 when data from 2017 are included.

Species	Year	N	Calibration	
			Factor	95% CI
Striped Bass	2015	21	0.52	0.40-0.83
	2015 and 2017	76	1.11	0.92-1.38
White Perch	2015	27	0.65	0.46-0.86
	2015 and 2017	75	0.85	0.69-1.04

**Appendix Table 2.** Catch of young-of-the-year Striped Bass per seine haul at index stations in 2020 summarized by drainage and river. The annual and reference period indices are calculated from observations from 180 seine hauls across 18 index stations in the James, Rappahannock, and York River watersheds. In 2020, due to Tropical Storm Isaias and precautions taken to protect researchers from COVID-19, researchers completed 145 seine hauls. With fewer hauls in 2020, the annual index may not be directly comparable to the reference period mean, which includes 180 hauls per year. Therefore, the reference period index was recalculated to include only the 145 seine hauls that were completed in 2020. Using the same methodology, reference period indices were also recalculated for the James, Rappahannock, and York drainages and rivers. Lastly, recruitment verdicts are given (below average, average, or above average). Below or above average verdicts indicate that the 2020 index falls outside the reference period mean 95% confidence intervals, and is significantly different from the reference period average. An average verdict indicates that the 2020 index falls within the reference period mean 95% confidence intervals, and is therefore not significantly different from the reference period mean.

Drainage River	<u>2020</u>				<u>Reference Period*</u> (1980-2009)				Verdict
	Total Fish	Index	C.I. (± 2 SE)	N (hauls)	Total Fish	Index	C.I. (± 2 SE)	N (years)	
JAMES	620	20.07	15.01-26.60	46	14,919	11.58	8.75-15.15	30	Average
James	436	19.12	12.61-28.47	30	8,571	10.40	7.52-14.13	30	Average
Chickahominy	184	21.96	15.74-30.32	16	6,348	14.00	10.19-18.98	30	Average
YORK	319	7.68	5.57-10.37	56	11,257	6.48	4.96-8.31	30	Average
Pamunkey	121	7.30	4.59-11.08	24	5,806	7.58	5.37-10.43	30	Average
Mattaponi	198	7.98	5.03-12.12	32	5,451	5.74	4.51-7.19	30	Average
RAPPAHANNOCK	897	19.20	11.43-31.39	43	12,947	9.12	6.44-12.63	30	Average
Overall	1,836	13.89	11.08-17.29	145	39,123	8.68	6.69-11.11	30	Average

\*Calculated using the sites, hauls, and rounds sampled during the 2020 survey.

**Appendix Table 3.** Catch of young-of-the-year Striped Bass at index stations in 2020 summarized by sampling round. The annual and reference period indices are calculated from observations from 180 seine hauls across 18 index stations in the James, Rappahannock, and York River watersheds. In 2020, due to Tropical Storm Isaias and precautions taken to protect researchers from COVID-19, researchers completed 145 seine hauls. With fewer hauls in 2020, the annual index may not be directly comparable to the reference period mean, which includes 180 hauls per year. Therefore, the reference period index was recalculated to include only the 145 seine hauls that were completed in 2020. Using the same methodology, reference period indices were also recalculated for each round of the survey. Lastly, recruitment verdicts are given (below average, average, or above average). Below or above average verdicts indicate that the 2020 index falls outside the reference period mean 95% confidence intervals, and is significantly different from the reference period average. An average verdict indicates that the 2020 index falls within the reference period mean 95% confidence intervals, and is therefore not significantly different from the reference period mean.

Month (Round)	N (hauls)	<u>2020</u>			<u>Reference Period (1980-2009)*</u>				Verdict
		Total Fish	Index	C.I. ( $\pm 2$ SE)	N (years)	Total Fish	Index	C.I. ( $\pm 2$ SE)	
July (1 <sup>st</sup> )	34	520	12.80	7.29-21.50	30	13,089	12.31	9.39-15.95	Average
(2 <sup>nd</sup> )	35	519	17.22	10.72-26.98	30	10,368	9.20	6.90-12.08	Average
Aug. (3 <sup>rd</sup> )	20	313	21.07	12.23-35.29	30	5,440	9.40	7.03-12.37	Average
(4 <sup>th</sup> )	35	287	10.02	6.46-15.02	26	6,848	7.02	5.22-9.25	Average
Sept. (5 <sup>th</sup> )	21	197	12.44	7.20-20.57	23	3,378	7.56	5.94-9.51	Average

\*Calculated using the sites, hauls, and rounds sampled during the 2020 survey.