

## ***Trollius austrosibiricus* (Ranunculaceae), a new species from South Siberia**

Andrey Erst<sup>1,2</sup>, Alexander Luferov<sup>3</sup>, Victoria Troshkina<sup>1</sup>, Dmitry Shauro<sup>1</sup>, Alexander Kuznetsov<sup>2</sup>, Kunli Xiang<sup>4,5</sup>, Wei Wang<sup>4,5</sup>

<sup>1</sup> Laboratory Herbarium (NS), Central Siberian Botanical Garden, Russian Academy of Sciences, Zolotodolinskaya st. 101, 630090, Novosibirsk, Russia;

<sup>2</sup> Laboratory Herbarium (TK), Tomsk State University, Lenin st. 36, 634050, Tomsk, Russia;

<sup>3</sup> I.M. Sechenov First Moscow State Medical University, Izmailovsky Boulevard, 8, 105043, Moscow, Russia;

<sup>4</sup> State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, Chinese Academy of Sciences, 100093, Beijing, China;

<sup>5</sup> University of Chinese Academy of Sciences, 19 Yuquan Road, Beijing 100049, China;

Corresponding author: Andrey Erst (erst\_andrew@yahoo.com)

### **Abstract**

*Trollius austrosibiricus* Erst & Luferov *sp. nov.*, a new species from Russian South Siberia is described and illustrated. This new species is endemic to Western and Central Siberia. Morphologically, it is close to the East Asian species *T. chinensis* and *T. macropetalus*. However, it differs from the aforementioned species due to the morphology of the rhizomes, aerial shoots, sepals and petals. This species is also distinguished from *T. asiaticus*, which is widespread in Russia (Western and Eastern Siberia), Mongolia, China, north-eastern Kazakhstan and in the northeast of the European part of Russia, in having a smaller number of sepals, longer persistent styles, and petals longer than sepals. In addition, identification key for all Russian species are given and all species have been discussed.

### **Keywords**

Ranunculaceae, *Trollius*, new species, South Siberia, Russia.

### **Introduction**

*Trollius* L. (Ranunculaceae) is distributed in temperate to arctic regions of the Northern Hemisphere and has two centres of diversity in SW China and the area ranging from Siberia, the Pamirs and Kashmir (Doroszewska 1974). The genus is characterised by conspicuous orange- or yellow-coloured flowers, similarly coloured petals, subscapose habit, ternate or deeply 3-lobed leaves (Kadota 1987). *Trollius* is distinguished by an unusual floral structure with petals divided into blade, pit (nectarostigma) and claw (Wang & al. 2010). Length ratio of nectaries to stamens has been considered to be one of the most important morphological characters for species delimitation (Shipczinsky 1937, Siplivinsky 1972, Kadota 1987, Tamura 1981, Luferov & al. 2018). Species belonging to the *T.* sect. *Longipetala* Dorosz. are characterised by linear, flat, thin petals, which are longer than the sepals or nearly equal to them. All representatives belonging to this group, except *T. asiaticus*, are common to the Far East part of Asia. All Siberian species are characterised by petals shorter or equal to sepals. When carrying out a revision of the genus Russia, we focused on specimens whose petals are

much longer than the sepals. Further revision of the herbarium material allowed us to describe a new species of *Trollius* from South Siberia.

## Methods

The revision of herbarium material was undertaken in the herbaria LE, MHA, ALTB, NS and NSK (Thiers 2017). The drawings of *Trollius austrosibiricus* are based on the images of the type specimens (NS-0013097). The photographs in the field were taken by a Nikon D90 camera. The morphological characters were measured using AxioVision 4.8. The flowering and fruiting periods and habitats are given as cited on the collector's labels. The IUCN Red List Category Criteria (IUCN 2016) were applied to assess the conservation status. All revised localities of *Trollius austrosibiricus* mentioned in the paper are shown on a map (Fig. 1) made with SimpleMappr (<http://www.simplemappr.net>).

## Taxonomy

*Trollius austrosibiricus* Erst & Luferov, sp. nov.

### Figs 2 & 3

**Type.** RUSSIA. Tuva Republic: Mongun-Tayga region, Tsagan-Shibetu ridge, the upper Barlyk river, valley of the right tributary, the lower part of the south-eastern slope (5°), forb-grass-sedge steppe meadow, 2350 alt., flowering, 22 Jul 1980, V. Khanminchun, M. Danilov & P. Enns without collector number (holotype: NS barcode 0013097!).

**Diagnosis.** *Trollius austrosibiricus* is morphologically close to *T. chinensis* Bunge and *T. macropetalus* (Regel) Fr.Schmidt. It differs from these species in simple rhizomes, shorter aerial shoots, smaller flowers and shorter persistent styles. The new species is distinguished from *T. asiaticus* L. by a smaller number of sepals, longer persistent styles, and petals longer than sepals.

**Description.** *Herbs* perennial, (20–)40–70 cm high. *Rhizomes* simple or slightly branching, short, erect or arched, with a bundle of adventitious roots. *Stems* straight, simple, less often weakly branched, slightly grooved. *Basal leaves* (1–)2–4, spirally-alternate, congested in a basal rosette; petioles 10–25(–35) cm long; blades 4–7 × 5–8 cm, rhomboid, 3–5(–7)-lobed, segments dissected almost to the midrib into lobes, ending sharply with edges dentate. *Cauline leaves* 2–5(–7), opposite, with short petioles or sessile, gradually smaller towards the apex. *Inflorescence* terminal, 1(–2)-flowered. *Flowers* 3.5–4.5(–5.5) cm diam.; pedicels 5–12 cm long, elongating in fruit up to 8–15 cm long; sepals 8–10(–14), 1.3–2.4 × 0.8–1.5 cm, rhombic-ovate or broadly elliptic, reddish-orange or yellow-orange; petals 9–18, 2–2.8 × 0.2–0.3 cm, oblong-lanceolate, slightly wider at middle, base narrow cuneate, apex acute, orange- or reddish-orange coloured, nectarostigma 2.5–3 mm from base; stamens more 10, filaments 7–11mm long, anthers 1.5–2(–2.5) mm long, linear; *Fruits* aggregate, with 9–14(17) follicles, 10–15 mm long, persistent style 1.5–3.0 mm long, slightly incurved.

**Specimens seen (paratypes).** RUSSIA. Tuva Republic: Western Sayan, Kurtushibinskiy ridge, the upper Mynas river, right tributary of the Hut river, forb-grass

meadow, 1080 alt., 6 Jul 1989, D. Shaulo & I. Kovaleva 4695 (NS barcode 0013098!); Mongun-Tayga region, valley of Tolayty river, yernik-sedge wetland meadow, 2500 alt., 26 Jul 1980, M. Danilov & A. Krytsin 932 (NS barcode 0013096!); the northern slope of the East Tannu-Ola ridge, the average flow of Elegest river, floodplain, No., 22 Jul 1973, V. Khanminchun & V. Dyukov 2270 (NS barcode 0013095!); **Kemerovo Oblast**: Tashtagol region, the village of Ust-Kabyrza 4 km above Pysas river, N 52°48', E 88°28', 450 alt., grass meadow, haymaking, 11 Sep 2000, I. M. Krasnoborov, A. I. Shmakov, D. Germann, S. Kostyukov, E. Antonuk, P. Kosachev & A. Vashchenko 207 (NS barcode 0013094!); **Krasnoyarsk Krai**: Abansky region, near Ustyanskoye village, the upper part of the eastern slope, upland meadow, 6 Jul 1956 T. Vagina (NS barcode 0013093!); Ermakovsky district, valley of the US river, elevation 852 alt., 52.2847°N, 93.2517°E, 03 Jul 2010, I.V. Khan & E.A. Balde 152 (NSK barcode 0028601!); **Khakassia Republic**: Abansky ridge, near Biskamzha station, southern slope, burning, N 53.25, E 089.30, 700 alt., 4 Jun 1991, E. Ankipovich (NS barcode 0013092!).

**Affinities.** *Trollius austrosibiricus* is morphologically close to *T. chinensis* Bunge and *T. macropetalus* (Regel) Fr.Schmidt. It is well distinguished by simple rhizomes (rather than by the multi-headed basal part of the plant, as in *T. chinensis* and *T. macropetalus*), shorter aerial shoots, smaller flowers and shorter persistent styles (Table 1). *Trollius chinensis* is an East Asian species occurring in Russia (Primorsky and Khabarovsk territories, Sakhalin), in the north and northeast of China and on the Korean peninsula (Schipczinsky 1937, Siplivinsky 1972, Doroczewska 1974, Kitagawa 1979, Woroshilov 1982, Lufarov 1991, 2004).

This new species is distinguished from *T. asiaticus* L. by a smaller number of sepals, longer persistent styles, and petals longer than sepals (Table 1). *Trollius asiaticus* grows mainly in extra-tropical Asia (Western and Eastern Siberia, Mongolia, northeast Kazakhstan and China), as well as in the northeast of European Russia (Schipczinsky 1937, Siplivinsky 1972, Doroczewska 1974, Borodina-Grabovskaya 2001, Friesen 2003).

**Phenology.** Flowering from April–early May; fruiting in May.

**Distribution.** *Trollius austrosibiricus* is endemic to mountainous areas of the southern part of Western and Central Siberia. Russia: Tuva Republic, Kemerovo district, Krasnoyarsk territory, Khakassia (Figure 1).

**Habitat and ecology.** *Trollius austrosibiricus* grows in subalpine and forest zones, in moist valleys at 350–2400 m elevation. It occurs in forest glades and fringes, in mixed-grass and mixed-grass-cereal dry and swampy meadows, along the banks of rivers, streams and small ponds with fresh water.

**Etymology.** The specific epithet of the new species is derived from the type locality, South Siberia, Russia.

**Preliminary conservation assessment.** No appropriate data on abundance and/or

distribution of the taxon are available. It can be included in the Not Evaluated (NE) category of IUCN Red List Categories (IUCN 2014) as it lacks adequate information to make a direct or indirect assessment of its risk of extinction based on its distribution and/or population status.

### Discussion

According to our data, 16 species of *Trollius* occur in Russia. The "Flora of the USSR" includes 11 species of *Trollius*, one of which are not found in Russia: *T. dschungaricus* Regel is confined to the east of Central Asia and China, (Schipczinsky 1937). The greatest species diversity is observed in Siberia (12 species: our data); 11 species of *Trollius* (Friesen 2012) were previously indicated for this region, of which we recognize 9, and *T. sayanensis* (Malyshev) Sipl. and *T. vicarius* Sipl. are synonyms to *T. asiaticus* L and *T. uniflorus* Sipliv. Nine species grow in the Far East (Luferov & al. 2018), 5 species of *Trollius* are found in the European part of Russia (Tzvelev 2001), and 1 species occurs in the Caucasus (Borodina-Grabovskaya 2012). Many species that have been described recently from Russia require more detailed investigation and evidence of species independence (Luferov & al. 2018). Many taxa of the genus have intermediate characteristics of the groups described by A. Doroszevska (1974) as sections. The groups considered in this monograph require verification, since many species exhibit intermediate features between sections. Molecular phylogeny has not yet been developed, but there are some works related to a small number of species that do not consider the classification of groups (Despres & al. 2003, Wang & al. 2010). Representatives of the *Trollius* section occur in Russia (petals 6–14 mm long, linear, apex rounded or spatulate, with no groove at the base; plants of Asia, Europe and the Caucasus): *T. europaeus*, *T. altaicus*, and *T. ramunculinus*; the *Longipetala* sections (petals 7–40 mm long, oblong, narrow, flat, thin, not thickened and dull, similar to sepals and usually orange): *T. macropetalus*, *T. ledebourii*, *T. austrosibiricus*, *T. asiaticus*, *T. kytmanovii*, and *T. apertus*; the *Insulaetrollius* sections: (petals 2–12 mm long, linear, extended or pyriform, grooved at the base; plants of the Okhotsk-Japan-Kamchatka region): *T. riederianus*, *T. sibiricus* *T. uniflorus*, *T. japonicus*, *T. miyabei*, *T. membranostylis*, and *T. chartosepalus*. It should be noted that many morphological characters used to distinguish between species and groups of species are not fully developed, do not cover the entire morphological diversity of the genus *Trollius*, and are very variable (many hybridogenic taxa have not yet been described, many species ecomorphs are not yet known). For proper identification, classification and understanding of groups in the genus *Trollius*, complex studies are needed, including micro- and macro-morphological, molecular phylogenetic, cytogenetic, etc. This paper presents the results of taxonomic studies that enabled development of the key to identification of the *Trollius* species based on the novel morphological and rhythmological features: for example, indication of the falling (in *T. chartosepalus*) and non-falling sepals (in other species), shape, the size of sepals, petals, persistent styles, their size ratio, the color of the flower elements, leaflets and stigmas.

According the newest data, the key for *Trollius* identification (from Russia) is provided by us.

#### Key to identification species of the genus *Trollius* from Russia

1. Flowers globular, closed due to overlapping sepal edges; petals, stamens and pistils not visible during flowering ..... *T. europaeus*

- Flowers bowl-shaped, saucer-shaped or, if globular, always open: sepal edges do not overlap; petals, stamens and pistils visible during flowering ..... 2
- 2. Petals longer than sepals and protrude from the flower ..... 3
  - Petals equal to sepals or shorter ..... 4
- 3. Sepals 5–7. Petals 30–40 mm long, 1.5–2 times longer than sepals. Persistent styles 3.5–5 mm long ..... *T. macropetalus*
  - Sepals 8–14. Petals 20–28 mm long, 1.2–1.5 times longer than sepals. Persistent styles 1.5–3 mm long ..... *T. austrosibiricus*
- 4. Petals 1.5–2 times longer than stamens ..... 5
  - Petals shorter ..... 8
- 5. Persistent styles and stigmas purplish black or often blackening. Persistent styles 2.5–3 mm long (Plants from the Altai Republic) ..... *T. altaicus*
  - Persistent styles and stigmas light green or yellow-green. Persistent styles less than 2 mm long ..... 6
- 6. Plants 80–100 cm high. Sepals 5–9 (12) ..... *T. ledebourii*
  - Plants 20–75 cm high. Sepals 10–20 ..... 7
- 7. Sepals reddish-orange. Persistent styles 0.5–1 mm long ..... *T. asiaticus*
  - Sepals yellow-orange or yellow. Persistent styles 1.5–2 mm long ..... *T. kytmanovii*
- 8. Plants bloom prior to leaf expansion. Sepals white or pale cream, finely dentate along the edge, do not fall long after flowering. Leaflets up to 25 mm long. Persistent styles 8–18 mm long, equal to or longer than the ovary, thin, straight or slightly curved ..... *T. chartosepalus*
  - Plants bloom with leaves developed. Sepals from pale yellow to orange, typically smooth-edged, fall at the end of flowering. Leaflets up to 15 (18) mm long. Persistent styles not more than 5 mm long, several times shorter than the ovary, more or less thickened, incurved ..... 9
- 9. Persistent styles 4–5 mm long, 3 times shorter than leaflets. Sepals golden-yellow ..... *T. ramunculimus*
  - Persistent styles less than 3 mm long, 5–7 times shorter than leaflets ..... 10
- 10. Petals equal to stamens, 1–3 mm longer or shorter ..... 11
  - Petals 2–2.5 times shorter than stamens ..... 15
- 11. Sepals 5–7 ..... 12
  - Sepals 9–12 ..... 14
- 12. Petals oblong-obovate, apex cuneate. Persistent styles 0.4–1.2 mm long ..... *T. apertus*
  - Petals obovate or spatulate, apex rounded. Persistent styles not less than 2 mm long ..... 13
- 13. Leaf blades round-pentagonal, 4–8 cm long, 5–10 cm wide, dentate with triangular acute and sharp teeth. Flowers 3–4 cm in diameter. Sepals yellow-orange or yellow. Petals are reddish-orange, equal to stamens or 1–3 mm longer. Leaflets with arcuate, unbreakable persistent styles 2–3 mm long ..... *T. riederianus*
  - Leaf blades are rounded- reniform, 8–14 cm long, 10–24 cm wide, dentate with narrow-triangular acute and sharp teeth. Flowers 2.5–3.5 cm in diameter. Sepals pale yellow or yellow-orange. Petals orange, 1–3 mm shorter than stamens. Leaflets straight or slightly arcuate, with longer (3–4.5 mm long), thin, brittle persistent

style ..... *T. japonicus*

14. Plants up to 40 cm high. Stem simple, with 1 flower. Sepals gray-yellow. Petals narrow-linear, acute, yellow-orange, 1–3 mm longer than stamens. Pedicels up to 10 cm long with fruits up to 20 cm. Leaflets are light green. (Plants from Siberia and the mainland of the Far East) ..... *T. sibiricus*

– Plants 70–120 cm high. Stems branched, with 2–5 flowers, less often simple. Sepals yellow-orange or golden-yellow. Petals obovate or spatulate, blunt, orange, equal to stamens in length. Pedicels 2–5 cm long with fruits up to 10 cm. Leaflets reddish-brown later blackening. (Plants from Sakhalin Island) ..... *T. miyabei*

15. Sepals 5–6. Persistent styles up to 1.4 mm long, subulate, straight or slightly curved ..... *T. uniflorus*

– Sepals 9–12. Persistent styles about 2 mm long, with flattened edges, webbed, arched ..... *T. membranostylis*

**Table 1. Morphological comparison between *Trollius austrosibiricus* and related species**

Characters	<i>T. austrosibiricus</i>	<i>T. chinensis</i>	<i>T. macropetalus</i>	<i>T. asiaticus</i>
Stem height, cm	(20)40–70	70–150 (180)	70–150 (180)	20–75
Undeground organs	simple rhizomes	multi-headed basal part	multi-headed basal part	multi-headed basal part
Length/width of the basal leaf blade	4–7/5–8	6–15/7–25	6–15/7–25	4–7/5–8
Number of flowers on shoot	1(2)	2–7(1)	2–7(1)	1–2(3)
Flower diameter, cm	3.5–4.5(5.5)	4–5(6)	(4)5–7(8)	3–4(5)
Number of sepals	8–10(14)	8–12	5–7	10–20
Petal length, mm	20–28	25–35	30–35 (40)	14–22
Petal apex	Acute, narrowed smoothly from the middle part of the petal	Acute, narrowed abruptly upwards	Acuminate, narrowed smoothly upwards	Rounded
Distance from nectary pit to petal base	2.5–3	3.5–4	3.5–4.5	1.5–2
Sepals/Petals length ratio	<1	<1	<1	≥1
Petal/Stamen length ratio	2–3.5/1	2–3/1	3.5–5/1	1.5–2/1
Follicle length, mm	10–15	14–18	14–22	8–12
Style length, mm	1.5–3	3–4	3.5–5	0.5–1(2)
Persistent style shape	Almost erect at the base, above: bent arc-like inwards	Bent outwards at the base, above: bent slightly arc-like towards the centre of the flower	Bent outwards at the base, above: bent slightly arc-like towards the centre of the flower	Bent sharply arc-like towards the centre of the flower

## Acknowledgements

We thank the staff of the Herbaria visited, I. Khan and E. Balde for the images of *Trollius* flowers, and Iuzhakova Maria for text correction. The research was supported by the Scientific programme AAAA-A17-117012610055-3 of the Central Siberian Botanical Garden, of SB RAS (revision in the herbarium NSK) and the Tomsk State University competitiveness improvement programme (revision in the herbarium NS) and the Russian Foundation for Basic Research, grant 18-04-0653 A (field work in South Siberia and DNA work). We would like to thank the reviewers for their valuable comments and suggestions.

## References

Borodina-Grabovskaya AE (2001) *Trollius* L. In: *Plantae Asiae Centralis*, Vol. 12, SPb.: Publ. SPKHFA, 17-21 [In Russian].

Borodina-Grabovskaya AE (2012) *Trollius* L. In: *Conspectus Florae Caucasi*. T. 3, SPb. - Moscow: 67 [In Russian].

Despres L, Gielly L, Redoutet B, Taberlet P (2003) Using AFLP to resolve phylogenetic relationships in a morphologically diversified plant species complex when nuclear and chloroplast sequences fail to reveal variability. *Molecular Phylogenetics and Evolution*. 27: 185–196.

Doroczewska A (1974) The genus *Trollius* L. A taxonomical study. *Monographiae botanicae*. Warszawa: Panstwowe wydawnictwo naukowe. 41. 184 p.

Friesen NV (2003) *Trollius* L. *Flora of Siberia*. Enfield (NH), USA; Plymouth, UK: Science Publ., Inc. 6: 98-103.

IUCN (2014) The IUCN Red List of threatened species: Version 2013.2. Cambridge U.K.: IUCN Red List Unit. Available from: <http://www.iucnredlist.org> (accessed: 17 May 2017).

Kitagawa M (1979) Neo-Lineamenta Florae Manshuricae. *Flora et vegetatio mundi*. Vaduz: Gantner. 4. 717 p.

Luferov AN (1991) New nomenclatural combinations in taxonomy of Far Eastern species of the family Ranunculaceae Juss. *Bulletin of Moscow Society of Naturalists, Biological series*, 96(5): 74-76. [In Russian].

Luferov AN (2004) A taxonomic synopsis of Ranunculaceae of the Far East of Russia. *Turczaninowia*. 7(1): 1-85 [In Russian].

Luferov AN, Erst AS, Luferov DN, Shmakov AI, Wang W (2018) The genus *Trollius* (Ranunculaceae) in the Russian Far East. *Turczaninowia*. 21(2): 110-116 [In Russian].



<https://doi.org/10.14258/turczaninowia.21.2>

Schipczinsky NV (1937) Genus globe-flower – *Trollius* L. In: Flora of the USSR. Moscow, Leningrad: Publishing House of the Academy of Sciences of the USSR. 7: 42-53 [In Russian].

Siplivinsky VN (1972) The genus *Trollius* L. on the North and East of Asia. Novitates systematicae plantarum vascularium. Leningrad. 9: 163-182 [In Russian with Latin description].

Thiers B (2017) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/> (accessed 1 July 2017).

Tzvelev N (2001) Globe-flower – *Trollius* L. In: Flora Europae Orientalis. T. 10. Petropoli: Mir i Semia, 2001: 40-43.

Wang W, Hu H, Xiang X-G, Yu S-X, Chen Z-D (2010) Phylogenetic placements of *Calathodes* and *Megaleranthis* (Ranunculaceae): Evidence from molecular and morphological data. *Taxon*. 59 (6): 1712-1720. <https://doi.org/10.2307/41059868>

Woroshilov VN (1982) Plants of the Soviet Far East. Moscow: Nauka, 672 pp. [In Russian].

#### Figure Captions


Figure 1. Distribution map of *Trollius austrosibiricus*.

Figure 2. Photograph of *Trollius austrosibiricus*. A – Flowering plant, B – Flower, C – Leaf laminae (Photographs by E. Balde and A. Erst).


Figure 3. *Trollius austrosibiricus*. A – General view, B – Sepal, C – Petal, D – Stamen, E – Fruit, F – Follicle. Scale bar: 1 cm (A-F).



**FIGURE 1.** Distribution map of *Trollius austrosibiricus*.



**FIGURE 2.** Photograph of *Trollius austrosibiricus*. **A.** Flowering plant; **B.** Flower; **C.** Leaf blade. (Photographs by E. Balde and A. Erst).



**FIGURE 3.** *Trollius austrosibiricus*. A – General view, B – Sepal, C – Petal, D – Stamen, E – Fruit, F – Follicle. Scale bar: 1 cm (A-F).