

Antioxidant Activity and Cytotoxicity of *Taraxacum hispanicum* Aqueous and Ethanolic Extracts on HepG2 Cells

Laranjeira C^{1*}, Nogueira A¹, Almeida R², Oliveira A I³, Oliveira R F^{3,4}, Pinho C³, Cruz A³

¹Escola Superior de Saúde (ESS), Instituto Politécnico do Porto (IPP), Vila Nova de Gaia, Portugal.

²Department of Botany, Faculty of Sciences, University of Porto, Porto, Portugal & CIBIO/InBIO Research Network in Biodiversity and Evolutionary Biology.

³Núcleo de Investigação e Intervenção em Farmácia (NIIF), Centro de Investigação em Saúde e Ambiente (CISA), Escola Superior de Saúde (ESS), Instituto Politécnico do Porto (IPP), Vila Nova de Gaia, Portugal.

⁴Secção Autónoma de Ciências da Saúde, Universidade de Aveiro, Aveiro, Portugal.

Received: 13th Nov, 16; Revised: 26th Nov, 16; Accepted: 6th Dec, 16; Available Online: 15th January, 2017

ABSTRACT

Introduction: Plants belonging to the genus *Taraxacum* have been used in traditional medicine. Nowadays, extracts of these plants have been reported for the treatment of diseases, including liver disorders. Increasing interest and research on these plants also revealed its potential for treating cancer. This study aims to evaluate the antioxidant activity and cytotoxic properties of crude extracts from aerial parts of *Taraxacum hispanicum* H.Lindb, against human hepatocarcinoma (HepG2). Material and methods: Evaluation of the antioxidant properties was performed using DPPH *in vitro* test, superoxide scavenging assay and Fe²⁺ chelating activity. MTT assay was used to determine metabolic activity, for 24 and 48 hours. Results: For antioxidant capacity of the ethanolic extract (overall the one with the best results), IC₅₀ values were 62.4 ± 6.7 µg/ml (DPPH radical scavenging activity) 53.9 ± 10.3 µg/ml (Fe²⁺ chelating activity) and 2.0 ± 0.3 µg/ml (superoxide scavenging assay). The aqueous and ethanolic extracts had different effects on HepG2 cell viability. Aqueous extract induced cell cytotoxicity in a time and dose-dependent manner, leaving only 52.6% viable cells at a concentration of 200 µg/ml, after 48 h. An increase in the cell viability was seen in the ethanolic extract, from 24 h to 48 h at higher concentrations. Conclusions: Ethanolic extract of *T. hispanicum* was the most promising, presenting anti-oxidative capacity and only the aqueous extract of the plant presented more relevant cytotoxicity over HepG2 cell line. These activities may be related with the extract phenolic content. However, further studies are needed to elucidate the main mechanisms responsible for these potential effects.

Keywords: Antioxidant, cytotoxicity, HepG2 cells, *Taraxacum hispanicum*.

INTRODUCTION

Plants have been used for centuries by people of all cultures for treating various ailments. It is estimated that 80% of the world's population living in developing countries rely on plants as a primary source of healthcare and traditional medical practice¹. The use of medicinal plants over the years has attracted the attention of scientists' worldwide². However, although some plants have promising potential and are widely used, many of them remain untested³.

Reactive oxygen species (ROS) are highly reactive molecules derived from the metabolism of oxygen. ROS induce lipid peroxidation, damage biomolecules, and affect cellular viability⁴. The uncontrolled generation of ROS often correlates directly with molecular markers of many diseases, including cancer⁵. The formation of cancer cells in the human body can be directly induced by free radicals⁶.

Antioxidants are substances capable of neutralize the excess of free radicals and protect cells against their toxic effects⁷. Natural antioxidants, found in plants, can protect

cells from oxidative stress by preventing the formation or detoxifying free radicals, resulting in the prevention of a variety of pathophysiological processes⁸. Therefore, the knowledge and identification of natural products that could limit ROS-mediated injuries and that could act as chemotherapeutic and chemopreventive agents has gained popularity.

Plants of the genus *Taraxacum* play an important role in Chinese, Arabian and Native American traditional medicine, and their leaves and roots are frequently used to treat lung, breast and uterine tumors as well as hepatitis and some digestive diseases^{9,10}.

Taraxacum officinale (L.) Weber ex F.H. Wigg (family Asteraceae), also known as dandelion, is the most studied species, and seems to be the most consumed both as food and in traditional medicine¹¹. The plant contains flavonoids, triterpenes, coumarins, and phytosterols¹². Some studies have revealed that extracts of *T. officinale* possess antioxidant, hepatoprotective, anti-inflammatory, and antitumoral activities^{4,12,13,14}. Although plants of the genus *Taraxacum*, including *T. officinale*, are often used

*Author for Correspondence: catialaranjeira7@gmail.com

in traditional medicine for the treatment of cancer, scientific evidence to support this effect is lacking. Hata et al.,¹⁵ found that upon screening a variety of compounds from wild plants, *T. officinale* was an effective inducer of differentiation in mouse melanoma cells. Koo et al.,¹⁶ showed that antitumor action has been demonstrated for aqueous *T. officinale* extracts, through TNF- α and IL-1 α secretion in HepG2 cells. In another study, Sigstedt et al.,¹⁷ also demonstrated that the crude extract of dandelion leaf decreased the growth of MCF-7/AZ breast cancer cells in an ERK-dependent manner. Chatterjee et al.,¹³ showed that dandelion root extract specifically and effectively induces apoptosis in human melanoma cells without inducing toxicity in noncancerous cells. More recently, Yoon et al.,¹⁸ confirmed that combination treatment of Huh7 cells with TNF-related apoptosis induced ligand (TRAIL) and *T. officinale* led to TRAIL-induced apoptosis.

Studies on the anti-oxidative capacity of *T. officinale* showed that both root and leaf extracts diminish lipid peroxidation and reduce cytochrome c¹⁹. The content of phenolic compounds in dandelion extracts was responsible for the inhibition of reactive oxygen species and nitric oxide induced damage²⁰.

The active phytochemicals of *T. officinale* (the most studied species of the genus) are found in both roots and leaves. Chemical constituents present in the dandelion leaves are bitter sesquiterpene lactones, several polyphenols and coumarins²¹. Other compounds include β -amyrin, taraxasterol and taraxerol as well as free sterols such as sitosterin, stigmasterin, and phytosterin⁹. Polyphenols are the subject of increasing scientific interest because of their beneficial effects on human health²². Some polyphenols like chicoric acid, caffeic acid, luteolin, luteolin 7-glucoside, rutin, and apigenin have been identified in the genus *Taraxacum*^{11,21}.

Due to genus morphology, determination of the correct *Taraxacum* species is very difficult for a non-expert. Furthermore, because *T. officinale* is not present in the South of Europe, other common *Taraxacum* species are being used as traditional medicine in these areas¹¹. To our knowledge, few studies have been reported addressing the cytotoxic effects of *T. hispanicum* extracts, on different cell lines. This study aims to evaluate whether dried aqueous and ethanolic extracts of aerial parts of the plant have an antioxidative and cytotoxic effect on HepG2 cells.

MATERIAL AND METHODS

Chemicals

Dulbecco's modified Eagle's medium (DMEM), antibiotic-antimycotic solution, trypsin solution, 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyl tetrazolium bromide (MTT), 2,2-diphenyl-1-picrylhydrazyl (DPPH), nitroblue tetrazolium (NBT), reduced form of nicotinamide adenine dinucleotide (NADH), N-phenylmethazonium methosulfate (PMS), quercetin, ferrozine, and all other not specified reagents were from Sigma-Aldrich (St. Louis, MO, USA). Fetal bovine serum (FBS) was purchased from Biochrom KG (Berlin, Germany). L(+)-

Ascorbic acid and pyruvate were purchased from Panreac (Barcelona, Spain). Ethylenediaminetetraacetic acid (EDTA) and dimethyl sulfoxide (DMSO) were purchased from VWR (Portugal). All other chemicals were of analytical reagent-grade.

Plant material and preparation of the extracts

The aerial parts of *T. hispanicum* used in this study were collected from Vila Nova de Gaia (Portugal) during May 2016, identified and authenticated by a specialist at Sciences Faculty, University of Porto, Portugal. A voucher specimen was kept in the herbarium of the institution (PO-V62372). The plant parts were washed under running tap water to remove dirt and soil and finally rinsed with distilled water.

The aqueous extract of *T. hispanicum* for *in vitro* screening was prepared as follows: 75 g of the dried leaves were soaked in water for 24 h at room temperature and in the darkness. The mixture was filtered through a paper filter (Whatman, No. 1), to remove particulate matter, lyophilized and the resulting powder was stored at 4°C for further use. Freeze-drying conditions were of 0.008 mBar, for 3 days, with a condenser surface temperature of -77°C.

For the ethanolic extract of *T. hispanicum* 75 g of the dried leaves were macerated with an ethanol-water solution (80:20, V/V), at room temperature and in the darkness for 4-5 days. The mixture was then filtered through a paper filter (Whatman, No. 1), and concentrated to obtain the crude extract. Stock solutions of test extracts were made in dimethylsulfoxide (DMSO) and aliquots were kept at -20°C for further use.

Antioxidant Activity

DPPH radical scavenging activity

The measurement of the *T. hispanicum* extract scavenger activity against the DPPH radical was performed in accordance with Lima et al.,²³ with modifications²⁰. DPPH[•] solution (90 μ M) was added to a medium containing different concentrations of the aqueous and ethanolic extract (1, 5, 10, 50, 100, and 200 μ g/mL) or vehicle (EtOH) and incubated in the dark, at room temperature, for 30 min. The reduction of DPPH absorption was measured at 515 nm using a plate reader spectrophotometer. Quercetin was used as a positive control. Radical scavenging ability was calculated using the following formula: radical scavenging ability (%) = 100 x [(AC-AS)/AC], where AC is the absorbance of the control, and AS is the absorbance of the sample (extract or standard). The concentration of the plant extract/control required to scavenge 50% of the total DPPH radicals (IC₅₀) available was calculated.

Superoxide radical scavenging activity

Superoxide radical scavenging activity was determined using the PMS-NADH nonenzymatic assay as previously described²⁴. Briefly, the reaction mixture in the sample wells consisted of NADH (166 μ M), NBT (43 μ M), *T. hispanicum* extracts and PMS (2.7 μ M) in 19 mM phosphate buffer, pH 7.4. The assay was performed at room temperature during 2 min and started with the addition of PMS. The absorbance was determined over time at 560 nm. Ascorbic acid was used as positive

control. Superoxide radical scavenging ability was calculated using the following formula: superoxide radical scavenging ability (%) = $100 \times [(AC-AS)/AC]$, where AC is the absorbance of the control, and AS is the absorbance of the sample (extract or standard). The sample concentration providing 50% inhibition (IC_{50}) was obtained by plotting the inhibition percentage against extracts concentrations.

Fe²⁺ chelating activity

Iron chelating activity of the *T. hispanicum* extracts was evaluated according to Russo et al.,²⁵ with some modifications. Briefly, 50 μ l of the extracts at different concentrations were added to a solution of 0.12 mM of ferrous sulphate ($FeSO_4$) (50 μ l) and 50 μ l of ferrozine (0.6 mM). The mixture was then shaken vigorously and left at room temperature for 10 min. Absorbance of the solution was then measured at 562 nm in a spectrophotometer. EDTA was used as a positive control. The ability of *T. hispanicum* extracts to chelate ferrous ion was calculated relative to the control using the formula: chelating activity (%) = $100 \times [(AC-AS)/AC]$, where AC is the absorbance of the control, and AS is the absorbance of the sample (extract or standard). The IC_{50} value (μ g/ml), which is the concentration of the extract/standard that chelates 50% of the ferrous ion, was calculated.

Cell culture

The human hepatocellular carcinoma (HepG2) cell line was obtained from the American Type Culture Collection (ATCC), and maintained in 25 cm² polystyrene flasks in DMEM supplemented with 10% FBS, 1% antibiotic-antimycotic solution, 1 mM sodium pyruvate, 10 mM HEPES and 1.5 g/l sodium bicarbonate. Cells were grown at 37°C in a humidified incubator with 5% CO₂. When the cells reached at >80% confluency, subculture was conducted at a cell density of 2.0×10^6 cells/well. The medium was refreshed three to four times a week.

Assay for cell viability

Cell viability of HepG2 cells was estimated using the MTT colorimetric assay²⁶. Briefly, cells were seeded (2.0×10^5 cells/well) into 24-multiwell culture plates and cultured at 37°C in a humidified atmosphere of 5% CO₂. The plate was incubated for a period of time to assure attachment and 40% to 60% confluency. Cells were then treated with increasing concentrations of each *T. hispanicum* extract (ranging from 1 to 200 μ g/ml), in fresh complete medium for 24 and 48 h.

After the exposure time, MTT solution (final concentration 0.5 mg/ml) was added to each well and incubated for more 1 h. Then, MTT-containing medium was removed gently and replaced with a 50:50 (V/V) DMSO:ethanol solution to mix the formazan crystals until dissolved. Presence of viable cells was visualized by the development of purple color due to formation of formazan crystals. The absorbance was measured at 570 nm using a 96-well microplate reader. The cytotoxicity was expressed as the percentage of control, which had no sample. For each extract tested, the IC_{50} (concentration of drug needed to inhibit cell growth by 50%) was generated from the dose-response curves for HepG2 cell line, using

the GraphPad Prism 5.0 (GraphPad Software, Inc., San Diego, USA).

Statistical Analysis

Data are presented as mean \pm SD of at least three independent experiments. In the *in vitro* cell assays One-way ANOVA was employed with Dunnett's multiple comparison test, when comparing each concentration against a control and unpaired t test when comparing two similar concentrations, using GraphPad Prism 5.0. Differences were considered to be significant at a level of $p < 0.05$. In the *in vitro* antioxidant assays unpaired Students t test was used. The IC_{50} was calculated from the dose-response curve obtained by plotting the percentage of inhibition versus the concentrations (in logarithm), using GraphPad Prism 5.0.

RESULTS

In this study, antioxidant activities of the aqueous and ethanolic extracts of *T. hispanicum* were evaluated by applying three common methods, DPPH, iron chelating assay and superoxide radical scavenging activity. All extracts presented different antioxidant activities (Table 1).

In the measurement of *T. hispanicum* extract scavenger activity against the DPPH radical, ethanolic extract showed significant better results ($IC_{50} = 62.4 \pm 6.7$ μ g/ml) when compared to aqueous extract ($p < 0.001$) ($IC_{50} = 124.5 \pm 2.3$ μ g/ml). According to the results, both plant extracts are not as good as the standard quercetin ($IC_{50} = 1.85 \pm 0.2$ μ g/ml).

Regarding iron chelating activity, best IC_{50} value was shown by aqueous extract ($IC_{50} = 49.7 \pm 11.1$ μ g/ml) when compared to ethanolic extract ($IC_{50} = 53.9 \pm 10.3$ μ g/ml). Again, and according to the results, both plant extracts are not as good as the standard EDTA (1.2 ± 0.0 μ g/ml).

Analyzing the results of the superoxide radical scavenging activity, best IC_{50} value was shown by ethanolic extract (2.0 ± 0.3 μ g/ml), being significantly lower ($p < 0.0001$), when compared with aqueous extract (48.8 ± 3.2 μ g/ml). In both cases, the IC_{50} values were lower than the positive control, ascorbic acid ($IC_{50} = 77.5 \pm 10.9$ μ g/ml), a well-known antioxidant.

To evaluate the cytotoxic activity of aqueous and ethanolic extracts of *T. hispanicum*, against HepG2 cell line, cells were incubated with different concentrations (ranging from 1 to 200 μ g/ml) of extract. After 24 and 48 h of incubation, cell viability was determined by MTT assay. For the aqueous extract, HepG2 cells experienced a significant decrease in viability at the highest concentrations of the extracts, after 24 and 48 h. The extract induced cell cytotoxicity in a time and dose-dependent manner, as presented in Figure 1. Based on metabolic activity of HepG2 cells, treatment with 200 μ g/ml of the aqueous extract resulted in 7.5% and 47.4% reduction in cell viability against control within 24 h and 48 h, respectively (Fig. 1).

The aqueous and ethanolic extracts had different effects on HepG2 cell viability, with aqueous extract being more toxic after 48 h of exposure, leaving only 52.6% viable at

Table 1: Antioxidant activities of studied plant extracts from *T. hispanicum*.

Plant Extract	IC ₅₀ value (µg/ml)			Fe ²⁺ chelating ability	Superoxide radical scavenging activity
	DPPH scavenging	free radical			
<i>Aqueous extract</i>	124.5 ± 2.3			49.7 ± 11.1	48.8 ± 3.2
<i>Ethanollic extract</i>	62.4 ± 6.7 [#]			53.9 ± 10.3	2.0 ± 0.3 [‡]

n = 3; results are showed as mean ± standard deviation

[#]significantly different (p < 0.001)

[‡]significantly different (p < 0.0001)

a concentration of 200 µg/ml (Fig. 1).

On the other hand, in the ethanolic extract the percentage of living HepG2 cells significantly increased from 79.1% (at 24 h) to 119.0% (at 48 h) at 200.0 µg/ml, as presented in Figure 2. An increase in the cell viability was seen only in the ethanolic extract, from 24 h to 48 h at higher concentrations (25 to 200 µg/ml).

DISCUSSION

In this study, we evaluate antioxidant activities of aqueous and ethanolic extracts of *T. hispanicum* by applying different methods, based on single electron transfer mechanisms (DPPH), metal ion chelation and free radical scavenging.

Antioxidant properties found in plant extracts have been attributed to polyphenols²⁷. Polyphenols act as antioxidants through several mechanisms, like radical scavenger, metal-chelating agents, hydrogen donor, or electron donor²⁸.

DPPH assay is a decolorization method capable of measuring the relative antioxidant abilities of natural compounds and extracts to scavenge free radicals. It is a commonly used antioxidant method due to excellent reproducibility, stability, commercial availability and by the fact that it is an easy measurement method²⁹. The effect of antioxidants on DPPH radical scavenging is thought to be due to their hydrogen-donating ability³⁰.

Superoxide radical is formed in almost all aerobic cells as a result of one oxygen electron donation, being relevant in the oxygen toxicity mechanism³¹. By itself, superoxide is a weak radical but it may cause severe damage to the cell by generating hydroxyl radical and singlet oxygen³². The scavenging activity of *T. hispanicum* extracts was established by the NBT reduction method, where the non-enzymatic PMS/NADH system generates superoxide radicals, which reduce the yellow NBT²⁺ in order to produce the purple NBT formazan. Antioxidants have the ability to inhibit purple NBT formation through competition with NBT for superoxide radical³¹.

Iron can stimulate lipid peroxidation by the Fenton reaction and by decomposing lipid hydroperoxides into peroxy and alkoxy radicals that can perpetuate chain of reaction. Metal chelating capacity is therefore significant since it reduces the concentration of the transition metal that catalyzes lipid peroxidation. Ferrozine produces a violet complex with Fe²⁺. In the presence of a chelating agent, complex formation is interrupted and as a result the violet color of the complex is decreased³³.

In this study, ethanolic extract showed better results in DPPH and superoxide radical scavenging activity, as

presented by IC₅₀ values (Table 1). In their study, Mingarro et al.,¹¹ showed that all extracts studied (*T. obovatum*, *T. marginellum*, *T. hispanicum*, *T. lambinonii*, and *T. lacistrum*) had a moderate capacity to scavenge the DPPH radical.

The best DPPH radical scavenging observed in ethanolic extract of *T. hispanicum* is thought to be due to their hydrogen donating ability³⁴. Hu & Kitts²⁰ showed that DPPH radical scavenging activity in the *T. officinale* flower extract was attributed to a reducing activity derived from the phenolic content of the plant. Also, in their study Hu & Kitts²⁰ found that the superoxide radical inhibition of dandelion flower extract was associated with a direct affinity to scavenge superoxide radical, and this observed inhibition could be attributed in part to the phenolic content. Polyphenolic compounds may react with the superoxide radical, via a one-electron transfer, or by a hydrogen abstraction mechanism³⁵.

Several plant extracts exert their antioxidant activity by chelating metals³⁶. The metal chelating ability of flavonoids is dependent on their unique phenolic structure and position of hydroxyl groups³⁷. In this study, both extracts did not show relevant Fe²⁺ chelating activity. The majority of compounds present in the extracts are, therefore, unlikely to have the require structure and/or to exist in a sufficient amount to exert this ability, explaining the results observed.

Phenolic compounds, such as luteolin, luteolin-7-glucoside, caffeic acid, chicoric acid, rutin and apigenin are found in extracts of *T. hispanicum*^{11,14}. Compared to roots, dandelion leaves are characterized by higher polyphenol contents²¹. The most abundant phenolic compounds in leaves and flowers are hydroxycinnamic acid derivatives^{21,38}. It has been shown that chicoric acid, a derivative of caffeic acid, has the most powerful antioxidant activity among reference compounds such as caffeic acid or rosmarinic acid³⁹. Various flavonoid glycosides such as 7-*O*-glycosides of apigenin, rutin, and luteolin were identified in *T. hispanicum* aerial parts¹¹. Therefore, the antioxidant activities of plant extracts demonstrated in this study, especially by the ethanolic extract, could be explained by the presence of these compounds, being therefore, related to solvent polarity and solubility of compounds in tested solvents⁴⁰.

Cellular viability is affected by oxidative stress, since the production of reactive species causes damage to the inner and outer mitochondrial membranes and opens the mitochondrial permeability transition pores, thereby inducing apoptosis⁴¹. In this study, the MTT assay was used to estimate the cell viability. The assay is based on




Figure 1: Cytotoxicity of different concentrations of aqueous extract, ranging from 1.0 to 200.0 µg/ml, from *T. hispanicum* in HepG2. Cell viability was determined by MTT assay. Data are presented as mean ± SD of at least three independent experiments, and significant differences as compared with the control group were established at ***p < 0.001; ###p < 0.001.




Figure 2: Cytotoxicity of different concentrations of ethanolic extract, ranging from 1 to 200 µg/ml, from *T. hispanicum* in HepG2. Cell viability was determined by MTT assay. Data are presented as mean ± SD of at least three independent experiments, and significant differences as compared with the control group were established at *p < 0.05, #p < 0.05, ##p < 0.01, ***p < 0.001, ###p < 0.001.

the capacity of mitochondrial dehydrogenase enzymes in living cells to convert the yellow water-soluble tetrazolium salt MTT to purple formazan crystals (insoluble in water) only by viable cells⁴². HepG2 cells, which are easy to handle, retain many of the morphological characteristics of liver parenchymal cells⁴³, and contain several enzymes responsible for the activation of various xenobiotics^{44,45}.

Aqueous extract of *T. hispanicum* significantly induced cell death of human hepatocellular carcinoma cells. Aqueous extracts of *T. officinale* have been reported to decrease the growth and invasive ability of breast cancer cells¹⁷. Similar results were presented by Koo et al.¹⁶ where it was found that aqueous *T. officinale* extract cause a time-dependent and partially dose-dependent reduction in cell viability. In their study, at 0.2 mg/ml and 48 h, cell viability of the *T. officinale* treated group was

74% (P < 0.05) of the control. The authors also found that *T. officinale* induced the secretion of TNF-α and IL-1α and then induced apoptosis of HepG2 cells. The inhibitory effect may result from phenolic compounds as dandelion leaves are characterized by higher polyphenolic acids and flavonoid contents⁴⁶. This is further supported by studies that describe the effect of polyphenols on cancer cell proliferation⁴⁷. Hu & Kitts¹⁴ demonstrated that the antioxidant and cytotoxic properties of dandelion flower extracts can partly be attributed to the presence of luteolin and luteolin 7-O-glucoside. The aqueous extract of *T. officinale* also has high contents of chicoric acid that may be responsible for the cytotoxic activity⁴⁸. For example, the biological properties of chicoric acid have been reported to include anticancer and antiviral^{49,50}.

Some important components of the *T. officinale* extract include sesquiterpene lactones and phenylpropanoids,

which are believed to have anti-cancer properties leading to diverse observed effects of dandelion extracts⁵¹. However, other components of the plants of *Taraxacum* genus have not been fully characterized and therefore their activities remain unknown^{46,52}.

The higher polyphenol content of the leaves compared to root extracts of *T. officinale*, also proved to be effective in various *in vitro* systems for the determination of antioxidant properties and radical scavenging capacity¹⁹. These antioxidant properties may be related to the increase in cell viability with time (from 24 to 48 h) observed with ethanolic extract in almost tested concentrations (Fig. 2), suggesting a potential hepatoprotective effect of this extract in HepG2 cells. In their study, Colle et al.⁵³ showed that ethanolic *T. officinale* leaves and root extract significantly attenuated marker enzymes of liver toxicity (aspartate transaminase and alanine transaminase), lipid peroxidation and oxidative stress induced by acetaminophen in mice. These protective effects of *T. officinale* have been suggested to the presence of phenolic compounds in the extract.

CONCLUSION

In this study, the antioxidant properties and cytotoxic properties of aqueous and ethanolic extracts from *T. hispanicum* aerial parts, against human hepatocarcinoma (HepG2) were evaluated. Ethanolic extract, in general, revealed better antioxidant and cytoprotective properties, in comparison with the aqueous extract. Aqueous extract proved to be more toxic, especially after 48 h incubated with HepG2 cells. The results observed seem to be related with the extracts phenolic content. However, further studies are needed to elucidate the main mechanisms responsible for these potential effects.

REFERENCES

- Bodeker C, Bodeker G, Ong CK, Grundy CK, Burford G, Shein K. 2005. WHO Global Atlas of Traditional, Complementary and Alternative Medicine. Geneva, Switzerland: World Health Organization.
- Falodun A, Okenroba LO, Uzoamaka N. Phytochemical screening and anti-inflammatory evaluation of methanolic and aqueous extracts of *Euphorbia heterophylla* Linn (Euphorbiaceae). African Journal of Biotechnology 2006; 5(6):529-531.
- Ekor M. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. Frontiers in Pharmacology 2014; 4:177. doi: 10.3389/fphar.2013.00177.
- Jeon HJ, Kang HJ, Jung HJ, Kang YS, Lim CJ, Kim YM, Park EH. Anti-inflammatory activity of *Taraxacum officinale*. Journal of Ethnopharmacology 2008; 115(1):82-8. doi.org/10.1016/j.jep.2007.09.006.
- Lee KW, Lee HJ. Biphasic effects of dietary antioxidants on oxidative stress-mediated carcinogenesis. Mechanisms of Ageing and Development 2006; 127(5):424-31. doi.org/10.1016/j.mad.2006.01.021.
- Wang Cheng-Yuan, Wu Tien-Chou, Hsieh Shu-Ling, Tsai Yung-Hsiang, Yeh Chia-Wen, Huang Chun-Yung. Antioxidant activity and growth inhibition of human colon cancer cells by crude and purified fucoidan preparations extracted from *Sargassum cristaefolium*. Journal of Food and Drug Analysis 2015; 23(4):766-777. doi.org/10.1016/j.jfda.2015.07.002.
- Pham-Huy LA, He H, Pham-Huy C. Free radicals, antioxidants in disease and health. International Journal of Biomedical Science 2008; 4(2):89-96.
- Mates JM, Sanchez-Jimenez FM. Role of reactive oxygen species in apoptosis: implications for cancer therapy. The International Journal of Biochemistry & Cell Biology 2000; 32(2):157-170. doi.org/10.1016/S1357-2725(99)00088-6.
- Gonzalez-Castejon M, Visioli F, Rodriguez-Casado A. Diverse biological activities of dandelion. Nutrition Reviews 2012; 70(9):534-547. doi:10.1111/j.1753-4887.2012.00509.x.
- Sweeney B, Vora M, Ulbricht C, Basch E. Evidence-based systematic review of dandelion (*Taraxacum officinale*) by natural standard research collaboration. Journal of Herbal Pharmacotherapy 2005; 5(1):79-93. doi: 10.1300/J157v05n01_09.
- Mingarro DM, Plaza A, Galán A, Vicente JA, Martínez MP, Acero N. The effect of five *Taraxacum* species on *in vitro* and *in vivo* antioxidant and antiproliferative activity. Food & Function 2015; 6(8):2787-93. doi: 10.1039/c5fo00645g.
- You Y, Yoo S, Yoon HG, Park J, Lee YH, Kim S, Oh KT, Lee J, Cho HY, Jun W. *In vitro* and *in vivo* hepatoprotective effects of the aqueous extract from *Taraxacum officinale* (dandelion) root against alcohol-induced oxidative stress. Food and Chemical Toxicology 2010; 48(6):1632-7. doi: 10.1016/j.fct.2010.03.037.
- Chatterjee SJ, Ovadje P, Mousa M, Hamm C, Pandey S. The Efficacy of Dandelion Root Extract in Inducing Apoptosis in Drug-Resistant Human Melanoma Cells. Evidence-based Complementary and Alternative Medicine 2011; 2011:129045. doi: 10.1155/2011/129045.
- Hu C, Kitts DD. Antioxidant, prooxidant, and cytotoxic activities of solvent-fractionated dandelion (*Taraxacum officinale*) flower extracts *in vitro*. Journal of Agricultural and Food Chemistry 2003; 51(1):301-10. doi: 10.1021/jf0258858.
- Hata K, Ishikawa K, Hori K, Konishi T. Differentiation-inducing activity of lupeol, a lupane-type triterpene from Chinese dandelion root (Hokouei-kon), on a mouse melanoma cell line. Biological & Pharmaceutical Bulletin 2000; 23(8):962-7.
- Koo HN, Hong SH, Song BK, Kim CH, Yoo YH, Kim HM. *Taraxacum officinale* induces cytotoxicity through TNF-alpha and IL-1alpha secretion in Hep G2 cells. Life Science 2004; 74(9):1149-57. doi.org/10.1016/j.lfs.2003.07.030.
- Sigstedt SC, Hooten CJ, Callewaert MC, Jenkins AR, Romero AE, Pullin MJ, Kornienko A, Lowrey TK, Slambrouck SV, Steelant WF. Evaluation of aqueous extracts of *Taraxacum officinale* on growth and

- invasion of breast and prostate cancer cells. *International Journal of Oncology*. 2008; 32(5):1085-90. doi: 10.3892/ijo.32.5.1085.
18. Yoon JY, Cho HS, Lee JJ, Lee HJ, Jun SY, Lee JH, Song HH, Choi S, Saloura V, Park CG, Kim CH, Kim NS. Novel TRAIL sensitizer *Taraxacum officinale* F.H. Wigg enhances TRAIL-induced apoptosis in Huh7 cells. *Molecular Carcinogenesis* 2016; 55(4):387-96. doi: 10.1002/mc.22288.
 19. Hagymási K, Blázovics A, Fehér J, Lugasi A, Kristó ST, Kéry T. The *in vitro* effect of dandelion antioxidants on microsomal lipid peroxidation. *Phytotherapy Research* 2000, 14(1):43-44. doi:10.1002/(SICI)1099-1573(200002)14:13.3.CO;2-H.
 20. Hu C, Kitts DD. Dandelion (*Taraxacum officinale*) flower extract suppresses both reactive oxygen species and nitric oxide and prevents lipid oxidation *in vitro*. *Phytomedicine* 2005; 12(8):588-597. doi.org/10.1016/j.phymed.2003.12.012.
 21. Williams CA, Goldstone F, Greenham J. Flavonoids, cinnamic acids and coumarins from the different tissues and medicinal preparations of *Taraxacum officinale*. *Phytochemistry* 1996; 42(1):121-7. doi:10.1016/0031-9422(95)00865-9.
 22. Pandey KB, Rizvi SI. Plant polyphenols as dietary antioxidants in human health and disease. *Oxidative Medicine and Cellular Longevity* 2009; 2(5):270-8. doi: 10.4161/oxim.2.5.9498.
 23. Lima CF, Valentao PC, Andrade PB, Seabra RM, Fernandes-Ferreira M, Pereira-Wilson C. Water and methanolic extracts of *Salvia officinalis* protect HepG2 cells from *t*-BHP induced oxidative damage. *Chemico-Biological Interactions*, 2007; 167(2):107-115. doi: 10.1016/j.cbi.2007.01.020.
 24. Valentão P, Fernandes E, Carvalho F, Andrade PB, Seabra RM, Bastos ML. Antioxidant activity of *Centaureum erythraea* infusion evidenced by its superoxide radical scavenging and xanthine oxidase inhibitory activity. *Journal of Agricultural and Food Chemistry* 2001; 49(7): 3476-3479. doi: 10.1021/jf001145s.
 25. Russo A, Cardile V, Lombardo L, Vanella L, Vanella A, Garbarino JA. Antioxidant activity and antiproliferative action of methanolic extract of *Geum quellyon* Sweet roots in human tumor cell lines. *Journal of Ethnopharmacology*, 2005; 100(3):323-332. doi: 10.1016/j.jep.2005.03.032.
 26. Lima CF, Pereira-Wilson C, Rattan SI. Curcumin induces heme oxygenase-1 in normal human skin fibroblasts through redox signaling: relevance for anti-aging intervention. *Molecular Nutrition & Food Research* 2011; 55(3):430-442. doi: 10.1002/mnfr.201000221.
 27. Peschel W, Sánchez-Rabaneda F, Diekmann W, Plescher A, Gartzia I, Jiménez D. An industrial approach in the search of natural antioxidants from vegetable and fruit wastes. *Food Chemistry* 2006; 97(1):137-150. doi.org/10.1016/j.foodchem.2005.03.033.
 28. Barreira JC, Ferreira IC, Oliveira MB, Pereira JA. Antioxidant activity and bioactive compounds of ten Portuguese regional and commercial almond cultivars. *Food and Chemical Toxicology* 2008; 46(6):2230-2235. doi: 10.1016/j.fct.2008.02.024.
 29. Silva BA, Dias ACP, Ferreres F, Malva JO, Oliveira C. Neuroprotective Effect of *H. perforatum* Extracts on β -Amyloid-induced Neurotoxicity. *Neurotoxicity Research* 2004; 6(2): 119-130. doi: 10.1007/BF03033214.
 30. Brighente IMC, Dias M, Verdi LG, Pizzolatti MG. Antioxidant activity and total phenolic content of some brazilian species. *Pharmaceutical Biology* 2007; 45(2):156-61. doi: 10.1080/13880200601113131.
 31. Magalhaes LM, Segundo MA, Reis S, Lima JL. Methodological aspects about *in vitro* evaluation of antioxidant properties. *Analytica Chimica Acta* 2008; 613(1):1-19. doi: 10.1016/j.aca.2008.02.047.
 32. Ahmad B, Khan MR, Shah NA, Khan RA. *In vitro* antioxidant potential of *dicliptera roxburghiana*. *BMC Complementary and Alternative Medicine* 2013; 13:140. doi: 10.1186/1472-6882-13-140.
 33. Hazra B, Biswas B, Mandal N. Antioxidant and free radical scavenging activity of *Spondias pinnata*. *BMC Complementary and Alternative Medicine* 2008; 8:63. doi: 10.1186/1472-6882-8-63.
 34. Soares JR, Dinis TC, Cunha AP, Almeida LM. Antioxidant activities of some extracts of *Thymus zygis*. *Free Radical Research* 1997; 26(5):469-78. doi: 10.3109/10715769709084484.
 35. Cui Y, Kim DS, Park KC. Antioxidant effect of *Inonotus obliquus*. *Journal of Ethnopharmacology* 2005; 96(1-2):79-85. doi: 10.1016/j.jep.2004.08.037.
 36. Niu XL, Ichimori K, Yang X, Hirota Y, Hoshiai K, Li M, Nakazawa H. Tanshinone II-A inhibits low density lipoprotein oxidation *in vitro*. *Free Radical Research* 2000; 33(3):305-312. doi: 10.1080/1071576000301471.
 37. Heim KE, Tagliaferro AR, Bobilya DJ. Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. *Journal of Nutritional Biochemistry* 2002; 13(10):572-584. doi.org/10.1016/S0955-2863(02)00208-5.
 38. Budzianowski J. Coumarins, caffeoyltartaric acids and their artefactual methyl esters from *Taraxacum officinale* leaves. *Planta Medica* 1997; 63(3):288. doi: 10.1055/s-2006-957681.
 39. Dalby-Brown L, Barsett H, Landbo AK, Meyer AS, Mølgaard P. Synergistic antioxidative effects of alkamides, caffeic acid derivatives, and polysaccharide fractions from *Echinacea purpurea* on *in vitro* oxidation of human low-density lipoproteins. *Journal of Agricultural and Food Chemistry* 2005; 53(24):9413-9423. doi: 10.1021/jf0502395.
 40. Xi J, Shen D, Zhao S, Lu B, Li Y, Zhang R. Characterization of polyphenols from green tea leaves using a high hydrostatic pressure extraction. *International Journal of Pharmaceutics* 2009; 382(1-2):139-143. doi: 10.1016/j.ijpharm.2009.08.023.

41. Sas K, Robotka H, Toldi J, Vécsei L. Mitochondria, metabolic disturbances, oxidative stress and the kynurenine system, with focus on neurodegenerative disorders. *Journal of the Neurological Sciences* 2007; 257(1-2):221-239. doi: 10.1016/j.jns.2007.01.033.
42. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity tests. *Journal of Immunological Methods* 1983; 65(1-2):55-63. doi:10.1016/0022-1759(83)90303-4.
43. Knowles BB, Howe CC, Aden DP. Human hepatocellular carcinoma cell lines secrete the major plasma proteins and hepatitis B surface antigen. *Science* 1980; 209(4455):497-499. doi: 10.1126/science.6248960.
44. Diamond L, Kruszewski F, Aden DP, Knowles BB, Baird WM. Metabolic activation of benzo[a]pyrene by a human hepatoma cell line. *Carcinogenesis* 1980; 1(10):871-875. doi: 10.1093/carcin/1.10.871.
45. Sassa S, Sugita O, Galbraith RA, Kappas A. Drug metabolism by the human hepatoma cell HepG2. *Biochemical and Biophysical Research Communications* 1987; 143(1):52-57. doi:10.1016/0006-291X(87)90628-0.
46. Schutz K, Carle R, Schieber A. *Taraxacum* - a review on its phytochemical and pharmacological profile. *Journal of Ethnopharmacology* 2006; 107(3):313-323. doi.org/10.1016/j.jep.2006.07.021.
47. Lambert JD, Hong J, Yang GY, Liao J, Yang CS. Inhibition of carcinogenesis by polyphenols: evidence from laboratory investigations. *The American Journal of Clinical Nutrition* 2005; 81(1):284S-291S.
48. Ivanov IG. Polyphenols content and antioxidant activities of *Taraxacum officinale* F. H. Wigg (Dandelion) Leaves. *International Journal of Pharmacognosy and Phytochemical Research* 2014; 6(4):889-93.
49. Charvat TT, Lee DJ, Robinson WE, Chamberlin AR. Design, synthesis, and biological evaluation of chicoric acid analogs as Inhibitors of HIV-1 integrase. *Bioorganic & Medicinal Chemistry* 2006; 14(13):4552-4567. doi: 10.1016/j.bmc.2006.02.030.
50. Tsai YL, Chiu CC, Yi-Fu Chen J, Chan KC, Lin SD. Cytotoxic effects of *Echinacea purpurea* flower extracts and cichoric acid on human colon cancer cells through induction of apoptosis. *Journal of Ethnopharmacology* 2012; 143(3):914-919. doi: 10.1016/j.jep.2012.08.032.
51. Yarnell E, Abascal K. Dandelion (*Taraxacum officinale* and *T mongolicum*). *Integrative Medicine* 2009; 8(2):35-38.
52. Ovadje P, Chatterjee S, Griffin C, Tran C, Hamm C, Pandey S. Selective induction of apoptosis through activation of caspase-8 in human leukemia cells (Jurkat) by dandelion root extract. *Journal of Ethnopharmacology* 2011; 133(1):86-91. doi:10.1016/j.jep.2010.09.005.
53. Colle D, Arantes LP, Gubert P, da Luz SC, Athayde ML, Teixeira Rocha JB, Soares FA. Antioxidant properties of *Taraxacum officinale* leaf extract are involved in the protective effect against hepatotoxicity induced by acetaminophen in mice. *Journal of Medicinal Food* 2012; 15(6):549-556. doi: 10.1089/jmf.2011.0282.