

Central Washington University
ScholarWorks@CWU

All Faculty Scholarship for the College of the
Sciences

College of the Sciences

1-19-2021

Excessive intake of iodine and low prevalence of goiter in school age children five years after implementation of national salt iodization in Shebedino woreda, southern Ethiopia


Elilta Elias
Hawassa University

Workneh Tsegaya
Oklahoma State University

Barbara J. Stoecker
Oklahoma State University

Tafere Gebreegziabher Belay
Central Washington University, tafere.bl@gmail.com

Follow this and additional works at: <https://digitalcommons.cwu.edu/cotsfac>

 Part of the [Endocrinology, Diabetes, and Metabolism Commons](#), [Human and Clinical Nutrition Commons](#), and the [Nutritional Epidemiology Commons](#)

Recommended Citation

Elias, E., Tsegaye, W., Stoecker, B. J., & Gebreegziabher, T. (2021). Excessive intake of iodine and low prevalence of goiter in school age children five years after implementation of national salt iodization in Shebedino woreda, southern Ethiopia. *BMC Public Health*, 21(1). <https://doi.org/10.1186/s12889-021-10215-y>

This Article is brought to you for free and open access by the College of the Sciences at ScholarWorks@CWU. It has been accepted for inclusion in All Faculty Scholarship for the College of the Sciences by an authorized administrator of ScholarWorks@CWU. For more information, please contact scholarworks@cwu.edu.

RESEARCH ARTICLE

Open Access



Excessive intake of iodine and low prevalence of goiter in school age children five years after implementation of national salt iodization in Shebedino woreda, southern Ethiopia

Elilta Elias¹, Workneh Tsegaye², Barbara J. Stoecker² and Tafere Gebreegziabher^{1,3*} 

Abstract

Background: Iodine is a trace element required for the synthesis of thyroid hormones. The multiple effects of iodine deficiency on human health are called iodine deficiency disorders (IDDs). IDDs have been common nutritional problems in Ethiopia. In 2012, Ethiopia launched a national salt iodization program to address IDDs. The objective of this study was to assess the effects of this program after 5 years by measuring urinary iodine concentration (UIC) and prevalence of goiter in school age children as well as household salt iodine concentration (SIC).

Methods: A school-based cross-sectional design was employed. After ethical approval, 408 children from eight randomly selected primary schools provided urine samples. UIC was analyzed by inductively coupled plasma mass spectrophotometry (ICP-MS). A 10 g salt sample was collected from each household of a sampled child. SIC was analyzed with a digital electronic iodine checker (WYD, UNICEF) and goiter was assessed by palpation.

Results: The mean (\pm SD) age of the children was 9 ± 2 years. The prevalence of goiter was 4.2% and no child had grade 2 goiter. The median (IQR) UIC was 518 (327, 704) μ g/L and UIC ranged from 3.1 to 2530 μ g/L. Of the salt samples, 15.6% were not adequately iodized (< 15 ppm), 39.3% were adequately iodized (≥ 15 to ≤ 40 ppm), and 45.1% were > 40 ppm. SIC ranged from 4.2 to 195 ppm. Of the mothers, 92% said iodized salt prevents goiter and 8% mentioned prevents mental retardation.

Conclusions: In 2017 iodine deficiency was no longer a public health problem in the study area. However, the high variability in UIC and SIC and excessive iodine intake are of great concern. It is vital to ensure that salt is homogeneously iodized at the production site before being distributed to consumers.

Keywords: Excessive iodine, Iodization, Goiter, School children, Southern Ethiopia

* Correspondence: Tafere.bl@gmail.com

¹School of Nutrition, Food Science and Technology, Hawassa University, P.O. Box 5, SNNPR, Hawassa, Ethiopia

³Department of Health Sciences, Central Washington University, 400 E University way, Ellensburg, WA 98926, USA

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Iodine is an essential trace element for human growth and development and is required for the synthesis of thyroid hormones [1]. Low levels of thyroid hormones result in hypothyroidism which could cause serious functional and developmental disorders collectively called iodine deficiency disorders (IDDs) [2]. These can have immediate negative effects on children's learning capacity and school performance. If the trend continues, IDDs can lead to lower economic productivity and poor quality of life [2].

Currently, the preferred way to alleviate iodine deficiency is to increase iodine intake through implementation of universal salt iodization (USI). USI is the iodization of all salt in the country for human and livestock consumption [3]. However, despite national and international efforts to increase iodine intake, primarily through salt iodization programs, about 35% of the world's population and nearly 30% of school-age children (241 million) were estimated in 2014 to have insufficient iodine intake. The majority (134 million) of these lived in South-East Asia and Africa [4, 5]. In 2017, iodine deficiency was still considered a global public health problem [6].

Although USI is the best strategy to alleviate IDDs, it requires a close monitoring system and may take time until there are changes in the prevalence of signs of iodine deficiency, particularly goiter. According to the Ethiopian Public Health Institute, in 2015, 3 years after launch of mandatory iodization of salt for humans, 47.5% of Ethiopian children had low UIC. The problem was worse in the southern parts of the nation where the percentage of school age children with urinary iodine of less than 100 µg/L was as high as 56.7%, and iodine deficiency remained a severe public health problem [7]. However, 5 years after the iodization program was launched, 29% of school children in North-West Ethiopia had excessive iodine intake, 62% had adequate iodine intake and the median urinary iodine concentration (UIC) was 235 µg/L. Prevalence of goiter was a severe public health problem as 34% of the school children had palpable or visible goiter [8].

Although there are several studies from Ethiopia before the salt iodization program, there is a lack of information about iodine status of school children following the iodization program, specifically those in the study area. This study was designed to assess iodine status and prevalence of goiter among school age (6–12 year old) children in Shebedino *woreda*, Sidama Zone, Southern Ethiopia 5 years after the national salt iodization program.

Methods and materials

Study design

This study employed a school-based cross-sectional design. Data were collected from May 10 to June 11, 2017.

Description of study area and study population

The study was conducted in Shebedino *woreda* (district), Sidama Zone, Southern Nations, Nationalities, and Peoples Region (SNNPR). Ninety-five percent of the population depend on mixed farming, 3.8% depend on trade and only 1.2% work for either governmental or non-governmental organizations (NGOs). All students in randomly selected primary schools of Shebedino *woreda* were the study population.

Inclusion criteria

All children (6–12 years old) whose parents/guardians gave consent for their children to be part of the study were eligible to be included. No child included in the study was more than 12 years old.

Exclusion criteria

Based on the exclusion criteria, children to be excluded were those who had attended the targeted schools for less than 6 months prior to the study and those who were chronically or acutely ill at the time of data collection. Because there were no students in these categories, no students were excluded from the study.

Sample size

A single proportion sample size calculation formula was used to determine optimal number for estimating the prevalence of iodine deficiency. A sample size of 408 school age children was computed based on an estimated 59.1% prevalence of goiter [9], a 95% confidence level, 5% margin of error, design effect of 1.5% and non-response rate of 10%.

Sampling technique

The *woreda* has 37 primary schools, out of which eight schools were selected by simple random sampling. The sample was allocated for each school by population proportional to size. The sampling frame was prepared by using registers that listed children 6–12 years of age in all selected schools. Finally, systematic random sampling was used to select the study participants from each school.

Data collection method

Mothers were interviewed individually in their homes using a pretested structured questionnaire. Questions about socio-demographic and economic data were adapted from the Ethiopia Demographic and Health Survey (EDHS) questionnaire [10], and food consumption patterns were assessed using a standardized food frequency questionnaire [11]. The questionnaire was prepared in English, translated into the local language (Sidamigna) and translated back into English before data collection to check for consistency and to keep its

contextual equivalence. Data were collected by four trained health professionals fluent in the local language. A 10 g salt sample was collected in a tightly sealed plastic container from each household of a sampled child during interview with the mother. The plastic containers were labeled with school code and children's identification number. The salt samples were stored in a dry place until analyzed for iodine concentration.

Each child provided a urine sample at school in disposable plastic cups, and urine was transferred to screw-capped plastic vials. The vials were labeled with the school code and children's identification numbers using both permanent marker and pencil. Plastic vials were sealed with Parafilm® to prevent leakage and evaporation. The urine samples were immediately placed in a thermo cool box and were transported to Hawassa University and stored at -20 °C until shipped frozen to the USA for analysis by ICP-MS.

Examination for goiter

Children were examined for goiter at school by a single public health officer experienced in palpation. Stage of goiter was based on WHO criteria: grade 0, no palpable or visible goiter; grade 1, palpable goiter but not visible when neck is in the normal position; grade 2, visible goiter when neck is in the normal position [2].

Laboratory analysis

Urinary iodine concentration was analyzed by inductively coupled plasma mass spectrometry (ICP-MS, Elan 9000, Perkin Elmer, Norwalk, CT) at Oklahoma State University, USA. Urine was diluted in 2% ammonium hydroxide and iodine was analyzed with tellurium as internal standard (Caldwell et al., 2003). Quality control samples for urinary iodine were measured every 10 samples to ensure stability of the instrument.

Salt iodine concentration (SIC) was analyzed with a digital electronic iodine checker (WYD, UNICEF) at Hawassa University, Ethiopia. All procedures were performed according to the instructions provided in the WYD manual. From a standard KIO₃ solution containing 1000 µg/mL of iodine, 5 ml was diluted to 500 mL to make a 10 µg/mL working standard solution. A calibration solution was then made by combining 5 mL of the working standard solution, 2 mL of KI-starch solution, 2 mL (1 mol/L) sulfuric acid and further diluting with distilled water to a total of 50 mL and shaking well. After calibrating a zero reading on the WYD instrument using distilled water, the instrument's liquid crystal display (LCD) reading was calibrated to 50.0 ± 0.1 using the calibration solution. For sample measurement, 1 g of well-mixed salt was dissolved in 10 mL of distilled water and the same quantities of KI-starch and sulfuric acid solutions were added with additional distilled water to make

a total volume of 50 mL and mixed by shaking well. Salt solution samples were individually placed in the cell of the WYD iodine checker and iodine concentration was read on the LCD screen.

Statistical analysis

Data were analyzed using SPSS, version 23 (SPSS Statistics Version 23, IBM Corp., Armonk, NY, USA). Descriptive analysis was used to report characteristics of the study participants.

Mean (SD) and median (IQR) were employed for descriptive statistics. Normality for distribution of continuous data was checked using a histogram with line curve for a visual test as well as the one sample Kolmogorov-Smirnov test. Natural log transformation was used to transform positively skewed data. Independent-sample t-test was used to compare means of UIC between male and female children.

Results

A total of 408 mother-child dyads participated in this study. The mean age (±SD) of the mothers was 35 ± 8 years and of the children was 9 ± 2 years (Table 1). Most of the women had no formal education.

Table 1 Socio-demographic characteristics of school age children and their mothers in Shebedino woreda, southern Ethiopia, 2017 (n = 408)

Variables	Percent
Sex of child	
- Female	50.5
- Male	49.5
Age of child (years)	
- 6–8	42.2
- 9–12	57.8
Family size	
- 2–4	21.3
- 5–7	62.9
- ≥ 8	15.8
Age of mothers (years)	
- 21–28	15.7
- 29–35	49.0
- 36–44	21.6
- ≥ 45	13.7
Education of mothers	
- No formal education	52.5
- Able to read and write	2.5
- Primary school	40.6
- Middle or high school	4.4

As indicated in Table 2 most of the children consumed corn and enset at least once a day, followed in frequency by kale and haricot bean. More than 95% of the children had never consumed sweet potato, cassava, tomato, papaya, meat, chicken, egg, fish or cheese in the last month. A majority of the children consumed potato or avocado at least once a month. Milk, one of the potential good sources of dietary iodine, was consumed more frequently.

The majority of the mothers said they had heard about iodine, iodized salt or IDD before and their primary source of information was health workers. More than 90% of the respondents said they knew about the benefits of iodized salt; however 92% of those said it prevents goiter but only 8% related lack of iodine to mental retardation (Table 3).

As shown in Fig. 1, prevalence of goiter in school children was very low. Only 12 female and 5 male

children had grade 1 goiter; the rest did not have goiter. This was consistent with their UIC. The median (IQR) for UIC of children was 518 (327, 704) µg/L. Mean (SD) UIC of male children was 581 (311) µg/L and female children was 518 (310) µg/L and this was significantly different ($p = 0.043$). Fewer than 5% of children had UIC less than 100 µg/L. More concerning was that even the 25th percentile for UIC was greater than 300 µg/L (Fig. 2).

Most salt from households contained ≥ 15 ppm iodine. Among the 408 samples collected from each household, none were non-iodized. Of the collected samples 15.6% were inadequately iodized (< 15 ppm), 39.3% were adequately iodized (15 ppm – 40 ppm) and 45.1% had more than 40 ppm iodine (Fig. 3). The median (IQR) iodine concentration of the salt was 37 (22, 55) ppm and the range was 4.2 to 195 ppm.

Table 2 Food consumption patterns of school age children in Shebedino woreda, Southern Ethiopia, 2017 ($n = 408$)

Food types	Frequency (%)			
	At least once/day	3-6x/week	At least once/month	Never
Cereals				
- Teff	0	3 (0.7)	23 (5.7)	382 (93.6)
- Wheat	4 (1)	3 (0.7)	39 (9.6)	362 (88.7)
- Corn	387 (94.9)	14 (3.4)	2 (0.5)	5 (1.2)
Legumes				
- Haricot bean	107 (26.2)	95 (23.2)	176 (43.2)	30 (7.4)
- Bean	4 (1)	26 (6.4)	77 (18.9)	301 (73.7)
Root crops & vegetables				
- Potato	11 (2.7)	71 (17.4)	242 (59.3)	84 (20.5)
- Sweet potato	0	3 (0.7)	13 (3.1)	392 (96.1)
- Cassava	3 (0.7)	3 (0.7)	14 (3.4)	388 (95.1)
- Enset	352 (86.3)	46 (11.2)	7 (1.7)	3 (0.7)
- Tomato	1 (0.2)	2 (0.4)	16 (3.9)	389 (95.3)
- Kale	202 (49.5)	184 (45.1)	18 (4.4)	4 (1)
Fruits				
- Avocado	12 (3)	34 (8.3)	270 (66.2)	92 (22.5)
- Papaya	0	0	16 (3.9)	392 (96)
- Banana	6 (1.4)	15 (3.7)	161 (39.5)	225 (55.1)
Meat, egg or fish				
- Meat (beef, sheep or goat)	0	0	16 (3.9)	392 (96.1)
- Chicken	0	0	5 (1.2)	403 (98.8)
- Egg	0	1 (0.2)	11 (2.7)	396 (97.1)
- Fish	0	0	3 (0.7)	405 (99.3)
Dairy products				
- Milk	14 (3.4)	17 (4.2)	239 (58.6)	138 (33.8)
- Cheese	3 (0.7)	0	10 (2.4)	395 (96.8)
- Butter-milk	24 (5.9)	48 (11.7)	259 (63.5)	77 (18.9)

Table 3 Knowledge of mothers about iodized salt and IDD in Shebedino woreda, southern Ethiopia, 2017 (n = 408)

	Percent
Heard about iodine or IDD	
- Yes	75.2
- No	24.8
Heard about iodized salt	
- Yes	91.7
- No	8.3
Source of information	
- Health worker	60
- Mass media	27.2
- Public announcement	12.8
Know benefit of iodized salt	
- Yes	90.7
- No	9.3
Benefits of iodized salt	
- Prevents goiter	92
- Prevents mental retardation	8

Discussion

Iodine deficiency had been a serious problem throughout Ethiopia, including the study region, for many years. Repeated studies conducted before the launch of the national salt iodization program in the country, particularly in SNNPR, showed that iodine deficiency was a serious public health problem [12–15]. In order to alleviate this long-standing problem, the national salt iodization program was launched in 2012 and iodized salt became available as a source of iodine in the study region [15, 16].

The current study showed high urinary iodine concentration (UIC) and low prevalence of goiter following national salt iodization. Prevalence of goiter in school age children in the study region in 2007 was 56.2% [12], and in the current study it was only 4.2% which indicates successful elimination of iodine deficiency as a major public health problem according to WHO/UNICEF/ICCIDD criteria [2]. Similarly, in a nine year prospective study in Iran, total goiter rate reportedly decreased from 44% in 1996 to 7.6% in 2001 and to 0.4% in 2007 following their national salt iodization program [17].

According to the WHO/UNICEF/ICCIDD recommended criteria, a population without iodine deficiency should have median UIC between 100 µg/L and 299 µg/L and UIC should not be lower than 50 µg/L in more than 20% of subjects [18]. In the current study the median UIC was 518 µg/L; more than 95% of the children had UIC above 100 µg/L and only 4.3% had UIC < 100 µg/L indicating that iodine deficiency has been eliminated for the population. However, 78% of the children had UIC > 300 µg/L which reflects a risk for excessive iodine intake and could be a cause for adverse health consequences including iodine-induced hyperthyroidism and autoimmune thyroid diseases [19]. Because ours was a cross-sectional study, we do not know how long the children had been consuming household salt iodized at > 40 ppm. Risk of adverse effects of excessive iodine intake is expected 5 to 10 years following introduction of iodized salt, and the consequences of excessive iodine intake are worse in places where iodine deficiency previously existed [2, 20, 21]. Risk of excessive iodine intake and possible adverse effects have been reported in Ghana following their universal salt iodization





Fig. 1 Prevalence of goiter in school age children in Shebedino woreda, southern Ethiopia, 2017 (n = 408)




program [22], and a study in China reported thyroid nodules in response to excess iodine intake [23].

The large variation in UIC observed in this study is of great concern. UIC ranged from 3.1 µg/L to 2530 µg/L. This could be due to the large variation observed in SIC, which varied from 4.2 to 195 ppm (mg/kg). Because salt had been iodized using knapsack sprayers [16], it is unlikely to be iodized homogeneously which could be a reason for high variability in amounts of iodine in salt. Although not as high as the current study, school age children in northwest Ethiopia in 2016 had median UIC of 235 (161, 320) µg/L and 29.1% of the children had UIC that suggested excessive iodine intake [8]. In an urban area of central Italy, high variation of UIC (33.2–

819.5 µg/L) was observed in children who used iodized salt [24]. Due to the potential health risks, iodine intake that produces UIC > 300 µg/L per day should be discouraged [2, 25]. Generally in the current study more than 45% of the salt samples were above 40 ppm, an indication of more than adequate iodization [2].

The reduced prevalence of iodine deficiency in the study area is promising. However, iodine deficiency was still a public health problem in some parts of the country [26–28]. A study conducted in the Robe district in 2015 showed that 57% of the participants had low UIC and the median UIC was 78 µg/L [29]. In another study, nearly 65% of the households in Southern Wollo reported inadequate use of iodized salt 6 years after the



launch of the national salt iodization program [30]. Additionally adequate iodine intake in school-aged children doesn't assure adequate intake in at risk populations even in a country with a salt iodization program. In Sweden for instance, school age children had adequate iodine intake while pregnant women had inadequate intakes [31].

The ultimate goal of a universal salt iodization program is to alleviate iodine deficiency. However, the process must be closely monitored to prevent both inadequate and excessive iodine intakes [32]. The marked regional variation in Ethiopia implies the need to study factors causing such disparities in availability of iodized salt at the household level.

The WHO recommends that 90% of households in a population should be able to obtain iodized salt with an iodine level > 15 ppm for the effective elimination of IDD [2]. In the current study, 84.3% of households used adequate or more than adequately iodized salt. Although the percentage of households is lower than the recommendation and despite the high variability, the use of iodized salt is a great achievement compared to the iodized salt coverage of only 3 to 5% of households shortly before implementation of the national program [33]. The concern is, however, that unless salt iodization programs are practiced as industry norms, government-funded projects are not sustainable. For example, in Vietnam a government-funded universal salt iodization program was not sustainable as household iodized salt coverage declined from 90% in 2005 to 45% in 2011 [34]. In Tunisia, only 50% of households used adequately iodized salt after two decades of salt iodization which shows the program failed to meet the goals of USI, that is > 90% of households should use adequately iodized salt [35].

Maternal knowledge and practices are key factors for addressing public health problems. Low level of knowledge was associated with increased prevalence of IDD and has been identified as a major constraint in eliminating IDD in some African countries [36, 37]. In the current study, most of the women had heard about IDD, iodine and iodized salt. This is a remarkable improvement compared to previously reported studies in the region [13, 38]. Handling and storage conditions of iodized salt could also affect its iodine content. According to WHO, 20% of iodine from salt may be lost between production and the household, and another 20% of the iodine could be lost during cooking [2, 38]. However, in this study more than 97% of the respondents said they stored their salt in a dry place with a covered container which helps retain iodine in salt.

Because schools and children were randomly selected for this study, we are confident that our sample was representative of the study community and local region.

However, Ethiopia is a large country with many agroecological zones. Because soil types and water sources contribute to diets of subsistence farming households [39], a study in a single region cannot be nationally representative. Furthermore, the current study did not analyze iodine concentration of consumed foods which was a limitation.

Conclusions

In conclusion, although prevalence of goiter was low, based on the median UIC the school children could be assumed to be at risk of excessive iodine consumption. The salt iodization process needs to be monitored strictly before the population suffers detrimental health effects from excessive intake. Nonetheless, since this analysis was done on spot UIC samples, repeated measures at different times and in different population groups are warranted.

Abbreviations

UIC: Urinary iodine concentration; SIC: Salt iodine concentration; IDD: Iodine deficiency disorders; ICP-MS: Inductively coupled plasma mass spectrometry; IQR: Interquartile range; USI: Universal salt iodization; WHO: World Health Organization; SNNPR: Southern Nations, Nationalities and Peoples Region; NGO's: Non-governmental organizations; EDHS: Ethiopia Demographic and Health Survey

Acknowledgements

We thank the study participants who took part in this study and all data collectors. Some of the data in this paper were included in a poster available at the virtual Micronutrient Forum held in November, 2020. "Belay GT, Elias E, Tsegaye W, Stoecker BJ. Increased urinary iodine concentration and decline in goiter prevalence in school children in southern Ethiopia six years after launch of the national salt iodization program". Micronutrient Forum 5th global conference, 2020. (<https://www.eventscribe.com/2020/MNF-CONNECTED/fsPopup.asp?PosterID=310580+&efp=QU9WU1BJREMxMzcwMQ+&mode=posterinfo&rnd=0.0558427>).

Authors' contributions

EE and TG, conception of the research, study design, data analysis, and drafting the manuscript. BJS was involved in analysis, revising the study design and the manuscript. WT was involved in lab analysis and revising the manuscript. All authors have read and approved the final manuscript.

Funding

The research was funded by Hawassa University, Ethiopia and the USDA Multistate project, W3002. The funders had no role in the design of the study and collection, analysis, and interpretation of data or in writing the manuscript.

Availability of data and materials

The dataset used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

Ethical approval was obtained from the review boards for Hawassa University and the Ministry of Science and Technology, Ethiopia. Written informed consent was obtained from the mothers.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹School of Nutrition, Food Science and Technology, Hawassa University, P.O. Box 5, SNNPR, Hawassa, Ethiopia. ²Department of Nutritional Sciences, Oklahoma State University, 421 Human Sciences, Stillwater, OK 74078, USA. ³Department of Health Sciences, Central Washington University, 400 E University way, Ellensburg, WA 98926, USA.

Received: 4 June 2020 Accepted: 11 January 2021

Published online: 19 January 2021

References

- Zimmermann MB. The role of iodine in human growth and development. *Semin Cell Dev Biol.* 2011;22(6):645–52.
- WHO/UNICEF/ICCIDD. Assessment of iodine deficiency disorders and monitoring their elimination. a guide for program managers. Third edition. World Health Organization 2007.
- WHO. Iodine status worldwide. WHO global database on iodine deficiency. Geneva: WHO; 2004. p. 1–58. (www.who.int/publication/2004/9241592001.pdf)
- WHO. Urinary iodine concentrations for determining iodine status deficiency in populations. Vitamin and mineral nutrition information system. Geneva: World Health Organization; 2013. (<http://www.who.int/nutrition/vmnis/indicators/urinaryiodine>, accessed May 05, 2019)
- WHO. Guideline: fortification of food-grade salt with iodine for the prevention and control of iodine deficiency disorders. Geneva: World Health Organization; 2014.
- Biban BG, Lichiardopol C. Iodine deficiency, still a global problem? *Curr Health Sci J.* 2017;43(2):103–11.
- Ethiopian Public Health Institute: Ethiopian national micronutrient survey report. Addis Ababa 2016.
- Wassie MM, Abebe Z, Tariku A, Gebeye E, Awoke T, Gete AA, Yesuf ME, Kebede Y, Biks GA, Zhou SJ. Iodine status five years after the mandatory salt iodization legislation indicates above requirement: a cross sectional study in Northwest Ethiopia. *BMC Nutr.* 2018;4(1):52.
- Mezgebu Y, Mossie A, Rajesh P, Beyene G. Prevalence and severity of iodine deficiency disorder among children 6–12 years of age in Shebe Senbo district, Jimma zone, southwest Ethiopia. *Ethiop J Health Sci.* 2012;22:196–204.
- CSA and ICF. Ethiopian Demographic and Health Survey. 2016. Addis Ababa and Rockville: CSA and ICF; 2016.
- Gibson RS. Principles of nutritional assessment, second ed: Oxford University press; 2005.
- Abuye C, Berhane Y, Akalu G, Getahun Z, Ersumo T. Prevalence of goiter in children 6 to 12 years of age in Ethiopia. *Food Nutr Bull.* 2007;28:391–7.
- Bogale A, Abebe Y, Stoecker BJ, Abuye C, Ketema K, Hambidge KM. Iodine status and cognitive function of women and their five year-old children in rural Sidama, southern Ethiopia. *East Afr J Public Health.* 2009;6:296–9.
- Ersino G, Tadele H, Bogale A, Abuye C, Stoecker BJ. Clinical assessment of goiter and low urinary iodine concentration depict presence of severe iodine deficiency in pregnant Ethiopian women: a cross-sectional study in rural Sidama, southern Ethiopia. *Ethiop Med J.* 2013;51(2):133–41.
- Gebreegziabher T, Teyike N, Mulugeta A, Abebe Y, Hambidge KM, Stoecker BJ. Lack of dietary sources of iodine and the prevalence of iodine deficiency in rural women from Sidama zone, southern Ethiopia. *Afr J Food Agric Nutr Dev.* 2013;13:8401–14.
- Ethiopian Public Health Institute: National salt iodization coverage towards prevention of iodine deficiency disorders in Ethiopia. 2014.
- Nouri Saeidlou S, Babaei F, Ayremlou P, Entezarmahdi R. Has iodized salt reduced iodine-deficiency disorders among school-aged children in north-West Iran? A 9-year prospective study. *Public Health Nutr.* 2018;21(3):489–96.
- UNICEF. Guidance on the monitoring of salt iodization programs and determination of population iodine status. UNICEF, 2018.
- Farebrother J, Zimmermann MB, Andersson M. Excess iodine intake: sources, assessment, and effects on thyroid function. *Ann N Y Acad Sci.* 2019;1446(1):44–65.
- Stanbury JB, Ermans AE, Bourdoux P, Todd C, Oken E, Tongle R, Vidor G, Braverman EL, Medeiros-Neto G. Iodine-induced hyperthyroidism: occurrence and epidemiology. *Thyroid.* 1998;8:83–100.
- Todd CH, Allain T, Gomo ZA, Hasler JA, Ndiweni M, Oken E. Increase in thyrotoxicosis associated with iodine supplements in Zimbabwe. *Lancet.* 1995;346(8989):1563–4.
- Abu BAZ, Oldewage-Theron W. Risks of excess iodine intake in Ghana: current situation, challenges, and lessons for the future. *Ann N Y Acad Sci.* 2019;1446(1):117–38.
- Du Y, Gao Y, Meng F, Liu S, Fan Z, Wu J, Sun D. Iodine deficiency and excess coexist in China and induce thyroid dysfunction and disease: a cross-sectional study. *PLoS One.* 2014;9(11):e111937.
- Coccaro C, Tuccilli C, Prinzi N, D'Armiento E, Pepe M, Del Maestro F, Cacciola G, Forlini B, Verdolotti S, Bononi M, et al. Consumption of iodized salt may not represent a reliable indicator of iodine adequacy: evidence from a cross-sectional study on schoolchildren living in an urban area of Central Italy. *Nutrition.* 2016;32(6):662–6.
- Farebrother J, Zimmermann MB, Abdallah F, Assey V, Fingerhut R, Gichohi-Wainaina WN, Hussein I, Makokha A, Sagno K, Untoro J, et al. Effect of excess iodine intake from iodized salt and/or groundwater iodine on thyroid function in nonpregnant and pregnant women, infants, and children: a multicenter study in East Africa. *Thyroid.* 2018;28(9):1198–210.
- Dessie G, Amare D, Dagne AB, Mulugeta H, Haile Kassa D, Negesse A, Kassa GM, Wagne F, Islam SMS, Burrows S. Prevalence of goiter among children in Ethiopia and associated factors: a systematic review and meta-analysis. *BMC Public Health.* 2019;19(1):1191.
- Bekele A, Adilo TM. Prevalence of goiter and its associated factors among primary school children in Chole District, Arsi zone, Ethiopia: a cross-sectional study. *BMC Nutr.* 2019;5:5.
- Hassen HY, Beyene M, Ali JH. Dietary pattern and its association with iodine deficiency among school children in Southwest Ethiopia; a cross-sectional study. *PLoS One.* 2019;14(8):e0221106.
- Hailu S, Wubshet M, Woldie H, Tariku A. Iodine deficiency and associated factors among school children: a cross-sectional study in Ethiopia. *Arch Public Health.* 2016;74:46.
- Mekonnen TC, Eshete S, Wasihun Y, Arefaynie M, Cherie N. Availability of adequately iodized salt at household level in Dessie and Combolcha towns, south Wollo, Ethiopia. *BMC Public Health.* 2018;18(1):1152.
- Manousou S, Andersson M, Eggertsen R, Hunziker S, Hulthen L, Nystrom HF. Iodine deficiency in pregnant women in Sweden: a national cross-sectional study. *Eur J Nutr.* 2019;59:2535–45.
- Pearce EN, Andersson M, Zimmermann MB. Global iodine nutrition: where do we stand in 2013? *Thyroid.* 2013;23(5):523–8.
- Gebreegziabher T, Girma M. Challenges and progress towards universal salt iodization (USI) in Ethiopia. *FASEB J.* 2012;26(10):21.
- Codling K, Quang NV, Phong L, Phuong do H, Quang ND, Begin F, Mathisen R. The rise and fall of universal salt iodization in Vietnam: lessons learned for designing sustainable food fortification programs with a public health impact. *Food Nutr Bull.* 2015;36(4):441–54.
- Doggui R, El Ati-Hellal M, Traissac P, Lahmar L, El Ati J. Adequacy assessment of a universal salt iodization program two decades after its implementation: a national cross-sectional study of iodine status among school-age children in Tunisia. *Nutrients.* 2016;9(1):6.
- Hessler J, Todd C, Mutamba J, Nyamandi T, Jumo D, Ropi F. Africa struggles for independence from IDD. *IDD News Letter.* 1997;13(2):17–27.
- Heidari Z, Arefhosseini SR, Hedayati M, Vaghef-Mehrabany E, Ebrahimi-Mameghani M. Iodine status, and knowledge about iodine deficiency disorders in adolescent school girls aged 14–19 years, 2016. *Health Promot Perspect.* 2019;9(1):77–84.
- Gebreegziabher T, Stoecker BJ. High variability of iodine in iodized salt and urine from rural households in Sidama zone, southern Ethiopia: a cross-sectional study. *Afr J Food Agric Nutr Dev.* 2020;20:15235–51.
- Watts MJ, Middleton DRS, Marriott AL. Source apportionment of micronutrients in the diets of Kilimanjaro, Tanzania and Counties of Western Kenya. *Sci Rep.* 2019;9(1):14447.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.