
GEOSUR: Mesozoic to Quaternary evolution of Tierra del Fuego and neighbouring austral regions II

M. MENICETTI^{|1|} and A. TASSONE^{|2|}

^{|1|} Istituto di Scienze della Terra. Università di Urbino.

Campus Scientifico Universitario. 61029 Urbino. Italy. E-mail: menichetti@uniurb.it

^{|2|} CONICET-Instituto de Geofísica "Daniel A. Valencio" (INGEODAV). Departamento de Geología.
Facultad de Ciencias Exactas y Naturales. Universidad de Buenos Aires

Ciudad Universitaria. Pabellón 2. 1428 Ciudad Autónoma de Buenos Aires. Argentina. E-mail: atassone@gl.fcen.uba.ar

This second thematic issue on Tierra del Fuego geology and geophysics, as the first one (Menichetti and Tassone, 2007) resulted from the new and profuse available information presented at the November 2004 **Geosur International Symposium** (Buenos Aires, Argentina). The aim of the organizing committee of **Geosur 2004** was to provide sets of synthetic papers supplying a comprehensive review of the geological and geophysical knowledge on the Tierra del Fuego region; and also more specific papers focused on subjects still under study and debate, including original contributions from Tierra del Fuego and some neighbouring areas of the Patagonian Andes and Scotia plate.

From the Chile trench to the transform boundaries of the western Scotia Sea, Tierra del Fuego Island shows a complex geological and structural architecture. Its geodynamic history includes a Mesozoic extensional phase in the southern Gondwanaland margin that was followed by the Late Cretaceous collisional events that led to the Andean Cordillera upbuilding and the Magallanes and Malvinas foreland basin generation. The opening of the Scotia Sea, with the separation from the Antarctic Peninsula and with superposition of important transform faults, influenced largely on the Cenozoic evolution of this orogenic segment. The last glacial events modelled and contributed to the present landscape.

The first thematic issue on Tierra del Fuego Mesozoic to Quaternary evolution contents contributions from a wide disciplinary scope (Geodesy, Paleomagnetism, Gravimetry, bathymetric studies and Geomorphology) by authors as Rappalini (2007), Bujalesky (2007), Chávez et al. (2007), Del Cogliano et al. (2007), Lodolo et al. (2007), Polonia and Torelli (2007). This second issue also presents contributions from a variety of disciplines including Petrology, Stratigraphy, Structural Geology and Seismostratigraphy.

Olivero and Malumián (this issue) provide an updated and exhaustive synthesis of the Mesozoic-Cenozoic Stratigraphy of the Fuegian Andes. Within the Cretaceous-Cenozoic record the authors recognize distinct sedimentary cycles in the northern cratonic areas of Tierra del Fuego and in the southern fold-and-thrust belt. In the cratonic areas four major sedimentary cycles evolved in a shallow marine platform setting mainly controlled by eustasy; whereas in the southern orogenic areas the propagation of the thrust front controlled the generation and migration of successive depocenters in the inner, central, and outer fold belt.

On the basis of new acquired and already published data Menichetti et al. (this issue) present a comprehensive synthesis of the structural geology of the Tierra del Fuego

Island. During the Mesozoic-Cenozoic evolution three complex tectonic stages are distinguished: extensional in the Upper Jurassic, compressional in the Late Cretaceous onward, and left lateral strike-slip from the Oligocene onwards, with the latter phase being superimposed and possibly reactivating the oldest structures. The structural characteristics and associated geodynamic events of each stage are exhaustively reviewed. At the same time, the authors detail a number of remaining topics related to Tierra del Fuego geology, which require further work.

The petrology, P-T evolution, geochronology and geodynamic meaning of the metamorphic complexes of the Patagonian and Fuegian Andes are presented in Hervè et al. (this issue). The authors distinguish metamorphic complexes located to the East of the Meso-Cenozoic South Patagonian Batholith and the so-called western complexes. The former correspond to the mostly low grade late Paleozoic Eastern Andes Metamorphic Complex (EAMC), which likely represents a passive margin metasedimentary sequence, later locally submitted to Late Jurassic high metamorphic conditions. The Cordillera Darwin Metamorphic Complex could be a higher grade equivalent of the EAMC. The western accretionary complexes point to the existence of a subduction zone environment near the Triassic - Jurassic boundary (Chonos Metamorphic Complex) up to the Middle Jurassic-Cretaceous (Diego de Almagro Complex) at least. The former comprises both allochthonous and autochthonous components, whereas the latter would postdate the southward drifting of Antarctic Peninsula.

On the basis of offshore multichannel seismic reflection profiles, combined with exploratory wells, the sedimentary architecture of the southern Malvinas foreland basin and the structural framework of its deformed inner and outer front are presented (Tassone et al., this issue). The basin infilling and the structure imprint record the main tectono-stratigraphic stages of the Fuegian Andes: the earliest Jurassic to Cretaceous extensional tectonics, the main compressional phase related to the Andean orogeny since Late Cretaceous, with the development of a thick-skinned tectonic style and the Mid-Tertiary to present influence of strike-slip activity, which has contributed to the subsidence of the foredeep. Restricted Neogene pull-apart basins formed by transcurrent motion occur superimposed to the Magallanes fold-and-thrust belt.

Rossello et al. (this issue) present a summary of the hydrocarbon exploration history and petroleum potential of Tierra del Fuego. Within the thick Late Cretaceous to Tertiary sedimentary and volcanoclastic sequence of the Austral-Magellan Basin, the sandy levels of the Springhill Formation host -in onshore as well as in offshore areas- productive hydrocarbon deposits.

Original information on the submerged stratigraphy and structure along the Magellan Strait, which bounds the Tierra del Fuego Island, is presented by Bartole et al. (this issue). On the basis of multichannel seismic profiles the authors distinguish three segments with specific sedimentary and tectonic architectures: the eastern Atlantic foreland domain characterized by nearly undeformed very thick Cretaceous to Tertiary sedimentary package, the central fold-and-thrust belt province with seismic evidence of deformations of the foreland units and the Cordillera province (Pacific segment) where the Magallanes-Fagnano transform fault exerts an important morpho-tectonic control that strongly conditions its bathymetric profile. Additionally, the seismic profiles also highlighted some depositional features of the glacial cycles in southernmost South America and present evidences of Miocene activity of the Magallanes-Fagnano fault in western Magellan Strait.

Finally, the paper by Waldmann et al. (this issue) includes the first seismic reflection profiles of Lago Fagnano that combined with sediment cores are presented in a preliminary succession of the depositional processes, which characterize the last deglaciation in Lago Fagnano. The authors recognize three major seismostratigraphic units: the oldest corresponds to glacially derived sediments; the overlying unit is interpreted as proglacial turbidite sequences that reflect sediment pulses released by the retreating Fagnano glacier during the last deglaciation. A major environmental change occurred during deposition of the uppermost unit when the pelagic style of sedimentation is interrupted by intercalated sequences of downslope mass flow events probably triggered by tectonic pulses along the Magallanes Fagnano system. The authors also provide important tools for the paleoclimate change analysis.

We would like to remind the readers that, as mentioned in the first Geosur thematic issue (Menichetti and Tassone, 2007), the terms Fuegian and Fueguian are synonymous and both are used as adjective meaning "of Tierra del Fuego" or "pertaining to Tierra del Fuego".

ACKNOWLEDGEMENTS

Most of the articles presented in this volume have resulted from researches carried out in the last decade under the sponsoring and the financial support of Antarctic national scientific programs jointly managed by Argentinean and Italian Institutions. The continuous scientific cooperation between Argentinean and Italian researchers was developed within the frame of several joint projects dealing with Tierra del Fuego geology and geophysics (PNRA - FORTE Project), onshore/offshore geophysical data acquisition (PNRA - TESAC Project) and the installation in

1994 of the broad-band seismological network in the Scotia Arc region, among other activities. We must express our deep gratitude to all the colleagues and friends that in many different ways helped us in Tierra del Fuego during these years. We particularly thank the Italian Embassy in Argentina for both the encouragement and the financial support in the organization of the Geosur 2004 International Symposium. The editors thank the following colleagues and friends who kindly helped with the reviewing of the papers submitted to this second *Geologica Acta* issue: M. E. Cerredo (Argentina), R. Coccioni (Italy), G. Fontolán (Italy), E. Lodolo (Italy), A. Massabie (Argentina), E. Oviedo (Argentina), E. Ramos (Spain), A. Rapalini (Argentina), N. Rubinstein (Argentina), F. Sàbat (Spain), J. Sellés Martínez (Argentina), B.L. Valero-Garcés (Spain), J. Vergés (Spain). Special acknowledgement is due to Lluís Cabrera and Ángeles Santos (Scientific and Managing Editors of *Geologica Acta*, respectively) and also the whole managing scientific committee of the journal for their constant support during the editorial process of this second thematic issue. Finally, thanks are also given to Prof. Miquel Canals who encouraged us to edit the sets of papers resulting from Geosur 2004 meeting.

REFERENCES

- Bartole, R., De Muro, S., Morelli, D., Tosoratti, F., 2008. Glacigenic features and Tertiary stratigraphy of the Magellan strait (Southern Chile). *Geologica Acta*, 6(1), 85-100.
- Bujalesky, G.G., 2007. Coastal geomorphology and evolution of Tierra del Fuego (Southern Argentina). *Geologica Acta*, 5(4), 337-362.
- Chávez, R.E., Flores-Márquez, E.L., Suriñach, E., Galindo-Zaldívar, J.G., Rodríguez-Fernández, J.R., Maldonado, A., 2007. Combined use of the GGSFT data base and on board marine collected data to model the Moho beneath the Powell Basin, Antarctica. *Geologica Acta*, 5(4), 323-335.
- Del Cogliano, D., Dietrich, R., Richter, A., Perdomo, R., Hormaechea, J.L., Liebsch, G., Fritsche, M., 2007. Regional geoid determination in Tierra del Fuego including GPS levelling. *Geologica Acta*, 5(4), 315-322.
- Hervé, F., Calderón, M., Faúndez, V., 2008. The metamorphic complexes of the Patagonian and Fuegian Andes. *Geologica Acta*, 6(1), 43-53.
- Lodolo, E., Lippai, H., Tassone, A., Zanolla, C., Menichetti, M., Hormaechea, J.L., 2007. Gravity map of the Isla Grande de Tierra del Fuego, and morphology of Lago Fagnano. *Geologica Acta*, 5(4), 307-314.
- Menichetti, M., Tassone, A. (eds.), 2007. GEOSUR 2004: Mesozoic to Quaternary evolution of Tierra del Fuego and neighbouring austral regions I. *Geologica Acta*, 5(4), 283-286.
- Menichetti, M., Lodolo, E., Tassone, A., 2008. Structural geology of the Fuegian Andes and Magallanes fold-and-thrust belt – Tierra del Fuego Island. *Geologica Acta*, 6(1), 19-42.
- Olivero, E.B., Malumián, N., 2008. Mesozoic-Cenozoic stratigraphy of the Fuegian Andes, Argentina. *Geologica Acta*, 6(1), 5-18.
- Polonia, A., Torelli, L., 2007. Antarctic/Scotia plate convergence off southernmost Chile. *Geologica Acta*, 5(4), 295-306.
- Rapalini, A.E., 2007. A paleomagnetic analysis of the Patagonian Orocline. *Geologica Acta*, 5(4), 287-294.
- Rossello, E.A., Haring, C.E., Cardinali, G., Suárez, F., Laffitte, G.A., Nevistic, A.V., 2008. Hydrocarbons and petroleum geology of Tierra del Fuego, Argentina. *Geologica Acta*, 6(1), 69-83.
- Tassone, A., Lodolo, E., Menichetti, M., Yagupsky, D., Caffau, M., Vilas, J.F., 2008. Seismostratigraphic and structural setting of the Malvinas basin and its southern margin. (Tierra del Fuego Atlantic offshore). *Geologica Acta*, 6(1), 55-67.
- Waldmann, N., Ariztegui, D., Anselmetti, F.S., Austin Jr., J.A., Dunbar, R., Moy, C.M., Recasens, C., 2008. Seismic stratigraphy of Lago Fagnano sediments (Tierra del Fuego, Argentina) - A potential archive of paleoclimatic change and tectonic activity since the Late Glacial. *Geologica Acta*, 6(1), 101-110.



Marco Menichetti received PhD in structural geology from the University of Perugia, Italy, in 1991. He spent the postdoctoral fellowship working on the crustal seismic survey project in Central Apennines. From 1995 to 1997 he was a consultant for oil and water companies.

He joined Urbino University, Italy, in 1998 where he is currently in charge of the degree courses on Structural geology and Tectonics. His research interest focus on field geology and structural concepts and methods for interpretation of mountain chains.



Alejandro A. Tassone graduated in 1989 from the Buenos Aires University (Argentina) and in 1998 obtained his PhD from Barcelona University (Spain). He is presently a researcher of the CONICET, the national council of scientific and technological research of Argentine, working at the the D. Valencio Institute of Geophysics of the Department of Geology

(Buenos Aires University). His main research interest is related to the application of several geophysical methodologies (seismic reflection, gravimetry, magnetometry, etc.) in deciphering the tectonostratigraphic and geodynamic evolution of the Patagonian and Antarctic continental margins.