



CHALMERS
UNIVERSITY OF TECHNOLOGY

Umbilical cord blood metabolome differs in relation to delivery mode, birth order and sex, maternal diet and possibly future allergy development in rural children

Downloaded from: <https://research.chalmers.se>, 2021-08-31 12:10 UTC

Citation for the original published paper (version of record):

Ross, A., Barman, M., Hartvigsson, O. et al (2021)

Umbilical cord blood metabolome differs in relation to delivery mode, birth order and sex, maternal diet and possibly future allergy development in rural children

PLoS ONE, 16(1 January)

<http://dx.doi.org/10.1371/journal.pone.0242978>

N.B. When citing this work, cite the original published paper.

RESEARCH ARTICLE

Umbilical cord blood metabolome differs in relation to delivery mode, birth order and sex, maternal diet and possibly future allergy development in rural children

Alastair B. Ross^{1,2}, Malin Barman¹, Olle Hartvigsson¹, Anna-Carin Lundell³, Otto Savolainen¹, Bill Hesselmar⁴, Agnes E. Wold^{3,5}, Ann-Sofie Sandberg^{1*}

1 Department of Biology and Biological Engineering, Chalmers University of Technology, Gothenburg, Sweden, **2** Food and Biobased Products Group, AgResearch, Lincoln, New Zealand, **3** Dept. of Rheumatology and Inflammation Research, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden, **4** Department of Paediatrics, Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden, **5** Clinical Bacteriology Section, Department of Infectious Diseases, University of Gothenburg, Gothenburg, Sweden

* Ann-Sofie.Sandberg@chalmers.se



OPEN ACCESS

Citation: Ross AB, Barman M, Hartvigsson O, Lundell A-C, Savolainen O, Hesselmar B, et al. (2021) Umbilical cord blood metabolome differs in relation to delivery mode, birth order and sex, maternal diet and possibly future allergy development in rural children. PLoS ONE 16(1): e0242978. <https://doi.org/10.1371/journal.pone.0242978>

Editor: Pal Bela Szecsi, Copenhagen University Hospital Holbæk, DENMARK

Received: May 13, 2020

Accepted: November 13, 2020

Published: January 25, 2021

Copyright: © 2021 Ross et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The metabolomics dataset is available via the MetaboLights database (www.ebi.ac.uk/metabolights), reference MTBLS2017.

Funding: This work was supported by the Swedish Research Council for Environmental, Agricultural Sciences and Spatial Planning (Formas), Grant number 222-2004-1958; the Swedish federal government under the LUA/ALF agreement; the

Abstract

Allergy is one of the most common diseases among young children yet all factors that affect development of allergy remain unclear. In a small cohort of 65 children living in the same rural area of south-west Sweden, we have previously found that maternal factors, including prenatal diet, affect childhood allergy risk, suggesting that *in utero* conditions may be important for allergy development. Here, we studied if metabolites in the umbilical cord blood of newborns may be related to development of childhood allergy, accounting for key perinatal factors such as mode of delivery, birth order and sex. Available umbilical cord blood plasma samples from 44 of the participants were analysed using gas chromatography-mass spectrometry metabolomics; allergy was diagnosed by specialised paediatricians at ages 18 months, 3 years and 8 years and included eczema, asthma, food allergy and allergic rhinoconjunctivitis. Nineteen cord blood metabolites were related to future allergy diagnosis though there was no clear pattern of up- or downregulation of metabolic pathways. In contrast, perinatal factors birth order, sex and mode of delivery affected several energy and biosynthetic pathways, including glutamate and aspartic acid—histidine metabolism ($p = 0.004$) and the tricarboxylic acid cycle ($p = 0.006$) for birth order; branched chain amino acid metabolism ($p = 0.0009$) and vitamin B₆ metabolism ($p = 0.01$) for sex; and glyoxylate and dicarboxylic acid metabolism ($p = 0.005$) for mode of delivery. Maternal diet was also related to some of the metabolites associated with allergy. In conclusion, the cord blood metabolome includes individual metabolites that reflect lifestyle, microbial and other factors that may be associated with future allergy diagnosis, and also reflects temporally close events/factors. Larger studies are required to confirm these associations, and perinatal factors such as birth order or siblings must be considered in future cord-blood metabolome studies.

Swedish Research Council (VR) under Grant number 521-2013-3154 and grant number 2019-01317.

Competing interests: The authors declare that they have no competing interests.

Introduction

Childhood allergy is one of the most prevalent diseases among children in countries with a Western lifestyle [1]. Causes of allergy are incompletely understood. The hygiene hypothesis proposes that a reduced exposure to microbes during early life results in allergy development [2,3], but other lifestyle factors, such as maternal and early life diet, may contribute [4]. Children who grow up on small family farms are at much lower risk of developing allergy than other children living in rural areas [5–8]. Both exposures of the infant and the pregnant mother to the farming environment are thought to contribute to protection against allergy development in the child [9–11].

The FARMFLORA birth cohort was established to investigate factors of the farming environment that contribute to the protection of allergy in farm children. We recently showed that farming mothers ate more butter and less margarine during pregnancy and lactation compared to non-farming mothers [12], and that a high maternal margarine intake was associated with more allergy in the offspring [12]. We also found that high proportions of the long-chain omega (n)-3 polyunsaturated fatty acids (PUFAs) docosahexaenoic acid (DHA) in breast milk [13] and eicosapentaenoic acid (EPA) in the serum of new born infants [14] and of four-month old infants were associated with a decreased risk of allergy development, although this was unrelated to growing up on a farm [13].

Mechanisms behind why growing up on a farm may be protective against allergy have been suggested, including increased exposure to farm animals and their associated microorganisms, consumption of unpasteurized milk [11] and dietary differences as outlined in our earlier work [12]. A new avenue for deciphering the mechanisms behind disease development is to use metabolomics, the profiling of a wide range of metabolites in a sample [15]. Metabolomics not only detects molecules that represent endogenous metabolism, it also contains metabolites derived from food and the environment which together can provide important information about external factors which may trigger allergy [16]. Several studies have used metabolomics to better understand mechanisms behind certain types of childhood allergic diseases, especially asthma [17,18]. Several studies on newborn infants have used metabolomics to study factors associated with foetal exposures such as maternal smoking [19], and gestational diabetes [20,21], as well as child health outcomes, including birth weight [21], and preterm delivery [19]. Even though maternal and *in utero* factors are thought to be related to development of allergy, there is little known about whether the metabolome at birth could reflect future allergy risk. We hypothesised that metabolites in cord blood are influenced by foetal sex, mode of delivery, birth order and maternal diet during pregnancy and may predict allergy development.

In order to explore whether metabolites in the cord blood of newborn infants born in farm or non-farm environments could be related to future risk of ‘any allergy’ (i.e. the child have one or more of: food allergy, eczema, asthma or allergic rhinoconjunctivitis), we performed metabolomics on cord blood serum from a small observational study [12]. We also investigated whether the metabolic profile of umbilical cord blood was associated with other factors associated with allergy development in the same cohort [12], such as diet, caesarean delivery, sex and being first-born.

Materials and methods

Subjects

The FARMFLORA birth-cohort study recruited 65 pregnant women at maternity clinics in the Skaraborg county in South-West Sweden between September 2005 and May 2008, 28 of whom lived on small dairy farms and 37 residing in the same rural areas, but not on farms.

Participants from farming environments other than dairy farms, or urban areas were excluded. Final inclusion in the cohort occurred at delivery, when children born between gestational week 36 and 42 were included. Children were followed up regularly from birth until around eight years of age.

Ethics statement

The study was conducted according to the relevant guidelines of the Declaration of Helsinki, and approved by the Regional Ethics Committee in Gothenburg (No. 363–05). Written informed consent was obtained from both parents. All participants were informed about their right to withdraw from the study at any point and to have their data removed.

Power analysis

All available samples were used in this study, and no statistical power analysis was carried out due to no pre-defined hypotheses about which metabolites would be likely to change, nor available data in a similar cohort. This is in line with the exploratory nature of this metabolomics study.

Sampling of umbilical cord blood

Cord blood was collected directly after delivery according to standard clinical routines at the hospital. The blood was allowed to clot for 30 min and centrifuged. Serum was removed, aliquoted and frozen. Frozen sera were stored at -80°C until analyzed.

Clinical examination

The children were examined clinically by pediatricians at 18 months, 3 years and around 8 years of age to diagnose food allergy, eczema, asthma and allergic rhinoconjunctivitis according to carefully standardized protocols as previously described [20,22,23]. All 65 children participated in the follow-up at 18 months, 63 children (97%) participated in the three year follow up and 48 of these children participated also in the 8-year follow up (76%, median age 8.3 years, range 6.5–9.4).

Allergic rhinoconjunctivitis (ARC) was defined as symptoms from eyes and/or nose upon exposure to pollen or animal dander, combined with demonstration of allergen-specific IgE to the corresponding inhalant allergen (Phadiatop, Phadia, Uppsala, Sweden).

Asthma was diagnosed at 18 months and 3 years of age based on ≥ 3 wheezing episodes combined either with a) eczema, allergic rhinoconjunctivitis or food allergy, or b) a response to leukotriene antagonists or inhaled glucocorticoids. At least one wheezing episode had to have occurred after 2 years of age for an asthma diagnosis at age 3 years. At 8 years of age asthma was defined as wheeze or heavy breathing in the last year together with a) bronchial obstruction reversible in response to β_2 -agonist, b) bronchial hyperresponsiveness on methacholine challenge or c) ongoing preventive asthma treatment with inhaled corticosteroids.

Food allergy was defined based on immediate or late-onset reactions after ingestion of a specific food that rapidly improved after allergen elimination. The diagnosis was supported by open food challenge tests and/or positive Fx5 Food Mix test (Phadia), followed by Immunocap tests (Phadia) for identification of the allergen.

Atopic eczema was diagnosed according to William's criteria or based on at least three major Hanifin and Rajka criteria [24].

Sensitization against common foods (Fx5 Food Mix) and inhalant allergens (Phadiatop) were assessed by blood tests (Phadia, Uppsala, Sweden) [20].

Due to the low number of subjects who developed allergy over the follow-up period, all allergies were grouped together in the statistical analyses rather than separating into different types of allergy. Allergy diagnoses at the three follow up visits are outlined in [S1 Table](#).

Metabolomics

The metabolite profile of umbilical cord blood plasma was analysed by gas chromatography-mass spectrometry using a previously published method [25]. In brief, 100 μ L of cord blood serum was extracted using 90% methanol with an internal standard mix to correct for variation in extraction and instrument response. Samples were derivatised using methoxymation and silylation and 1 μ L was injected onto a Shimadzu GCMS8030 gas chromatograph-tandem mass spectrometer (GC-MS) [25]. Data were processed to identify compounds based on their retention index and spectrum against the Swedish Metabolomics Centre library. Only identified compounds were used for further data analysis.

Maternal diet

Maternal diet data was collected using a semi-quantitative food frequency questionnaire (FFQ) adapted from the validated Northern Sweden 84-item questionnaire [26] extended with food items reflecting fat-containing foods [27]. The FFQ was sent out shortly after delivery and asked about intake frequencies the past year, hence covering the intake during the whole pregnancy. The methodology and results from the direct association between these assessments of maternal diet during pregnancy and allergy have been reported previously [12].

Data analysis

The initial analysis of the cord blood metabolome focused on differences between children were born to parents living on dairy farms or not (the basis for the design of the study), and differences in cord blood metabolites between subsequently allergic and non-allergic children (controls) based on a diagnosis determined at 18 months, 3 years and 8 years of age. Children who were not allergic at 3 years or 8 years, but had been diagnosed as being allergic at an earlier time-point were not included in the non-allergic control group for those follow up ages. Statistical significance was tested using Mann-Whitney tests due to non-normal distribution of metabolite data. Further, the relation between cord metabolic profiles and three predefined factors that were previously determined to be associated with allergy were investigated, namely delivery mode (caesarean or vaginal) [28], being first born [2], and being of male sex [29]. The relationship between metabolite pattern and allergy diagnosis was also adjusted for these three factors using logistic regression. Sensitivity analyses were performed where difference in cord blood metabolites between allergic and non-allergic children, at 18 months and 3 years, were analyzed in the non-farm group only by excluding all children in the farm group from the analysis. Statistical analyses were carried out using R version 3.5.1.

As we have previously found maternal diet during pregnancy to be associated with childhood allergy in this cohort [12], maternal food frequency scores were correlated with umbilical cord metabolites that were found to be associated with future allergy development at any time-point using Spearman's rank correlation. Correlations >0.4 or <-0.4 are reported. Due to the low number of subjects, maternal diet was not included as a factor in the logistic regression models.

Principal Components Analysis (PCA) was used to identify any outliers due to analytical issues (Simca 15, Umetrics AB, Sweden).

The potential effects of the metabolite differences on metabolic pathway upregulation was tested using the Pathway Analysis tool in MetaboAnalyst [30]. Separate analyses for allergy at

18 months and 3 years and 8 years of age, exposure to farming conditions, mode of delivery, having siblings or not and sex were carried out. All identified cord blood metabolites with corresponding Kyoto Encyclopaedia of Genes and Genomes (KEGG) metabolite identification numbers were included in this analysis (133 out of 155 metabolites), and data were log-transformed prior to analysis and mean centred prior to analysis. The data were matched to the KEGG human metabolite pathway library (Kyoto Encyclopaedia of Genes and Genomes, www.genome.jp).

Differences between conditions for metabolites, and pathway enrichment were considered significant at $P < 0.05$, with no correction for multiple testing in line with the exploratory nature of this study.

Results

Characteristics of the 44 included families are shown in [Table 1](#). The mean maternal age of the women were 33 years (range 21–42 years). The infants were born between gestational week 36 and 42 with a mean of 39 weeks. The mean birth weight was 3512 grams (range 2730–4435). One infant was born large for gestational age (LGA) while no infants were born small for gestational age (SGA) according to ultrasound-based growth curves [31]. Four infants were born preterm, i.e. before 37 full weeks of gestation. Around half of the infants were of male sex, 6 were born with caesarean section, 26 had older siblings and 4 were part of a twin pair ([Table 1](#)). The mothers of the children in this cohort were not undergoing any specialist medical care during their pregnancy so maternal health has not been included as a factor in subsequent analyses.

A total of 155 metabolites were identified in cord blood plasma after matching for retention index and mass spectra, and after visual inspection of the mass spectrum and chromatogram for the aligned peak result. No trends due to analysis were observed, and nor was there any clear grouping for allergy, farming, delivery mode, birth order, siblings or sex with PCA ([S1 Fig](#)).

Effect of farm residence

Twelve metabolites were associated with perinatal exposure to a farming environment, but none of them very strongly ([Table 2](#)). Infants born to mothers living on dairy farms had higher

Table 1. Demographic data of the 44 included mother/new-born dyads.

	Mean or n	SD(%)
Maternal age	33	(4.3)
Gestational age, weeks	39	(1.6)
Gestational age, days	278	(11)
Birth weight	3512	(403)
SGA	0	(0%)
LGA	1	(2%)
Preterm birth	4	(10%)
Male sex	21	(48%)
Twin pairs	2	(5%)
Caesarean section	6	(14%)
Older siblings	26	(59%)

Small for gestational age (SGA) and large for gestational age (LGA) status according to ultrasound-based growth curves [31]. Preterm birth is defined as delivery before 37 full weeks of gestation.

<https://doi.org/10.1371/journal.pone.0242978.t001>

Table 2. Metabolic differences in umbilical cord blood between farm and non-farm infants.

	<i>Farm, n = 21</i>		<i>Non-farm, n = 23</i>		P-value	Fold change
	Median	IQR	Median	IQR		
Anhydro-D-glucitol	52506	23677	62409	27438	0.013	-0.42
Lactose	4998	3850	2706	2823	0.013	0.55
N-acetyl-L-cysteine	447895	340228	273604	149632	0.020	0.72
Cellobiose	4751	3484	2602	2283	0.028	0.51
Ursodeoxycholic acid	29686	24368	55408	39445	0.034	-0.64
Lactulose	5852	3635	3126	2523	0.040	0.47
Pyruvic acid	123734	48985	107303	40785	0.040	0.41
Glycolic acid	37458760	10366464	32018982	7180247	0.042	0.27
Rhamnose	2696570	1824822	1747505	943766	0.042	0.57
Aconitic acid	138590	64230	111944	55014	0.045	0.28
Lyxose	4019	2054	3274	1609	0.045	0.37
Hydroxybutyric acid	18851	21051	21709	28178	0.048	-0.64

Data are GC-MS peak areas. Differences between farm and non-farm infants were analysed with Mann-Whitney U-test.

<https://doi.org/10.1371/journal.pone.0242978.t002>

levels of lactose, n-acetyl-l-cysteine, cellobiose, lactulose, pyruvic acid, glycolic acid, rhamnose, aconitic acid and lyxose while infants born to non-farming mothers had higher levels of anhydro-d-glucitol, ursodeoxycholic acid and hydroxybutyric acid. There were no metabolic pathways impacted by farm vs non-farm residence.

The effect of delivery mode

The three predefined factors that are known to be associated with allergy, i.e. mode of delivery (caesarean or vaginal delivery) [28], whether the child was the first born or had siblings [2], and sex [29], were all found to be strongly associated with metabolic differences. Fourteen metabolites differed between children delivered vaginally and through caesarean section, covering several types of metabolites including amino acids, carbohydrates and one-carbon metabolism (Table 3; S2 Fig). Pathway analysis of the identified cord metabolites found that glyoxylate and dicarboxylic acid pathways were significantly altered based on whether children were born by vaginal or caesarean section delivery ($p = 0.005$). Of the six caesarean sections, one was acute, four were planned while no information was available for one.

The effect of birth order

The effect of being first born or having older siblings was strongly imprinted in the umbilical cord metabolome, with 25 metabolites differing between these two groups (Table 4; S3 Fig). Birth order affected several metabolic pathways: glutamate and aspartic acid-histidine metabolism ($p = 0.004$), tricarboxylic acid (TCA) cycle ($p = 0.006$), fructose and mannose metabolism ($p = 0.01$) and butanoate metabolism ($p = 0.01$).

The effect of sex

Ten metabolites differed between male and female infants (Table 5; S4 Fig). Notably, all metabolites that differed between sexes were higher in girls except for erythritol and 2-oxoisocaproic acid.

The sex of the children was reflected in differences between several pathways; branched chain amino acid metabolism ($p = 0.0009$), vitamin B₆ metabolism ($p = 0.01$), taurine and

Table 3. Significant metabolic differences between mode of delivery (caesarean section or vaginal delivery).

	Caesarean section, n = 6		Vaginal delivery, n = 38		p value	Fold change
	Median	IQR	Median	IQR		
Lyxose	2345	937	3669	1852	0.0021	-0.83
Quinic acid	43781	19374	79624	30289	0.0130	-0.66
Uridine	30185	8908	25170	9019	0.0145	0.39
Ribose-5-phosphate	19612	5896	11732	7992	0.0222	0.51
Alanine	44880	17508	58163	27489	0.0246	-0.52
Fructose	4705	1965	11450	17622	0.0271	-1.80
Sorbose	4403	2143	11344	18241	0.0271	-1.81
Creatinine	44201	29212	29821	11690	0.0299	0.54
Homocysteine	24301	13183	12929	9345	0.0299	0.60
Nigerose	2463	1799	4496	4064	0.0299	-0.96
Valine	275398	14795	338292	80070	0.0299	-0.27
Homoserine	25773	13377	35253	16044	0.0361	-0.53
Glucosamine	173120	262624	373811	323590	0.0434	-0.94
Isoleucine	73896	21988	87316	28365	0.0434	-0.40

Data are GC-MS peak areas. Differences between the two groups were analysed with Mann-Whitney U-test.

<https://doi.org/10.1371/journal.pone.0242978.t003>

hypotaurine metabolism ($p = 0.01$), pyruvate metabolism ($p = 0.017$), terpenoid backbone biosynthesis ($p = 0.017$), cysteine and methionine metabolism ($p = 0.018$), glyoxylate and dicarboxylate metabolism ($p = 0.024$), and nicotinate and nicotinamide metabolism ($p = 0.026$).

Prediction of future allergy by cord blood metabolic pattern

Eight metabolites were associated with allergy development when all follow up timepoints were considered (“allergy at any time-point”, Table 6). However, of these eight, only ursodeoxycholic acid was predictive for allergy development at any one of the individual time points, being higher in infants diagnosed with allergy at 18 months of age. Cord blood metabolites that were associated with future allergy diagnosis differed according to the time point at which allergy was diagnosed (Table 6 and S5–S7 Figs) and no single metabolite was associated with allergy at all time points, though α -ketoglutaric acid was lower in the cord blood of children who would go on to be diagnosed with allergy at 18 months and 3 years. Notably, there was a stronger association between cord blood metabolite composition and allergy at age 8 years, than with allergy at the earlier time points (Table 6). The sample size was too small to allow further investigation based on the type of allergy diagnosis. Adjustment for being first-born, male sex and caesarean delivery attenuated approximately half of the unadjusted analysis differences between allergic and non-allergic children (Table 6).

Some of the metabolites that were associated with allergy status were also associated with a farming environment. For example, ursodeoxycholic was lower in infants from farming families than from non-farming families (Table 2) and higher in allergic compared to non-allergic infants at 18 months of age as well as being allergic at any time point during follow up (Table 6). Conversely, disaccharides lactose and cellobiose were higher in the cord blood of children born to farmers (Table 2), and lower in children who developed allergy at any point during follow up (Table 6). Accordingly, allergy was much more prevalent in non-farming than farming families at the two first diagnostic time points [12] and for the 18 months and 3 years follow ups, growing up on a farm or not is a major confounding factor for the allergy analysis.

Table 4. Significant metabolic differences between children with or without siblings.

	Have older siblings, n = 26		No older siblings, n = 18		p value	Fold change
	Median	IQR	Median	IQR		
Oxalic acid	159634	46356	215444	91995	0.00003	-0.69
Citric acid	11302	4506	18922	8789	0.00124	-0.57
Inositol	46903	22119	34617	14392	0.00124	0.71
N-acetyl glucosamine	1707288	821658	1406169	531941	0.00214	0.72
Phenylpyruvic acid	64297	16030	79925	20230	0.00214	-0.32
Sorbitol	3249	2773	5449	5676	0.00256	-0.87
Lactic acid	298432	51095	366236	90623	0.00279	-0.23
Isoerythritol	375786	133672	498367	195137	0.00424	-0.35
α -tocopherol	161923	91978	245157	75151	0.00738	-0.28
Glutamic acid	51698	26943	33843	11331	0.00796	0.46
N-acetylmannosamine	166880	67235	132443	54499	0.00796	0.53
Fumaric acid	223664	101344	305973	141690	0.01428	-0.41
Xylose	3313	1354	4395	2440	0.01428	-0.40
Ornithine	33095	16235	20638	14835	0.01532	0.49
Malic acid	2967379	1341621	3740148	1756233	0.01641	-0.41
Nigerose	3604	2447	4951	4616	0.01758	-0.62
Cellobiose	2795	3111	4665	4679	0.01881	-0.67
N-acetylornithine	74127	54777	107498	57345	0.02012	-0.55
Allothreonine	79426	28318	71611	17009	0.02297	0.26
Isocitric acid	35225	14285	43186	15557	0.02297	-0.27
Aspartic acid	1654723	604487	1383135	267027	0.02787	0.25
Lactose	2651	3098	4884	4441	0.02970	-0.62
Xylitol	19956	14046	33050	12069	0.03161	-0.46
Lactulose	3089	3138	5899	4173	0.04284	-0.57
Phenylalanine	1464103	549524	1372503	293543	0.04284	0.22
Arabinose	74523	39838	98879	37775	0.04544	-0.43

Data are GC-MS peak areas. Differences between the two groups were analysed with Mann-Whitney U-test.

<https://doi.org/10.1371/journal.pone.0242978.t004>

Table 5. Significant metabolic differences between male and female sex.

	Male, n = 21		Female, n = 23		p value	Fold change
	Median	IQR	Median	IQR		
Amiloride	39878	5597	45074	15684	0.009	-0.28
Leucine	31319	9779	36673	23121	0.010	-0.50
Isoleucine	80114	26489	98224	50301	0.013	-0.40
Galactitol	1029	2346	3146	4555	0.031	-1.32
Hydroxybutyric acid	17053	21402	24599	19108	0.032	-0.66
Uric acid	51282	47882	65892	93350	0.033	-0.90
Erythritol	2221990	1257668	1595929	926665	0.036	0.54
Sucrose	8834	5910	13690	20589	0.040	-1.21
Mannose	3416	6329	5442	15173	0.042	-0.05
2-oxoisocaproic acid	27838	14608	22864	12665	0.045	0.28

Data are GC-MS peak areas. Differences between the two groups were analysed with Mann-Whitney U-test.

<https://doi.org/10.1371/journal.pone.0242978.t005>

Table 6. Significant differences in cord blood metabolites between subsequently allergic and non-allergic children based on diagnosis of allergy at different ages.

Allergy at any time point	Allergic, n = 14		Not allergic, n = 30		p values					Fold change
	Median	IQR	Median	IQR	unadjusted	Adjusted*				
						BO	S	DM	BO S DM	
Ophthalmic acid	23843	14351	36824	34405	0.016	0.048	0.044	0.054	0.049	-0.522
Ursodeoxycholic acid	55213	47987	31957	34161	0.019	0.010	0.008	0.018	0.010	0.726
δ-tocopherol	29378	21603	16356	13392	0.027	0.027	0.026	0.034	0.027	0.622
Glyceric acid	1824	872	2633	1697	0.034	0.076	0.069	0.088	0.080	-0.411
Lactose	2682	2612	4002	4051	0.035	0.105	0.108	0.082	0.119	-0.690
Cellobiose	2558	2474	3640	4200	0.038	0.108	0.113	0.084	0.124	-0.701
Sorbitol	4245	3702	3950	7878	0.040	0.111	0.141	0.094	0.125	-0.798
Nigerose	3564	2321	4649	3978	0.047	0.127	0.141	0.097	0.145	-0.607
18 months	Allergic, n = 10		Non-allergic, n = 34							
Ursodeoxycholic acid	57662	34659	32062	33350	0.021	0.012	0.029	0.010	0.013	0.674
Pyroglutamic acid	607144	108081	664308	116454	0.026	0.034	0.031	0.031	0.042	-0.148
α-ketoglutaric acid	25078	2136	31390	9669	0.041	0.112	0.080	0.087	0.135	-0.245
3 years	Allergic, n = 6		Non-allergic, n = 31							
Oxoisocaproic acid	16762	3623	27249	13368	0.019	0.053	0.049	0.040	0.062	-0.549
Uridine	31445	5643	25161	7826	0.022	0.042	0.045	0.028	0.040	0.423
Benzoic acid	30115	3808	35220	7458	0.025	0.109	0.106	0.093	0.108	-0.292
Putrescine	12761	13004	9209	4980	0.035	0.041	0.039	0.034	0.044	0.804
α-ketoglutaric acid	24001	1716	31714	9364	0.044	0.145	0.109	0.111	0.146	-0.303
Isocitric acid	30102	11344	42209	15486	0.048	0.037	0.050	0.034	0.045	-0.360
8 years	Allergic, n = 7		Non-allergic, n = 24							
E-Octadecenoic acid	3599571	432707	2700887	889046	0.003	0.023	0.016	0.044	0.017	0.391
Z-Octadecenoic acid	3213916	306038	2602898	748100	0.011	0.051	0.046	0.127	0.045	0.231
Aspartic acid	1284643	246623	1513347	420257	0.016	0.031	0.040	0.032	0.040	-0.318
Cholic acid	1420	196	857	498	0.022	0.141	0.095	0.122	0.139	0.544
Hydroxyphenylacetic acid	11028	5790	7579	4389	0.033	0.099	0.161	0.070	0.097	0.665
Hexadecanoic acid	225387	38680	194448	39945	0.041	0.030	0.031	0.049	0.028	0.216
N-acetylmethionine	109336	51788	72155	44892	0.041	0.057	0.070	0.066	0.068	0.611
Uric acid	78720	115274	46950	33403	0.041	0.224	0.133	0.128	0.230	1.113

Data are GC-MS peak areas. Differences between the two groups were analysed with Mann-Whitney U-test.

* Adjusted for: Birth Order (BO), Sex (S) and Delivery mode (DM).

<https://doi.org/10.1371/journal.pone.0242978.t006>

To account for the difference in prevalence of allergy in the non-farming and farming families, we performed sensitivity analyses on difference in cord blood metabolites between allergic and non-allergic children, at 18 months and 3 years, in the non-farm group only (Table 7). None of the three metabolites that were associated with allergy at 18 months of age (ursodeoxycholic acid, pyroglutamic acid, alpha-ketoglutaric acid) were associated with allergy when only children in the non-farm group were included in the analyses between allergic and non-allergic individuals. Instead four other metabolites were significantly different between allergic and non-allergic individuals (δ-tocopherol, dodecanoic acid, glucose and myristoleic acid). However, when examining allergy at 3 years of age three of the six metabolites that differed between allergic and non-allergic individuals in the whole cohort were still significantly different when looking only in the non-farm group: uridine, benzoic acid and alpha-ketoglutaric acid. At 8 years of age, the distribution of allergy in the farm and non-farm group did not differ statistically ($p = 0.43$) with 7 allergic children in the non-farm group and 3 allergic

Table 7. Significant metabolic differences between allergic and non-allergic children based on diagnosis at 18 months and 3 years in the non-farm group only.

	Allergic, n = 8		Non-allergic, n = 15		p value	Fold change
	Median	IQR	Median	IQR		
18 months						
δ-tocopherol	18667	18009	17498	17895	0.026	0.62
Dodecanoic acid	56684	32315	78308	41385	0.034	-0.09
Glucose-6-phosphate	1432	816	1867	1458	0.045	-0.87
Myristoleic acid	193299	108112	229996	182648	0.047	-0.58
3 years	Allergic, n = 5		Non-allergic, n = 14			
α-ketoglutaric acid	26937	10347	29692	9288	0.026	-0.36
Uridine	25733	4878	23070	6345	0.026	0.45
Monostearoylglycerol	33446	13958	44943	106907	0.044	-0.48
Adenosine	14825	13161	20101	12181	0.044	0.76
Asymmetrical-N-N-dimethylarginine	276868	114167	269977	158079	0.044	-0.37
Benzoic acid	31213	2741	36317	12814	0.044	-0.39
Glucose-6-phosphate	1392	718	1884	1239	0.044	-1.05

Data are GC-MS peak areas. Differences between the two groups were analysed with Mann-Whitney U-test. Statistical analyses were not adjusted for any covariables due to the low number of participants when only including non-farm children.

<https://doi.org/10.1371/journal.pone.0242978.t007>

children in the farm group. Hence, no sensitivity analyses based on farming environment were performed for allergy at this age.

Pathway analysis did not find any significant differences between cord blood metabolome at birth in allergic and non-allergic children at 18 months, 3 years or 8 years of age, suggesting that of the metabolites that were mapped in metabolic pathways, those that differed did not represent a concerted difference in metabolic status and instead likely reflect the impact of genetic, lifestyle and other environmental differences.

To probe this, we tested the association between maternal diet and metabolites associated with allergy in the overall cohort. Of the 23 metabolites that were associated with allergy overall or any time point, 19 were also correlated with one or more foods in the maternal diet during pregnancy (S2 Table).

Discussion

In this small, but well controlled study we found that metabolites in umbilical cord blood were strongly associated both with delivery mode, sex and birth order, with potentially important impacts on metabolic pathways. Cord blood metabolites were also associated with prenatal exposure to maternal diet and a farming environment and later allergy development but were not indicative of consistent systemic metabolic effects.

Our aim was to identify pre- and peri-natal factors associated with subsequent allergy development, but we did not find any relationship with perturbation of metabolic pathways, though both low statistical power and scattered metabolite coverage may have hindered finding differences in biochemical pathways related to future allergy.

The strongest relationships with the cord blood metabolome were events that were in close temporal proximity to the sample being collected (delivery mode) or of fundamental biological importance (sex and birth order). While sex imprinting on the metabolome of infants has been frequently described [32,33], as has the difference in delivery mode [34–36], we were surprised to find that birth order (i.e. first born or having older siblings) was also reflected in the metabolome. Having older siblings or not impacted on several related metabolic pathways, with the TCA cycle, butanoate metabolism, aspartic acid and glutamate metabolism all being

linked, especially through fatty acid metabolism and acetyl-CoA. Possible reasons are speculative, though practical aspects such as the mother having already experienced giving birth leading to a different level of psychological and physiological preparedness, or the need to look after another child during pregnancy may play a role. Time may also be a factor, and although time from rupture of membranes to delivery was not recorded in this study, in general first deliveries take longer than subsequent deliveries [37]. In support of the hypothesis that use of energy, length of delivery and overall effort being greater for first delivery is that the concentration of energy metabolism end product lactic acid was higher in cord blood of children with no siblings, as were TCA-cycle metabolites. Further, lactic acid, an end product of muscle energy metabolism, was borderline lower in cord blood after caesarean delivery ($p = 0.061$), suggesting that muscular contractions do impact on the cord blood metabolome and is supported by earlier work comparing delivery mode using metabolomics [34,35]. As a crude measure of the relative importance of having siblings versus other factors, the number of metabolites related to having siblings or not, was more than twice that of either sex or delivery mode (26 compared to 12 and 14 respectively). Most metabolites that differed between children with siblings or no siblings were higher in those with no siblings (20 out of 26).

Mode of delivery also led to several differences in umbilical cord metabolites, notably a higher concentration of homocysteine in caesarean delivered children compared to vaginally delivered children. Homocysteine, an intermediate of one carbon metabolism and DNA methylation, has been linked to birth complications [38], and has previously been found to be higher in women having caesarean sections under general anaesthesia [39]. The elevation of homocysteine in the cord blood of caesarean delivered infants may reflect differences in requirements for methyl-groups and increased use of methyl-donors and related enzyme cofactors such as folate and vitamin B₁₂. We did not have information about the time between rupture of membranes and delivery, and this may also be informative in future studies if delivery mode has an impact on the cord blood metabolome. There did not appear to be any clear metabolic connection between factors associated with delivery mode, siblings or sex that could be clearly linked to metabolic factors that were related to allergy risk.

The metabolites that differed between children who went on to develop allergy compared to those who did not covered a wide range of compound classes, though notably included several that are related to lipid and energy metabolism including bile acids (ursodeoxycholic acid and cholic acid), fatty acids (octadecenoic acids 9Z and 9E, eicosapentaenoic acid, and hexadecenoic acid), amino acid derivatives (pyroglutamic acid, 2-oxoisocaproic acid, aspartic acid, acetylornithine and putrescine) and Krebs Cycle intermediates (α -ketoglutaric acid and isocitric acid). Earlier work in this cohort on both the fatty acid profile of the infants [13] and maternal intake of dietary fat [12] have found that both have an impact on allergy risk. The finding that bile acids were elevated in the cord blood of infants that went on to develop allergy supports the role of dietary lipids in allergy development in this cohort. Bile acids have been associated with immune regulation and allergy [40,41], though given the potential confounding from different fat intakes and endogenous fatty acid profiles, it is not possible to establish if there is a causative relationship between elevated bile acids and allergy in this study.

The metabolites associated with allergy development differed when comparing allergy status at different ages. Symptomatic allergy is variable with age, often seen with children 'growing out of' allergy, while more severe and longer-term allergy is observed in older children, reflected in the eight year allergy diagnosis follow up, a phenomenon called 'the atopic march' [42]. Only three umbilical cord metabolites differed between children who were diagnosed as being allergic at 18 months, while increasing numbers were different in relation to allergy diagnosis at the later follow up visits (six and eleven metabolites at 3 years and 8 years respectively). Only α -ketoglutaric acid was associated with allergy at two time points (18 months and 3

years). Metabolites related to protein breakdown and central energy metabolism were related to 3 year diagnosis, while several lipids differed between allergic and non-allergic children at 8 years old, which were not at all related to allergy diagnosis at 18 months and 3 years. Metabolites related to inflammatory pathways were not covered in this method, so it is not possible to state whether the umbilical cord concentrations of metabolites in these pathways with a clear relationship to allergy were in anyway related to future allergy. The possibility that later life allergy is imprinted already at birth is intriguing, and this would support the idea that conditions around *in utero* development and birth, along with genetic and general lifestyle factors may be associated with long-term allergy development. However, there were no overall effects on metabolic pathways detected, while such effects were detected for factors that would be expected to impact on the metabolome at the time of sample collection (e.g. sex, delivery mode). This also raises the possibility that the metabolites related to future allergy instead reflect lifestyle and familial factors that are related to future allergy risk, rather than being directly related biochemically to future allergy development. We investigated this in a further analysis on the association between maternal diet and cord blood metabolites related to allergy.

The role of maternal diet during pregnancy in the development of allergy is of great interest as it presents a potential modifiable factor in allergy risk [12,43]. In the current study we found that the metabolites that were associated with allergy risk were associated with many, and diverse, dietary factors. We did not model these dietary factors within the metabolite-allergy model due to the low overall number of subjects. We have earlier found that maternal diet during pregnancy influenced allergy in this cohort, with margarine intake being associated with higher allergy risk [22]. It was notable that many of the foods associated with allergy-related metabolites were also related to dietary fats. Higher intake of margarine or soya oil overall or in cooking was associated with metabolites that were predictive of allergy at 3 and 8 years (E- and Z-octadecanoic acid, hydroxyphenylacetic acid and uric acid), while the use of olive oil, rapeseed oil and butter on sandwiches were negatively associated with metabolites that predicted allergy at 3 and 8 years (hydroxyphenylacetic acid). In the case of 2-oxoisocaproic acid, a metabolite associated with lower risk of allergy, higher concentrations were measured with higher butter intake, and lower concentrations with higher margarine intake. These findings further support the importance of choice of dietary fat in the maternal diet during pregnancy for future allergy risk in this population. Due to the small size of this cohort we have not sought to interpret the potential-diet-cord metabolome-allergy interaction, only to highlight that there are many relationships between maternal diet and the cord blood metabolome and that this should be of future interest.

The children included in this study were from the FARMFLORA study, a small but well-controlled birth cohort following families living on dairy farms and families living in the same rural area but not on farms, with the aim to determine what in the farming environment could be responsible for the apparent protection against developing childhood allergy. We have previously found that eating habits among farmers and their infants differ from the families that do not live on farms [12,13,44]. For example pregnant mothers not living on dairy farms eat more margarine which was also associated with allergy in their infants [12]. Sensitivity analysis when non-farm raised children were removed from the model suggested that the differences between cord blood metabolites and allergy diagnoses at 18 months and 3 years could not be disentangled from whether children were raised on farms or not. This may further explain the difference between metabolites associated with allergy at the two first follow ups and the 8 year follow up, as the impact of maternal environment plays less of a role as the children grow older. Other metabolites, including uridine, benzoic acid and α -ketoglutaric acid were consistent in their association with allergy in both non-farm children and the entire cohort.

Only three metabolites were associated with both farming environment and allergy; ursodeoxycholic acid, lactose and cellobiose. Ursodeoxycholic acid, a bile acid, suggests a difference in lipid metabolism which matches earlier results on differences in maternal fat intake [12], and infant blood fatty acid composition [13]. The two disaccharides lactose and cellobiose may also be related to differences in dietary intake, with lactose being found in substantial quantities in milk, and cellobiose potentially coming from plant-based foods. In both cases it would be expected that most lactose and cellobiose is broken down during maternal absorption, though both compounds have been reported previously as being detected in plasma. As bile acids interact with the intestinal microbiota and are modified by conjugation [45], and both lactose and cellobiose represent fermentable substrate that has avoided breakdown in the large intestine, it is possible that these plasma metabolite differences are also indicative of a diet-environment-microbiome interaction playing a role in allergy development, though more detailed analyses on both faecal samples and microbe-related metabolites will be needed to prove this.

There are a number of weaknesses in this study and approach used which should be accounted for when interpreting the data. We did not use correction for multiple testing, which increases the likelihood of false positive results, due to the low number of subjects in the study. While this restricts the possible statistical power and generalisability of the outcomes, it has the advantage of allowing the cohort to be geographically well defined, and to measure many exposures in relation to doctor diagnosed allergy outcomes in a prospective setting and starting during pregnancy. The rate of allergy among those children in the study living on dairy farms could be a confounding factor in this study as the markers may reflect lifestyle rather than allergy risk. However all children in the study lived in the same rural environment, including while *in utero*, so while exposure to microorganisms from animals would be expected to be higher among children born to dairy farmers, other exposures such as weather and pollen are likely to be similar. Sensitivity analysis suggests that growing up on a farm or not is an important factor for metabolites related to childhood allergy in this study. Not all possible factors that have previously been related to allergy were controlled for, for example season of birth [46,47] or maternal physical activity and other lifestyle factors [48]. This study also had a moderately high loss to follow up at 3 and 8 years, representing 18% of the total children diagnosed with allergy. This number of participants dropping out of the study is not unexpected, especially at 8 years where families had moved from the region where the study was located. This does reduce the statistical power at the 8 year follow up, and the number of allergic children is reduced from 15 at 18 months to 11 at 3 years and 10 at 8 years. However this also follows the expected decline in allergy prevalence as children grow older. Although we have not stratified this analysis by type of allergy, it is also notable that eczema is the most prevalent allergy diagnosis at 18 months (73% of diagnoses including eczema, declining to 50% at 8 years). Of the four lost to follow up at 8 years, allergy diagnoses at earlier time points included eczema, asthma, food allergy and allergic rhinoconjunctivitis, suggesting that there was no impact of the type of diagnosis on participants dropping out of the study. The age of the children at the third follow up varied (range 6.5–9.4 years). This may confound comparisons of allergy diagnoses at either end of this range which spans 3 years. This age range is due to the practicalities of getting the children to the third follow up visit with the same paediatrician, meaning that it was not possible to have all children at exactly the same age when this was done. When detecting the impact of delivery mode on the cord blood metabolome, we have grouped together all types of caesarean deliveries, which themselves can differ widely, especially between elective and emergency caesarean delivery. With only six cases of caesarean delivery we did not have sufficient statistical power to further divide the cohort by type of caesarean delivery. Similarly, there was only one child who was born large for gestational age, and

two twin pairs and four children born pre-term in the cohort, so no sub-analyses were carried out for these factors.

Analytical methods are crucial to account for in metabolomics as the type of approach used dictates what metabolites are included in the final analysis. In this study we have used GC-MS for metabolomics, an instrument which has the advantage of allowing rapid identification of compounds based on mass spectral and retention index matching, but does not have the breadth of metabolite coverage of liquid chromatography-mass spectrometry when several different chromatographic modes are used. Not all identifications made using spectral and retention time matching could be confirmed, and the identity of one metabolite strongly associated with allergy at eight years could not be confirmed based on spectral matching and hence not included in the overall analysis. Additional metabolomics methods should be used to get wider coverage of the metabolome to discover other metabolites that are associated with allergy development.

Conclusions

Results from this work suggests that the umbilical cord blood metabolome is strongly influenced by sex, delivery mode and birth order, with apparent impacts on several biochemical pathways. There may also be imprinting of future allergy risk, though consistent changes related to mapped biochemical pathways were not evident and these metabolites may instead reflect lifestyle and familial factors that are in turn related to allergy risk. Maternal diet during pregnancy appears to be reflected in cord blood metabolites associated with allergy, supporting the hypothesis that maternal lifestyle may have an impact on future childhood allergy. These results underline that the metabolome is highly responsive to a myriad of factors, and that important but often overlooked covariables in health research such as siblings or delivery mode need to be accounted for in studies analysing samples collected at birth. Further work needs to be done to follow up on these relationships in larger cohorts [49] to determine what wider role metabolic status at birth may influence future allergy risk.

Supporting information

S1 Fig. PCA plot of all data together. PC1 explains 72.5% of total variation. Data points are coloured according to farmer (1) or non-farmer (0) status. There are no clear analysis-related trends in the data.

(DOCX)

S2 Fig. Box plots of the cord blood metabolites that differed ($p < 0.05$) at birth between children born via vaginal birth and children born via caesarean section. Children born via vaginal birth are represented by the dark grey boxes and children born via caesarean section by the light grey boxes.

(DOCX)

S3 Fig. Box plots of the cord blood metabolites that differed ($p < 0.05$) at birth between children with siblings and children without siblings at birth. Children without siblings are represented by the dark grey boxes and children with siblings by the light grey boxes.

(DOCX)

S4 Fig. Box plots of the cord blood metabolites that differed ($p < 0.05$) between sexes. Females are represented by the dark grey boxes and males by the light grey boxes.

(DOCX)

S5 Fig. Box plots of the cord blood metabolites that differed ($p < 0.05$) at birth between non-allergic children and children diagnosed with allergy at 18 months.

Non-allergic children are represented by the dark grey boxes and allergic children by the light grey boxes.

(DOCX)

S6 Fig. Box plots of the cord blood metabolites that differed ($p < 0.05$) at birth between non-allergic children and children diagnosed with allergy at 3 years.

Non-allergic children are represented by the dark grey boxes and allergic children by the light grey boxes.

(DOCX)

S7 Fig. Box plots of the cord blood metabolites that differed ($p < 0.05$) at birth between non-allergic children and children diagnosed with allergy at 8 years old.

Non-allergic children are represented by the dark grey boxes and allergic children by the light grey boxes.

(DOCX)

S1 Table. Distribution of types of allergy among allergic children followed up at 18 months, 3 and 8 years.

(DOCX)

S2 Table. Correlation (Spearman's rho) between metabolites associated with allergy development and maternal diet.

(PDF)

Acknowledgments

We thank Associate Professor Carl Brunius for helpful discussions on aspects related to data analysis. We also acknowledge the study nurses Helen Andersson and Anders Nordberg for their work in coordinating sampling and questionnaires. In addition, we acknowledge the paediatricians Margareta Ceder, Gunhild Lindhagen, Stefan Stentoft and Carl-Johan Törnåge, who diagnosed the children, and all the families for their participation in the FARMFLORA birth cohort.

Author Contributions

Conceptualization: Malin Barman, Bill Hesselmar, Agnes E. Wold, Ann-Sofie Sandberg.

Formal analysis: Alastair B. Ross, Malin Barman, Olle Hartvigsson.

Funding acquisition: Ann-Sofie Sandberg.

Investigation: Malin Barman, Anna-Carin Lundell, Bill Hesselmar, Ann-Sofie Sandberg.

Methodology: Alastair B. Ross, Otto Savolainen.

Project administration: Malin Barman, Ann-Sofie Sandberg.

Resources: Anna-Carin Lundell, Ann-Sofie Sandberg.

Supervision: Alastair B. Ross, Malin Barman, Agnes E. Wold, Ann-Sofie Sandberg.

Writing – original draft: Alastair B. Ross.

Writing – review & editing: Alastair B. Ross, Malin Barman, Olle Hartvigsson, Anna-Carin Lundell, Otto Savolainen, Bill Hesselmar, Agnes E. Wold, Ann-Sofie Sandberg.

References

1. Haanpää L, af Ursin P, Nermes M, Kaljonen A, Isolauri E. Association of allergic diseases with children's life satisfaction: population-based study in Finland. *BMJ Open*. 2018; 8(3):e019281. <https://doi.org/10.1136/bmjopen-2017-019281> PMID: 29602839
2. Strachan D. Hay fever, hygiene, and household size. *British Medical Journal*. 1989; 299:1259–60. <https://doi.org/10.1136/bmj.299.6710.1259> PMID: 2513902
3. Wold AE. The hygiene hypothesis revised: is the rising frequency of allergy due to changes in the intestinal flora? *Allergy*. 1998; 53(46 Suppl):20–5. <https://doi.org/10.1111/j.1398-9995.1998.tb04953.x> PMID: 9825991.
4. Black PN, Sharpe S. Dietary fat and asthma: is there a connection? *Eur Respir J*. 1997; 10(1):6–12. <https://doi.org/10.1183/09031936.97.10010006> PMID: 9032484.
5. Braun-Fahrlander C, Gassner M, Grize L, Neu U, Sennhauser FH, Varonier HS, et al. Prevalence of hay fever and allergic sensitization in farmer's children and their peers living in the same rural community. SCARPOL team. Swiss Study on Childhood Allergy and Respiratory Symptoms with Respect to Air Pollution. *Clin Exp Allergy*. 1999; 29(1):28–34. <https://doi.org/10.1046/j.1365-2222.1999.00479.x> PMID: 10051699.
6. Remes ST, Koskela HO, Iivanainen K, Pekkanen J. Allergen-specific sensitization in asthma and allergic diseases in children: The study on farmers' and non-farmers' children. *Clinical and Experimental Allergy*. 2005; 35(2):160–6. <https://doi.org/10.1111/j.1365-2222.2005.02172.x> PMID: 15725186
7. Riedler J, Eder W, Oberfeld G, Schreuer M. Austrian children living on a farm have less hay fever, asthma and allergic sensitization. *Clin Exp Allergy*. 2000; 30(2):194–200. <https://doi.org/10.1046/j.1365-2222.2000.00799.x> PMID: 10651771.
8. Von Ehrenstein OS, Von Mutius E, Illi S, Baumann L, Bohm O, von Kries R. Reduced risk of hay fever and asthma among children of farmers. *Clin Exp Allergy*. 2000; 30(2):187–93. <https://doi.org/10.1046/j.1365-2222.2000.00801.x> PMID: 10651770.
9. Douwes J, Cheng S, Travier N, Cohet C, Niesink A, McKenzie J, et al. Farm exposure in utero may protect against asthma, hay fever and eczema. *European Respiratory Journal*. 2008; 32(3):603–11. <https://doi.org/10.1183/09031936.00033707> PMID: 18448493
10. Ege MJ, Bieli C, Frei R, van Strien RT, Riedler J, Åcøblagger E, et al. Prenatal farm exposure is related to the expression of receptors of the innate immunity and to atopic sensitization in school-age children. *Journal of Allergy and Clinical Immunology*. 2006; 117(4):817–23. <https://doi.org/10.1016/j.jaci.2005.12.1307> PMID: 16630939
11. Riedler J, Braun-Fahrlander C, Eder W, Schreuer M, Waser M, Maisch S, et al. Exposure to farming in early life and development of asthma and allergy: a cross-sectional survey. *Lancet*. 2001; 358(9288):1129–33. [https://doi.org/10.1016/S0140-6736\(01\)06252-3](https://doi.org/10.1016/S0140-6736(01)06252-3) PMID: 11597666.
12. Jonsson K, Barman M, Moberg S, Sjoberg A, Brekke HK, Hesselmar B, et al. Fat intake and breast milk fatty acid composition in farming and nonfarming women and allergy development in the offspring. *Pediatric research*. 2016; 79(1–1):114–23. Epub 2015/09/22. <https://doi.org/10.1038/pr.2015.187> PMID: 26389822.
13. Jonsson K, Barman M, Moberg S, Sjoberg A, Brekke HK, Hesselmar B, et al. Serum fatty acids in infants, reflecting family fish consumption, were inversely associated with allergy development but not related to farm residence. *Acta Paediatr*. 2016; 105(12):1462–71. Epub 2016/09/18. <https://doi.org/10.1111/apa.13592> PMID: 27637371.
14. Barman M, Rabe H, Hesselmar B, Johansen S, Sandberg A-S, Wold AE. Cord Blood Levels of EPA, a Marker of Fish Intake, Correlate with Infants' T- and B-Lymphocyte Phenotypes and Risk for Allergic Disease. *Nutrients*. 2020; 12(10). Epub Sep 30. <https://doi.org/10.3390/nu12103000> PMID: 33007868
15. Dessì A, Cesare Marincola F, Masili A, Gazzolo D, Fanos V. Clinical metabolomics and nutrition: the new frontier in neonatology and pediatrics. *Biomed Res Int*. 2014; 2014:981219–. Epub 2014/08/27. <https://doi.org/10.1155/2014/981219> PMID: 25247199.
16. Agache I, Miller R, Gern JE, Hellings PW, Jutel M, Muraro A, et al. Emerging concepts and challenges in implementing the exposome paradigm in allergic diseases and asthma: a Practall document. *Allergy*. 2019; 74(3):449–63. Epub 2018/12/06. <https://doi.org/10.1111/all.13690> PMID: 30515837.
17. Turi KN, Romick-Rosendale L, Ryckman KK, Hartert TV. A review of metabolomics approaches and their application in identifying causal pathways of childhood asthma. *Journal of Allergy and Clinical Immunology*. 2018; 141(4):1191–201. <https://doi.org/10.1016/j.jaci.2017.04.021> PMID: 28479327
18. Chawes BL, Giordano G, Pirillo P, Rago D, Rasmussen MA, Stokholm J, et al. Neonatal Urine Metabolic Profiling and Development of Childhood Asthma. *Metabolites*. 2019; 9(9):185. <https://doi.org/10.3390/metabo9090185> PMID: 31527391

19. Gil AM, Duarte D. Biofluid Metabolomics in Preterm Birth Research. *Reprod Sci*. 2018; 25(7):967–77. Epub 2018/02/15. <https://doi.org/10.1177/1933719118756748> PMID: 29439621.
20. Strombeck A, Rabe H, Lundell AC, Andersson K, Johansen S, Adlerberth I, et al. High proportions of FOXP3(+) CD25(high) T cells in neonates are positively associated with allergic sensitization later in childhood. *Clin Exp Allergy*. 2014; 44(7):940–52. Epub 2014/02/18. <https://doi.org/10.1111/cea.12290> PMID: 24528482; PubMed Central PMCID: PMC4215110.
21. Robinson O, Keski-Rahkonen P, Chatzi L, Kogevinas M, Nawrot T, Pizzi C, et al. Cord Blood Metabolic Signatures of Birth Weight: A Population-Based Study. *Journal of Proteome Research*. 2018; 17(3):1235–47. <https://doi.org/10.1021/acs.jproteome.7b00846> PMID: 29401400
22. Lundell AC, Hesselmar B, Nordstrom I, Adlerberth I, Wold AE, Rudin A. Higher B-cell activating factor levels at birth are positively associated with maternal dairy farm exposure and negatively related to allergy development. *Journal of Allergy and Clinical Immunology*. 2015; 136(4):1074–+. Epub 2015/05/06. <https://doi.org/10.1016/j.jaci.2015.03.022> WOS:000362976300026. PMID: 25936566
23. Strombeck A, Nordstrom I, Andersson K, Andersson H, Johansen S, Maglio C, et al. Allergic disease in 8-year-old children is preceded by delayed B cell maturation. *Clinical and Experimental Allergy*. 2017; 47(7):918–28. <https://doi.org/10.1111/cea.12922> WOS:000404962400007. PMID: 28295779
24. Williams HC. Diagnostic criteria for atopic dermatitis. *Lancet*. 1996; 348(9038):1391–2. Epub 1996/11/16. [https://doi.org/10.1016/S0140-6736\(05\)65466-9](https://doi.org/10.1016/S0140-6736(05)65466-9) PMID: 8918311.
25. Savolainen OI, Sandberg AS, Ross AB. A Simultaneous Metabolic Profiling and Quantitative Multimetabolite Metabolomic Method for Human Plasma Using Gas-Chromatography Tandem Mass Spectrometry. *J Proteome Res*. 2016; 15(1):259–65. Epub 2015 Nov 30. <https://doi.org/10.1021/acs.jproteome.5b00790> PMID: 26615962
26. Johansson I, Hallmans G, Wikman A, Biessy C, Riboli E, Kaaks R. Validation and calibration of food-frequency questionnaire measurements in the Northern Sweden Health and Disease cohort. *Public health nutrition*. 2002; 5(3):487–96. Epub 2002/05/11. <https://doi.org/10.1079/phn2001315> S136898000200068X [pii]. PMID: 12003662.
27. Brekke HK, Jansson PA, Mansson JE, Lenner RA. Lifestyle changes can be achieved through counseling and follow up in first-degree relatives of patients with type 2 diabetes. *J Am Diet Assoc*. 2003; 103(7):835–43. [https://doi.org/10.1016/s0002-8223\(03\)00378-x](https://doi.org/10.1016/s0002-8223(03)00378-x) WOS:000183926400013. PMID: 12830021
28. Bager P, Wohlfahrt J, Westergaard T. Caesarean delivery and risk of atopy and allergic disease: meta-analyses. *Clinical & Experimental Allergy*. 2008; 38(4):634–42. <https://doi.org/10.1111/j.1365-2222.2008.02939.x> PMID: 18266879
29. Larsson L. Incidence of asthma in Swedish teenagers: relation to sex and smoking habits. *Thorax*. 1995; 50(3):260. <https://doi.org/10.1136/thx.50.3.260> PMID: 7660339
30. Chong J, Yamamoto M, Xia J. MetaboAnalystR 2.0: From Raw Spectra to Biological Insights. *Metabolites*. 2019; 9(3):57. <https://doi.org/10.3390/metabo9030057> PMID: 30909447
31. Marsal K, Persson PH, Larsen T, Lilja H, Selbing A, Sultan B. Intrauterine growth curves based on ultrasonically estimated foetal weights. *Acta Paediatr*. 1996; 85(7):843–8. Epub 1996/07/01. <https://doi.org/10.1111/j.1651-2227.1996.tb14164.x> PMID: 8819552.
32. Ruoppolo M, Scolamiero E, Caterino M, Mirisola V, Franconi F, Campesi I. Female and male human babies have distinct blood metabolomic patterns. *Mol Biosyst*. 2015; 11(9):2483–92. Epub 2015/07/04. <https://doi.org/10.1039/c5mb00297d> PMID: 26140445.
33. Lau C-HE, Siskos AP, Maitre L, Robinson O, Athersuch TJ, Want EJ, et al. Determinants of the urinary and serum metabolome in children from six European populations. *BMC Medicine*. 2018; 16(1):202. <https://doi.org/10.1186/s12916-018-1190-8> PMID: 30404627
34. Hashimoto F, Nishiumi S, Miyake O, Takeichi H, Chitose M, Ohtsubo H, et al. Metabolomics analysis of umbilical cord blood clarifies changes in saccharides associated with delivery method. *Early Hum Dev*. 2013; 89(5):315–20. Epub 2012/11/28. <https://doi.org/10.1016/j.earlhumdev.2012.10.010> PMID: 23178109.
35. Perng W, Rifas-Shiman SL, McCulloch S, Chatzi L, Mantzoros C, Hivert M-F, et al. Associations of cord blood metabolites with perinatal characteristics, newborn anthropometry, and cord blood hormones in project viva. *Metabolism*. 2017; 76:11–22. Epub 2017/07/17. <https://doi.org/10.1016/j.metabol.2017.07.001> PMID: 28987236.
36. Shokry E, Marchioro L, Uhl O, Bermúdez MG, García-Santos JA, Segura MT, et al. Investigation of the impact of birth by cesarean section on fetal and maternal metabolism. *Archives of Gynecology and Obstetrics*. 2019; 300(3):589–600. <https://doi.org/10.1007/s00404-019-05213-w> PMID: 31201538
37. Abalos E, Oladapo OT, Chamillard M, Díaz V, Pasquale J, Bonet M, et al. Duration of spontaneous labour in 'low-risk' women with 'normal' perinatal outcomes: A systematic review. *European Journal of*

- Obstetrics and Gynecology and Reproductive Biology. 2018; 223:123–32. <https://doi.org/10.1016/j.ejogrb.2018.02.026> PMID: 29518643
38. Braekke K, Ueland PM, Harsem NK, Karlsen A, Blomhoff R, Staff AC. Homocysteine, Cysteine, and Related Metabolites in Maternal and Fetal Plasma in Preeclampsia. *Pediatric research*. 2007; 62:319. <https://doi.org/10.1203/PDR.0b013e318123fba2> PMID: 17622947
 39. Zanardo V, Caroni G, Burlina A. Higher homocysteine concentrations in women undergoing caesarean section under general anesthesia. *Thromb Res*. 2003; 112(1–2):33–6. Epub 2004/03/12. <https://doi.org/10.1016/j.thromres.2003.11.004> PMID: 15013270.
 40. Nakada EM, Bhakta NR, Korwin-Mihavics BR, Kumar A, Chamberlain N, Bruno SR, et al. Conjugated bile acids attenuate allergen-induced airway inflammation and hyperresponsiveness by inhibiting UPR transducers. *JCI Insight*. 2019; 4(9). <https://doi.org/10.1172/jci.insight.98101> PMID: 31045581
 41. Sipka S, Bruckner G. The immunomodulatory role of bile acids. *Int Arch Allergy Immunol*. 2014; 165(1):1–8. Epub 2014/10/04. <https://doi.org/10.1159/000366100> PMID: 25277277.
 42. Spergel JM. From atopic dermatitis to asthma: the atopic march. *Ann Allergy Asthma Immunol*. 2010; 105(2):99–106; quiz 7–9, 17. Epub 2010/08/03. <https://doi.org/10.1016/j.anaai.2009.10.002> PMID: 20674819.
 43. Baiz N, Just J, Chastang J, Forhan A, de Lauzon-Guillain B, Magnier A-M, et al. Maternal diet before and during pregnancy and risk of asthma and allergic rhinitis in children. *Allergy, Asthma & Clinical Immunology*. 2019; 15(1):40. <https://doi.org/10.1186/s13223-019-0353-2> PMID: 31285746
 44. Jonsson K, Green M, Barman M, Sjöberg A, Brekke HK, Wold AE, et al. Diet in 1-year-old farm and control children and allergy development: results from the FARMFLORA birth cohort. *Food Nutr Res*. 2016; 60:32721. Epub 2016/08/19. <https://doi.org/10.3402/fnr.v60.32721> PMID: 27534847; PubMed Central PMCID: PMC4989273.
 45. Staley C, Weingarden AR, Khoruts A, Sadowsky MJ. Interaction of gut microbiota with bile acid metabolism and its influence on disease states. *Appl Microbiol Biotechnol*. 2017; 101(1):47–64. Epub 2016/11/25. <https://doi.org/10.1007/s00253-016-8006-6> PMID: 27888332.
 46. Vassallo MF, Banerji A, Rudders SA, Clark S, Mullins RJ, Camargo CA Jr. Season of birth and food allergy in children. *Annals of allergy, asthma & immunology: official publication of the American College of Allergy, Asthma, & Immunology*. 2010; 104(4):307–13. <https://doi.org/10.1016/j.anaai.2010.01.019> PMID: 20408340.
 47. Ramesh M, Cohen L, Kachirayan M, Rajan TV. Relationship Of Date Of Birth And Seasonal Allergies. *Journal of Allergy and Clinical Immunology*. 2010; 125(2):AB32. <https://doi.org/10.1016/j.jaci.2009.12.159>
 48. Chen Y, Lyu J, Xia Y, Zhu J, Tong S, Yin Y, et al. Effect of Maternal Sleep, Physical Activity and Screen Time during Pregnancy on the Risk of Childhood Respiratory Allergies: a Gender-specific Study. *Research Square*; 2020.
 49. Barman M, Murray F, Bernardi AI, Broberg K, Bölte S, Hesselmar B, et al. Nutritional impact on Immunological maturation during Childhood in relation to the Environment (NICE): a prospective birth cohort in northern Sweden. *BMJ Open*. 2018; 8(10). <https://doi.org/10.1136/bmjopen-2018-022013> PMID: 30344169