

Response to: 'Correspondence on 'Concomitant use of oral glucocorticoids and proton pump inhibitors and risk of osteoporotic fractures among patients with rheumatoid arthritis: a population-based cohort study" by Zheng

Citation for published version (APA):

Abtahi, S., Driessen, J. H. M., Burden, A. M., Souverein, P. C., van den Bergh, J. P., van Staa, T. P., Boonen, A., & de Vries, F. (2021). Response to: 'Correspondence on 'Concomitant use of oral glucocorticoids and proton pump inhibitors and risk of osteoporotic fractures among patients with rheumatoid arthritis: a population-based cohort study" by Zheng. *Annals of the Rheumatic Diseases*. <https://doi.org/10.1136/annrheumdis-2021-219856>

Document status and date:

E-pub ahead of print: 08/02/2021

DOI:

[10.1136/annrheumdis-2021-219856](https://doi.org/10.1136/annrheumdis-2021-219856)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Download date: 06 Oct. 2022

Response to: 'Correspondence on 'Concomitant use of oral glucocorticoids and proton pump inhibitors and risk of osteoporotic fractures among patients with rheumatoid arthritis: a population-based cohort study' by Gong and Zhang and Lin *et al*

We read with much interest the two correspondences on our recently published paper by Lin *et al* and Gong and Zhang in the *Annals of the Rheumatic Diseases*.¹⁻³ We conducted a retrospective cohort study to evaluate the association between concomitant use of two commonly used medications in patients with rheumatoid arthritis (RA), that is, oral glucocorticoids (GCs) and proton pump inhibitors (PPIs), and risk of osteoporotic (OP) fractures. We used data from one of the largest primary care databases in the world, the Clinical Practice Research Datalink (CPRD). While this database is a rich source, it does have some inherent limitations.

The first question of both correspondences addresses one of these limitations of the CPRD, that is, the absence of any direct measure of RA disease activity in the database, such as the Disease Activity Score in 28 joints. The values of C reactive protein (CRP) were also not available for all patients, and if available, not for the whole follow-up period. Therefore, we decided not to include CRP in the analyses, considering our on-treatment study design and importance of having information on covariates during all person-times from patients for the entire follow-up. We were aware of this limitation and mentioned this in the paper. We also reported our strategy to overcome this by using proxies of RA disease activity, such as use of conventional synthetic disease-modifying antirheumatic drugs (csDMARDs), and five analgesics (including non-selective non-steroidal anti-inflammatory drugs, selective cyclooxygenase-2 inhibitors, paracetamol, opioids and tramadol). Based on the European Alliance of Associations for Rheumatology and the UK National Institute for Health and Care Excellence guidelines on RA management,^{4,5} we assume that patients with RA would have been prescribed csDMARDs and analgesics according to the severity of the inflammation and pain that they had experienced. Thus, by considering these six covariates in the Cox regression model, we aimed to adjust (at least partly) for RA disease severity in our analyses.

Continuing with the effect of disease severity indicators on our risk estimates, questioned by Gong and Zhang, the frequency of use of some types of analgesics, such as opioids and tramadol, was higher among concomitant GC and PPI users at baseline versus non-users (data shown in table 1 in the manuscript). Nevertheless, we had an on-treatment study design and adjusted our final Cox regression model for use of analgesics and csDMARDs during follow-up, so any confounding effect from these proxies of disease severity have been taken into account in the fully adjusted analyses.

We agree with Lin *et al* that postmenopausal status is an important predisposing factor to primary osteoporosis and the resulting fractures. But there is little evidence to think that a postmenopausal status is associated with higher use of oral GCs or PPIs, as by definition a true confounder should be associated with both the exposure and outcome and does not lie in the causal pathway of this association.^{6,7} We included hormone replacement therapy (HRT) in our primary list of covariates as an indirect measure of postmenopausal status. Not so many patients were on HRT at baseline (4.3% of concomitant users and 2.6%

of non-users), and our univariate Cox regression models showed that HRT was not associated with a significant change in the beta coefficient of the main association (ie, >5%).

The association between statin use and fracture risk was not uniform in the literature. Meta-analyses of randomised controlled trials (RCTs) showed a neutral effect from statins on fracture risk.⁸ But, many observational studies in the past two decades claimed surprisingly beneficial effects of statins on various adverse outcomes such as cancers, diabetes, respiratory diseases and fractures, which has been later found out to be explained by time-related biases or healthy-user bias.⁹⁻¹² Based on the stronger evidence from RCTs, we did not feel a necessity to add statins as a covariate in our study.

Regarding the last point by Lin *et al* on our inclusion and exclusion criteria for establishing our patient population, we should clarify that we used Thomas algorithm (updated by Muller *et al*) for selecting our study population within the CPRD and we only included those patients from the CPRD who had a definite RA diagnosis based on Thomas algorithm. The 17 111 patients who were excluded in the second step, as shown in figure 1 in the manuscript, were either those without a definite RA diagnosis or those who had a previous OP fracture or used oral GCs and/or PPIs in the 1-year look-back period.

Answering the last question by Gong and Zhang on GC dose changes and stratification of GC use, we can explain that the average duration of drug use was estimated based on the time between first and last drug exposure during the follow-up, and for this we did not consider the drug holidays or intervals. However, we carefully calculated the average daily use of oral GCs and PPIs, cumulative use of oral GCs and continuous duration of PPIs during follow-up, while accounting for drug dose changes. The average daily dose of oral GCs was calculated at each current GC use interval. When the average daily dose was ≤ 7.5 mg prednisolone equivalent dose (PED)/day, an interval (not patient) was classified as low GC use, when this was 7.6–14.9 mg PED/day, an interval was classified as medium GC use, and when the average daily dose was ≥ 15 mg PED/day, an interval was classified as high GC use. So, patients could move between the groups, depending on what was the most recent average daily GC exposure.

Shahab Abtahi ^{1,2,3}, Johanna H M Driessen,^{1,2,3,4} Andrea M Burden ^{1,5}, Patrick C Souverein,² Joop P van den Bergh,^{6,7,8} Annelies Boonen ^{6,9}

¹Department of Clinical Pharmacy and Toxicology, Maastricht University Medical Centre, Maastricht, The Netherlands

²Division of Pharmacoepidemiology and Clinical Pharmacology, Utrecht Institute for Pharmaceutical Sciences, Utrecht University, Utrecht, The Netherlands

³Cardiovascular Research Institute Maastricht (CARIM), Maastricht University, Maastricht, The Netherlands

⁴NUTRIM School for Nutrition and Translational Research in Metabolism, Maastricht University Medical Centre, Maastricht, The Netherlands

⁵Institute of Pharmaceutical Sciences, Department of Chemistry and Applied Biosciences, ETH Zurich, Zurich, Switzerland

⁶Department of Internal Medicine, Division of Rheumatology, Maastricht University Medical Center, Maastricht, The Netherlands

⁷Faculty of Medicine and Life Sciences, Hasselt University, Hasselt, Belgium

⁸Department of Internal Medicine, VieCuri Medical Centre, Venlo, The Netherlands

⁹Care and Public Health Research Institute (CAPHRI), Maastricht University, Maastricht, The Netherlands

Correspondence to Dr Johanna H M Driessen, Division of Pharmacoepidemiology and Clinical Pharmacology, Utrecht Institute for Pharmaceutical Sciences, Utrecht University, PO BOX 80082, 3508 TB, Utrecht, The Netherlands; j.h.m.driessen@uu.nl

Handling editor Josef S Smolen

Twitter Shahab Abtahi @Sh_Abtahi and Andrea M Burden @ETH_PharmEpi

Contributors SA wrote the manuscript. All authors were involved in editing or quality control, are accountable for their own contribution, in addition to all aspects of the manuscript, and have approved the submitted version of the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests JvdB reports grants from Amgen, Eli Lilly and UCB, outside the submitted work. AB received research grants for her department from Celgene and AbbVie and speakers' fees or honoraria from Sandoz, UCB, Lilly, Novartis, Biogen and Gilead, not related to the current study. SA, JD, AMB, and PCS have no competing interests.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Abtahi S, Driessen JHM, Burden AM, *et al.* *Ann Rheum Dis* Epub ahead of print: [please include Day Month Year]. doi:10.1136/annrheumdis-2021-220477

Received 15 May 2021

Accepted 17 May 2021



► <http://dx.doi.org/10.1136/annrheumdis-2021-220453>

► <http://dx.doi.org/10.1136/annrheumdis-2021-220494>

Ann Rheum Dis 2021;**0**:1–2. doi:10.1136/annrheumdis-2021-220477

ORCID iDs

Shahab Abtahi <http://orcid.org/0000-0003-0482-5563>

Andrea M Burden <http://orcid.org/0000-0001-7082-8530>

Annelies Boonen <http://orcid.org/0000-0003-0682-9533>

REFERENCES

- 1 Abtahi S, Driessen JHM, Burden AM, *et al.* Concomitant use of oral glucocorticoids and proton pump inhibitors and risk of osteoporotic fractures among patients with rheumatoid arthritis: a population-based cohort study. *Ann Rheum Dis* 2021;**80**:423–31.
- 2 Lin TK, Pan LF, Jong GP. Correspondence on 'Concomitant use of oral glucocorticoids and proton pump inhibitors and risk of osteoporotic fractures among patients with rheumatoid arthritis: a population-based cohort study'. *Ann Rheum Dis* 2021.
- 3 Gong Z, Zhang F. Correspondence on 'Concomitant use of oral glucocorticoids and proton pump inhibitors and risk of osteoporotic fractures among patients with rheumatoid arthritis: a population-based cohort study'. *Ann Rheum Dis* 2021.
- 4 Smolen JS, Landewé RBM, Bijlsma JWJ, *et al.* EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Ann Rheum Dis* 2020;**79**:685–99.
- 5 Allen A, Carville S, McKenna F, *et al.* Diagnosis and management of rheumatoid arthritis in adults: summary of updated NICE guidance. *BMJ* 2018;**362**:k3015.
- 6 Strom BL, Kimmel SE, Hennessy S. Chapter 2, Study designs available for pharmacoepidemiologic studies. In: *Textbook of pharmacoepidemiology*. 2nd ed. West Sussex, UK: Wiley-Blackwell, 2013: 18–20.
- 7 European Medicines Agency. The European network of centres for pharmacoepidemiology and pharmacovigilance (ENCePP). guide on methodological standards in pharmacoepidemiology (revision 8), 2010. Available: http://www.encepp.eu/standards_and_guidance
- 8 Wang Z, Li Y, Zhou F, *et al.* Effects of statins on bone mineral density and fracture risk: a PRISMA-compliant systematic review and meta-analysis. *Medicine* 2016;**95**:e3042.
- 9 Suissa S, Dell'Aniello S. Time-Related biases in pharmacoepidemiology. *Pharmacoepidemiol Drug Saf* 2020;**29**:1101–10.
- 10 Suissa S, Dell'aniello S, Vahey S, *et al.* Time-Window bias in case-control studies: statins and lung cancer. *Epidemiology* 2011;**22**:228–31.
- 11 de Vries F, de Vries C, Cooper C, *et al.* Reanalysis of two studies with contrasting results on the association between statin use and fracture risk: the general practice research database. *Int J Epidemiol* 2006;**35**:1301–8.
- 12 Klop C, Driessen JHM, de Vries F. Statin use and reduced cancer-related mortality. *N Engl J Med* 2013;**368**:574.