



Title	Intrauterine infection with Mycobacterium avium subsp. paratuberculosis in pregnant cattle diagnosed with Johne's disease
Author(s)	Taniguchi, Yukiko; Konnai, Satoru; Sakakibara, Shinichi; Yamamoto, Ayano; Okagawa, Tomohiro; Maekawa, Naoya; Murata, Shiro; Ohashi, Kazuhiko
Citation	Japanese Journal of Veterinary Research, 69(1), 51-55
Issue Date	2021-02
DOI	10.14943/jjvr.69.1.51
Doc URL	http://hdl.handle.net/2115/80620
Type	bulletin (article)
File Information	JJVR69-1_51-55_YukikoTaniguchi.pdf



[Instructions for use](#)

Intrauterine infection with *Mycobacterium avium* subsp. *paratuberculosis* in pregnant cattle diagnosed with Johne's disease

Yukiko Taniguchi¹⁾, Satoru Konnai^{2,3,*}, Shinichi Sakakibara¹⁾,
Ayano Yamamoto¹⁾, Tomohiro Okagawa³⁾, Naoya Maekawa³⁾,
Shiro Murata^{2,3)} and Kazuhiko Ohashi^{2,3)}

¹⁾Hokkaido Tokachi Livestock Hygiene Service Center, Obihiro, 089-1182, Japan

²⁾Department of Disease Control, Faculty of Veterinary Medicine, Hokkaido University, Sapporo, 060-0818, Japan

³⁾Department of Advanced Pharmaceutics, Faculty of Veterinary Medicine, Hokkaido University, Sapporo, 060-0818, Japan

Received for publication, September 17, 2020; accepted, October 22, 2020

Abstract

Johne's disease (JD) is caused by infection with *Mycobacterium avium* subsp. *paratuberculosis* (MAP). We confirmed the intrauterine infection of MAP in 22 pregnant cattle diagnosed with JD in Hokkaido, Japan. MAP was isolated from the umbilical cord (3/22: 13.6%) or caruncle (6/22: 27.3%) derived from the pregnant dams. Furthermore, dams with MAP from which MAP was isolated were also found to have a high amount of MAP or detected bacterial load in their feces. Fetuses of the tested dams indicated positive polymerase chain reaction (PCR) results for the MAP gene (17/22:77.3%) in several tissues. MAP was isolated from the PCR-positive sites of dams detected with high levels of bacteria (6/22: 27.3%). These results indicate that MAP infection in pregnant cattle must be prevented as it is important for JD control.

Key Words: *Mycobacterium avium* subsp. *paratuberculosis*, intrauterine infection, vertical transmission

Johne's disease (JD) is caused by infection with *Mycobacterium avium* subsp. *paratuberculosis* (MAP). JD causes chronic enteritis, resulting in decreased milk production, wasting, and eventual death in ruminants, including cattle⁶⁾. Recently, the number of cattle with JD has been increasing in Japan, especially in Hokkaido (https://www.maff.go.jp/j/syouan/douei/kansi_densen/kansi_densen.html). Basically, MAP is transmitted orally among the cattle herd via contaminated materials, such as feces and milk

from infected animals. Therefore, based on the MAP transmission route and because there is no effective vaccine for MAP infection⁵⁾, recommended disease control measures include prompt removal of infected animals and pasteurization of milk that is fed to calves.

In cases of MAP-infected pregnant dams, MAP is vertically transmitted to newborn calves and fetuses in the dam¹¹⁾. To date, two possible routes of vertical MAP transmission have been recognized. One is oral bacterial acquisition in

* Corresponding author: Satoru Konnai

Mailing address: Department of Disease Control, Faculty of Veterinary Medicine, Hokkaido University, Sapporo, Hokkaido 060-0818, Japan

Phone: (81) 11 706 5216, Fax: (81) 11 706 5217, E-mail: konnai@vetmed.hokudai.ac.jp

doi: 10.14943/jjvr.69.1.51

the newborn via the maternal colostrum or milk containing MAP-infected cells, and another is intrauterine infection of the fetus in the MAP-infected pregnant dam³⁾. Among reports on the vertical transmission of MAP, fetal infection was first reported by Alexejeff-Goloff²⁾. The presence of MAP was confirmed in fetal membranes, blood, liver, and other fetal tissues from a clinically affected cow with JD²⁾. Similar findings concerning fetal infection have been reported since the first report¹¹⁾.

Thus, in addition to the prevention of oral MAP transmission, the recognition of risk of intrauterine MAP infection in pregnant cattle also is important to control the incidence of JD. We investigated 22 fetuses from infected pregnant dams with a high bacterial load in Hokkaido to gather further evidences concerning the intrauterine transmission of MAP.

In the Tokachi District of Hokkaido, between July 2016 and March 2017, 22 pregnant cattle (14 Japanese Black cattle and 8 Holstein Friesian cattle) that were legislatively diagnosed with JD (>0.001 pg/well [2.5 µL] of MAP DNA) were investigated. The rectal stool, umbilical cord, caruncle, umbilical blood, and amniotic fluid were collected as maternal specimens. The fetal specimens were carefully collected from the heart, lung, kidney, liver, spleen, ilea, jejunum, colon, and meconium. All specimens except the maternal rectal feces were collected immediately at rendering plants after euthanasia. The collected maternal and fetal samples were used for MAP detection and isolation.

Briefly, the bacterial load of MAP was confirmed using real-time polymerase chain reaction (PCR), Johne-spin (FASMAC, Atsugi, Japan) for DNA extraction, and Johne's Gene test kit "KS" (Kyoritsu Seiyaku Co., Tokyo, Japan) targeting the MAP-specific gene *IS900* according to the manufacturer's instructions. MAP was further confirmed via bacterial isolation using a Harold's medium with mycobactin (Kyoritsu Seiyaku Co.) according to the manufacturer's instructions.

MAP was isolated from six (27.3%) maternal specimens of caruncle and umbilical cord (Dam IDs 1, 3, 4) or either caruncle or umbilical cord (Dam IDs 2, 10, 11) among the 22 tested pregnant dams (Table 1). All MAP infections isolated from the maternal specimens of pregnant dams showed extremely high bacterial loads in the feces. Detection was easier and the detected colony-forming units (cfu) were higher in the caruncle than in the umbilical cord. However, 17 (77.3%) of 22 fetuses derived from the pregnant dams showed positive PCR results for the *MAP* gene in any tested tissues, and MAP was isolated in six (27.3%) fetuses. All MAP-isolated fetal tissues were from PCR-positive sites. Interestingly, three of six MAP-isolated fetuses were derived from dams (Dam IDs 1, 3, 4) with high bacterial loads and MAP isolation from the caruncle plus umbilical cord. For the youngest fetus (4 months, from pregnant dam ID 3), MAP was isolated from three tissues (liver, spleen, and kidney). The remaining 3 MAP-isolated fetuses were obtained from dams (Dam IDs 7, 8, 9) with high bacterial loads in the feces; however, MAP was not isolated from the caruncle or umbilical cord. On the other hand, MAP was not detected and isolated in the umbilical blood and amniotic fluid derived from pregnant dams and meconium in this study (data not shown).

To date, there are many reports on the intrauterine transmission of MAP in cattle with JD. However, it is likely that these reports varied on the rarity and risk of intrauterine transmission of MAP in the dam¹¹⁾. Indeed, Whittington *et al.* (2009)¹¹⁾ reported the prevalence of fetal infection with MAP to be as high as 39% of cows with clinical signs of JD, which is higher than 8.6% of fetuses in cows without signs of the disease. Seitz *et al.* (1989) reported a high rate (26.4%) of intrauterine infection in a dairy herd⁷⁾, whereas Adaska *et al.* (2012) reported a low rate (4.3%)¹¹⁾.

In addition to the prevalence of MAP-infected animals in the herd¹¹⁾, the difference in findings may be attributable to different bacterial loads or disease progression in the infected cattle¹¹⁾. Indeed,

Table 1. Detection of *M. paratuberculosis* in the infected dams and fetus.

Dam ID	Breed	Age (years)	Bacterial load (pg/well)	Bacteria isolation (+(cfu)/-)			Fetus									
				Feces	Feces	Umbilical cord	Caruncle	Fetal age (months)	Bacteria isolation (+/-)	PCR (+/-)	Details of the detection (Tested tissue (Bacteria isolation(+/(cfu)/-) / PCR(+/-))					
											Lung	Heart	Liver	Spleen	Kidney	Intestines
1	J.B	11	185.000	+(UC)	+(1)	+(60)	5	+	+	-/+	-/+	+(1)/+	-/+	-/+	-/+	
2	J.B	11	119.000	+(UC)	-	+(12)	8	-	+	-/-	-/+	-/-	-/-	-/-	-/+	
3	J.B	4	26.900	+(UC)	+(7)	+(24)	4	+	+	-/+	-/-	+(1)/+	+(1)/+	+(1)/+	-/+	
4	J.B	5	8.010	+(UC)	+(1)	+(2)	9	+	+	-/+	-/-	+(1)/+	-/-	-/+	-/+	
5	J.B	9	7.980	NT	-	-	6	-	-	-/-	-/-	-/-	-/-	-/-	-/-	
6	J.B	11	3.740	+(UC)	-	-	8	-	+	-/-	-/-	-/+	-/-	-/-	-/+	
7	Hol	4	0.110	+(UC)	-	-	7	+	+	-/-	-/-	+(1)/+	-/-	-/-	-/+	
8	J.B	3	0.027	+(UC)	-	-	8	+	+	-/+	+(1)/+	-/-	-/-	-/+	-/+	
9	J.B	6	0.017	+(7)	-	-	6	+	+	-/-	-	-/-	-/-	-/-	+(4)/+	
10	Hol	5	0.013	+(UC)	-	+(1)	7.5	-	+	-/+	-/+	-/+	-/-	-/+	-/+	
11	J.B	10	0.013	+(30)	-	+(4)	9	-	+	-/-	-/+	-/-	-/-	-/-	-/+	
12	Hol	4	0.010	+(13)	-	-	4	-	+	-/-	-/-	-/-	-/-	-/-	-/+	
13	J.B	3	0.008	+(8)	-	-	6	-	-	-/-	-/-	-/-	-/-	-/-	-/-	
14	Hol	4	0.007	NT	-	-	7	-	+	-/-	-/-	-/-	-/+	-/-	-/-	
15	Hol	1	0.007	-	-	-	6	-	+	-/-	-/-	-/-	-/+	-/+	-/-	
16	J.B	9	0.005	-	-	-	4.5	-	+	-/-	-/+	-/-	-/-	-/-	-/+	
17	Hol	8	0.004	+(6)	-	-	5	-	+	-/-	-/-	-/-	-/-	-/+	-/-	
18	Hol	5	0.002	+(16)	-	-	6	-	-	-/-	-/-	-/-	-/-	-/-	-/+	
19	J.B	13	0.001	NT	-	-	7	-	+	-/+	-/-	-/-	-/-	-/-	-/-	
20	J.B	9	0.001	NT	-	-	4.5	-	-	-/-	-/-	-/-	-/-	-/-	-/-	
21	Hol	6	0.001	-	-	-	5	-	+	-/+	-/+	-/-	-/+	-/-	-/-	
22	J.B	8	0.001	NT	-	-	7	-	-	-/-	-/-	-/-	-/-	-/-	-/-	
Total					3/22	6/22		6/22	17/22							
n=22					(13.6%)	(27.3%)		(27.3%)	(77.3%)							

UC: uncountable, NT: not tested,

Sweeney *et al.* (1992) reported that MAP was isolated from fetal tissues in 5 of 58 (17.8%) cattle and all 5 positive fetuses were from dams with heavy fecal shedding⁹. In our study, intrauterine infection was confirmed in pregnant cattle with different bacterial loads. As expected, the risk of MAP intrauterine transmission tended to be higher in infected dams with high bacterial loads as shown in previous reports^{9,11}. Sonawane *et al.* (2009) reported that the isolation rate of MAP from blood increased in infected cattle with severe lesions⁸. Therefore, MAP might migrate to the blood of infected cattle with the progression of the disease state, infecting the fetus hematogenously through the umbilical cord⁴. Indeed, in our study, MAP was isolated from the umbilical cord or caruncle as noted in previous reports¹¹. However, although hematogenous infection of MAP through

the umbilical cord is possible, the bovine placenta is classified histologically as a syndesmochorial placenta with a strong junction. Therefore, the possibility of hematogenous infection of MAP through the umbilical cord must be clarified. In addition to this issue, although a high level of MAP DNA was detected (77.3%) in the fetal sample, the intrauterine infection was established when MAP was isolated from the fetal specimen because the positive PCR results included those derived from dead cells and gene fragments. However, it is possible for the pregnant cattle to have an intrauterine infection with a low bacteria load. Further clarification is required to determine the accurate risk of intrauterine infection in MAP-infected cattle.

In conclusion, our study presents direct evidence of intrauterine MAP infection. In

Hokkaido, Japan, the number of cattle with JD is increasing because there is no effective treatment or vaccine for MAP infection. Recently, we reported that the shedding level of MAP DNA in feces was related to the histopathologic classification of disease progression in cattle with JD¹⁰. As mentioned, such cattle with high MAP shedding or clinical signs could be at a high risk for intrauterine infection^{9,11}. Furthermore, it is notable that fetal infection with MAP in utero tended to demonstrate high shedding with clinical signs¹¹. Thus, Whittington and Windsor explained the possibility of intrauterine infection with a high risk of progressive MAP infection because the uncontrolled infection could retard the success of disease control programs for JD based on pasteurization of milk and removal of infected cattle¹¹. According to a report, the incidence of intrauterine transmission depends on within-herd prevalence of MAP infection¹. Although intrauterine transmission of MAP could not be prevented, the removal of suspected cattle with JD is the most important issue to prevent vertical transmission of MAP. These results indicated that prevention of MAP in the pregnant cattle is important for JD control.

Acknowledgements

This work was supported by JSPS KAKENHI grant number 19KK0172 [to S.K.], grants from the Project of the NARO, Bio-oriented Technology Research Advancement Institution (Research Program on Development of Innovative Technology 26058 BC [to S.K.] and Special Scheme Project on Regional Developing Strategy, Grant 16817557 [to S.K.]), and Regulatory research projects for food safety, animal health and plant protection (JPJ008617.17935709) funded by the Ministry of Agriculture, Forestry and Fisheries of Japan. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- 1) Adaska JM, Whitlock RH. Low rate of detectable *in utero* transmission of *Mycobacterium avium* subspecies *paratuberculosis* in a dairy herd with a low prevalence of Johne's disease. *J Vet Diagn Invest* 24, 153-155, 2012.
- 2) Alexejeff-Goloff NA. Zur Frage der Pathogenese und Bazillenausscheidung bei Rinderparatuberkulose. *Zeitschrift für Infektionskrankheiten, Parasitaerkrankheiten und Hygiene der Haustiere* 36, 313-317, 1929. Abstracted in *Journal of Comparative Pathology and Therapeutics* 48, 81-82, 1935.
- 3) Barkema HW, Orsel K, Nielsen SS, Koets AP, Rutten VPMG, Bannantine JP, Keefe GP, Kelton DF, Wells SJ, Whittington RJ, Mackintosh CG, Manning EJ, Weber MF, Heuer C, Forde TL, Ritter C, Roche S, Corbett CS, Wolf R, Griebel PJ, Kastelic JP, De Buck J. Knowledge gaps that hamper prevention and control of *Mycobacterium avium* subspecies *paratuberculosis* infection. *Transbound Emerg Dis* 65, 125-148, 2018.
- 4) Chiodini RJ, Van Kruningen HJ, Merkal RS. Ruminant paratuberculosis (Johne's disease): the current status and future prospects. *Cornell Vet* 74, 218-262, 1984.
- 5) Clarke CJ. The pathology and pathogenesis of paratuberculosis in ruminants and other species. *J Comp Pathol* 116, 217-261, 1997.
- 6) Ott SL, Wells SJ, Wagner BA. Herd-level economic losses associated with Johne's disease on US dairy operations. *Prev Vet Med* 40, 179-192, 1999.
- 7) Seitz SE, Heider LE, Hueston WD, Bech-Nielsen S, Rings DM, Spangler L. Bovine fetal infection with *Mycobacterium paratuberculosis*. *J Am Vet Med Assoc* 194, 1423-1426, 1989.
- 8) Sonawane GG, Tripathi BN. Comparison of a quantitative real-time polymerase chain reaction (qPCR) with conventional PCR, bacterial culture and ELISA for detection of

- Mycobacterium avium* subsp. *paratuberculosis* infection in sheep showing pathology of Johne's disease. Springerplus 2, 45, 2013.
- 9) Sweeney RW, Whitlock RH, Rosenberger AE. *Mycobacterium paratuberculosis* isolated from fetuses of infected cows not manifesting signs of the disease. Am J Vet Res 53, 477-480, 1992.
 - 10) Taniguchi Y, Sakakibara SI, Fujihara M, Yagi A, Fujiyoshi S. The association between detection of *Mycobacterium avium* subsp. *paratuberculosis* DNA in feces and histopathological classification. J Vet Med Sci 82, 541-545, 2020.
 - 11) Whittington RJ, Windsor PA. In utero infection of cattle with *Mycobacterium avium* subsp. *paratuberculosis*: A critical review and meta-analysis. Vet J 179, 60-69, 2009.