

Publ. Mat. **48** (2004), 309–333

RIESZ TRANSFORMS ON GENERALIZED HEISENBERG GROUPS AND RIESZ TRANSFORMS ASSOCIATED TO THE CCR HEAT FLOW

FRANÇOISE LUST-PIQUARD

Abstract

Let $1 < q < \infty$. We prove that the Riesz transforms $R_k = X_k L^{-\frac{1}{2}}$ on a generalized Heisenberg group G satisfy $\left\| \left(\sum_{k=1}^K |R_k(f)|^2 \right)^{\frac{1}{2}} \right\|_{L^q(G)} \leq C(q, J) \|f\|_{L^q(G)}$ where K, J are respectively the dimensions of the first and second layer of the Lie algebra of G . We prove similar inequalities on Schatten spaces $S^q(H)$, with dimension free constants, for Riesz transforms associated to commuting inner $*$ -derivations D_k and a suitable substitute of the square function. An example is given by the derivations associated to n commuting pairs of operators (P_j, Q_j) on a Hilbert space H satisfying the canonical commutation relations $[P_j, Q_j] = iI_H$.

General introduction

This paper is divided in two parts: we solve similar problems in two different settings, using similar methods inspired by the first part of [P1], which contains a proof of the following classical inequalities [S]: for $1 < q < \infty$ and $f \in \mathcal{D}(\mathbb{R}^n)$,

$$D_q \|f\|_{L^q(\mathbb{R}^n)} \leq \left\| \left(\sum_{k=1}^n |R_k(f)|^2 \right)^{\frac{1}{2}} \right\|_{L^q(\mathbb{R}^n)} \leq C_q \|f\|_{L^q(\mathbb{R}^n)}$$

where

$$R_k(f) = \frac{\partial}{\partial x_k} L^{-\frac{1}{2}}(f), \quad L = - \sum_{k=1}^n \frac{\partial^2}{\partial x_k^2}$$

and the constants do not depend on n .

2000 *Mathematics Subject Classification.* 43A80, 46L50, 46L57.

Key words. Heat operator, Riesz transforms, H-groups, commuting $*$ -inner derivations.

The first part deals with Riesz transforms acting on $L^q(G)$, where G is a generalized Heisenberg group, and owes a lot to [CMZ].

The second part deals with Riesz transforms acting on the Schatten space $\mathcal{S}_q(H)$, associated to commuting $*$ -inner derivations on the algebra $K(H)$ of compact operators on a Hilbert space H ; an example is given by the inner derivations defined by $(P_j, Q_j)_{j=1}^n$, where P_j, Q_j satisfy the canonical commutation relation $[Q_j, P_j] = iI_H$ and the other commutators are zero.

We already used Pisier's method in other settings, see e.g. [LP2], [LP3]. However the difficulties arise at different steps in different applications.

Since the two settings we consider are very different, we present more precise introductions in each part.

Acknowledgement. We thank V. Georgescu for fruitful discussions on the setting of the second part. We also thank the referee for detecting a gap in the proof of Theorem 1.b in an earlier version of the paper.

1. Riesz transforms on generalized Heisenberg groups

1.1. Introduction.

For any stratified Lie group G , we denote by X_1, \dots, X_K a basis of the top layer of the Lie algebra \mathcal{G} of G , by

$$L = - \sum_{j=1}^K X_j^2$$

the subelliptic Kohn Laplacian on G , by

$$R_k = X_k L^{-\frac{1}{2}}, \quad 1 \leq k \leq K,$$

the Riesz transforms. The boundedness of each R_k on $L^q(G)$, $1 < q < \infty$, is known: the classical proof uses the homogeneity of its kernel and the singular integral results of [FS, Chapter 6]; see also Lemma 2 below. Our interest in this paper is to look for dimension free inequalities involving

$$\left\| \left(\sum_{k=1}^K |R_k(f)|^2 \right)^{\frac{1}{2}} \right\|_{L^q(G)}.$$

We first consider the case where G is a Heisenberg group \mathbb{H}_n and give a simpler proof of the main result of [CMZ]. We extend this result to the generalized Heisenberg groups $\mathbb{H}_{K,J}$ defined by Kaplan [K]; they are particular step two stratified Lie groups, and K, J denote respectively the dimensions of the first and second layer in the Lie algebra of $\mathbb{H}_{K,J}$. The Heisenberg group \mathbb{H}_n is the same as $\mathbb{H}_{2n,1}$.

Theorem 1. *Let $1 < q < \infty$, $\frac{1}{q} + \frac{1}{q'} = 1$.*

- a) **[CMZ]** *There exist constants C_q such that, for every $n \in \mathbb{N}^*$ and every $f \in \mathcal{D}(\mathbb{H}_n)$,*

$$C_{q'}^{-1} \|f\|_{L^q(\mathbb{H}_n)} \leq \left\| \left(\sum_{k=1}^{2n} |R_k(f)|^2 \right)^{\frac{1}{2}} \right\|_{L^q(\mathbb{H}_n)} \leq C_q \|f\|_{L^q(\mathbb{H}_n)}.$$

- b) *The same holds for generalized Heisenberg groups $\mathbb{H}_{K,J}$, with constants $C(q, J)$ which depend on q and J but not on K .*

It is a standard fact (see e.g. **[CMZ]**) that, in the above formula, for any Lie group G , the lefthand side inequality for q is an easy consequence of the righthand side one for q' , $\frac{1}{q} + \frac{1}{q'} = 1$.

Theorem 1 relies on Christ's study **[C]** of Hilbert transforms along curves for homogeneous nilpotent Lie groups, and on the use of dilations δ_t in order to get an expression of the convolution operator $e^{-\frac{1}{2}t^2L}$ involving the (heat) kernel p of $e^{-\frac{1}{2}L}$, namely (see Lemma 2 a))

$$(1) \quad e^{-\frac{1}{2}t^2L}(f)(\gamma) = \int_G f(\gamma\delta_t g^{-1})p(g) dg.$$

Theorem 1 also uses a formula (Lemma 2 b) below)

$$(2) \quad \sqrt{2\pi}XL^{-\frac{1}{2}}(f)(\gamma) = \int_G F(\gamma, g)(Xp)(g) dg$$

which holds on every stratified group G , X lying in the first layer of \mathcal{G} , F being a Hilbert transform of f . These ingredients are already in **[CMZ]**; they are reminiscent of the method of **[P1]**.

In the proof of Theorem 1 a), our improvement upon **[CMZ]** is that we do not use the explicit formula of p and avoid computations. Denoting $p = p^{(n)}$ when $G = \mathbb{H}_n$, we use only the following properties:

- (i) **[FS]** p is a positive function lying in $\mathcal{S}(G)$, and $\int_G p dg = 1$,
- (ii) $p^{(n)}(x_1, \dots, y_n, u)$ is radial with respect to (x_1, \dots, y_n) , i.e. depends on (r, u) ,
- (iii) $p^{(n)}(x_1, \dots, y_n, u) = p^{(1)}(x_1, y_1, u) *_u \dots *_u p^{(1)}(x_n, y_n, u)$,

where $*_u$ denotes convolution in \mathbb{R} with respect to the variable u . Property (ii) is used through the observation that

$$X_j p^{(n)} = x_j \frac{1}{r} \frac{\partial p^{(n)}}{\partial r} + 2y_j \frac{\partial p^{(n)}}{\partial u}.$$

In the proof of Theorem 1 b) we use the analogue for $\mathbb{H}_{K,J}$.

We recall that the heat kernels (i.e. the kernels of e^{-tL}) on \mathbb{H}_n and the non isotropic Heisenberg groups, or rather their Fourier transform with respect to u , were first computed in [G] and [H], then in several subsequent papers, by rather complicated methods; the heat kernels on general step two stratified groups were computed in [CY]. A more explicit formula for generalized Heisenberg groups $\mathbb{H}_{K,J}$ is given in [R], using the result for \mathbb{H}_n . Let us mention our computation of the heat kernels for isotropic or non isotropic Heisenberg groups and the free step two stratified groups $N_{n,2}$ [LP1], which is simpler than the previous ones and relies on the common starting point of [CMZ] and the present paper, namely formula (1).

1.2. Notation.

For a background on stratified groups (which are particular homogeneous groups) we refer to [FS, Chapter I]. We consider stratified Lie groups G equipped with their Haar measure, denoted by dg or $d\gamma$, which is Lebesgue measure on the underlying space \mathbb{R}^d . $\mathcal{D}(G)$ denotes the space of C^∞ compactly supported functions on G , $\mathcal{S}(G)$ denotes the Schwartz class. The convolution of two functions f, h lying in $\mathcal{S}(G)$ is defined by

$$f * h(\gamma) = \int_G f(\gamma g^{-1})h(g) dg.$$

The Lie algebra of left invariant vector fields on G is denoted by \mathcal{G} . For $X \in \mathcal{G}$,

$$(3) \quad X(f * p) = f * Xp.$$

The first layer of \mathcal{G} is the linear subspace which spans \mathcal{G} as a Lie algebra. We denote by σ the automorphism of G , corresponding to the automorphism σ of \mathcal{G} whose action on the first layer is

$$\sigma: X \longrightarrow -X.$$

G is equipped with a dilation $\delta_t, t > 0$, corresponding to the automorphism of \mathcal{G} whose action on the first layer is

$$(4) \quad \delta_t X = tX.$$

The induced action on functions $f: G \rightarrow \mathbb{R}$ is denoted by $(\delta_t f)(g) = f(\delta_t g)$, and [FS, I C]

$$(5) \quad X\delta_t f = t\delta_t(Xf).$$

A stratified Lie group G is said to be step two if the central layer of \mathcal{G} is the second one; denoting by X_1, \dots, X_K a basis of the first layer, and by U_1, \dots, U_J a basis of the second layer, it means that all

commutators $[X_j, X_k]$ belong to the linear span of the U_j 's and the other commutators are zero. Every $g \in G$ is defined in a unique way by $g = \exp\left(\sum_{k=1}^K x_k X_k + \sum_{j=1}^J u_j U_j\right)$ and we denote $g = (x, u) = (x_1, \dots, x_K, u_1, \dots, u_J)$. In this setting, the Haar measure on G , i.e. the Lebesgue measure on \mathbb{R}^{K+J} , is also denoted by $dx du$. In particular, $\sigma g = \sigma(x, u) = (-x, u)$, hence $\sigma g^{-1} = (x, -u)$, and $\delta_t(x, u) = (tx, t^2u)$.

We will use the following easy, but crucial, result which is standard when $J = 0$ and ψ is the gaussian density on \mathbb{R}^K .

Lemma 2. *Let ψ be a measurable function: $\mathbb{R}^{K+J} \rightarrow \mathbb{R}$ such that $\int_{\mathbb{R}^J} |\psi(x, u)| du$ only depends on $|x|$, $x = (x_k)_{k=1}^K \in \mathbb{R}^K$. Then*

a) *if $1 \leq q < \infty$, $a_k \in \mathbb{C}$,*

$$\left\| \sum_{k=1}^K a_k x_k \right\|_{L^q(|\psi| dx du)} = \left(\sum_{k=1}^K |a_k|^2 \right)^{\frac{1}{2}} \|x_1\|_{L^q(|\psi| dx du)};$$

b) *if $1 < q < \infty$, $\frac{1}{q} + \frac{1}{q'} = 1$, for every $h \in L^{q'}(|\psi| dx du)$,*

$$\left(\sum_{k=1}^K \left| \int_{\mathbb{R}^{K+J}} x_k h \psi dx du \right|^2 \right)^{\frac{1}{2}} \leq \|x_1\|_{L^{q'}(|\psi| dx du)} \|h\|_{L^q(|\psi| dx du)}.$$

Proof: For short, we write $L^q(|\psi|)$ instead of $L^q(|\psi| dx du)$. We use polar coordinates in \mathbb{R}^K , namely $a = (a_1, \dots, a_K) = |a| w$, $x = |x| v = r v$, with $w, v \in \Sigma_K$, $v = (v_1, \dots, v_K)$, here Σ_K denotes the unit sphere of \mathbb{R}^K and $d\sigma_K$ is the uniform measure on it, with total mass the area of Σ_K .

a) Since $\int_{\mathbb{R}^J} |\psi| du$ only depends on r ,

$$\begin{aligned} \left\| \sum_{k=1}^K a_k x_k \right\|_{L^q(|\psi|)}^q &= |a|^q \int_0^\infty r^q \left(\int_{\mathbb{R}^J} |\psi| du \right) r^{K-1} dr \int_{\Sigma_K} |\langle w, v \rangle|^q d\sigma_K(v) \\ &= |a|^q \int_0^\infty r^q \left(\int_{\mathbb{R}^J} |\psi| du \right) r^{K-1} dr \int_{\Sigma_K} |v_1|^q d\sigma_K(v) \\ &= |a|^q \|x_1\|_{L^q(|\psi|)}^q. \end{aligned}$$

b) This follows from a): indeed, by Hölder inequality,

$$\begin{aligned} \left(\sum_{k=1}^K \left| \int x_k h \psi \, dx \, du \right|^2 \right)^{\frac{1}{2}} &= \sup_{|a|=1} \left| \int \left(\sum_{k=1}^K a_k x_k \right) h \psi \, dx \, du \right| \\ &\leq \|h\|_{L^q(|\psi|)} \sup_{|a|=1} \left\| \sum_{k=1}^K a_k x_k \right\|_{L^{q'}(|\psi|)} \\ &= \|h\|_{L^q(|\psi|)} \|x_1\|_{L^{q'}(|\psi|)}. \quad \square \end{aligned}$$

Let G be a stratified Lie group; to $g \in G$ we associate the curve: $\mathbb{R} \rightarrow G$

$$\begin{aligned} g(t) &= \delta_t g \quad \text{for } t \geq 0, \\ g(t) &= \delta_{|t|} \sigma g \quad \text{for } t < 0. \end{aligned}$$

In particular, if G is step two, the curve associated to $g = (x, u)$ is

$$g(t) = (tx, t^2u), \quad t \in \mathbb{R}.$$

The Hilbert transform of f along the curve $g(t)$ is

$$\begin{aligned} F(\gamma, g) &= \text{pv} \int_{-\infty}^{\infty} f(\gamma g(t)^{-1}) \frac{dt}{t} \\ &= \lim_{\varepsilon \rightarrow 0^+} \int_{\varepsilon < |t| < \varepsilon^{-1}} f(\gamma g(t)^{-1}) \frac{dt}{t} \\ &= \lim_{\varepsilon \rightarrow 0^+} F(\gamma, g, \varepsilon). \end{aligned}$$

The function $F(\gamma, g)$ is well defined on $G \times G$ because $f \in \mathcal{D}(G)$. $F(\gamma, g, \varepsilon)$ is called the truncated Hilbert transform.

The map $h_t^g: f \rightarrow f(\cdot g(t)^{-1})$, $t \in \mathbb{R}$, $L^\infty(G) \rightarrow L^\infty(G)$, is a $*$ -homomorphism of the $*$ -algebra $L^\infty(G)$, but $\{h_t^g\}_{t \in \mathbb{R}}$ is not a one parameter group. This makes an important difference with the settings of [LP1], [LP2], [LP3] and the second part of this paper.

The next result comes from [CMZ, Proof of Lemmas 1 and 5]. For the sake of completeness we give a more precise proof.

Lemma 3. *Let G be a stratified Lie group and $f \in \mathcal{D}(G)$. Then*

$$(1) \quad e^{-\frac{1}{2}t^2L}(f)(\gamma) = \int_G f(\gamma\delta_t g^{-1})p(g) dg, \quad t > 0,$$

and, for X in the first layer of \mathcal{G} , $d\gamma$ a.s.,

$$(2) \quad \sqrt{2\pi}XL^{-\frac{1}{2}}(f)(\gamma) = \int_G F(\gamma, g)(Xp)(g) dg,$$

where $F(\gamma, g)$ is the Hilbert transform of f along the curve $g(t)$, $g \in G$.

For $1 < q < \infty$, the Riesz transforms satisfy

$$\left\| \sqrt{2\pi}XL^{-\frac{1}{2}}(f) \right\|_{L^q(G)} \leq h_q \|Xp\|_{L^1(G)} \|f\|_{L^q(G)}.$$

Proof: Formula (1) holds true for $t = 1$ by definition of p . By (4) (see also [LP1]),

$$e^{-\frac{1}{2}t^2L} = \delta_{t^{-1}}e^{-\frac{1}{2}L}\delta_t,$$

hence, by (5),

$$(1) \quad e^{-\frac{1}{2}t^2L}(f)(\gamma) = \delta_{t^{-1}}[(f \circ \delta_t) * p(\gamma)] = \int_G f(\gamma\delta_t g^{-1})p(g) dg.$$

By (5), (1) and (3),

$$Xe^{-\frac{1}{2}t^2L}(f) = t^{-1}\delta_{t^{-1}}X[(f \circ \delta_t) * p] = t^{-1}\delta_{t^{-1}}[(f \circ \delta_t) * (Xp)].$$

The automorphism σ maps L to L , so $p = p \circ \sigma$ and

$$Xp = X(p \circ \sigma) = -(Xp) \circ \sigma.$$

For $h \in \mathcal{D}(G)$, since $\sigma^2g = g$ and σ is measure preserving,

$$\begin{aligned} h * (Xp)(\gamma) &= \int_G h(\gamma g^{-1})(Xp)(g) dg \\ &= - \int_G h(\gamma g^{-1})(Xp)(\sigma g) dg \\ &= - \int_G h(\gamma \sigma g^{-1})(Xp)(g) dg. \end{aligned}$$

In particular, for $f \in \mathcal{D}(G)$,

$$2Xe^{-\frac{1}{2}t^2L}(f)(\gamma) = t^{-1} \int_G [f(\gamma\delta_t g^{-1}) - f(\gamma\delta_t \sigma g^{-1})](Xp)(g) dg.$$

Since

$$\sqrt{\frac{\pi}{2}}XL^{-\frac{1}{2}}(f) = \int_0^\infty Xe^{-\frac{1}{2}t^2L}(f) dt$$

we get

$$\sqrt{2\pi}XL^{-\frac{1}{2}}(f)(\gamma) = \int_0^\infty \left(\int_G [f(\gamma\delta_t g^{-1}) - f(\gamma\delta_t \sigma g^{-1})](Xp)(g) dg \right) \frac{dt}{t}.$$

Since $p \in \mathcal{S}(G)$, $Xp \in L^1(G)$, and formula (2) now comes from the subsequent Lemma 4 b).

By (2) and Hölder inequality with $\frac{1}{q} + \frac{1}{q'} = 1$, $d\gamma$ a.s.,

$$\sqrt{2\pi} \left| XL^{-\frac{1}{2}}(f)(\gamma) \right| \leq \|F(\gamma, \cdot)\|_{L^q(|Xp| dg)} \|Xp\|_{L^1(G)}^{\frac{1}{q'}}.$$

By the subsequent Lemma 4 a),

$$\|F(\gamma, g)\|_{L^q(d\gamma, L^q(|Xp| dg))} \leq h_q \|f\|_{L^q(G)} \|Xp\|_{L^1(G)}^{\frac{1}{q}},$$

which ends the proof. □

The next lemma comes from [C]; it is already used in this setting in [CMZ], see [CMZ, Lemma 5].

Lemma 4. *Let G be a stratified Lie group and $f \in \mathcal{D}(G)$. For $g \in G$ let $F(\gamma, g)$ be the Hilbert transform of f along the curve $g(t)$ and let $\psi \in L^1(G)$. Then*

a) *for $1 < q < \infty$, there exists a constant h_q such that*

$$\|F\|_{L^q(d\gamma \otimes |\psi| dg)} \leq h_q \|f\|_{L^q(d\gamma)} \|\psi\|_1^{\frac{1}{q}}.$$

b)

$$\begin{aligned} & \int_G F(\gamma, g)\psi(g) dg \\ &= \int_0^\infty \left[\int_G (f(\gamma\delta_t g^{-1}) - f(\gamma\delta_t \sigma g^{-1}))\psi(g) dg \right] \frac{dt}{t}, \quad d\gamma \text{ a.s.} \end{aligned}$$

Proof: a) When g runs through G , the corresponding family of curves $g(t)$ satisfies the assumptions of [C, pp. 579 and 594]. By the main result of [C], if $1 < q < \infty$, there exists a constant h_q such that, for every $g \in G$,

$$\|F(\cdot, g)\|_{L^q(G)} \leq h_q \|f\|_{L^q(G)}.$$

This proves a) by integration with respect to g and Fubini theorem.

b) For every $g \in G$, the truncated Hilbert transform $F(\gamma, g, \varepsilon)$ along the curve $g(t)$ satisfies [C, Lemma 6.3]

$$\|F(\cdot, g) - F(\cdot, g, \varepsilon)\|_{L^2(G)} \xrightarrow{\varepsilon \rightarrow 0^+} 0$$

and

$$\sup_{\varepsilon > 0} \|F(\cdot, g, \varepsilon)\|_{L^2(G)} \leq h_2 \|f\|_{L^2(G)}.$$

Hence, since $\psi \in L^1(G)$, the dominated convergence theorem implies

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0^+} \left\| \int_G (F(\gamma, g) - F(\gamma, g, \varepsilon)) \psi(g) dg \right\|_{L^2(G)} \\ & \leq \lim_{\varepsilon \rightarrow 0^+} \int_G \|F(\cdot, g) - F(\cdot, g, \varepsilon)\|_{L^2(G)} |\psi(g)| dg = 0. \end{aligned}$$

This implies b), because, by Fubini theorem, $d\gamma$ a.s.,

$$\begin{aligned} & \int_G F(\gamma, g, \varepsilon) \psi(g) dg \\ & = \int_{\varepsilon < t < \varepsilon^{-1}} \left[\int_G (f(\gamma \delta_t g^{-1}) - f(\gamma \delta_t \sigma g^{-1})) \psi(g) dg \right] \frac{dt}{t}. \quad \square \end{aligned}$$

Proof of Theorem 1: We treat Heisenberg groups first because the proof in this case is simpler and the idea is more apparent.

a) When $G = \mathbb{H}_n$, we denote $p = p^{(n)}$. The group law on \mathbb{H}_n is

$$\begin{aligned} \gamma g &= (x_1, y_1, \dots, x_n, y_n, u)(x'_1, y'_1, \dots, x'_n, y'_n, u') \\ &= \left(x_1 + x'_1, y_1 + y'_1, \dots, x_n + x'_n, y_n + y'_n, u + u' + 2 \sum_{j=1}^n (y_j x'_j - x_j y'_j) \right). \end{aligned}$$

By definition, for $g = (x_1, \dots, y_n, u)$,

$$\begin{aligned} (X_j p^{(n)})(g) &= \frac{\partial p^{(n)}}{\partial x_j}(g) + 2y_j \frac{\partial p^{(n)}}{\partial u}(g) \\ (Y_j p^{(n)})(g) &= \frac{\partial p^{(n)}}{\partial y_j}(g) - 2x_j \frac{\partial p^{(n)}}{\partial u}(g). \end{aligned}$$

The Laplacian is given by

$$-L = \sum_{j=1}^n \left[\frac{\partial^2}{\partial x_j^2} + \frac{\partial^2}{\partial y_j^2} + 4(x_j^2 + y_j^2) \frac{\partial^2}{\partial u^2} + 4 \left(x_j \frac{\partial^2}{\partial y_j \partial u} - y_j \frac{\partial^2}{\partial x_j \partial u} \right) \right]$$

hence commutes with rotations on (x_1, \dots, y_n) . It follows that $p^{(n)}$ is radial with respect to (x_1, \dots, y_n) , hence so are $\frac{\partial p^{(n)}}{\partial u}$ and the function

$$\frac{1}{r} \frac{\partial p^{(n)}}{\partial r} = \frac{1}{x_j} \frac{\partial p^{(n)}}{\partial x_j} = \frac{1}{y_j} \frac{\partial p^{(n)}}{\partial y_j},$$

where $r = \left(\sum_{j=1}^n x_j^2 + y_j^2 \right)^{\frac{1}{2}}$. Rewriting

$$(X_j p^{(n)})(g) = x_j \frac{1}{r} \frac{\partial p^{(n)}}{\partial r}(g) + 2y_j \frac{\partial p^{(n)}}{\partial u}(g)$$

we have, by (2), $d\gamma$ a.s.,

$$\sqrt{2\pi} X_j L^{-\frac{1}{2}}(f)(\gamma) = \int_G x_j F(\gamma, g) \frac{1}{r} \frac{\partial p^{(n)}}{\partial r}(g) dg + 2 \int_G y_j F(\gamma, g) \frac{\partial p^{(n)}}{\partial u}(g) dg$$

and a similar formula for $Y_j L^{-\frac{1}{2}}(f)$. Denoting for $1 \leq j \leq n$

$$\begin{aligned} R_j &= X_j L^{-\frac{1}{2}} \\ R_{n+j} &= Y_j L^{-\frac{1}{2}} \end{aligned}$$

by triangular inequality in l_{2n}^2 and Lemma 2 applied on \mathbb{R}^{2n+1} , $d\gamma$ a.s.,

$$\begin{aligned} (6) \quad \sqrt{2\pi} \left(\sum_{k=1}^{2n} |R_k(f)(\gamma)|^2 \right)^{\frac{1}{2}} &\leq \|F(\gamma, \cdot)\|_{L^q\left(\frac{1}{r} \left| \frac{\partial p^{(n)}}{\partial r} \right| dg\right)} \|x_1\|_{L^{q'}\left(\frac{1}{r} \left| \frac{\partial p^{(n)}}{\partial r} \right| dg\right)} \\ &\quad + 2 \|F(\gamma, \cdot)\|_{L^q\left(\left| \frac{\partial p^{(n)}}{\partial u} \right| dg\right)} \|x_1\|_{L^{q'}\left(\left| \frac{\partial p^{(n)}}{\partial u} \right| dg\right)}. \end{aligned}$$

Since L is the sum of $-(X_j^2 + Y_j^2)$, $1 \leq j \leq n$, which act on different set of variables except for the central u [CMZ, p. 371], (see also the proof of b) below)

$$p^{(n)}(x_1, \dots, y_n, u) = p^{(1)}(x_1, y_1, u) *_u \dots *_u p^{(1)}(x_n, y_n, u).$$

Since $p^{(n)} \geq 0$ for every n ,

$$\frac{1}{r} \left| \frac{\partial p^{(n)}}{\partial r} \right| = \left| \frac{1}{x_1} \frac{\partial p^{(n)}}{\partial x_1} \right| \leq \left| \frac{1}{x_1} \frac{\partial p^{(1)}}{\partial x_1} \right| *_u p^{(n-1)}.$$

Since $\int_{\mathbb{H}_{n-1}} p^{(n-1)}(g) dg = 1$, for $q \geq 0$,

$$\int_{\mathbb{H}_n} |x_1|^q \left| \frac{1}{r} \frac{\partial p^{(n)}}{\partial r} \right| dg \leq \int_{\mathbb{R}^3} |x_1|^q \left| \frac{1}{x_1} \frac{\partial p^{(1)}}{\partial x_1} \right| dx_1 dy_1 du = A_q.$$

Since $p^{(1)}$ belongs to $\mathcal{S}(\mathbb{R}^3)$, A_q is obviously finite for $q \geq 1$, and also for $q = 0$. Indeed

$$\left\| \frac{1}{x_1} \frac{\partial p^{(1)}}{\partial x_1} \right\|_{L^1(\mathbb{R}^3)} = 2\pi \int_{\mathbb{R}^+ \times \mathbb{R}} \left| \frac{\partial p^{(1)}}{\partial r} \right| dr du$$

and $\left| \frac{\partial p^{(1)}}{\partial r} \right| = \left(\left| \frac{\partial p^{(1)}}{\partial x_1} \right|^2 + \left| \frac{\partial p^{(1)}}{\partial y_1} \right|^2 \right)^{\frac{1}{2}}$. Similarly

$$\int_{\mathbb{H}_n} |x_1|^q \left| \frac{\partial p^{(n)}}{\partial u} \right| dg \leq \int_{\mathbb{R}^3} |x_1|^q \left| \frac{\partial p^{(1)}}{\partial u} \right| dx_1 dy_1 du = B_q$$

and B_q is finite for $q \geq 0$.

Integrating (6) with respect to γ , we get by Lemma 4 a)

$$\begin{aligned} \sqrt{2\pi} \left\| \left(\sum_{k=1}^{2n} |R_k(f)|^2 \right)^{\frac{1}{2}} \right\|_{L^q(\mathbb{H}_n)} &\leq A_{q'}^{\frac{1}{q'}} \|F\|_{L^q(d\gamma \otimes |\frac{1}{r} \frac{\partial p^{(n)}}{\partial r}| dg)} \\ &+ 2B_{q'}^{\frac{1}{q'}} \|F\|_{L^q(d\gamma \otimes |\frac{\partial p^{(n)}}{\partial u}| dg)} \leq h_q \|f\|_{L^q(\mathbb{H}_n)} (A_{q'}^{\frac{1}{q'}} A_0^{\frac{1}{q}} + 2B_{q'}^{\frac{1}{q'}} B_0^{\frac{1}{q}}) \end{aligned}$$

which gives the righthand side inequality in the statement of the theorem. Denoting

$$\mathcal{R}: f \longrightarrow (R_k(f))_{k=1}^K,$$

this means that \mathcal{R} is bounded: $L^q(G) \rightarrow L^q(G, l_{2n}^2)$, and so is

$$\mathcal{R}^*: (h_k)_{k=1}^{2n} \longrightarrow \sum_{k=1}^{2n} R_k^*(h_k), \quad L^{q'}(G, l_{2n}^2) \longrightarrow L^{q'}(G).$$

Since $f = \sum_{k=1}^{2n} R_k^* R_k(f) = \mathcal{R}^* \mathcal{R}(f)$ and $\|\mathcal{R}^*\|_{q \rightarrow q} = \|\mathcal{R}\|_{q' \rightarrow q'}$, we get in the standard way the lefthand side inequality:

$$\|f\|_{L^q(G)} \leq \|\mathcal{R}^*\|_{q \rightarrow q} \left\| \left(\sum_{k=1}^{2n} |R_k(f)|^2 \right)^{\frac{1}{2}} \right\|_{L^q(G)}.$$

b) When $G = \mathbb{H}_{K,J}$, we recall [K], [CDKR] that its Lie algebra \mathcal{G} is equipped with a scalar product, so that the first layer \mathcal{V} is a real vector space with dimension K , orthogonal to the second layer \mathcal{Z} with dimension J . We denote by $|X|$ the norm on \mathcal{G} defined by the scalar product.

The specific property is that, for every X with norm 1 in \mathcal{V} , the operator ad_X defined on \mathcal{V} by

$$\text{ad}_X: Y \longrightarrow [X, Y]$$

is an isometry from $E = (\ker \text{ad}_X)^\perp \subset \mathcal{V}$ onto \mathcal{Z} . In particular, E has dimension J and $|\text{ad}_X(Y)| = |P_E(Y)|$ where P_E is the orthogonal projection onto E .

Let us now precise the structure of $\mathcal{V} + \mathcal{Z}$ both as a Hilbert space and a Lie algebra. For $U \in \mathcal{Z}$ let $\Phi_U: \mathcal{V} \rightarrow \mathcal{V}$ be the linear operator defined by

$$\langle \Phi_U(X), X' \rangle = \langle U, [X, X'] \rangle.$$

In particular, $(\Phi_U)^* = -\Phi_U$, so that, if $F \subset \mathcal{V}$ is an invariant subspace for Φ_U , F^\perp is also invariant. The operator $\Phi: \mathcal{Z} \rightarrow B(\mathcal{V})$ defined by $U \rightarrow \Phi_U$ satisfies

$$(*) \quad \Phi_U \Phi_{U'} + \Phi_{U'} \Phi_U = -2 \langle U, U' \rangle \text{Id}_{\mathcal{V}}.$$

In particular, if U_1, \dots, U_J is an orthonormal basis of \mathcal{Z} , $(\Phi_{U_j})_{j=1}^J$ are unitary skew-adjoint anticommuting operators on \mathcal{V} . In more sophisticated words (see [BTV, 3.1.2]), denoting by q the quadratic form defined on \mathcal{Z} by $q(U) = -\langle U, U \rangle$, Φ induces a representation of the real Clifford algebra \mathcal{C}_J built on (\mathcal{Z}, q) into $B(\mathcal{V})$. In \mathcal{C}_J one has $U^2 = -\text{Id}$ for every $U \in \mathcal{Z}$ with norm 1, so that \mathcal{C}_J is the linear span of Id and $U_{i_1} U_{i_2} \dots U_{i_n}$, $1 \leq i_1 < i_2 < \dots < i_n \leq J$, and has dimension 2^J . Φ is a direct sum of irreducible representations of \mathcal{C}_J . By the classification of the Clifford algebras \mathcal{C}_J , their irreducible representations are as follows (see e.g. [Hu, Chapter 11] or [ABS, Part I]):

If $J \not\equiv 3 \pmod{4}$, there is only one (up to equivalence) irreducible representation of \mathcal{C}_J . Hence \mathcal{V} must be splitted as a hilbertian sum $\mathcal{V} = \mathcal{V}_1 \oplus \dots \oplus \mathcal{V}_N$ where the spaces \mathcal{V}_l have the same dimension K_J and are invariant under all Φ_U , $U \in \mathcal{Z}$. In particular, the \mathcal{V}_l 's are commuting copies with isomorphic Lie structure. We may choose orthonormal basis $(X_{(l-1)K_J+i})_{i=1}^{K_J}$ of the \mathcal{V}_l 's such that $[X_{(l-1)K_J+i}, X_{(l-1)K_J+h}]$ does not depend on l , for $1 \leq i, h \leq K_J$.

If $J \equiv 3 \pmod{4}$, there are two non equivalent irreducible representations of \mathcal{C}_J , with the same dimension K_J . \mathcal{V} must be splitted as a hilbertian sum $\mathcal{V}_1 \oplus \dots \oplus \mathcal{V}_k \oplus \mathcal{V}'_{k+1} \oplus \dots \oplus \mathcal{V}'_N$ where the \mathcal{V}_l 's (resp. the \mathcal{V}'_l 's) are commuting copies with isomorphic Lie structure, and the \mathcal{V}_l 's commute with the the \mathcal{V}'_l 's. We choose orthonormal basis of the \mathcal{V}_l 's (resp. the \mathcal{V}'_l 's) as above.

Conversely, by [K], if \mathcal{Z}, \mathcal{V} are real finite dimensional Hilbert spaces, from a linear isometry $\Phi: \mathcal{Z} \rightarrow B(\mathcal{V})$ satisfying (*), one can build a structure of Lie algebra on $\mathcal{V} + \mathcal{Z}$. If $J \equiv 3 \pmod{4}$, there are several non isomorphic groups $\mathbb{H}_{K,J}$ for every admissible $K > K_J$ and only one $\mathbb{H}_{K_J,J}$ [BTV, 3.1.2].

The value of K_J is computed as follows. By [K], the couple (K, J) satisfies

$$J < \rho(K) = 8\alpha + 2^\beta \text{ where } K = m2^{4\alpha+\beta}, \quad m \text{ odd}, \quad 0 \leq \beta \leq 3, \quad \alpha, \beta \in \mathbb{N}.$$

In particular, K must be even. Let ρ_J be the smallest integer such that

$$\rho_J = 8\alpha_J + 2^{\beta_J} > J, \quad 0 \leq \beta_J \leq 3, \quad \alpha_J, \beta_J \in \mathbb{N}$$

and let

$$K_J = 2^{4\alpha_J + \beta_J}, \quad \text{hence} \quad \rho(K_J) = \rho_J.$$

This ensures the existence of $\mathbb{H}_{K_J, J}$ and K_J is the minimal possible dimension of \mathcal{V} , i.e. the dimension of each \mathcal{V}_l . In particular $J < K_J$.

For a given $\mathbb{H}_{K, J}$ we choose an orthonormal basis X_1, \dots, X_K of \mathcal{V} as above. We denote by $g = (x, u)$ an element of $\mathbb{H}_{K, J}$, where $x = (x_k)_{k=1}^{NK_J}$ are the coordinates corresponding to X_1, \dots, X_K , $u = (u_j)_{j=1}^J$ are the coordinates corresponding to U_1, \dots, U_J , and by $y_l \in \mathbb{R}^{K_J}$ the coordinates of x corresponding to the basis of \mathcal{V}_l or of \mathcal{V}'_l , $1 \leq l \leq N$, so that $x = (y_1, \dots, y_N)$.

By definition [BTV, 3.1.5], $U_j = \frac{\partial}{\partial u_j}$, $1 \leq j \leq J$, and

$$X_i = \frac{\partial}{\partial x_i} + \frac{1}{2} \sum_{j=1}^J \left\langle \left[\sum_{k=1}^K x_k X_k, X_i \right], U_j \right\rangle \frac{\partial}{\partial u_j}, \quad 1 \leq i \leq K.$$

In particular, if X_i belongs to \mathcal{V}_l or to \mathcal{V}'_l , it only depends on y_l .

The heat kernel $p = p^{(K, J)}$ has properties analogous to those of $p^{(n)}$. First, $p^{(K, J)}$ is radial both with respect to x and u and depends only on (K, J) . This follows from the following formula established by J. Randall [R, Proof of Lemma 1.3.3]:

$$(\mathcal{F}_u p^{(K, J)})(x, \lambda) = (\mathcal{F}_v p^{(\frac{K}{2})})(x, |\lambda|), \quad \lambda \in \mathbb{R}^J,$$

where \mathcal{F}_u denotes the Fourier transform on \mathbb{R}^J with respect to u , \mathcal{F}_v the Fourier transform on \mathbb{R} with respect to $v \in \mathbb{R}$, and $p^{(\frac{K}{2})}$ denotes the heat kernel on the Heisenberg group $\mathbb{H}_{\frac{K}{2}}$. In particular, there is a unique $p^{(K, J)}$ for any given J .

We claim that

$$p^{(K, J)}(x, u) = p^{(K_J, J)}(y_1, u) *_u \dots *_u p^{(K_J, J)}(y_N, u)$$

where convolution is on \mathbb{R}^J , with respect to u . This follows from Randall's formula and the fact that

$$(\mathcal{F}_v p^{(\frac{K}{2})})(x, |\lambda|) = \prod_{i=1}^N (\mathcal{F}_v p^{(\frac{K_J}{2})})(y_i, |\lambda|).$$

It can also be proved directly. Indeed, let us denote by q the convolution product on the right hand side. Since $p^{(K_J, J)}$ is a positive function with norm 1 in $L^1(\mathbb{R}^{K_J+J})$, q is a positive function with norm 1 in $L^1(\mathbb{R}^{K+J})$. As well known (see e.g. [FS]), the heat kernel $p_t^{(K, J)}$ is the only positive function with norm 1 in $L^1(\mathbb{R}^{K+J})$ satisfying $\frac{\partial f}{\partial t} = -\frac{1}{2}Lf$; moreover,

$p_t^{(K,J)} = t^{-(\frac{K}{2}+J)} p^{(K,J)} \circ \delta_{t^{-\frac{1}{2}}}$. Hence $p^{(K,J)}$ is the only positive function with norm 1 in $L^1(\mathbb{R}^{K+J})$ satisfying

$$(K \text{ Id} - L)p + \sum_{k=1}^K x_k X_k p = -2 \sum_{j=1}^J U_j(u_j p)$$

(see e.g. [LP1, Lemmas 2, 3]). So we only have to verify that q satisfies this equation. The left term can be splitted as $\sum_{l=1}^N D_l(p)$ where

$$D_l = K_J \text{ Id} + \sum_{i=1}^{K_J} -X_{i+(l-1)K_J}^2 + x_{i+(l-1)K_J} X_{i+(l-1)K_J}.$$

By our choice of coordinates, D_l only acts on the l^{th} factor of q . For the right term, we note that

$$\begin{aligned} & u_j q(y_1, \dots, y_N, u) \\ &= \sum_{l=1}^N p^{(K_J,J)}(y_l, u) *_{u} \dots *_{u} (u_j p^{(K_J,J)})(y_l, u) *_{u} p^{(K_J,J)}(y_{l+1}, u) *_{u} \dots, \end{aligned}$$

which is easily verified by Fourier transform with respect to u . So it remains to verify that, for $1 \leq l \leq N$,

$$(D_l p^{(K_J,J)})(y_l, u) = -2 \sum_{j=1}^J U_j(u_j p^{(K_J,J)})(y_l, u).$$

These are the heat equations corresponding to each Lie algebra $\mathcal{V}_l + \mathcal{Z}$ or $\mathcal{V}'_l + \mathcal{Z}$. Hence they are all satisfied by $p^{(K_J,J)}$ as we saw above (this is obvious when there are only \mathcal{V}'_l 's).

For $g = (x, u) \in \mathbb{H}_{K,J}$ let $r = |x|$, $\rho = |u|$. As in a), we may rewrite for $1 \leq i \leq K$,

$$X_i p^{(K,J)} = \frac{x_i}{r} \frac{\partial p^{(K,J)}}{\partial r} + \frac{1}{2} \left\langle \left[\sum_{k=1}^K x_k X_k, X_i \right], \sum_{j=1}^J u_j U_j \right\rangle \frac{1}{\rho} \frac{\partial p^{(K,J)}}{\partial \rho},$$

and

$$\begin{aligned} (7) \quad & \sqrt{2\pi} \left(\sum_{i=1}^K |R_i(f)(\gamma)|^2 \right)^{\frac{1}{2}} \leq \left(\sum_{i=1}^K \left| \int_G x_i F(\gamma, g) \frac{1}{r} \frac{\partial p^{(K,J)}}{\partial r}(g) dg \right|^2 \right)^{\frac{1}{2}} \\ & + \frac{1}{2} \left(\sum_{i=1}^K \left| \int_G F(\gamma, g) \left\langle \left[\sum_{k=1}^K x_k X_k, X_i \right], \sum_{j=1}^J u_j U_j \right\rangle \frac{1}{\rho} \frac{\partial p^{(K,J)}}{\partial \rho}(g) dg \right|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

The $L^q(d\gamma)$ norm of the first term in the righthand side of (7) is estimated as in a), replacing $p^{(1)}$ by $p^{(K,J)}$; it is less than

$$\begin{aligned} h_q \|x_1\|_{L^{q'}(\mathbb{R}^{K+J}, |\frac{1}{x_1} \frac{\partial p^{(K,J)}}{\partial x_1}| dx du)} &\left\| \frac{1}{x_1} \frac{\partial p^{(K,J)}}{\partial x_1} \right\|_{L^1(\mathbb{R}^{K+J})}^{\frac{1}{q}} \|f\|_{L^q(\mathbb{H}_{K,J})} \\ &= C_1(q, J) \|f\|_{L^q(\mathbb{H}_{K,J})} \end{aligned}$$

and the constant is finite as in a) because $p^{(K,J)} \in \mathcal{S}(\mathbb{H}_{K,J})$.

The second term is rewritten as

$$\begin{aligned} \sup_{|a|=1} \int_G F(\gamma, g) \left\langle \left[\sum_{k=1}^K x_k X_k, \sum_{i=1}^K a_i X_i \right], \sum_{j=1}^J u_j U_j \right\rangle \frac{1}{\rho} \frac{\partial p^{(K,J)}}{\partial \rho}(g) dg \\ \leq A_{q'} \|F(\gamma, \cdot)\|_{L^q(\frac{1}{\rho} |\frac{\partial p^{(K,J)}}{\partial \rho}(g)| dg)} \end{aligned}$$

where

$$A_{q'} = \sup_{|a|=1} \left\| \sum_{j=1}^J u_j \left\langle \left[\sum_{k=1}^K x_k X_k, \sum_{i=1}^K a_i X_i \right], U_j \right\rangle \right\|_{L^{q'}(\frac{1}{\rho} |\frac{\partial p^{(K,J)}}{\partial \rho}(g)| dg)}.$$

Hence, by Lemma 4 a), the $L^q(d\gamma)$ norm of the second term in the righthand side of (7) is estimated by

$$\begin{aligned} A_{q'} h_q \left\| \frac{1}{u_1} \frac{\partial p^{(K,J)}}{\partial u_1} \right\|_{L^1(\mathbb{R}^{K+J})}^{\frac{1}{q}} \|f\|_{L^q(\mathbb{H}_{K,J})} \\ \leq A_{q'} h_q \left\| \frac{1}{u_1} \frac{\partial p^{(K,J)}}{\partial u_1} \right\|_{L^1(\mathbb{R}^{K+J})}^{\frac{1}{q}} \|f\|_{L^q(\mathbb{H}_{K,J})}. \end{aligned}$$

We now estimate $A_{q'}$. By Lemma 2 a) applied to u_1, \dots, u_J and fixed x ,

$$\begin{aligned} \left\| \sum_{j=1}^J u_j \left\langle \left[\sum_{k=1}^K x_k X_k, \sum_{i=1}^K a_i X_i \right], U_j \right\rangle \right\|_{L^{q'}(\frac{1}{\rho} |\frac{\partial p^{(K,J)}}{\partial \rho}| du)} \\ = B_a(x) \|u_1\|_{L^{q'}(\frac{1}{\rho} |\frac{\partial p^{(K,J)}}{\partial \rho}| du)} \end{aligned}$$

where, since $\left| \sum_{i=1}^K a_i X_i \right| = 1$,

$$\begin{aligned} B_a(x) &= \left| \left[\sum_{k=1}^K x_k X_k, \sum_{i=1}^K a_i X_i \right] \right| = \left| \text{ad}_{\sum_{i=1}^K a_i X_i} \left(\sum_{k=1}^K x_k X_k \right) \right| \\ &= \left| P_{E_a} \left(\sum_{k=1}^K x_k X_k \right) \right| \end{aligned}$$

with $E_a = (\ker \text{ad}_{\sum_{i=1}^K a_i X_i})^\perp$. By rotation on the x variables, we may suppose that E_a is the span of X_1, \dots, X_J . Since $\frac{\partial p^{(K,J)}}{\partial \rho}$ is radial with respect to x

$$\begin{aligned} A_{q'}^{q'} &= \sup_{|a|=1} \int_{\mathbb{R}^{K+J}} |B_a(x)|^{q'} |u_1|^{q'} \frac{1}{\rho} \left| \frac{\partial p^{(K,J)}}{\partial \rho}(x, u) \right| dx du \\ &= \int_{\mathbb{R}^{K+J}} \left(\sum_{k=1}^J |x_k|^2 \right)^{\frac{q'}{2}} |u_1|^{q'-1} \left| \frac{\partial p^{(K,J)}}{\partial u_1}(x, u) \right| dx du \\ &\leq \int_{\mathbb{R}^{K+J}} \left(\sum_{k=1}^J |x_k|^2 \right)^{\frac{q'}{2}} |u_1|^{q'-1} \left| \frac{\partial p^{(K,J)}}{\partial u_1}(x, u) \right| dx du \end{aligned}$$

where the inequality is verified as in a), replacing $p^{(1)}$ by $p^{(K,J)}$, since $J < K_J$. Finally

$$C_2(q, J) = h_q \left\| \left(\sum_{k=1}^J |x_k|^2 \right)^{\frac{1}{2}} |u_1| \right\|_{L^{q'(|\frac{1}{u_1} \frac{\partial p^{(K,J)}}{\partial u_1}| dx du)} \left\| \frac{1}{u_1} \frac{\partial p^{(K,J)}}{\partial u_1} \right\|_{L^1(\mathbb{R}^{K+J})}^{\frac{1}{q}}.$$

This constant is finite because $p^{(K,J)} \in \mathcal{S}(\mathbb{H}_{K,J})$ and we get

$$\left\| \left(\sum_{i=1}^K |R_i(f)(\gamma)|^2 \right)^{\frac{1}{2}} \right\|_{L^q(\mathbb{H}_{K,J})} \leq \frac{1}{\sqrt{2\pi}} \left(C_1(q, J) + \frac{1}{2} C_2(q, J) \right) \|f\|_{L^q(\mathbb{H}_{K,J})}.$$

This implies as in a) the lefthand side inequality. □

Remark. The above proof of a) does not seem to extend to the setting of non isotropic Heisenberg groups, because it uses in a crucial way the radiality of p ; so does the proof in [CMZ]. However, it is stated in [BDJ, p. 59], without explanation, that the proof in [CMZ] can be extended to the non isotropic case.

2. Riesz transforms associated to commuting inner *-derivations

2.1. Introduction and notation.

We now show how the classical result on Riesz transforms on $L^q(\mathbb{R}^n, dx)$ can be extended to Riesz transforms on Schatten spaces $\mathcal{S}^q(H)$, defined by inner (bounded or unbounded) commuting *-derivations acting on the C^* algebra $K(H)$ of compact operators on the Hilbert space H .

We denote by γ_n the standard gaussian density on \mathbb{R}^n and by Π the orthogonal projection onto the linear span of the coordinates y_j in $L^2(\gamma_n)$. As well known, Π extends as a bounded operator: $L^q(\gamma_n) \rightarrow L^q(\gamma_n)$, $1 \leq q < \infty$, which we still denote by Π .

The Fourier transform on \mathbb{R} is defined by $\widehat{f}(u) = \int_{\mathbb{R}} e^{-iyu} f(y) dy$.

We recall that $\mathcal{S}^q(H)$ is the space of compact operators X on H such that $\text{tr}(X^*X)^{\frac{q}{2}} < \infty$, $1 \leq q < \infty$. $B(H)$ is the dual space of $\mathcal{S}^1(H)$ which is itself the dual space of $K(H)$. $\mathcal{S}^2(H)$ is the Hilbert space of Hilbert Schmidt operators on H . From now on, we assume H is separable.

We denote by $|X|_s^2 = X^*X + XX^*$ the symmetrized modulus of $X \in K(H)$.

Let h be a *-automorphism of $K(H)$, hence h^{**} is a *-automorphism of $B(H)$. By [Pe, Theorem 8.9.2] there exists a unitary operator U on H such that $h(X) = UXU^*$, $X \in K(H)$. Let $(h_y)_{y \in \mathbb{R}}$ be a strongly continuous one parameter group of *-automorphisms of $K(H)$; then there exists a one parameter group of unitaries $(U_y)_{y \in \mathbb{R}}$ such that $h_y(X) = U_y X U_y^*$, $X \in K(H)$ (see e.g. [V, Theorem 11.1] for an actually stronger result, or [Par, pp. 86–87]); by Stone's theorem $U_y = e^{yA}$, where iA is self-adjoint on H . Hence the generator D of $(h_y)_{y \in \mathbb{R}}$ is an inner *-derivation, in general unbounded, defined by $D(X) = [A, X]$. In particular $(h_y)_{y \in \mathbb{R}}$ is a group of isometries of $\mathcal{S}^2(H)$ (and a group of isometries of every $\mathcal{S}^q(H)$, $1 \leq q < \infty$). By [RS, Theorem VIII.9], $(h_y)_{y \in \mathbb{R}}$ is strongly continuous on $\mathcal{S}^2(H)$. Its generator is naturally induced by D , and iD is self-adjoint on $\mathcal{S}^2(H)$ by Stone's theorem.

Let us now recall some facts about joint functional calculus for “commuting” *-inner derivations. We consider n strongly continuous one

parameter groups of $*$ -automorphisms of $K(H)$, with respective generators D_j , $1 \leq j \leq n$, which we denote respectively by $(e^{y_j D_j})_{y_j \in \mathbb{R}}$. We assume that

- (i) their restrictions commute on $\mathcal{S}^2(H)$
- (ii) the only $X \in \mathcal{S}^2(H)$ which is invariant under every $e^{y_j D_j}$ is $X = 0$.

In particular, by (i),

$$y \longrightarrow U(y) = e^{\sum_{j=1}^n y_j D_j}$$

is a strongly continuous map of \mathbb{R}^n into the unitary operators on $\mathcal{S}^2(H)$, satisfying $U(y+z) = U(y)U(z)$, $y, z \in \mathbb{R}^n$ and $U(0) = I$. Then [RS, Theorem VIII.12], there is a projection valued measure E on \mathbb{R}^n such that

$$(8) \quad \left\langle e^{\sum_{j=1}^n y_j D_j}(X), Y \right\rangle = \int_{\mathbb{R}^n} e^{i\langle y, \lambda \rangle} d\langle E_\lambda(X), Y \rangle, \\ X, Y \in \mathcal{S}^2(H), \quad y \in \mathbb{R}^n.$$

By bounded functional calculus and Fubini theorem, for every $F \in \mathcal{S}(\mathbb{R}^n)$, as bounded operators on $\mathcal{S}^2(H)$ [RS, p. 272],

$$(9) \quad \widehat{F}(iD_1, \dots, iD_n) = \int_{\mathbb{R}^n} e^{\sum_{k=1}^n y_k D_k} F(y) dy.$$

Moreover, let $h: \mathbb{R}^n \rightarrow \mathbb{R}$ be a Borel measurable function and let $X \in \mathcal{S}^2(H)$ be such that $\int_{\mathbb{R}^n} |h(\lambda)|^2 d\langle E_\lambda(X), X \rangle < \infty$; then the formula

$$\langle h(iD_1, \dots, iD_n)(X), Y \rangle = \int_{\mathbb{R}^n} h(\lambda) d\langle E_\lambda(X), Y \rangle, \quad Y \in \mathcal{S}^2(H)$$

defines an operator $h(iD_1, \dots, iD_n)$ which is densely defined and self-adjoint on $\mathcal{S}^2(H)$. This holds in particular for

$$L = - \sum_{k=1}^n D_k^2.$$

By assumption (ii), the projection $E_{\{0\}}$ is null. Indeed, let $Z \in \mathcal{S}^2(H)$ lying in its range. Since the measure $E - \delta_0 \otimes E_{\{0\}}$ is valued in the set of orthogonal projections P on $\mathcal{S}^2(H)$ such that $PE_{\{0\}} = 0$, hence $P(Z) = 0$, the scalar measure $1_{\{\mathbb{R}^n \setminus \{0\}\}}(\lambda) d\langle E_\lambda(Z), Y \rangle$ is zero for every $Y \in \mathcal{S}^2(H)$; then, by (8),

$$\left\langle Y, e^{\sum_{j=1}^n y_j D_j}(Z) \right\rangle = \left\langle e^{i\langle y, \lambda \rangle}, \delta_0 \right\rangle \langle Y, E_{\{0\}}Z \rangle = \langle Y, Z \rangle,$$

which, by (ii), implies $Z = 0$.

The Riesz transforms are defined by

$$R_j = D_j L^{-\frac{1}{2}}, \quad 1 \leq j \leq n,$$

so they are contractions on their domain in $\mathcal{S}^2(H)$. Since $E_{\{0\}} = 0$, R_j is actually defined on $\mathcal{S}^2(H)$ by

$$(10) \quad \langle R_j(X), Y \rangle = \int_{\mathbb{R}^n \setminus \{0\}} \frac{i\lambda_j}{|\lambda|} d\langle E_\lambda(X), Y \rangle.$$

The main result of this part is the following theorem:

Theorem 5. *Let H be a separable Hilbert space and let $(e^{y_j D_j})_{y_j \in \mathbb{R}}$, $1 \leq j \leq n$ be strongly continuous one parameter groups of $*$ -automorphisms of $K(H)$ satisfying the above conditions (i), (ii). Let $1 < q < \infty$.*

a) *Then, for every $X \in K(H)$ and $t \in \mathbb{R}$,*

$$e^{-\frac{1}{2}t^2 L}(X) = \int_{\mathbb{R}^n} e^{t \sum_{j=1}^n y_j D_j} (X) \gamma_n(y) dy;$$

$e^{-\frac{1}{2}t^2 L}$ is a completely positive contraction: $K(H) \rightarrow K(H)$ and a contraction of every $\mathcal{S}^q(H)$.

b) *The operator $\mathcal{R} = \sum_{j=1}^n y_j R_j$ is a (completely) bounded operator: $\mathcal{S}^q(H) \rightarrow L^q(\gamma_n(y) dy, \mathcal{S}^q)$ which satisfies*

$$\sqrt{2\pi} \mathcal{R}(X) = (\Pi \otimes I_{\mathcal{S}^q}) \left(pv \int_{-\infty}^{\infty} e^{t \sum_{j=1}^n y_j D_{A_j}} (X) \frac{dt}{t} \right).$$

c) *For $X \in \mathcal{S}^q$, $\|X\|_{\mathcal{S}^q}$ is respectively equivalent, with constants which depend only on q , to*

$$(i) \quad \left\| \left(\sum_{j=1}^n |R_j(X)|_s^2 \right)^{\frac{1}{2}} \right\|_{\mathcal{S}^q}, \quad 2 \leq q < \infty,$$

and to

$$(ii) \quad \inf \left\{ \left\| \left(\sum_{j=1}^n B_j B_j^* \right)^{\frac{1}{2}} \right\|_{\mathcal{S}^q} + \left\| \left(\sum_{j=1}^n C_j^* C_j \right)^{\frac{1}{2}} \right\|_{\mathcal{S}^q} \right\}, \quad 1 < q < 2,$$

where the infimum is taken over all decompositions $R_j(X) = B_j + C_j$ in $\mathcal{S}^q(H)$.

Note that, on H , the operators $R_j(X)$, $1 \leq j \leq n$, do not commute in general, and $R_j(X)^* = R_j(X^*)$ does not commute in general with $R_j(X)$.

Proof of Theorem 5: The strategy is similar to Pisier's in the first part of [P1] and to the one we used in the first part of this paper. On one hand, things are much easier than for Heisenberg groups because we deal with a one parameter group; on the other hand, some difficulties arise from the setting of non commutative L^q 's.

a) Formula (9) defines bounded operators on $K(H)$ and $\mathcal{S}^q(H)$, $1 \leq q < \infty$, with norm less than $\|F\|_{L^1(dy)}$, because $e^{\sum_{k=1}^n y_k D_k}$ is an isometry of $K(H)$ and $\mathcal{S}^q(H)$ for every y .

This formula gives a Stinespring factorization (see e.g. [Pa]) of $\widehat{F}(iD_1, \dots, iD_n)$ acting on $K(H)$ because

$$X \longrightarrow e^{\sum_{j=1}^n y_j D_j}(X)$$

is a *-homomorphism: $K(H) \rightarrow L^\infty(d\gamma_n, B(H)) \subset B(L^2(d\gamma_n, H))$.

Applying this to $F(y) = \gamma_n(y)$ and tD_1, \dots, tD_n proves assertion a).

b) α) By (10) and Fubini theorem, for $X \in \mathcal{S}^2(H)$,

$$\begin{aligned} \sqrt{\frac{\pi}{2}} \langle R_j(X), X \rangle &= \sqrt{\frac{\pi}{2}} \int_{\mathbb{R}^n} \frac{i\lambda_j}{|\lambda|} d \langle E_\lambda(X), X \rangle \\ &= \int_{\mathbb{R}^n} i\lambda_j \left(\int_0^\infty e^{-\frac{1}{2}t^2|\lambda|^2} dt \right) d \langle E_\lambda(X), X \rangle \\ &= \int_0^\infty \left(\int_{\mathbb{R}^n} i\lambda_j e^{-\frac{1}{2}t^2|\lambda|^2} d \langle E_\lambda(X), X \rangle \right) dt \\ &= \int_0^\infty \langle D_j e^{-\frac{1}{2}t^2 L}(X), X \rangle dt. \end{aligned}$$

β) Let

$$F(y) = y_j \gamma_n(y), \quad \text{hence} \quad \widehat{F}(u) = -iu_j e^{-\frac{1}{2}|u|^2}.$$

By (9) applied to F and tD_1, \dots, tD_n , by symmetry with respect to t , for $X \in \mathcal{S}^q(H)$,

$$\begin{aligned} tD_j e^{-\frac{1}{2}t^2 L}(X) &= \int_{\mathbb{R}^n} e^{t \sum_{k=1}^n y_k D_k}(X) y_j \gamma_n(y) dy \\ &= \frac{1}{2} \int_{\mathbb{R}^n} (e^{t \sum_{k=1}^n y_k D_k}(X) - e^{-t \sum_{k=1}^n y_k D_k}(X)) y_j \gamma_n(y) dy. \end{aligned}$$

Hence, by α), for $X \in \mathcal{S}^2(H)$,

$$\begin{aligned}
 \sqrt{2\pi}R_j(X) &= \int_0^\infty \left[\int_{\mathbb{R}^n} (e^{t \sum_{k=1}^n y_k D_k}(X) \right. \\
 (11) \quad &\quad \left. - e^{-t \sum_{k=1}^n y_k D_k}(X)) y_j \gamma_n(y) dy \right] \frac{dt}{t} \\
 &= \int_{-\infty}^\infty \left[\int_{\mathbb{R}^n} e^{t \sum_{k=1}^n y_k D_k}(X) y_j \gamma_n(y) dy \right] \frac{dt}{t}.
 \end{aligned}$$

γ) We claim that, for every $X \in \mathcal{S}^q(H)$, $y \in \mathbb{R}^n$, $1 < q < \infty$,

$$F(y, X) = \text{pv} \int_{-\infty}^\infty e^{t \sum_{k=1}^n y_k D_k}(X) \frac{dt}{t}$$

is well defined as the norm limit in $\mathcal{S}^q(H)$ of

$$F(y, X, \varepsilon) = \int_{\varepsilon < |t| < \varepsilon^{-1}} e^{t \sum_{k=1}^n y_k D_k}(X) \frac{dt}{t}$$

and satisfies

$$(12) \quad \|F(y, X)\|_{\mathcal{S}^q} \leq h_q \|X\|_{\mathcal{S}^q}.$$

Indeed, $\mathcal{S}^q(H)$ is UMD for $1 < q < \infty$ [BGM, Theorem 6.1] and, for fixed y , $(e^{t \sum_{k=1}^n y_k D_k})_{t \in \mathbb{R}}$ is a strongly continuous one parameter group of isometries of $\mathcal{S}^q(H)$. Hence [BGM, Theorems 5.12 and 5.16] prove the claim.

δ) It follows that, for the $\mathcal{S}^2(H)$ norm, by Fubini theorem and (11),

$$\begin{aligned}
 \int_{\mathbb{R}^n} F(y, X) y_j \gamma_n(y) dy &= \lim_\varepsilon \int_{\mathbb{R}^n} F(y, X, \varepsilon) y_j \gamma_n(y) dy \\
 &= \lim_\varepsilon \int_{\varepsilon < |t| < \varepsilon^{-1}} \left(\int_{\mathbb{R}^n} e^{t \sum_{k=1}^n y_k D_k}(X) y_j \gamma_n(y) dy \right) \frac{dt}{t} \\
 &= \sqrt{2\pi}R_j(X).
 \end{aligned}$$

Hence, for $X \in \mathcal{S}^q(H) \cap \mathcal{S}^2(H)$,

$$\sqrt{2\pi}\mathcal{R}(X) = \sum_{j=1}^n y_j \int_{\mathbb{R}^n} F(y, X) y_j \gamma_n(y) dy = (\Pi \otimes I_{\mathcal{S}^q})(F(y, X)),$$

and, by (12),

$$\|F(y, X)\|_{L^q(\gamma_n dy, \mathcal{S}^q)} \leq h_q \|X\|_{\mathcal{S}^q}.$$

ε) By [P2, Remark 8.4.6], $\Pi \otimes I_{\mathcal{S}^q}$ is bounded on $L^q(\gamma_n dy, \mathcal{S}^q)$, $1 < q < \infty$, and its norm depends only on q . By δ) this proves the boundedness of \mathcal{R} on $\mathcal{S}^q(H)$ and ends the proof of b).

Actually, Π is completely bounded, hence so is $\Pi \otimes I_{\mathcal{S}^q}$, and the same argument as above, applied to $(e^{t \sum_{j=1}^n y_j D_{A_j}} \otimes I_{\mathcal{S}^q})_{t \in \mathbb{R}}$ shows that, for fixed y , $\text{pv} \int_{-\infty}^{\infty} e^{t \sum_{j=1}^n y_j D_{A_j}} \frac{dt}{t}$ is completely bounded: $\mathcal{S}^q(H) \rightarrow \mathcal{S}^q(H)$.

c) By [P2, Theorem 8.4.1], the norms in the statement are equivalent, with constants which depend only on q , to

$$\left\| \sum_{j=1}^n y_j R_j(X) \right\|_{L^q(d\gamma_n, \mathcal{S}^q)} = \|\mathcal{R}(X)\|_{L^q(d\gamma_n, \mathcal{S}^q)}.$$

Hence, we have to show that $\|X\|_{\mathcal{S}^q}$ is equivalent to $\|\mathcal{R}(X)\|_{L^q(d\gamma_n, \mathcal{S}^q)}$. One inequality has been proved in b). For the other one, we notice that $\mathcal{R}^* \mathcal{R} = \text{Id}$ on $\mathcal{S}^2(H)$ because

$$\begin{aligned} \langle \mathcal{R}(X), \mathcal{R}(Y) \rangle &= \int_{\mathbb{R}^n} \left\langle \sum_{j=1}^n y_j R_j(X), \sum_{k=1}^n y_k R_k(Y) \right\rangle \gamma_n(y) dy \\ &= \sum_{j=1}^n \langle R_j(X), R_j(Y) \rangle = \sum_{j=1}^n \langle R_j^* R_j(X), Y \rangle = \langle X, Y \rangle. \end{aligned}$$

Hence, for $\frac{1}{q} + \frac{1}{q'} = 1$,

$$\|X\|_{\mathcal{S}^q} \leq \|\mathcal{R}^*\|_{q \rightarrow q} \|\mathcal{R}(X)\|_{L^q(\gamma_n dy, \mathcal{S}^q)} = \|\mathcal{R}\|_{q' \rightarrow q'} \|\mathcal{R}(X)\|_{L^q(\gamma_n dy, \mathcal{S}^q)}. \quad \square$$

Example 6. The CCR heat flow on $K(L^2(\mathbb{R}^n))$.

The assumptions of Theorem 5 are satisfied by the following example, taken from [A], where assertion a) is proved in this special case.

The operator $P = -i \frac{d}{dx}$ and the operator Q of multiplication by x are well defined: $\mathcal{S}(\mathbb{R}) \rightarrow \mathcal{S}(\mathbb{R})$ and satisfy the Canonical Commutation Relation

$$[Q, P] = iI.$$

P, Q are formally selfadjoint unbounded operators on $L^2(\mathbb{R})$. They generate two one parameter unitary groups of operators on $L^2(\mathbb{R})$, respectively e^{isP} (translation by $-s$) and e^{itQ} (multiplication by e^{itx}), satisfying

$$(13) \quad e^{isP} e^{itQ} = e^{ist} e^{itQ} e^{isP}, \quad s, t \in \mathbb{R}.$$

We may define the inner $*$ -derivations

$$D_P(X) = i[P, X],$$

$$D_Q(X) = i[Q, X]$$

for operators X on $L^2(\mathbb{R})$ which are defined by kernels $k(x, y) \in \mathcal{S}(\mathbb{R}^2)$. Note that the kernels of $D_P(X)$, $D_Q(X)$ are respectively $\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y}\right)k$ and $(x - y)k$. D_P , D_Q are generators of groups of $*$ -automorphisms of $K(L^2(\mathbb{R}))$, namely

$$e^{sD_P}(X) = e^{isP} X e^{-isP},$$

$$e^{tD_Q}(X) = e^{itQ} X e^{-itQ}$$

and, owing to (13), e^{sD_P} and e^{tD_Q} commute on $K(L^2(\mathbb{R}))$. Obviously, only $X = 0$ is stable under these groups.

On $\mathcal{S}(\mathbb{R}^n)$ we consider in the same way $P_j = -i\frac{\partial}{\partial x_j}$, Q_j the operator of multiplication by x_j . The $*$ -automorphisms $e^{s_j D_{P_j}}$ and $e^{t_k D_{Q_k}}$, $1 \leq j, k \leq n$ all commute on $K(L^2(\mathbb{R}^n))$. Denoting

$$L = -\sum_{j=1}^n D_{P_j}^2 + D_{Q_j}^2,$$

the semigroup e^{-tL} , $t \geq 0$, acting on $K(L^2(\mathbb{R}^n))$, is called the CCR heat flow. The $2n$ Riesz transforms $D_{P_j} L^{-\frac{1}{2}}$, $D_{Q_k} L^{-\frac{1}{2}}$, $1 \leq j, k \leq n$ satisfy the conclusion of Theorem 5.

References

- [A] W. ARVESON, The heat flow of the CCR algebra, *Bull. London Math. Soc.* **34(1)** (2002), 73–83.
- [ABS] M. F. ATIYAH, R. BOTT AND A. SHAPIRO, Clifford modules, *Topology* **3**, suppl. 1 (1964), 3–38.
- [BDJ] C. BERENSTEIN, DER-CHEN CHANG AND JINGZHI TIE, “Laguerre calculus and its applications on the Heisenberg group”, AMS/IP Studies in Advanced Mathematics **22**, American Mathematical Society, Providence, RI; International Press, Somerville, MA, 2001.
- [BGM] E. BERKSON, T. A. GILLESPIE AND P. S. MUHLY, Abstract spectral decompositions guaranteed by the Hilbert transform, *Proc. London Math. Soc.* (3) **53(3)** (1986), 489–517.

- [BTV] J. BERNDT, F. TRICERRI AND L. VANHECKE, “*Generalized Heisenberg groups and Damek-Ricci harmonic spaces*”, Lecture Notes in Mathematics **1598**, Springer-Verlag, Berlin, 1995.
- [C] M. CHRIST, Hilbert transforms along curves. I. Nilpotent groups, *Ann. of Math. (2)* **122(3)** (1985), 575–596.
- [CMZ] T. COULHON, D. MÜLLER AND J. ZIENKIEWICZ, About Riesz transforms on the Heisenberg groups, *Math. Ann.* **305(2)** (1996), 369–379.
- [CDKR] M. COWLING, A. H. DOOLEY, A. KORÁNYI AND F. RICCI, H -type groups and Iwasawa decompositions, *Adv. Math.* **87(1)** (1991), 1–41.
- [CY] J. CYGAN, Heat kernels for class 2 nilpotent groups, *Studia Math.* **64(3)** (1979), 227–238.
- [FS] G. B. FOLLAND AND E. M. STEIN, “*Hardy spaces on homogeneous groups*”, Mathematical Notes **28**, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1982.
- [G] B. GAVEAU, Principe de moindre action, propagation de la chaleur et estimées sous elliptiques sur certains groupes nilpotents, *Acta Math.* **139(1–2)** (1977), 95–153.
- [H] A. HULANICKI, The distribution of energy in the Brownian motion in the Gaussian field and analytic-hypoellipticity of certain subelliptic operators on the Heisenberg group, *Studia Math.* **56(2)** (1976), 165–173.
- [Hu] D. HUSEMOLLER, “*Fibre bundles*”, Third edition, Graduate Texts in Mathematics **20**, Springer-Verlag, New York, 1994.
- [K] A. KAPLAN, Fundamental solutions for a class of hypoelliptic PDE generated by composition of quadratic forms, *Trans. Amer. Math. Soc.* **258(1)** (1980), 147–153.
- [LP1] F. LUST-PIQUARD, A simple minded computation of heat kernels on Heisenberg groups, *Colloq. Math.* **97(2)** (2003), 233–249.
- [LP2] F. LUST-PIQUARD, Dimension free estimates for discrete Riesz transforms on products of abelian groups, *Adv. Math.* (to appear).
- [LP3] F. LUST-PIQUARD, Riesz transforms associated with the number operator on the Walsh system and the fermions, *J. Funct. Anal.* **155(1)** (1998), 263–285.
- [Par] K. R. PARTHASARATHY, “*An introduction to quantum stochastic calculus*”, Monographs in Mathematics **85**, Birkhäuser Verlag, Basel, 1992.

- [Pa] V. I. PAULSEN, “*Completely bounded maps and dilations*”, Pitman Research Notes in Mathematics Series **146**, Longman Scientific & Technical, Harlow; John Wiley & Sons, Inc., New York, 1986.
- [Pe] G. K. PEDERSEN, “*C*-algebras and their automorphism groups*”, London Mathematical Society Monographs **14**, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London-New York, 1979.
- [P1] G. PISIER, Riesz transforms: a simpler analytic proof of P.-A. Meyer’s inequality, in: “*Séminaire de Probabilités, XXII*”, Lecture Notes in Math. **1321**, Springer, Berlin, 1988, pp. 485–501.
- [P2] G. PISIER, Non-commutative vector valued L_p -spaces and completely p -summing maps, *Astérisque* **247** (1998), 131 pp.
- [R] J. RANDALL, The heat kernel for generalized Heisenberg groups, *J. Geom. Anal.* **6(2)** (1996), 287–316.
- [RS] M. REED AND B. SIMON, “*Methods of modern mathematical physics. I. Functional analysis*”, Second edition, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1980.
- [S] E. M. STEIN, Some results in harmonic analysis in \mathbb{R}^n , for $n \rightarrow \infty$, *Bull. Amer. Math. Soc. (N.S.)* **9(1)** (1983), 71–73.
- [V] V. S. VARADARAJAN, “*Geometry of quantum theory*”, Vol. II, The University Series in Higher Mathematics, Van Nostrand Reinhold Co., New York-Toronto, Ont.-London, 1970.

Département de Mathématiques
Université de Cergy
2 Av. A. Chauvin
95302 Cergy
France
E-mail address: Francoise.Piquard@math.u-cergy.fr

Primera versió rebuda el 20 de març de 2003,
darrera versió rebuda el 2 de març de 2004.

