

ACTA GEOLOGICA HISPANICA, v. 35 (2000), nº 1-2, p. 135-147

Checklist of South American Cenozoic Insects

J.F. PETRULEVICIUS⁽¹⁾ and R.G. MARTINS-NETO⁽¹⁾

(1) *Departamento Científico Paleozoología Invertebrados, Museo de La Plata, Paseo del Bosque s/n, 1900 La Plata, Argentina.*

CONICET. E-mail: levicius@museo.fcnym.unlp.edu.ar

(2) *Laboratório de Paleontologia, FFCL - USP Campus de Ribeirão Preto, Av. Bandeirantes, 3900,*

CEP 14040-901 Ribeirão Preto, SP, Brazil. E-mail: mtmsneto@ffclrp.usp.br

ABSTRACT

This checklist is a compilation of literature about South American Cenozoic insects. The list include 73 named insects belonging to 11 orders. Records are from the Oligocene and Pleistocene of south-eastern Brazil, late Paleocene of north-western Argentina and Paleocene-Eocene of southern Argentina.

Keywords: Fossil insects. Cenozoic. Checklist. South America.

RESUMEN

Esta lista es una compilación de literatura acerca de los insectos del Cenozoico de América del Sur. La lista contiene 73 insectos nominados pertenecientes a 11 órdenes. Los hallazgos provienen del Oligoceno y Pleistoceno del sudeste de Brasil, del Paleoceno tardío del noroeste de Argentina y del Paleoceno-Eoceno del sur de Argentina.

Palabras clave: Insectos fósiles. Cenozoico. Lista de especies nominadas. América del Sur.

INTRODUCTION

South American Cenozoic insects are poorly known, since there are only 73 named species belonging to 11 orders. Findings come from three distinct regions of the subcontinent: south-eastern Brazil, north-western and south Argentina. In the Oligocene of Southeast Brazil, 31 species were discovered; 29 from Tremembé Formation, one from Fonseca Formation, one from Pirassununga Formation and other from the Pleistocene. In Northwest

Argentina 38 species were described from the late Paleocene Maíz Gordo Formation, whereas in southern Argentina, one and two species were discovered from the Paleocene-Eocene Laguna del Hunco and Ventana Formations, respectively. The aim of this work is to list the described genera and species from the South American Cenozoic arranged by orders and families. This checklist also provides information about the type localities, stratigraphic levels and ages; some remarks are also included. To emphasize the dubious nature of some generic as-

signments, these names have been enclosed in quotation marks.

In the present paper, we follow the wing venation nomenclature of Kukalová-Peck (1991).

SYSTEMATIC PALEONTOLOGY

Order: Odonata FABRICIUS

Superfamily: Libelluloidea RAMBUR

Family: Palaeomacromiidae PETRULEVICIUS,
NEL AND MUZÓN

GENUS *Palaeomacromia* PETRULEVICIUS, NEL
AND MUZÓN

Palaeomacromia multicellulata PETRULEVICIUS,
NEL AND MUZÓN 1999: pp. 678-680, figs. 1-2.

Type locality: stratigraphic level and age: La Mendieta, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on a 3/4 forewing.

Order: Blattodea LATREILLE

Family: Blattidae STEPHENS

GENUS “*Amazonina*”HEBARD

Amazonina purpurae PINTO 1991: p. 90, figs. 3-4.

Type locality and age: Mateus Leme County, Minas Gerais, Brazil. Pleistocene?.

Remarks: Age attribution based on vegetal remains.

Order: Orthoptera OLIVIER

Superfamily: Grylloidea LATREILLE

GENUS “*Gryllus*”LINNÉ

Gryllus vociferans COCKERELL 1925b: p. 712,
fig. 2.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on a basal fragment of tegmina. This specimen was placed by Cockerell (1925b) in a broad sense into the genus *Gryllus* LINNÉ, then Cockerell (1936) allocated the species to the collective group *Gryllites* GERMAR. New material collected from the same formation presents similarities with this specimen, both probably belonging to a new genus. However the narrow costal area and subcostal area without cross veins removes this species of the genus *Gryllus* (Martins-Neto and Petrulevicius, in prep.).

GENUS *Gryllites* GERMAR

Gryllites vocalis COCKERELL 1936: p. 9, fig. 19.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on a basal fragment of tegmina. This specimen was allocated by Cockerell (1925b) to the genus *Gryllites* GERMAR. The species is available only for the purposes of the Principle of Homonymy (ICZN 1985-41, Art. 20). New material collected from the same formation presents similarities with this specimen, both probably belonging to a new genus (Martins-Neto and Petrulevicius, in prep.).

Order: Dermaptera DE GEER

Family: Labiduridae VERHOEFF

GENUS “*Carcinophora*”SCUDDER

Psalis pachyura COCKERELL 1925b: p. 712,
fig. 3.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on forceps. Cockerell (1925b) placed the species into the genus *Psalis* ‘in a broad sense’, a latter synonymous of *Carcinophora* SCUDDER (Carpenter, 1992).

Order: Isoptera BRULLÉ

Family: Mastotermitidae DESNEUX

GENUS *Spargotermes* EMERSON

Spargotermes costalimai EMERSON 1965: pp. 19-24, fig. 4.

Type locality; stratigraphic level and age: Minas Gerais State, Brazil; Fonseca Basin, Fonseca Formation, Oligocene.

Remarks: Species description is based on isolated wings. Emerson (1965) mentioned an Eocene age for the remains, updated by Martins-Neto (1997b) to the Oligocene.

Order: Hemiptera LINNÉ
Suborder: Homoptera LEACH
Family: Cercopidae WESTWOOD

GENUS *Parafitopterix* MARTINS-NETO

Parafitopterix duateae MARTINS-NETO 1989: pp. 376-378, pls. I A-B, fig. 2 J.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Paraná Basin, Pirassununga Formation, Oligocene.

Remarks: Species description is based on a tegmina.

Superfamily: Fulgoroidea LATREILLE

GENUS *Hipocixius* COCKERELL

Hipocixius oblitescens COCKERELL 1926a: pp. 501-502, fig. 1.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on a middle part of a tegmina. Placed in Fulgoroidea (“apparently related to Cixiidae”) by Cockerell (1926a) and in Homoptera uncertain by Carpenter (1992). This species could not be assigned to any family of Fulgoroidea because of the lack of the apical part of CuP and AA₃₊₄, but seems not to be related to Cixiidae because the AA₃₊₄ seems to finish at the same time with CuP.

Family: Flatidae SPINOLA

GENUS “*Ormenis*” STÅL

Ormenis devinctus COCKERELL 1926a: pp. 502-503, fig. 2.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on a costal and distal tegmina fragment.

Family: Aetalionidae *sensu* EVANS

GENUS *Tremembaetalion* MARTINS-NETO

Tremembaetalion mimum MARTINS-NETO 1998a: p. 61, fig. 1 N.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen without legs.

Family: Cicadellidae LATREILLE

GENUS *Taubocicadellina* MARTINS-NETO

Taubocicadellina breviptera MARTINS-NETO 1998a: p. 60, fig. 1 B.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen without legs.

GENUS *Tremembellina* MARTINS-NETO

Tremembellina micropcellata MARTINS-NETO 1998a: p. 60, fig. 1 M.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen without legs.

GENUS *Trulaxia* MARTINS-NETO

Trulaxia primula MARTINS-NETO 1998a: p. 60, fig. 1 L.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen without legs.

Suborden: Heteroptera LATREILLE
Family: Veliidae DOUGLAS AND SCOTT

GENUS *Paleohebrus* MARTINS-NETO

Paleohebrus tremembeensis MARTINS-NETO 1997b: pp. 67-68, pl. B, fig. 2.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen. Originally described as Hebridae by Martins-Neto (1997b).

Family: Pentatomidae LEACH

GENUS *Taubatecoris* MARTINS-NETO

Taubatecoris quadntiformis MARTINS-NETO 1997b: p. 67, pl. A, fig. 1.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen.

Family: Corixidae LEACH

GENUS *Taubarixa* MARTINS-NETO

Taubarixa macocelata MARTINS-NETO 1998a: p. 59, pl. 1 A, fig. 1 A.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen without legs.

GENUS *Tauborixella* MARTINS-NETO

Tauborixella santosae MARTINS-NETO 1998a: p. 59, pl. 1 B, figs. 1 C-G.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen without legs.

GENUS *Tauborixiellopsis* MARTINS-NETO

Tauborixiellopsis bεviclavata MARTINS-NETO 1998a: p. 60, pl. 1 C, fig. 1 H-K.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen.

Family: Coreidae LEACH

GENUS “*Corizus*” FALLÉN

Corizus deflagratus COCKERELL 1926a: p. 503, fig. 3.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on a nearly complete (ventral view) specimen.

Order: Coleoptera LINNÉ
Family: Carabidae LATREILLE

GENUS *Tremembecarabus* MARTINS-NETO

Tremembecarabus rotundus MARTINS-NETO 1998a: p. 61, pl. 1 D, fig. 2 A.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen without head and legs.

Family: Psephenidae LACORDAIRE

GENUS *Psephenella* MARTINS-NETO

Psephenella ferrera MARTINS-NETO 1998a: p. 62, pl. 2 B, fig. 2 B.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen.

Family: Meloidae GUILLENHALL

GENUS *Microbasis* MARTINS-NETO

Microbasis longinota MARTINS-NETO 1998a: p. 62, pl. 2 A, fig. 2 C.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen.

Family: uncertain

GENUS *Carabites* HEER

Carabites harringtoni COCKERELL 1936: p. 4; fig. 2.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: This specimen was allocated by Cockerell (1925b) to the collective group Carabites HEER 1852. The species is available only for the purposes of the Principle of Homonymy (ICZN 1985-41, Art. 20). At present knowledge the species could not be assigned to any family because of the lack of the synapomorphic characters in the specimen.

Carabites schueli COCKERELL 1926b: p. 321, fig. 11.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on an isolated elytra. Same nomenclatural and taxonomic considerations as in Carabites harringtoni COCKERELL.

GENUS *Cerambycites* COCKERELL

Cerambycites wilmatae COCKERELL 1925a: p. 712, fig. 1.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on an isolated elytra. This specimen was allocated by Cockerell (1925b) to the collective group Cerambycites. The species is available only for the purposes of the Principle of Homonymy (ICZN 1985-41, Art. 20). At present knowledge the species could not be assigned to any family because of the lack of the synapomorphic characters in the specimen.

GENUS *Chrysomelites* HEER

Chrysomelites danielis COCKERELL 1926b: p. 321, fig. 10.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on an isolated elytra. This specimen was allocated by Cockerell (1925b) to the collective group Chrysomelites HEER. The species is available only for the purposes of the Principle of Homonymy (ICZN 1985-41, Art. 20). At present knowledge the species could not be assigned to any family be-

cause of the lack of the synapomorphic characters in the specimen.

GENUS *Curculionites* HEER

Curculionites angustior COCKERELL AND WAGNER, Cockerell, 1936: p. 6, fig. 11.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on an isolated elytra. This specimen was allocated by Cockerell (1925b) to the collective group *Curculionites* HEER. The species is available only for the purposes of the Principle of Homonymy (ICZN 1985-41, Art. 20). At present knowledge the species could not be assigned to any family because of the lack of the synapomorphic characters in the specimen.

Curculionites epistictus COCKERELL AND WAGNER, Cockerell, 1936: p. 8, fig. 8.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Same nomenclatural and taxonomic considerations as in *Curculionites angustior* COCKERELL AND WAGNER.

Curculionites eustictus COCKERELL AND WAGNER, Cockerell, 1936: p. 8, fig. 9.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Same nomenclatural and taxonomic considerations as in *Curculionites angustior* COCKERELL AND WAGNER.

Curculionites harringtoni COCKERELL 1925a: p. 4, pl. 1, fig. 5.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Same nomenclatural and taxonomic considerations as in *Curculionites angustior* COCKERELL AND WAGNER.

Curculionites jujuyensis COCKERELL 1925a: p. 4, pl. 1, fig. 6.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Same nomenclatural and taxonomic considerations as in *Curculionites angustior* COCKERELL AND WAGNER.

Curculionites latiusculus COCKERELL AND WAGNER, Cockerell, 1936: p. 6, fig. 8.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Same nomenclatural and taxonomic considerations as in *Curculionites angustior* COCKERELL AND WAGNER.

Curculionites magdalinus COCKERELL AND WAGNER, Cockerell, 1936: p. 9, fig. 6.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Same nomenclatural and taxonomic considerations as in *Curculionites angustior* COCKERELL AND WAGNER.

Curculionites megastictus COCKERELL AND WAGNER, Cockerell, 1936: p. 9, fig. 17.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Same nomenclatural and taxonomic considerations as in *Curculionites angustior* COCKERELL AND WAGNER.

Curculionites microstictus COCKERELL AND WAGNER, Cockerell, 1936: p. 8, fig. 10.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Same nomenclatural and taxonomic considerations as in *Curculionites angustior* COCKERELL AND WAGNER.

***Curculionites pamstictus* COCKERELL AND WAGNER**, Cockerell, 1936: p. 9; fig. 13.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Same nomenclatural and taxonomic considerations as in *Curculionites angustior* COCKERELL AND WAGNER.

***Curculionites stebingeri* COCKERELL** 1926: pp. 318-319, fig. 6.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Same nomenclatural and taxonomic considerations as in *Curculionites angustior* COCKERELL AND WAGNER.

***Curculionites sunchalicus* COCKERELL AND WAGNER**, Cockerell, 1936: p. 6, fig. 14.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Same nomenclatural and taxonomic considerations as in *Curculionites angustior* COCKERELL AND WAGNER.

***Curculionites wielandi* COCKERELL** 1925a: p. 5, pl. 1, fig. 8.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Same nomenclatural and taxonomic considerations as in *Curculionites angustior* COCKERELL AND WAGNER.

GENUS *Elaterites* HEER

***Elaterites bruchi* COCKERELL** 1926b: pp. 320-321, fig. 9.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on an isolated elytra. This specimen was allocated by Cockerell (1925b) to the collective group *Elaterites* HEER. Carpenter (1992) placed the species in Coleoptera suborder and family uncertain. The species is available only for the purposes of the Principle of Homonymy (ICZN 1985-41, Art. 20). At present knowledge the species could not be assigned to any family because of the lack of the synapomorphic characters in the specimen.

***Elaterites micostictus* COCKERELL** 1926b: p. 320, fig. 8.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Same nomenclatural considerations as *Elaterites bruchi* COCKERELL.

GENUS *Otiorhynchites* HAUPT

***Otiorhynchites aterrimus* COCKERELL** 1925a: p. 3, pl. 1, fig. 1.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on an isolated elytra. This specimen was allocated by Cockerell (1925b) to the genus *Otiorhynchites* HAUPT. The species is available only for the purposes of the Principle of Homonymy (ICZN 1985-41, Art. 20). At present knowledge the species could not be assigned to any family because of the lack of the synapomorphic characters in the specimen.

***Otiorhynchites crassus* COCKERELL** 1936: p. 6, fig. 12.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Same nomenclatural and taxonomic considerations as in *Otiorhynchites aterrimus* COCKERELL.

GENUS *Tenebrionites* COCKERELL

Tenebrionites inclinans COCKERELL 1925a: p. 5, pl. 1, fig. 7.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on an isolated elytra. This specimen was allocated by Cockerell (1925b) to the collective group *Tenebrionites* COCKERELL. Carpenter (1992) placed the species into Tenebrionidae. The species is available only for the purposes of the Principle of Homonymy (ICZN 1985-41, Art. 20). At present knowledge the species could not be assigned to any family because of the lack of the synapomorphic characters in the specimen.

Species doubtfully assigned to recent genera

GENUS “*Anthicus*” PAYKULL

Anthicus (s. l.) *sepultulus* COCKERELL 1926b: p. 318, fig. 5.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on a nearly complete specimen without antennae and legs. Allocated by Cockerell (1926b) to the genus *Anthicus* in a broad sense. Carpenter (1992) placed this species in “recent genera of Coleoptera with doubtfully assigned species”.

GENUS “*Anthonomus*” GERMAR

“*Anthonomus*” (?) *sunchalensis* COCKERELL 1925a: p. 4, pl. 1, figs. 3-4.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on an isolated elytra. Carpenter (1992) placed this species in “recent genera of Coleoptera with doubtfully assigned species”.

GENUS “*Cossonus*” CLAIRVILLE AND SCHELLEMBERG

Cossonus devoratus COCKERELL 1925a: p. 3, pl. 1, fig. 2.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on an isolated elytra. Carpenter (1992) placed this species in “recent genera of Coleoptera with doubtfully assigned species”.

GENUS “*Cryptophagus*” HERBST

Cryptophagus sunchalensis COCKERELL 1926b: p. 317, fig. 4.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on a nearly complete specimen without legs.

GENUS “*Haruspex*”

Haruspex (?) *defectus* COCKERELL 1926a: p. 503, fig. 4.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on an isolated elytra. Carpenter (1992) placed this species in “recent genera of Coleoptera with doubtfully assigned species”.

GENUS “*Lebia*” LATREILLE

Lebia harrelli COCKERELL 1936: pp. 4-5, fig. 3.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on an isolated elytra. Carpenter (1992) placed this species in “recent genera of Coleoptera with doubtfully assigned species”.

GENUS “*Podabrus*” WESTWOOD

Podabrus (?) *santaritensis* COCKERELL 1936: p. 5, fig. 5.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on an isolated elytra. Carpenter (1992) placed this species in “recent genera of Coleoptera with doubtfully assigned species”.

GENUS “*Trichodes*” HERBST

Trichodes stebingeri COCKERELL 1936: p. 5, fig. 4.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on an isolated elytra. Carpenter (1992) placed this species in “recent genera of Coleoptera with doubtfully assigned species”.

Order: Diptera LINNÉ

Family: Hybotidae *sensu* CHVÁLA

Subfamily: Trachydromiinae MEIGIN

GENUS *Archaeodrapetiops* MARTINS-NETO, VIEIRA, KUCERA-SANTOS AND FRAGOSO

Archaeodrapetiops (?) *elongata* MARTINS-NETO, VIEIRA, KUCERA-SANTOS AND FRAGOSO 1992: pp. 37-38, pl. 2, figs. 3-4, figs. 1 E, 2 C.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen including chaetotaxy.

Archaeodrapetiops mezzalini MARTINS-NETO, VIEIRA, KUCERA-SANTOS AND FRAGOSO 1992: p. 36, pl. 1, figs. 1-3, 6, figs. 1 G, 2 F.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen.

Archaeodrapetiops nefra MARTINS-NETO, VIEIRA, KUCERA-SANTOS AND FRAGOSO 1992: pp. 34-36, pl. 1, figs. 4-5, 7; figs. 1 A, 2 E.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen.

Archaeodrapetiops transversa MARTINS-NETO, VIEIRA, KUCERA-SANTOS AND FRAGOSO 1992: p. 37, pl. 1, fig. 8; pl. 2, figs. 5-6; figs. 1 C, 2 H.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen.

Subfamily: Hybotinae *sensu* CHVÁLA

GENUS *Eternia* MARTINS-NETO, VIEIRA, KUCERA-SANTOS AND FRAGOSO

Eternia papaweroi MARTINS-NETO, VIEIRA, KUCERA-SANTOS AND FRAGOSO 1992: p. 40, pl. 2, fig. 2; pl. 3, figs. 2, 4; figs. 1 F, 2 I.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen.

GENUS *Tremabella* MARTINS-NETO, VIEIRA, KUCERA-SANTOS AND FRAGOSO

Tremabella gacilis MARTINS-NETO, VIEIRA, KUCERA-SANTOS AND FRAGOSO 1992: pp. 38-39, pl. 3, figs. 1, 3; figs. 1 B, 2 H.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen.

Family: Tabanidae LEACH

GENUS *Tabanus* LINNÉ

Tabanus tremembeensis MARTINS-NETO 1997a: pp. 53-54, pl. 1, fig. 1.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen.

Order: Trichoptera KIRBY

Family: Limnephilidae *sensu* FLINT

GENUS *Indusia* BOSC

Indusia suguioi MARTINS-NETO 1989: pp. 378-379, pl.1, fig. F.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a caddis-case.

Family: uncertain

GENUS “*Molanna*” CURTIS

Molanna (?) *derosa* COCKERELL 1925a: pp. 2-3, fig. 1.

Type locality; stratigraphic level and age: Sunchal, Jujuy Province, Argentina; Maíz Gordo Formation, late Paleocene.

Remarks: Species description is based on a forewing fragment. Placed in Molannidae by Cockerell (1925a).

Order: Lepidoptera LINNÉ

Family: Nepticulidae *sensu* OPLER

GENUS “*Nepticula*” OPLER

Nepticula (?) *almeidae* MARTINS-NETO 1989: p. 381, pl. 1, fig. C.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on leaf-mines. Generic attribution was merely tentative.

Family: Gracillaridae *sensu* OPLER

GENUS “*Phyllonoryctes*” HUBNER

Phyllonoryctes (?) *oliveirae* MARTINS-NETO 1989: pp. 381-382, pl. 1, fig. D.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on leaf-mines. Generic attribution was merely tentative.

Family: Cossidae LEACH

GENUS *Kleopatra* MARTINS-NETO

Kleopatra nemogypsia MARTINS-NETO 1998b: p. 76, pl. 1, fig. B; figs. 1 C, 2.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a forewing.

Kleopatra noctodiva MARTINS-NETO 1998b: p. 75, pl. 1, figs. B; figs. 1 A, 2.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a forewing.

Family: Pyralidae LINNÉ
Subfamily: Chrysauginae LEDERER

GENUS ***Petisca*** MARTINS-NETO

Petisca dryellina MARTINS-NETO 1998b: pp. 63-64, pl. 3, fig. C; fig. 2 D.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a forewing.

Family: Nymphalidae *sensu* RYCHARD AND DAVIES

Subfamily: Danainae *sensu* RYCHARD AND DAVIES

GENUS ***Archaeolycorea*** MARTINS-NETO

Archaeolycorea ferreirai MARTINS-NETO 1989: p. 380, fig. 4 A.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a forewing.

Subfamily: Satyrinae *sensu* KRISTENSEN

GENUS ***Neorinella*** MARTINS-NETO, KUCERA-SANTOS, VIEIRA AND FRAGOSO

Neorinella garciae MARTINS-NETO, KUCERA-SANTOS, VIEIRA AND FRAGOSO 1993: pp. 6-8, pl. 1-2, figs. 2-3.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen well preserved including colour pattern.

Family: Noctuidae *sensu* KRISTENSEN

GENUS ***Philodarchia*** MARTINS-NETO

Philodarchia cigana MARTINS-NETO 1998b: pp. 77-78, pl.1, fig. C; fig. 1 C.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a complete specimen.

Order: Hymenoptera LINNÉ
Family: Ichneumonidae LATREILLE

GENUS ***Paratilgidopsis*** MARTINS-NETO

Paratilgidopsis paecursora MARTINS-NETO 1998: p. 63, pl. 3, fig. B, fig. 3 B.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete fore wing.

GENUS ***Taubatehymen*** MARTINS-NETO

Taubatehymen minuta MARTINS-NETO 1998: pp. 62-63, pl. 3, fig. A, fig. 3 A.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete hind wing.

Family: Siricidae BILLBERG
Subfamily: Siricinae BILLBERG

GENUS *Urocerus* GEOFFROY

Urocerus patagonicus FIDALGO AND SMITH 1987: pp. 63-65, figs. 1-3.

Type locality; stratigraphic level and age: São Paulo State, Brazil; Taubaté Basin, Tremembé Formation, Oligocene.

Remarks: Species description is based on a nearly complete specimen.

Family: Formicidae LATREILLE

GENUS *Ameghinoia* VIANA AND HAEDO-ROSSI

Ameghinoia piazzi VIANA AND HAEDO-ROSSI 1957: pp. 109-113, pl. 1-3, figs. 1-3.

Type locality; stratigraphic level and age: Pichileufú, Río Negro Province, Argentina; Ventana Formation, Paleocene-Eocene.

Remarks: Species description is based on a nearly complete specimen. This species was assigned by Viana and Haedo Rossi (1957) to the Subfamily Ponerinae. Naumann (1991), allocated the species to the Subfamily Myrmeciinae.

GENUS *Polanskiella* ROSSI DE GARCÍA

Polanskiella smekali ROSSI DE GARCÍA 1983: pp. 19-20, pl. 1, figs. 1-3, 5.

Type locality; stratigraphic level and age: Confluencia, Río Negro Province, Argentina; Ventana Formation, Paleocene-Eocene.

Remarks: Species description is based on a nearly complete specimen. This species was assigned by Rossi de García (1983) to the Subfamily Ponerinae. Probably this species has to be placed into the Subfamily Myrmeciinae, after the examination of the il-

lustrated material, but we think that a revision of the type material is necessary to arrive to a more precise decision.

ACKNOWLEDGEMENTS

Funds for this research were provided in part by CONICET. Thanks are also due to Drs. T. Deuve, A. Nel, A. Riccardi and C. Digiani for valuable help and discussion. This paper has been much improved by the comments and suggestions of one anonymous referee.

REFERENCES

- Carpenter, F.M., 1992. Superclass Hexapoda. In: L. Kaesler et al. (eds.). Treatise of Invertebrate Paleontology, part R, Arthropoda, 4(3-4), 1-655, Boulder and Lawrence, The Geological Society of America Inc. and University of Kansas.
- Cockerell, T.D.A., 1925a. Tertiary Insects from Argentina. Proc. U. S. Nat. Mus., 68(1), 1-5.
- Cockerell, T.D.A., 1925b. Tertiary fossil Insects from Argentina. Nature, 116, 711-712.
- Cockerell, T.D.A., 1926a. Tertiary fossil Insects from Argentina. Amer. Jour. Sci. (Series 5), 11, 501-504.
- Cockerell, T.D.A., 1926b. Some Tertiary fossil Insects. Ann. Mag. Nat. Hist. (Series 9), 18, 313-324.
- Cockerell, T.D.A., 1936. The fauna of the Sunchal (or Margas Verdes) Formation, Northern Argentina. Amer. Mus. Novit., 886, 1-10.
- International Code of Zoological Nomenclature., Adopted by the XX General Assembly of the International Union of Biological Sciences, 1985. International Trust for Zoological Nomenclature, 338 pp., London.
- Emerson, A.E., 1965. Review of the Mastotermitidae (Isoptera) including a new fossil genus from Brazil. Am. Mus. Nov., 2236, 1-45.
- Fidalgo, P., Smith, D.R., 1987. A fossil Siricidae (Hymenoptera) from Argentina. Ent. News, 98(2), 63-66.
- Kukalová-Peck, J., 1991. Fossil history and the evolution of hexapod structures. In: I.D. Naumann (ed.). The insects of Australia, A textbook for students and research workers, vol. 1, 141-179. Melbourne University Press, Melbourne.
- Martins-Neto, R.G., 1989. Novos Insetos terciários do Estado de São Paulo. Rev. bras. Geoc., 19(3), 375-386.
- Martins-Neto, R.G., 1997a. Dípteros (Insecta) da Formação Tremembé, Bacia de Taubaté, Oligoceno do Estado de São Paulo. III - Família Tabanidae. Acta Geol. Leopoldensia, 20(44), 51-57.
- Martins-Neto, R.G., 1997b. A Paleontomofauna da Formação Tremembé (Bacia de Taubaté) Oligoceno do Estado de São

- Paulo: descrição de novos hemípteros (Insecta). Rev. Universidade Guarulhos, Sér. Geociências, 2(6), 66-69.
- Martins-Neto, R.G., 1998a. A Paleontomofauna da Formação Tremembé (Bacia do Taubaté) Oligoceno do Estado de São Paulo: novos Hemiptera, Auchenorrhyncha, Hymenoptera, Coleoptera e Lepidoptera (Insecta). Rev. Universidade Guarulhos, Sér. Geociências, 3(6), 58-70.
- Martins-Neto, R.G., 1998b. A Paleontomofauna da Formação Tremembé (Bacia do Taubaté) Oligoceno do Estado de São Paulo: descrição de novos lepidópteros (Insecta). Acta Geol. Leopoldensia, 21(46-47), 75-82.
- Martins-Neto, R.G., Kucera-Santos, J.C., Vieira, F.R.M., Fragoso, L.M.C., 1993. Nova espécie de Borboleta (Lepidoptera: Nymphalidae: Satirynae) da Formação Tremembé, Oligoceno do Estado de São Paulo. Acta Geol. Leopoldensia, 16(37), 5-16.
- Martins-Neto, R.G., Vieira, F.R.M., Kucera-Santos, J.C., Fragoso, L.M.C., 1992. Dípteros (Insecta, Empidoidea) da Formação Tremembé, Bacia do Taubaté, Oligoceno do Estado de São Paulo. I - Família Hybotidae. Acta Geol. Leopoldensia, 15(36), 31-48.
- Naumann, I.D., 1991. Hymenoptera. In: I.D. Naumann et al. (eds.). The insects of Australia, vol. 1, 916-1000, Carlton, Melbourne University Press.
- Petrulevicius, J., Nel, A., Muzón, J. 1999. A new Libelluloid family from the Upper Paleocene of Argentina. Palaeontology, 42(4), 667-682.
- Pinto, I.D., 1991. The fossil Blattoid genus *Amazonina*. Taxonomy and geographical distribution. Pesquisas, 18(1), 88-92.
- Rossi de García, E., 1983. Insectos de la Formación Ventana (Eoceno). Provincia de Neuquén. Rev. Asoc. Geol. Arg., 38(1), 17-23.
- Viana, M.J., Haedo-Rossi, J.A., 1957. Primer hallazgo en el hemisferio sur de Formicidae extinguidos y catálogo mundial de los Formicidae fósiles. Primera parte. Ameghiniana, 1(1-2), 108-113.