

Eusyllinae, Exogoninae and Autolytinae (Syllidae, Annelida, Polychaeta) from the Chafarinas Islands (Alboran Sea, W Mediterranean)

E. López & G. San Martín

López, E. & San Martín, G., 1997. Eusyllinae, Exogoninae and Autolytinae (Syllidae, Annelida, Polychaeta) from the Chafarinas Islands (Alboran Sea, W Mediterranean). *Misc. Zool.*, 20.2: 101-111.

Eusyllinae, Exogoninae and Autolytinae (Syllidae, Annelida, Polychaeta) from the Chafarinas Islands (Alboran Sea, W Mediterranean).—A study on the benthic invertebrate communities of the Chafarinas Islands was carried out recently, paying special attention to some groups, including polychaetes. In this work the results of the study of subfamilies Eusyllinae, Exogoninae, and Autolytinae (Annelida, Polychaeta, Syllidae) are given, citing 48 species. One of these, *Amblyosyllis inmatura* Langerhans, 1879, is recorded for first time in the Mediterranean Sea and compared with previously known Mediterranean species of the genus. Some variations from typical form are described for two *Grubeosyllis* Verrill, 1900 species: *G. clavata* (Claparède, 1863) and *G. vieitezi* (San Martín, 1984). Specimens of a new species of genus *Pionosyllis* (Eusyllinae) and other of genus *Autolytus* (Autolytinae) were found.

Key words: Polychaeta, Syllidae, Western Mediterranean.

(Rebut: 20 I 97; Acceptació condicional: 1 IV 97; Acc. definitiva: 2 VII 97)

E. López & G. San Martín, Depto. de Biología, Unidad de Zoología, Lab. de Biología Marina e Invertebrados, Fac. de Ciencias, Univ. Autónoma de Madrid, Campus de Canto Blanco, 28049 Madrid, España (Spain).

E-mail: eduardo.lopez@uam.es

This study was financially supported by Comisión Interministerial de Ciencia y Tecnología (CICYT), as part of the research project "Invertebrados marinos bentónicos de sustratos duros de las Islas Chafarinas" (PB89-0204-C02).

Introduction

The Chafarinas Islands are a small Spanish archipelago close to the Mediterranean coast of Morocco. Although there is a wide body of literature on the distribution of polychaete species in the Western Mediterranean, few works have studied the polychaete fauna of the Mediterranean coast of North-western Africa (AMOREUX, 1976 in the Gibraltar Strait; WESTHEIDE, 1972; CANTONE et al., 1978 and ZGHAL & BEN AMOR, 1980 in the Tunisian Coast; SAN MARTÍN & PARAPAR, 1990; PARAPAR & SAN MARTÍN, 1992, and PARAPAR et al., 1993 in the vicinity of Ceuta).

From February 1991 to July 1993 a number of field studies were carried out by research staff of the Universidad de Alcalá de Henares, the Universidad Autónoma de Madrid, and the Universidad de Valencia.

The main objective was the study of benthic marine invertebrates. Study of the polychaetous annelid fauna was emphasized. Ecological and biogeographical data of the species of these subfamilies in the Chafarinas Islands are discussed. This paper follows a previous one (LÓPEZ et al., 1996) in which a brief description of the zone is given.

Material and methods

The list of localities and samples can be found in LÓPEZ et al. (1996). In the same paper, the used methods are described. Descriptions and references on all common species can be found in previous general works (DAY, 1967; WESTHEIDE, 1974; CAMPOY, 1982; SAN MARTÍN, 1984, 1990; SARDÀ, 1984a; UEBELACKER, 1984; ALÓS, 1988; NÚÑEZ et al., 1992). For newly described or poorly known species, this information can be found in more specific works: MARTÍN & ALÓS (1987) for *Autolytus sardai*, RUSSELL (1989) and OLANO et al. (in press) for *Sphaerosyllis belizensis*, MARTÍN et al. (1990) for *Miscellania dentata*, LANERA et al. (1994) for *Exogone gambiae* and, LÓPEZ et al. (1997) for *Pionosyllis serratisetosa* and *Autolytus longoprimicirratus*.

For each species, samples where it has been found and number of specimens per sample (in brackets) are given; samples are characterized by a capital letter (field station) and a number (for quantitative samples) or another capital letter (for qualitative samples). A brief list of references, comments and the known world distribution are also given for some interesting species.




Fig. 1. Anterior end of pharynx: A. *Amblyosyllis dorsigera* Claparède, 1863; B. *Amblyosyllis inmatura* Langerhans, 1879; C. *Amblyosyllis madeirensis* Langerhans, 1879. (Scales A, B: 20 µm; C: 48 µm.)

Extremo anterior de la faringe: A. *Amblyosyllis dorsigera* Claparède, 1863; B. *Amblyosyllis inmatura* Langerhans, 1879; C. *Amblyosyllis madeirensis* Langerhans, 1879. (Escalas A, B: 20 µm; C: 48 µm.)

Results

About 20,700 specimens belonging to the subfamilies Eusyllinae, Exogoninae, and Autolytinae were studied. Forty eight species were recorded.

Family Syllidae Grube 1850

Subfamily Eusyllinae Rioja, 1925

Genus *Amblyosyllis* Grube, 1857

Amblyosyllis dorsigera Claparède, 1864
(fig. 1A)

Material examined: G/1 (1).

Amblyosyllis inmatura Langerhans, 1879
(fig. 1B)

Amblyosyllis inmatura Langerhans, 1879.
LANGERHANS (1879): 561, Taf. XXXII, fig. 20.

Material examined: K/3 (1).

Remarks: this species is recorded for the first time since its original description. A previous record of this species as *Amblyosyllis cf. inmatura* (CAMPYO, 1982) was actually based on two specimens of *A. dorsigera*. Its most characterizing feature is the pharynx mouth (fig. 1B), with six small and unidentate teeth, very different to those of *A. dorsigera* Claparède, 1864 (fig. 1A) and *A. madeirensis* Langerhans, 1879 (fig. 1C); both species commonly found on the Spanish Mediterranean coasts. *A. inmatura* could be a juvenile stage of another *Amblyosyllis* species, but the study of more abundant and complete material is needed.

World distribution: Madeira.

First record for Mediterranean and Spanish fauna.

Amblyosyllis madeirensis Langerhans, 1879
(fig. 1C)

Material examined: C/1 (1); C/2(1); C/3 (1); F/1(1); F/A (3); H/D (1); L/2 (1); M/1 (1); O/1 (3); X/A (5).

Genus *Ehlersia* Quatrefages, 1865 enmend.
Langerhans, 1879

Ehlersia ferrugina Langerhans, 1881

Material examined: A/1 (2); A/B (1); E/1 (1); F/2 (5); F/A (1); G/1 (2); G/A (1); G/B (8); H/1 (1); I/1 (1);

J/1 (27); J/1' (1); J/3 (1); J/A (10); J/A (1); N/1 (1); O/1 (8); X/A (28).

Remarks: this species was formerly considered as belonging to the subfamily Syllinae. In a recent work the change of its taxonomical position was proposed (NÚÑEZ & SAN MARTÍN, 1996).

Genus *Eusyllis* Malmgren, 1867

Eusyllis assimilis Marenzeller, 1875

Material examined: A/1 (1); C/1 (1); D/1 (3); F/1 (1); F/A (1); G/1 (1); L/A (1); N/1 (6); O/1 (1).

Eusyllis lamelligera Marion & Bobretzky, 1875

Material examined: A/1 (2); A/2 (7); B/2 (3); C/2 (3); D/1 (5); F/1 (20); F/2 (2); F/A (7); G/1 (1); G/A (1); G/B (3); H/A (1); J/1 (1); L/A (6); L/B (1); L/D (2); N/1 (3); N/B (1); O/1 (1).

Genus *Miscellania* Martín, Alós & Sardà, 1990

Miscellania dentata Martín, Alós & Sardà, 1990

Material examined: B/2 (14); X/A (2).

Genus *Odontosyllis* Claparède, 1863

Odontosyllis ctenostoma Claparède, 1868

Material examined: A/1 (1); A/2 (3); B/2 (1); B/A (1); C/2 (1); C/3 (2); E/1 (4); F/2 (2); H/1 (1); H/D (2); I/1 (4); J/2 (1); K/3 (1); L/1 (2); L/A (1); M/1 (1); N/2 (1); N/B (1); O/1 (2); X/A (2); X/B (17).

Odontosyllis fulgorans (Audouin & Milne-Edwards, 1834)

Material examined: A/1 (3); B/1 (2); B/2 (1); D/2 (2); E/1 (3); F/1 (10); F/2 (5); H/1 (2); H/D (2); I/1 (1); J/2 (5); K/2 (1); L/A (1); N/2 (1); O/1 (16); X/A (5).

Odontosyllis gibba Claparède, 1863

Material examined: A/1 (3); A/2 (3); C/1 (3); C/2 (2); C/3 (1); D/1 (6); F/1 (11); F/A (3); G/1 (7); G/A (1); K/1 (4); L/A (7); L/C (5); N/1 (3).

Genus *Pionosyllis* Malmgren, 1867

Pionosyllis lamelligera Saint-Joseph, 1886

Material examined: A/1 (42); A/2 (26); A/A (1); A/B (2); B/1 (1); B/2 (11); C/1 (8); C/2 (33); C/3 (1);

C/4 (2); D/1 (3); E/1 (1); F/1 (91); F/2 (64); F/A (4); G/1 (7); G/2 (3); G/3 (8); G/A (20); G/B (49); H/1 (32); H/D (8); I/1 (5); J/1 (18); J/2 (2); J/3 (2); J/A (1); L/2 (5); L/A (12); L/C (3); L/D (2); M/1 (2); N/1 (7); N/2 (1); N/B (5); O/1 (55); X/A (134); X/D (2); X/E (1).

Pionosyllis longocirrata Saint-Joseph, 1886

Material examined: A/1 (1); C/2 (1); D/1 (1); D/2 (1); F/1 (5); F/A (5); G/1 (1); I/1 (1); J/1 (3); L/A (1); K/1 (2); N/1 (1); O/1 (5).

Pionosyllis pulligera (Krohn, 1852)

Material examined: A/1 (16); A/2 (41); C/1 (6); C/2 (11); C/3 (139); C/4 (15); D/1 (11); D/2 (14); D/3 (14); E/1 (15); F/1 (27); F/2 (3); F/A (4); G/1 (1); G/2 (1); G/3 (5); H/1 (6); H/2 (17); I/3 (7); J/2 (24); K/1 (11); K/2 (11); L/1 (1); L/2 (8); L/A (4); L/C (1); M/1 (100); N/1 (28); N/2 (23); O/1 (2); X/A (2); X/D (2).

Pionosyllis serratisetosa López, San Martín & Jiménez, 1997

Material examined: A/1 (1); B/2 (1); C/2 (2); F/1 (1); F/2 (5); G/B (3).

Genus *Syllides* Oersted, 1845

Syllides edentatus (Westheide, 1974)

Material examined: I/1 (3).

Syllides fulvus (Marion & Bobretzky, 1875)

Material examined: A/2 (3); B/1 (2); C/2 (2); C/3 (1); C/4 (1); D/1 (1); E/1 (2); F/1 (8); F/2 (1); F/A (2); G/1 (1); G/2 (1); G/A (2); G/B (4); H/D (4); J/1 (3); J/A (1); L/1 (1); L/2 (1); L/A (1); L/C (1); M/1 (1); X/A (21).

Subfamily Exogoninae Rioja, 1925

Genus *Brania* Quatrefages, 1865

Brania arminii (Langerhans, 1881)

Material examined: E/1 (2); F/A (1); J/1 (2); J/A (2).

Brania pusilla (Dujardin, 1839)

Material examined: A/1 (6); A/2 (4); A/B (1); C/1 (3); C/4 (7); D/3 (1); E/1 (9); F/1 (5); F/2 (1); G/2 (11); G/3 (10); G/A (3); G/B (8); J/A (11); H/2 (24); I/1 (15); J/1 (4); J/1' (1); J/2 (2); J/3 (6); K/2 (3); K/3 (7); L/2 (10); L/A (3); L/D (5); N/1 (7); N/B (3); O/1 (61); O/D (1); X/A (156); X/B (3); X/C (1).

Genus *Exogone* Oersted, 1845

Exogone (Parexogone) gambiae Lanera, Sordino & San Martín, 1994

Material examined: J/1 (3); J/3 (1); J/A (1); M/1 (1); X/A (1).

Exogone (Exogone) cf. dispar (Webster, 1879)

Material examined: A/B (1)

Remarks: this specimen is different from those typical of *E. dispar* (Webster, 1879) only by having a central antenna of the same length as the lateral antennae; this could be due to a loose of the same and the later regeneration process.

Exogone (Exogone) naidina Oersted, 1845

Material examined: A/1 (162); A/2 (578); A/B (11); B/1 (66); B/2 (13); B/A (10); C/1 (49); C/2 (127); C/3 (454); C/4 (18); D/1 (402); D/2 (596); D/3 (5); E/1 (137); F/1 (1435); F/2 (450); F/A (30); G/1 (80); G/2 (194); G/3 (3); G/A (26); G/B (12); H/1 (142); H/2 (33); H/D (2); I/1 (21); J/1 (32); J/1' (5); J/2 (242); J/3 (62); J/A (27); K/1 (44); K/2 (20); K/3 (11); L/1 (31); L/2 (12); L/A (84); L/B (3); L/C (6); L/D (6); M/1 (7); N/1 (141); N/2 (33); N/B (6); X/A (955); X/B (1); X/D (22).

Exogone (Exogone) rostrata Naville, 1933

Material examined: A/1 (1); J/1 (1); J/3 (1); X/A (78); G/B (31).

Exogone (Exogone) verugera Claparède, 1868

Material examined: A/1 (2); A/B (3); B/2 (1); B/A (1); C/1 (1); C/2 (2); C/3 (1); D/1 (2); D/3 (1); E/1 (30); F/1 (4); F/2 (1); F/A (1); G/1 (1); G/B (1); H/1 (1); I/1 (9); J/3 (3); J/A (2); L/A (6); L/B (1); M/1 (1); N/1 (2); N/B (1); O/1 (6); X/A (15); X/D (1).

Genus *Grubeosyllis* Verrill, 1900

Grubeosyllis euritmica (Sardà, 1984)

Material examined: G/3 (24); N/2 (1).

Grubeosyllis limbata (Claparède, 1868)

Material examined: A/1 (31); A/2 (66); A/B (1); B/2 (19); B/A (1); C/1 (36); C/2 (12); C/3 (11); C/4 (27); D/1 (3); E/1 (4); F/1 (79); F/2 (51); F/A (17); G/1 (48); G/2 (7); G/A (21); H/1 (41); H/2 (30); I/1 (51); J/1 (7); J/1' (1); J/2 (2); J/3 (5); J/A (1); K/1 (3); K/2 (7); K/3 (115); L/1 (2); L/2 (3); L/A (23); L/B (2); M/1 (1);

N/1 (86); N/2 (5); N/B (3); O/1 (29); X/A (148); X/D (1).

Grubeosyllis clavata (Claparède, 1863) (fig. 2)

Grubea clavata. FAUVEL (1923): 296, fig. 114 a-e.

Brania clavata. CAMPOY (1982): 253.

Pseudobrania clavata. SAN MARTÍN (1984): 167, lám. 33-34; ALÓS (1988): 275, figs. 49-54.

Grubeosyllis clavata. SAN MARTÍN (1991): 718, fig. 2 a-b. Material examined: A/1 (34); A/2 (118); B/1 (10); B/2 (2); C/2 (34); D/1 (93); D/2 (151); C/1 (13); C/3 (102); C/4 (21); D/3 (6); E/1 (92); F/1 (108); F/2 (22); F/A (7); G/1 (6); G/2 (49); G/3 (6); G/A (5); G/B (3); H/1 (29); H/2 (19); I/1 (57); J/2 (171); J/3 (57); K/1 (30); K/2 (38); K/3 (123); L/1 (30); L/2 (35); L/A (18); M/1 (3); N/1 (25); N/2 (5); N/B (2); O/1 (27); X/A (96); X/D (8).

Remarks: besides the typical form (figs. 2A-D), with broad pharynx and pharyngeal tooth, subdistal tooth of composite seta blades smaller than the distal one, and scarcely developed blade spinulation, three other forms were identified:

- form A (figs. 2E, 2F), previously recorded by Alós (1988) on Catalonian coasts. It is characterized by enlarged subdistal teeth in midbody and posterior parapodia. This feature can be confused with *G. euritmica* (Sardà 1984) (SARDÀ, 1984b) but blade and pharynx shapes, the same as in typical *G. clavata*, is sufficient to distinguish the two species.

- form B (fig. 2G, 2H), also recorded by Alós (1988). Its blades are proportionally longer and thinner than those with the typical form.

- form C (fig. 2I, 2J), characterized by a well-developed spinulation on blades, similar to that of *G. vieitezi*. However, it has a broad pharynx and pharyngeal tooth and its blade serration is clearly straight.

World distribution: cosmopolitan.

Grubeosyllis vieitezi (San Martín, 1984) (fig. 3)

Pseudobrania vieitezi San Martín, 1984. SAN MARTÍN (1984): 160-164, lám. 31.

Grubeosyllis vieitezi. SAN MARTÍN (1991): 718, fig. 2 e-f. Material examined: A/1 (24); A/2 (106); C/1 (1); C/2 (6); C/3 (50); C/4 (111); D/2 (17); D/3 (72); E/1 (105); F/1 (26); F/2 (1); G/1 (9); G/2 (48); G/3 (65); G/A (2); G/B (1); H/1 (8); H/2 (175); I/1 (49); J/1 (3); J/2 (67); J/3 (64); K/1 (1); K/2 (93); K/3 (424); L/1 (21); L/2 (193); L/A (7); M/1 (48); N/1 (18); N/2 (31); O/1 (79); X/A (6); X/C (11); X/D (2).

Remarks: the typical form of this species

has a long narrow pharynx (fig. 3C) with a thin tooth. Its aciculae (fig. 3D) are provided with long mucros, and its composite seta blades (figs. 3A, 3B) are more curved than those of *Grubeosyllis clavata*. On the dorsal setae, there are very characteristic long upward curved spines. There is a different form, form A, characterized by straight spinulation (figs. 3E, 3F) similar to *G. clavata* form C. The two can be distinguished by the narrower pharynx (fig. 3G), thinner pharyngeal tooth, the aciculum shape (fig. 3H), and the more curved blades of *G. vieitezi*.

World distribution: Western Mediterranean. Atlantic (Iberian peninsula, Canary Islands, Cuba).

Genus *Parapionosyllis* Fauvel, 1923

Parapionosyllis brevicirra Day, 1954

Material examined: A/B (3); L/B (2); M/1 (1); X/A (19); X/D (1).

Genus *Sphaerosyllis* Claparède, 1863

Sphaerosyllis austriaca Banse, 1959

Material examined: A/1 (4); A/2 (43); A/B (4); B/1 (21); B/A (1); C/1 (6); C/2 (4); C/3 (20); C/4 (39); D/1 (6); D/2 (15); E/1 (45); F/1 (4); F/2 (5); G/1 (3); G/2 (24); G/3 (1); G/A (2); G/B (2); H/1 (11); H/2 (9); I/1 (252); J/3 (28); K/2 (39); K/3 (381); L/1 (1); L/2 (5); L/A (1); L/B (1); N/1 (1); N/2 (21); O/1 (39); X/A (30).

Sphaerosyllis belizensis Russell, 1989

Material examined: A/B (4); J/A (1).

Sphaerosyllis campoyi San Martín, Acero, Contonente & Gómez, 1982

Material examined: A/2 (1); F/1 (8); J/1 (1); L/B (1).

Sphaerosyllis cryptica Ben-Eliahu, 1977

Material examined: A/B (2); C/2 (1); D/1 (1); F/1 (39); F/2 (14); F/A (5); G/1 (2); H/1 (1); J/1 (11); J/A (1); X/A (32).

Sphaerosyllis hystrix Claparède, 1863

Material examined: A/1 (66); A/2 (199); A/A (2); A/B (49); B/1 (25); B/2 (3); B/A (8); C/1 (49); C/2 (32); C/3 (107); C/4 (77); D/1 (16); D/2 (49); E/1 (34); F/1 (127); F/2 (76); F/A (5); G/1 (34); G/2 (64);




Fig. 2. *Grubeosyllis clavata* (Claparède, 1868): A. Pharynx and proventriculum; B. Aciculum; C. Composite setae of anterior parapodium (typical form); D. Composite setae of midbody parapodium (typical form); E. Composite setae of anterior parapodium (form A); F. Composite setae of midbody parapodium (form A); G. Composite setae of anterior parapodium (form B); H. Composite setae of midbody parapodium (form B); I. Composite setae of anterior parapodium (form C); J. Composite setae of midbody parapodium (form C). (Scales A: 0.2 mm; B-J: 20 µm.)

G/3 (37); G/A (20); G/B (22); H/1 (87); H/2 (9); H/D (4); I/1 (113); J/1 (10); J/1' (1); J/2 (43); J/3 (29); J/A (38); K/1 (2); K/2 (45); K/3 (1); L/1 (12); L/2 (38); L/A (20); L/B (1); L/C (1); L/D (1); M/1 (5); N/1 (15); N/2 (54); N/B (8); O/1 (76); O/B (1); X/A (418).

Sphaerosyllis pirifera Claparède, 1868

Material examined: A/1 (88); A/2 (148); A/A (1); A/B (26); B/1 (93); B/2 (9); B/A (16); C/1 (22); C/2 (34); C/3 (96); C/4 (21); D/1 (72); D/2 (49); D/3 (3); E/1 (94); F/1 (229); F/2 (100); F/A (25); G/1 (77); G/2 (106); G/3 (28); G/A (27); G/B (109); H/1 (102); H/2 (11); H/D (4); I/1 (56); J/1 (100); J/1' (2); J/2 (5); J/3 (17); J/A (21); K/1 (9); K/2 (17); K/3 (25); L/1 (21); L/2 (27); L/A (68); L/B (3); L/C (14); L/D (2); M/1 (3); N/1 (34); N/2 (52); N/A (1); N/B (23); O/1 (145); X/A (480); X/B (1); X/C (2); X/D (6).

Sphaerosyllis taylori Perkins, 1981

Material examined: A/1 (2); A/2 (7); A/B (7); B/1 (3); B/2 (4); C/1 (5); C/2 (1); C/3 (3); D/1 (14); D/2 (7); E/1 (6); F/1 (56); F/2 (8); F/A (10); G/1 (9); G/2 (2); H/1 (4); I/1 (4); J/1 (9); J/3 (1); J/A (1); K/1 (1); L/2 (1); L/A (3); L/B (3); L/C (1); N/1 (2); O/1 (3); X/A (22).

Sphaerosyllis xarifae Hartmann-Schröder, 1960

Material examined: F/1 (22); F/2 (14); F/A (10); E/1 (1); H/1 (4); J/1 (10); J/1' (1); J/A (1); L/D (2); O/1 (1); X/A (36).

Subfamily Autolytinae Rioja, 1925

Genus *Autolytus* Grube, 1850

Autolytus benazzi Cognetti, 1953

Material examined: J/3 (1); X/C (3).

Autolytus brachycephalus (Marenzeller, 1874)

Material examined: A/2 (3); B/1 (5); C/1 (1); C/3 (7); D/2 (1); E/1 (14); F/1 (6); G/1 (1); G/2 (2); H/2 (1); H/D (3); I/1 (22); J/3 (2); L/1 (36); L/A (2); L/C (1); L/D (1); O/1 (1); X/A (6).

Autolytus convolutus Cognetti, 1953

Material examined: A/2 (4); C/3 (7); C/4 (7); D/2 (4); D/3 (1); E/1 (19); F/2 (12); G/1 (2); G/2 (21); H/2 (7); I/1 (3); J/1 (2); J/2 (4); J/3 (2); K/1 (1); K/2 (32); K/3 (1); L/1 (1); L/C (1); N/2 (5); O/1 (12); O/B (1); X/A (1).

Autolytus edwardsi Saint-Joseph, 1887

Material examined: A/A (2); C/3 (2); C/4 (4); D/3 (6); E/1 (2); F/1 (1); F/2 (2); F/A (2); G/3 (7); G/B (7); H/2 (4); I/1 (2); J/1' (1); J/2 (2); J/3 (4); J/A (3); K/1 (2); K/2 (5); K/3 (4); L/1 (1); L/2 (6); L/A (1); M/1 (3); N/2 (2); O/1 (1); X/B (1).

Autolytus longoprimicirratus López, San Martín & Jiménez, 1997

Material examined: G/B (3).

Autolytus prolifer (Müller, 1788)

Material examined: A/B (5); G/A (3).

Autolytus quindecimdentatus Langerhans, 1879

Material examined: A/1 (6); A/2 (9); B/1 (5); B/2 (3); B/A (7); C/1 (1); C/2 (2); C/3 (17); C/4 (7); D/1 (3); D/3 (4); E/1 (29); F/1 (8); F/2 (2); F/A (1); G/1 (6); G/2 (27); G/3 (4); G/A (2); H/1 (10); H/2 (13); H/D (6); I/1 (5); J/1 (3); J/2 (6); J/3 (8); J/A (1); K/2 (27); K/3 (3); L/1 (8); L/2 (17); L/A (1); L/C (1); M/1 (2); N/1 (15); N/2 (1); N/B (1); O/1 (3); X/A (35); X/D (4).

Autolytus sardai Martín & Alós, 1987

Material examined: A/1 (4); A/2 (2); F/1 (2); F/A (1); G/1 (1); G/2 (2); G/B (7); N/1 (3); X/A (1).

Grubeosyllis clavata (Claparède, 1868): A. Faringe y proventrículo; B. Acícula; C. Sedas compuestas del parápodio anterior (forma típica); D. Sedas compuestas del parápodio intermedio (forma típica); E. Sedas compuestas del parápodio anterior (forma A); F. Sedas compuestas del parápodio intermedio (forma A); G. Sedas compuestas del parápodio anterior (forma B); H. Sedas compuestas del parápodio intermedio (forma B); I. Sedas compuestas del parápodio anterior (forma C); J. Sedas compuestas del parápodio intermedio (forma C). (Escalas A: 0,2 mm; B-J: 20 µm.)




Fig. 3. *Grubeosyllis vieitezi* (San Martín, 1984): A. Composite setae of anterior parapodium (typical form); B. Composite setae of midbody parapodium (typical form); C. Pharynx and proventriculum (typical form); D. Aciculum (typical form); E. Composite setae of anterior parapodium (form A); F. Composite setae of midbody parapodium (form A); G. Pharynx and proventriculum (form A); H. Aciculum (form A). (Scales A-B, D-F, H: 20 µm; C, G: 0.2 mm.)

Genus *Myrianida* Milne-Edwards, 1841*Myrianida pinnigera* (Montagu, 1808)

Material examined: C/4 (1); G/3 (1); I/1 (1).

Genus *Proceraea* Ehlers, 1864*Proceraea aurantiaca* Claparède, 1868

Material examined: A/2 (2); B/1 (2); B/A (1); C/2 (3); F/1 (6); F/A (1); J/2 (1); K/2 (1); L/B (1); L/C (1); N/1 (7); X/A (2).

Proceraea picta Ehlers, 1864

Material examined: A/2 (2); F/1 (12); F/2 (2); F/A (1); G/3 (2); G/A (3); G/B (2); J/1 (1); N/1 (2); N/B (2); X/A (6).

Genus *Umbellisyllis* Sars, 1869*Umbellisyllis clavata* (Langerhans, 1879)

Material examined: A/1 (4); A/2 (4); C/1 (5); C/2 (2); C/3 (3); F/1 (10); F/2 (3); G/A (1); G/1 (2); K/1 (1).

Discussion

Most polychaete species recorded in this work were collected in substrata on which their presence is well known from previous papers: photophilic and sciophilous algae (SARDÀ, 1991; ABBIATI et al., 1992), calcareous algae (MARTÍN, 1987a, 1987b; SAN MARTÍN & AGUIRRE, 1991), *Posidonia oceanica* meadows (COLOGNOLA, et al. 1984; SAN MARTÍN & VIÉTEZ, 1984; GIANGRANDE, 1985; SAN MARTÍN et al., 1990). CAMPOY (1982) and ALÓS (1988) gave complete compilations of substrata where many of these species were recorded. On the contrary, some others species were found for the first time on some substrata that have

been less studied in the past, among them, animal substrata. Thus, some polychaete species are cited for the first time as epibiontic on some gorgonians such as *Paramuricea clavata* (*Ehlersia ferrugina*, *Odontosyllis ctenostoma*, *Pionosyllis lamelligera*, *Brania pusilla*, *Exogone naidina*, *E. verugera*, *Grubeosyllis limbata*, *G. clavata*, *Sphaerosyllis hystrix*, *S. pirifera*, *Autolytus quindecimdentatus*, *Proceraea picta*), *Eunicella verrucosa* (*Pionosyllis lamelligera*, *Sphaerosyllis hystrix*, *S. pirifera*, *Autolytus edwardsi*), or *Ellisella parplexauroides* (*Pionosyllis lamelligera*, *P. pulligera*, *Exogone naidina*, *E. verugera*, *Grubeosyllis limbata*, *G. clavata*, *G. vieitezii*, *Parapionosyllis brevicirra*, *Sphaerosyllis pirifera*, *Autolytus quindecimdentatus*). Another poorly known animal substratum is the hidroids facies, where *Eusyllis lamelligera*, *Odontosyllis gibba*, *Syllides fulvus*, *Exogone verugera*, *Sphaerosyllis austriaca*, *S. campoyi*, *S. taylori*, *S. xarifae*, *Proceraea aurantiaca*, *P. picta*, and *Umbellisyllis clavata* are recorded for the first time.

For biogeographical study, the division in groups stated for syllids by SAN MARTÍN (1984) was followed. In most cases, the species belonging to Eusyllinae, Exogoninae, and Autolytinae found at Chafarinas Islands are those of Atlantic-Mediterranean distribution (52.1% of the species number), of which, only *Exogone rostrata* can be considered as endemic in the Mediterranean. The second group in importance are cosmopolitan or widely distributed species (29.1% of the species number). In despite of this, as they are a very abundant species, in number of specimens, this is the most important group (60.7% of specimens versus 38.4% of Atlantic-Mediterranean specimens). Only a few species (*Amblyosyllis dorsigera*, *A. madeirensis*, *Brania arminii*, *Sphaerosyllis belizensis*, and *S. xarifae*) are typical in warm or tropical seas, showing a

Grubeosyllis vieitezii (San Martín, 1984): A. Sedas compuestas del parápodio anterior (forma típica); B. Sedas compuestas del parápodio intermedio (forma típica); C. Faringe y proventrículo (forma típica); D. Acícula (forma típica); E. Sedas compuestas del parápodio anterior (forma A); F. Sedas compuestas del parápodio intermedio (forma A); G. Faringe y proventrículo (forma A); H. Acícula (forma A). (Escalas A-B, D-F, H: 20 µm; C, G: 0,2 mm.)

slight influence of faunae from this kind of waters in polychaete fauna from Chafarinas. Some species (*Exogone gambiae*, *Autolytus sardai*) are recorded for the first time since their original description; their known distribution is therefore enlarged although they remain as Mediterranean endemisms.

Acknowledgements

We wish to express our gratitude to all other members of the research staff of the project PB89-0204-C02, to ICONA, and to the Ministerio de Defensa for their kind help.

Resumen

Eusyllinae, Exogoninae y Autolytinae (Syllidae, Annelida, Polychaeta) de las Islas Chafarinas (Mar de Alborán, Mediterráneo occidental)

Recientemente fue realizado un estudio sobre las comunidades de invertebrados bentónicos de las Islas Chafarinas, prestándose especial atención a determinados grupos, entre los que se encontraban los poliquetos. En este trabajo se ofrecen los resultados referidos a las subfamilias Eusyllinae, Exogoninae y Autolytinae (Syllidae, Annelida, Polychaeta), citándose 48 especies de las cuales una, *Amblyosyllis inmatura* Langerhans, 1879 (fig. 1), se cita por primera vez para la fauna del Mar Mediterráneo, siendo comparado con las especies del género previamente conocidas en dicho mar. Se describen algunas variaciones sobre la forma típica de dos especies del género *Grubeosyllis* Verrill, 1900: *G. clavata* (Claparède, 1863) (fig. 2) y *G. vieitezi* (San Martín, 1984) (fig. 3). Se encontraron ejemplares de *Pionosyllis serratisetosa* (Eusyllinae) y de *Autolytus longoprimicirratus* (Autolytinae); sus descripciones se han realizado recientemente (LÓPEZ et al., 1997).

References

- ABBIATI, M., BIANCHI, C. N., CASTELLI, A., GIANGRANDE, A. & LARDICCI, C., 1992. Distribution of Polychaetes on Hard Substrates of the Midlittoral-Infralittoral Transition Zone, Western Mediterranean. *Ophelia*, Suppl. 5: 421-423.
- ALÓS, C., 1988. Anélidos Poliquetos del Cabo de Creus (Alt Empordá). Tesis doctoral, Universidad Central de Barcelona.
- AMOUREUX, L., 1976. Annélides Polychètes recoltées par J. Stirn, sur les côtes marocaiens du détroit de Gibraltar. *Cuad. C. Biol.*, 52: 5-33.
- CAMPOY, A., 1982. Fauna de España. Fauna de Anélidos Poliquetos de la Península Ibérica. EUNSA (Ediciones de la Universidad de Navarra S. A.), Serie Biológica, Pamplona.
- CANTONE, G., FASSARI, G. & BRIGANDI, S., 1978. Ricerche sui policheti della Tunisia. *Animalia*, 5(1/3): 51-78.
- COLOGNOLA, R., GAMBI, M. C. & CHESSA, L. A., 1984. Polychaetes of the *Posidonia oceanica* (L.) Delile foliar stratum: comparative observations. *International Workshop on Posidonia oceanica Beds*, 1: 12-15.
- DAY, J. H., 1967. A monograph on the Polychaeta of Southern Africa. Trustees of the British Museum (Natural History) XXIX, Londres.
- FAUVEL, P., 1923. Faune de France. 5: Polychètes Errantes. Chevalier Ed., Paris.
- GIANGRANDE, A., 1985. Policheti dei rizomi di *Posidonia oceanica* (L.) Delile (Helobinae, Potamogetonaceae) di una prateria dell'isola di Ischia (Napoli). *Atti Soc. Tosc. Sci. Nat. Mem. Serie B*, 92: 195-206.
- LANERA, P., SORDINO, P. & SAN MARTÍN, G., 1994. *Exogone (Parexogone) gambiae*, a new species of Exogoninae (Polychaeta, Syllidae) from the Mediterranean Sea. *Boll. Zool.*, 61: 235-240.
- LANGERHANS, P., 1879. Die Würmfauna von Madeira. *Zeitschr. wiss. Zool.*, 32: 513-592.
- LÓPEZ, E., SAN MARTÍN, G. & JIMÉNEZ, M., 1996. Syllinae (Syllidae, Annelida, Polychaeta) from Chafarinas Islands (Alborán Sea; Western Mediterranean). *Misc. Zool.*, 19.1: 105-118.
- 1997. Two new species of syllids (Syllidae, Polychaeta) from Chafarinas Islands (Alborán Sea, Southwestern Mediterranean). *Bull. Mar. Sci.*, 60(2): 293-299.
- MARTÍN, D., 1987a. La comunidad de Anélidos Poliquetos de las concreciones de algas calcáreas del litoral catalán. Caracterización de las especies. *Publ. Dpt. Zool.*, 13: 45-54.

- 1987b. Anélidos Poliquetos asociados a concreciones de algas calcáreas del litoral catalán. *Misc. Zool.*, 11: 61-75.
- MARTÍN, D. & ALÓS, C., 1987. *Autolytus sardai* n. sp. Nueva especie de Autolytinae (Syllidae, Polychaeta) en el Mediterráneo español. *Cah. Biol. Mar.*, 28: 579-584.
- MARTÍN, D., ALÓS, C. & SARDÀ, R., 1990. *Miscellania dentata* gen. et sp. n. (Polychaeta: Syllidae) from the Spanish Mediterranean coast. *Zool. Scr.*, 19(2): 169-172.
- NÚÑEZ, J. & SAN MARTÍN, G., 1996. Anélidos poliquetos de las islas Canarias: Familia Syllidae, I. Subfamilias Eusyllinae y Autolytinae. *Actas del Primer Congreso de Oceanografía y Recursos Marinos en el Atlántico Centro-Oriental*: 199-222.
- NÚÑEZ, J., SAN MARTÍN, G. & BRITO, M. C., 1992. Exogoninae (Polychaeta: Syllidae) from the Canary Islands. *Sci. Mar.*, 56(1): 43-52.
- OLANO, M. L., LÓPEZ, E. & SAN MARTÍN, G., in press. Dos nuevas especies de Sílidos (Polychaeta, Syllidae, Exogoninae) para el litoral ibérico: *Sphaerosyllis belizensis* Russell, 1989 y *Exogone (Parexogone) cognetti* Castelli, Badalamenti & Lardicci, 1987. *Bol. R. Soc. Esp. Hist. Nat.*, 94(1-2).
- PARAPAR, J. & SAN MARTÍN, J., 1992. *Grubeosyllis celiae*, a new species of Exogoninae (Syllidae, Polychaeta) from off Ceuta, Spain. *Proc. Biol. Soc. Wash.*, 105(1): 112-115.
- PARAPAR, J., SAN MARTÍN, G., URGORRI, V. & BESTEIRO, C., 1993. Anélidos poliquetos mesop-sámmicos sublitorales de la costa de Ceuta (España). *Cah. Biol. Mar.*, 34: 363-381.
- RUSSELL, D., 1989. Three new species of *Sphaerosyllis* (Polychaeta, Syllidae) from mangrove habitats in Belize. *Zool. Scr.*, 18(3): 375-380.
- SAN MARTÍN, G., 1984. Estudio biogeográfico, faunístico y sistemático de los Poliquetos de la familia Sílidos (Syllidae: Polychaeta) en Baleares. Publicaciones de la Universidad Complutense de Madrid 187/84, Madrid.
- 1990. Eusyllinae (Syllidae, Polychaeta) from Cuba and Gulf of Mexico. *Bull. Mar. Sci.*, 46(3): 590-619.
- 1991. *Grubeosyllis* and *Exogone* (Exogoninae, Syllidae, Polychaeta) from Cuba, the Gulf of Mexico, Florida, and Puerto Rico, with a revision of *Exogone*. *Bull. Mar. Sci.*, 49(3): 715-740.
- SAN MARTÍN, G. & AGUIRRE, O., 1991. Ciclo anual de los poliquetos asociados al alga *Mesophyllum lichenoides* (Ellis) en una playa mediterránea. *Bol. Inst. Esp. Oceanogr.*, 7(1): 157-170.
- SAN MARTÍN, G., ESTAPÉ, S., GARCÍA-OCEJO, A., GÓMEZ, C. & JIMÉNEZ, P., 1990. Estudio de la taxocenosis de Anélidos Poliquetos de rizomas de *Posidonia oceanica* en las costas de Almería. *Bol. Inst. Esp. Oceanogr.*, 6(1): 41-58.
- SAN MARTÍN, G. & PARAPAR, J., 1990. *Exogone acerata* (Exogoninae: Syllidae: Polychaeta), a new species without antennae from the Mediterranean Sea. *Proc. Biol. Soc. Wash.*, 103(3): 687-690.
- SAN MARTÍN, G. & VIÉITEZ, J. M., 1984. Anélidos Poliquetos de los rizomas de *Posidonia oceanica* en las costas de Cabo Palos (Murcia, España). *International Workshop on Posidonia oceanica Beds*, 1: 149-157.
- SARDÀ, R., 1984a. Estudio sobre la fauna de Anélidos Poliquetos de las zonas medio-litoral e infralitoral en la región del Estrecho de Gibraltar. Tesis doctoral, Universidad Central de Barcelona.
- 1984b. La subfamilia Exogoninae (Polychaeta: Syllinae) de Gibraltar con la descripción de *Pseudobranchia euritmica*. *Publ. Dpt. Biol.*, 10: 7-13.
- 1991. Polychaete Communities Related to Plant Covering in the Midlittoral and Infralittoral Zones of the Balearic Islands (Western Mediterranean). *Mar. Ecol.*, 12 (4): 341-360.
- UEBELACKER, J. M., 1984. Familiy Syllidae. In: *Taxonomic Guide to the Polychaetes of the Northern Gulf of Mexico*, Vol. IV, Chapter 30: 1-151 (J. M. Uebelacker & P. G. Johnson, Eds.). Barry A. Vitor & Associated, Mobile.
- WESTHEIDE, W., 1972. La faune des Polychètes et des Archiannélides dans les plages sableuses à ressac de la côte méditerranée de la Tunisie. *Bull. Inst. Océanogr. Pêche, Salammbô*, 2(3): 449-467.
- 1974. Interstitielle Fauna von Galapagos. XI. Pisionidae, Hesionidae, Pilargidae, Syllidae (Polychaeta). *Mikrofauna Meeresbodens*, 44: 1-146.
- ZGHAL, F. & BEN AMOR, Z., 1980. Annélides Polychètes de Tunisie. *Bull. Off. natn. Pêch. Tunisie*, 4(1): 1-11.