



Scientific basis for the industrialization of traditionally used plants of the *Rosaceae* family

P. Garcia-Oliveira^a, M. Fraga-Corral^{a,b}, A.G. Pereira^{a,b}, C. Lourenço-Lopes^a, C. Jimenez-Lopez^{a,b}, M.A. Prieto^{a,*}, J. Simal-Gandara^{a,*}

^a Nutrition and Bromatology Group, Analytical and Food Chemistry Department, Faculty of Food Science and Technology, University of Vigo, Ourense Campus, E-32004 Ourense, Spain

^b Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Campus de Santa Apolonia, 5300-253 Bragança, Portugal

ARTICLE INFO

Keywords:

Traditionally used plants
Ethnobotanic
Rosaceae family
Bioactive compounds

ABSTRACT

Plants have been traditionally used for the treatment of different types of illness, due to biomolecules with recognised benefits. *Rosaceae* family is used in traditional Galician medicine. The following plants *Agrimonia eupatoria*, *Crataegus monogyna*, *Filipendula ulmaria*, *Geum urbanum*, *Potentilla erecta* and *Rosa canina* are usually found in treatments. The aim of this study is to perform an ethnobotanical review about the bioactive compounds of these plants and their different bioactivities, both studied *in vitro* and *in vivo*. The nature of the bioactive compounds is varied, highlighting the presence of different phenolic compounds, such as phenolic acids, flavonoids or tannins. Understanding the beneficial effects of the administration of the whole plant or target tissues from *A. eupatoria*, *C. monogyna*, *F. ulmaria*, *G. urbanum*, *P. erecta* and *R. canina* as well as those from their individual compounds could lead to the development of new drugs based on the use of natural ingredients.

1. Oral tradition and the knowledge based on trial and error

1.1. History of traditionally used Plants: Worldwide to Galicia (NW Spain)

Since the ancient times, people instinctively sought the remedies for pain and diverse illness in the surrounding nature (Sengupta, Gaurav, & Tiwari, 2018; Srivastava, 2018). All knowledge was based on experience, as there was no information about the reasons for the disease or which plant and its form of consumption could be useful for its treatment. After thousands of years of trial and error throughout human history, specific plants consumed via oral administration were identified for the treatment of specific diseases, therefore known as medicinal plants (Srivastava, 2018). Plants may contain diverse chemical compounds, named active or bioactive compounds, such as alkaloids or essential oils (EO) (Pérez, 1998). The correct administration of these biomolecules may have beneficial effects on health. In fact, some of these bioactive compounds are the foundation of modern medicine drugs and they represent an alternative source of innovative ingredients with further applications in food and cosmetic industries. Along the history of the traditionally used plants, at first, information was passed orally from generation to generation, until the start of writing. Numerous texts describe the use of different plants in ancient civilizations,

such as the Sumerian, Egyptian, Indian, Chinese, Greek, Roman or Arab (Sengupta et al., 2018; Srivastava, 2018). The *Ebers Papyrus* contains descriptions of different treatments, based on plants, to treat skin, urinary, ophthalmologic and other infections (Lennihan, 2014; Srivastava, 2018). Some of the treatments from the Ayurveda Indian medical system and Traditional Chinese medicine have similar basis and the species they described are still used today (Lennihan, 2014; Srivastava, 2018). Initially, plants were consumed raw, and later, people added them to hot water (W) to create soup and tea. As knowledge grew, different herbs were combined to elaborate more complex treatises (Lennihan, 2014). Ancient Greeks are considered the parents of modern medicine, as they combined traditional practices with scientific thinking. Hippocrates, Galen and Dioscorides, among other authors, showed their interest in the deeper application of plants by writing vast treatises (Srivastava, 2018). In *Corpus Hippocraticum*, different herbal medicines and their forms of treatment were described for each disease (Lennihan, 2014; Pahlow, 1994). Galen differentiated between the use of plants “at natural” (meaning in powder form) and the galenic preparations: bioactive compounds were concentrated in W, ethanol (Et) or vinegar, then used in ointment, plasters or other galenic forms (Pahlow, 1994). Dioscorides wrote *De Materia Medica*, with descriptions, modes of collection, preparation and therapeutic effects of

* Corresponding authors.

E-mail addresses: mprieto@uvigo.es (M.A. Prieto), jsimal@uvigo.es (J. Simal-Gandara).

<https://doi.org/10.1016/j.foodchem.2020.127197>

Received 9 February 2020; Received in revised form 16 March 2020; Accepted 29 May 2020

Available online 08 June 2020

0308-8146/ © 2020 Elsevier Ltd. All rights reserved.

Abbreviations			
<i>Generic</i>		Ha	Hydroalcoholic
		M	Methanol
		W	Water
<i>Compounds</i>		<i>Antioxidant Essays</i>	
AP	Aerial parts	ABTS	2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid
EMA	European Medicines Agency	DPPH	2,2-diphenyl-1-picrylhydrazyl
FL	Flowers	FRAP	Ferric Reducing Antioxidant Power assay
FR	Fruits	ORAC	Oxygen radical absorbance capacity assay
L	Leaves	TEAC	Trolox Equivalent Antioxidant Capacity
NA	Not analysed	<i>Cell Lines</i>	
P	Petals	Caco-2	Heterogeneous human epithelial colorectal adenocarcinoma
R	Roots	HaCaT	Human aneuploid immortal keratinocyte
RH	Rose hips	HeLa:	Cervical carcinoma
		HepG2:	Hepatocellular carcinoma
		HL-60	Human promyelocytic leukemia
		HT-22	Mouse hippocampal neuronal cell line
		HT-29	Human colon cancer
		MCF-7:	Breast adenocarcinoma
		MRC-5	Human foetal lung
		NCI-H460:	Non-small cell lung cancer
		RB	Human rhabdomyosarcoma
		SH-SY5Y	Human neuroblastoma
		SMMC-7221	Human hepatoma
		VSMCs	Vascular smooth muscle
		WiDr	Human colon adenocarcinoma
<i>Solvents & Extractions</i>		<i>Anti-inflammatory Compounds</i>	
Ac	Acetone	12-HETE	12-Hydroxyeicosatetraenoic acid
AE	Alkaline extraction	15-LOX	Human 15 Lipoxygenase
CE	Commercial extract	AChE	Acetylcholinesterase
Ch	Chloroform	COX	Cyclooxygenase
CO ₂ s	CO ₂ supercritical	IL-1β	Interleukin 1 beta
DMSO	Dimethylsulfoxide	IL-6	Interleukin 6
EA	Ethyl acetate	iNOS	Inducible nitric oxide synthase
Er	Ether	NF-κB	Nuclear Factor kappa-light-chain-enhancer of activated B cells
Et	Ethanol	PGE2	Prostaglandin E2
H	Hexane	PPARs	Peroxisome proliferator-activated receptors
		TLRs	Toll-like receptors
		TNF-α	Tumor necrosis factor alpha

more than 600 MP (Srivastava, 2018). The Greek knowledge was extended by the Romans and Arabians. A scientific regression was suffered during the Middle Ages, as science and magic were mixed. Owing to their knowledge of Latin and Greek, clerics conserved the ancient knowledge (Pahlow, 1994). Until the 16th century, with the advent of iatochemistry (chemical medicine), plants have been the major source of treatments and relief to some of the most diverse diseases, such as digestive problems, microbial infections, cancer and even neurodegenerative illness (Sengupta et al., 2018; Srivastava, 2018). Synthetic drugs were available for people with more economic resources, while remedies based on plants remained the option to treat poor people (Lennihan, 2014; Srivastava, 2018). In the 18th century, urban citizens preferred pharmaceutical drugs rather than traditional medicine. However, drug side effects became known and medicinal plans resurfaced rapidly (Lennihan, 2014). In early 19th century, the development of chemical methods allowed the discovery and isolation of alkaloids

(such as morphine from *Papaver somniferum* or quinine from *Cinchona officinalis*), tannins (T), vitamins, hormones, saponins (Sa), among other bioactive compounds from vegetal origin starting scientific pharmacy (Srivastava, 2018).

Regarding traditional Spanish medicine, it has the heritage of hundreds of years of different people that influenced the country: diverse Iberian people, Greeks, Romans, Arabs and Orientals as well as German traditions (Verde, Rivera, Fajardo, Obón, & Cebrián, 2008). Some Roman authors described the traditional use of plants by Iberian tribes (de Santayana, Morales, Tardío, Aceituno, & Molina, 2014). During Middle Ages, Ibn al-Baytar, an Andalusian doctor and botanist) wrote the *Tratado de simples*, where he described more than 1400 species with beneficial properties (de Santayana et al., 2014). However, as the Christian religion increased its influence, the human body was considered a finite recipient for a soul, which aspired to divine salvation. In these ages, people were suspicious of medicine, and healers

were often considered heretics (Verde et al., 2008). Until the early 19th century, the Spanish Inquisition pursued and convicted many healers, men and women, of heresy (Verde et al., 2008). Little by little, science overcome religion, and two differentiated figures appeared: the physician, who carried out the diagnosis, and the pharmacist, who prepared the drugs (Mosquera Paans, 2013). After the Spanish Flu pandemic of 1918, which caused a dramatic number of deaths, traditional medicine started to grow, particularly among the population with less economic resources to obtain chemical drugs (Mosquera Paans, 2013). The botanist Pio Font Quer wrote *Plantas medicinales. El Dioscórides renovado*, which is the most complete text about application of Iberian plants and is still a basic reference (de Santayana et al., 2014). It includes botanic, chemical and pharmacologic data, names and local uses. He is considered one of the fathers of ethnobotany in Spain (de Santayana et al., 2014). The roots of traditional medicine lead to a new golden age of natural ingredients obtained from plants, whose use is gradually legislated more exhaustively and scientific research is favoured (Mosquera Paans, 2013).

Galicia is a region located in the NW of Spain. Not long ago, in case of ailment, the people had to seek the experience of the elders, especially in the rural areas (Rodríguez González, 2008). In these zones, visiting a doctor or buying a drug was a luxury. Healers (“menciñeiros” in Galician) practiced “shamanism” and the Christian religion, integrating magic and religion. These healers used their knowledge of MP combined with spells, magic symbolism and also Christian traditions (Mosquera Paans, 2013; Rodríguez González, 2008). Actually, they were important members of the society in the rural areas, since the medical advances took a long time to arrive. Regarding the important figures, the conservation of Galician traditional medicine, Father Sarmiento (18th century) stands out. He understood the necessity to understand, recover and preserve Galician tradition and popular culture (Mosquera Paans, 2013).

As in other regions, Galician traditional medicine has almost been replaced by modern medicine. However, some people have learned and continue transmitting this ancient knowledge, thereby conserving it (González-Hernández, Romero, Rodríguez-Guitián, & Rigueiro, 2004). As popular wisdom is concentrated on in the elder population and in the

rural areas, thousands of years of experience are at risk of disappearing when these people pass away. Preserving the ancient knowledge is exceedingly important, as the study of the traditional use of certain plant species could lead to the re-discovery of active compounds (initially found by our ancestors) with different applications, including medicine, nutrition, cosmetics, etc. (Franco, Guitián, & Resúa, 2013).

1.2. Current uses of natural ingredients extracted from plants

1.2.1. Food

Nowadays, the consumption of vegetal-based nutrients has been prompted by different reasons. In most of the developed countries where nutrition does not represent an issue for most of the population, consumers’ demands are focused on the improvement of the quality of the food. Some of these demands include the replacement of synthetic or artificial ingredients, additives or colorants for natural ones which are not considered to have side-effects. Other demand is to increase the presence of organic and fresh products in the market, as well as, the increment of vegetal-based products which may be included in alternative diets such as vegan or vegetarian ones. However, another important concern is about people with low economic inputs whose nutrition levels are scarce which may induce to the ingestion of unsafe products and thus cause diseases that might become chronic and even fatal. The use of vegetal-based diets has been demonstrated to be an economical approach to feed people. In this sense, since the 50’s the rise on the global population has represented a challenge due to the parallel increment of animal protein intake. Plants have been described to represent an alternative source of protein which may help to balance or even reduce the animal protein production (Asioli et al., 2017; Day, 2013; Flood, 2010). Traditionally many plants have been cultivated since ancient times, due to their nutritional and beneficial properties. Nowadays, it is very common to include some of them in our diet, as it is the case of different spices, such as thyme, rosemary, parsley, coriander; vegetables like tomatoes, garlic, pumpkins, onions, or leeks; and different fruit trees for example (López et al., 2018; Pahlow, 1994; Pinela et al., 2016; Vieira et al., 2017). Plants can be eaten raw, as in salads, cooked or in infusions and can be used in different ways: to




Fig. 1. Different traditional and industrial uses of Rosaceae plants.

garnish, like food ingredients, aromatize, or they can be consumed for their beneficial properties. Some herbal products are sold in pharmacies as dietary supplements, teas, capsules, extracts or powders used in self-medication to prevent diseases, soothe pains, relax, improve wellbeing, etc. Several studies have proven the efficacy of these products in health (Martín Ortega & Segura Campos, 2019; Srivastava, 2018).

Whole plants and some of their compounds could be included in foods or beverages, obtaining a “nutraceutical” food (also called nutraceutical), which, in addition to nutritional function, has different health benefits, helping in the prevention and treatment of diseases (Gulati, Ottaway, Jennings, Coppens, & Gulati, 2019; Lubbe & Verpoorte, 2011). In fact, this tendency to implement the properties of the products and the wide use of functional foods has commenced to blur the limits between pharma and nutrition (Corbo, Bevilacqua, Petruzzi, Casanova, & Sinigaglia, 2014). Currently, different nutraceuticals are available for consumers. One of the most well-known examples are dairy products (e.g. milk, yogurts and margarines) enriched with phytosterols, which have been proven to reduce the intestinal absorption of cholesterol (Tolve et al., 2019). Another example is folic acid, extracted from leafy green vegetables and legumes, commonly used to fortify cereal and their derivatives. This vitamin is considered to help in the prevention of central nervous system diseases (Samaniego-Vaesken, Alonso-Aperte, & Varela-Moreiras, 2016). In addition to compounds, there are also examples of products supplemented with the plant itself. For example, different plant materials are used to fortify flour, including *Chenopodium quinoa*, *Avena sativa* or *Hordeum vulgare*, due to the high quantity of bioactive compounds of these species (Cardoso, Fernandes, González-Paramás, Barros, & Ferreira, 2019).

The toxicological and carcinogenic effects associated with synthetic ingredients have triggered the replacement with natural ones and therefore food industries and academia have prompted their deeper study. Along the last decades many applications have been developed based on the use of natural biocompounds, such as the antioxidants that are present under different chemical structures. Carotenoids, anthocyanins and betalains extracted from plants have been proven to be effective natural additives for foods with antioxidants and have less toxic effects than synthetic ones (Roriz, Barros, Prieto, Morales, & Ferreira, 2017).

The previous food applications of different species of plants are summarized in Fig. 1 and in Table 1.

1.3. Pharmacological

Pharmacological applications are the most known application of these raw materials (Pérez, 1998). Nowadays, the synthetic drugs of modern medicine have almost completely replaced the use of natural plants in developed nations and they are used mostly in traditional medicine in Third world and developing countries. In developed countries, traditional medicine is categorized as “alternative medicine” and is used not for the treatment of serious diseases, but for minor disorders (Martín Ortega & Segura Campos, 2019; Srivastava, 2018).

Natural ingredients obtained from plants can be described as being applied in different ways such as being used as direct agents or as raw materials in the manufacture of drugs. Some examples of these drugs are aspirin, which derives from willow; opium-derived morphine, obtained from poppies; or the anti-tumour drug Taxol, which derives from the Pacific yew (Lennihan, 2014; Martín Ortega & Segura Campos, 2019). In fact, it is estimated that 25–28% of drugs used in modern medicine are derived from MP, directly or indirectly (Lubbe & Verpoorte, 2011; Martín Ortega & Segura Campos, 2019). Regarding pharmacological applications, diverse compounds have been proved to be related to the health benefits of biomolecules extracted from plants. Some examples are described below and summarized in Fig. 1 and in Table 1.

Plants from the Asteraceae, Rutaceae, Fabaceae, Lamiaceae, Zygophyllaceae, Rhamnaceae, Liliaceae and Zingiberaceae families are

Table 1
Current use of plants in different industries.

Industry	Application	Examples	Compounds	Mechanisms	Reference
Food	Foodstuff, nutraceuticals, animal feeding	Spices (thyme, rosemary, parsley); vegetables (garlic, pumpkin, onion), cereals (quinoa, oat, barley) fruits (apples, oranges), etc. Tea, supplements, enriched foods	PC, vitamins, phytoestersols, etc.	Diverse beneficial effects on health	(Cardoso et al., 2019; Delaveau et al., 1981; Samaniego-Vaesken et al., 2016; Tolve et al., 2019; Verde et al., 2008)
		<i>M. suawolens</i> , <i>T. zygis</i> or <i>F. vulgare</i>	PC, vitamins	Interruption of reactive oxygen reactions	(Stanković et al., 2017).
Drugs	Antioxidant	<i>Cymbopogon citratus</i> , <i>Chenopodium ambrosioides</i> , <i>Euphorbia stenoclada</i> , <i>Geranium mexicanum</i> , <i>Gnaphalium oxyphyllum</i> , <i>Helianthemum glomeratum</i> , <i>Larrea tridentate</i>	Terpens (e.g. carvacol, geraniol) and PC	Their antioxidant and antimicrobial prevent the spoilage of food by food borne pathogens	(Ortega-Ramirez et al., 2014)
	Food additives	Asteraceae, Rutaceae, Fabaceae, Lamiaceae, Zygophyllaceae, Rhamnaceae, Liliaceae and Zingiberaceae families	Pp, F, coumarins, etc.	Inhibit the ADP pathway, suppress TXA ₂ formation, reduce intracellular Ca ²⁺ mobilization	(El Haouari & Rosado, 2016)
Cosmetics	Thrombotic diseases (anti-platelet activity)	<i>Passiflora incarnata</i> , <i>Piper methysticum</i> , <i>Valeriana officinalis</i>	Sa, alkaloids, Pp, triterpenoid, EO, fatty acids or F	Increase of serotonin, dopamine or norepinephrine levels	(Martins & Brijesh, 2018)
	Anxiety and depression	<i>Cimicifuga racemosa</i> , <i>Hypericum perforatum</i> or <i>Melissa officinalis</i>	Alkaloids, PC, Te and proteins	Interrupt the life cycle of HIV as well as act as enhance the immune system	(Lailla et al., 2019)
	Against HIV	<i>Rheum palmatum</i> , <i>Rheum officinale</i> , <i>Trigonostem oxyphyloides</i> , <i>Vatica astrotricha</i> , <i>Vernonia amygdalina</i>	PC, Te	Disruption cell wall. Inhibition metabolic processes.	(Pandey & Agnihotri, 2015)
	Antimicrobials	<i>Taxus sp.</i> , <i>Ginkgo biloba</i> or <i>Artemisia annua</i>	PC, alkaloids, terpens, and Sa	Induction of apoptosis, autophagy, cell cycle arrest. Alteration different signalling pathways	(Benarba & Pandiella, 2018; Oyenihni & Smith, 2019)
Others	Anticancer	<i>Taxus brevifolia</i> , <i>Artemisia sieversiana</i> , <i>Chamaecyparis obtusa</i> , <i>Gurcuma longa</i> or <i>Zingiber officinal</i>	PC, vitamins, carotenes, Ps and fatty acids	Anti-aging, antioxidant, anti-inflammation, calming, relaxing	(Aburjai & Natsheh, 2003; Ganesan & Choi, 2016; Lubbe & Verpoorte, 2011)
	Beauty products	Aloe vera, cucumber, ginseng, tea, chamomileEOs (e.g. citron, lavender, eucalyptus)	Numerous bioactivity such as antistress, immunostimulant and antiparasitic	Cheaper and more sustainable treatment.	(Reverter et al., 2017)
Others	Aquaculture				

well known for their antiplatelet activity. Several bioactive compounds of these plants, specially polyphenols (Pp), flavonoids (F) and coumarins, has been proved to inhibit the ADP pathway, suppress TXA₂ formation and reduce intracellular Ca²⁺ mobilization, among other activities (El Haouari & Rosado, 2016).

MP are also a great source of bioactivity compounds to treat anxiety and depression. In this case, compounds such as Sa, alkaloids, Pp, F, triterpenoids, EO, and fatty acids are involved. *Passiflora incarnata*, *Piper methysticum*, *Valeriana officinalis*, *Cimicifuga racemose*, *Hypericum perforatum* or *Melissa officinalis* are examples of plants used to treat these disorders (Martins & Brijesh, 2018). After the administration of most of the plants considered to possess medicinal properties, the antidepressant activity is related to the increase of monoamines (serotonin, dopamine or norepinephrine) levels in the brain, inhibiting their receptor systems and transporters and also reducing the activity of monoamine oxidase enzymes, which break the monoamines (Martins & Brijesh, 2018).

Nowadays, several adverse effects of the current medication against HIV have been described. To reduce these side effects, scientists are searching for safer and more effective compounds. Several plants have shown promising anti-HIV activity (both *in vitro* and *in vivo*), such as *Rheum palmatum*, *R. officinale*, *Trigonostem oxyphylloides*, *Vatica astrotricha*, *Vernonia amygdalina*, *Hypoxias pelargonium*, *Sidoides hemerocallidea* and *Sutherlandia frutescens*. The mechanism of action of most of them has not been fully disclosed, but different compounds (alkaloids, phenolic compounds (PC), terpenoids (Te) and proteins) have been proven to interrupt the life cycle of HIV and enhance the immune system (Laila et al., 2019). Other species, for instance *Sambucus nigra*, *Caesalpinia pulcherrima* and *Hypericum connatum*, are also useful against other viruses like the herpes simplex virus, influenza, hepatitis, and coxsackie virus (Akram et al., 2018).

Antimicrobials may be obtained from plant matrixes, being alkaloids, PC and Te the responsible compounds of the activity. Generally, these compounds act through the disruption of the cell membrane and inhibition of different metabolic processes, such as the biosynthesis of nucleic acids or proteins. Species of genus *Taxus*, *Ginkgo biloba* or *Artemisia annua* have shown antimicrobial properties. *A. annua* also contains the anti-malaria compound artemisinin (Pandey & Agnihotri, 2015).

Several compounds are currently used in the formulation of drugs to treat cancer, including Te, alkaloids, Sa, glycosides (G), and PC. Briefly, these compounds act through the induction of apoptosis, autophagy or cell cycle arrest of cancer cells and also through the modification of signalling pathways (Benarba & Pandiella, 2018; Oyenih & Smith, 2019). To give some examples, *T. brevifolia*, *Artemisia sieversiana*, *Chamaecyparis obtusa*, *Curcuma longa* or *Zingiber officinale* are MP with anti-proliferative effects *in vitro* and *in vivo* models (Benarba & Pandiella, 2018; Martín Ortega & Segura Campos, 2019; Oyenih & Smith, 2019).

Numerous studies have demonstrated that thousands of plants contain potent antioxidant compounds, especially PC and vitamins, due to their redox properties, the ability to chelate metals and quench singlet oxygen reactive species (Borneo, León, Aguirre, Ribotta, & Cantero, 2009; Jaberian, Piri, & Nazari, 2013; Ortega-Ramirez et al., 2014; Roleira et al., 2015). Their efficiency has been assessed through *in vitro* assays and also with *in vivo* supplementation studies in animal and human models (Oyenih & Smith, 2019). *Mentha suaveolens*, *Thymus zygis* or *Foeniculum vulgare* are just a few examples of plants with antioxidant properties (Stanković, Radić, Blanco-Salas, Vázquez-Pardo, & Ruiz-Téllez, 2017). These compounds may be used to treat several diseases, such as diabetes, cancer, ageing and other health problems (Oyenih & Smith, 2019).

1.3.1. Cosmetics

Throughout the centuries, plants have been used in the treatment of different beauty problems and disorders, such as dry skin, inflammation, eczema, psoriasis, hair loss and others. Nowadays, numerous

cosmetic products contain extracts or plants-derived compounds in their formulation, such as sunscreens, moisturizers, skin cleansers, body lotions, lipsticks, etc. (Aburjai & Natsheh, 2003; Ganesan & Choi, 2016). Regarding skin products, aloe vera and cucumber are widely used, due to their anti-inflammatory, cooling and healing properties. Other examples of plants used in cosmetics are the ginseng (anti-aging properties), tea (several bioactivities, including antioxidant, photo-protective and anti-inflammatory effects), chamomile (anti-inflammatory activity) or onion (antiallergic anti-inflammatory and antimicrobial effects) (Aburjai & Natsheh, 2003). In hair care, plants such as *Ginkgo biloba*, aloe vera, sage and rosemary have been used, especially to promote hair growth (Aburjai & Natsheh, 2003). EOs are also an important cosmetic product, due to their beneficial effects for skin and hair care and relaxing properties. For instance, chamomile, lavender, eucalyptus and citron EOs are appreciated in cosmetic applications for their antioxidant and anti-inflammatory properties, among others (Lubbe & Verpoorte, 2011).

In many cases, bioactive compounds are included in the formulation of cosmetics, such as PC (e.g. quercetin, gallic acid, luteolin or catechins (Ca)), vitamins, carotenes, polysaccharides (Ps) and fatty acids. Several beneficial activities of these compounds have been described, including anti-ageing, antioxidant, anti-inflammatory and anti-microbiological (Ganesan & Choi, 2016). The previous food applications of different species of plants are summarized in Fig. 1 and in Table 1

1.3.2. Others

Currently, many consumers have a bad opinion of synthetic additives used to preserve foods. Thus, the food industry is searching new natural compounds to prevent the spoilage of food produced by pathogenic microorganisms and also the oxidation of lipids. Terpens (e.g. geraniol, citral or carvacrol) and PC extracted from plants have been demonstrated to be effective as food additives due to their antimicrobial and antioxidant properties. Some plant compounds have been tested as food additives, such as the terpenes of *Cymbopogon citratus* and *Chenopodium ambrosioides* and the PC of *Euphorbia stenoclada*, *Helianthemum glomeratum*rich and *Gnaphalium oxyphyllum*rich (Ortega-Ramirez et al., 2014).

The use of plants has been also considered a promising tool for aquaculture. Microbial diseases cause high economic losses, while the inappropriate use of antibiotics could lead to negative side effects appearing in the organisms and also in the environment. Several plants have been demonstrated to exert beneficial effects on the aquatic organism and also reduction of the pathogens present in the W (Van Hai, 2015). Therefore, plants are a cheaper and more sustainable treatment, since they have numerous interesting bioactivity such as promoting anti-stress and being immune-stimulants and anti-parasitic (bacterial, fungus, virus and ectoparasites) (Reverter, Tapissier-Bontemps, Sasal, & Saulnier, 2017).

2. Properties of traditionally used plants from *Rosaceae* family

As mentioned before, the information related to traditional uses of plants is in serious danger. To record this ancient knowledge, several ethnobotanical and ethnopharmaceutical studies have been performed in different Iberian Peninsula regions, including Galicia (Franco et al., 2013; González-Hernández et al., 2004). Most of these studies are focused on the different uses of plants and their forms of consumption/application but no extensive information about the active compounds responsible of the medicinal action has been collected. Investigating the phytochemistry is necessary to understand the relationship between their traditional uses (the result of hundreds of years of trial and error) and their bioactive compounds. In addition, this study allows the exploration of possible future uses of these plants.

In ethnobotanical studies, *Rosaceae* family is found among the most frequently used plant families (Franco et al., 2013). In the present study, *Agrimonia eupatoria*, *Crataegus monogyna*, *Filipendula ulmaria*,

Geum urbanum, *Potentilla erecta* and *Rosa canina* were chosen to perform an ethnobotanical review, because they are among the most utilised plants of this genus, and scientific studies have been carried out on their active compounds and bioactivities. All the six species mentioned can be found in the northwestern part of the Iberian Peninsula. However, the distribution of *F. ulmaria*, *G. urbanum*, and *P. erecta* have a lower density in the northwest, even though they constitute a representative flora from the northern part of the Peninsula (Castroviejo, 1986; Ministerio De Agricultura Alimentación y Medio Ambiente, 2019). Additionally, all the species can be found in all four Galician provinces: Lugo, Ourense, A Coruña and Pontevedra (Romero, 2008). The traditional uses of the selected plants (parts used, mode of application and infections treated) are described in the following paragraphs and summarized in Table 2.

2.1. *Agrimonia eupatoria*

2.1.1. Traditional importance

A. eupatoria is an indigenous plant from the middle and northern areas of Europe and also the temperate regions from Asia and North America (Al-snafi, 2015). Its aerial parts (AP) are usually collected during the flowering period, spring and summer, to be applied in different ways. Traditional uses reported along the Iberian Peninsula comprised diverse application modes, such as, infusion, decoction, dry extraction, wet extraction (using aqueous, oil or alcoholic solvents) and tincture (de Santayana et al., 2014; Escudero, 1999; Febrer, Blanquer, & Pí, 2001; Romero Martín, 1998; Verde, 2008). *A. eupatoria* has several medicinal properties: it is an astringent, hypotensive, anti-inflammatory, antiemetic, analgesic and healing. It has been used as mouthwash, to gargle and for larynges infections, to clean varicose ulcers in renal or hepatic infections and to control diarrhea and menstruation pain. Additional uses includes its application to detoxify blood, eliminate pimples and rashes, control headaches, reduce muscular pain, or as anthelmintic, among others (de Santayana et al., 2014; de Santayana, Morales, Tardío, Aceituno, & Molina, 2018; Escudero, 1999; Febrer et al., 2001; Romero Martín, 1998; Verde, 2008).

2.1.2. Scientific studies

Agrimonia eupatoria, whose phytochemical composition includes compounds of interest such as carbohydrates (Carb), T, terpenes and their derivatives, PC, vitamins and oils, among other substances, has been linked to very promising biological functions at the pharmacological level, probably associated with the presence of that compounds. Some of the bioactivities *A. eupatoria* produces are antitumor, antiviral, antibacterial, antioxidant, antidiabetic, analgesic, immunomodulatory, anti-inflammatory, anti-obesity, hepatoprotective, neuroprotective and healing activities, as demonstrated in numerous experimental studies, both *in vitro* and *in vivo* (Table 3, Table 4).

Starting with the antioxidant activity and the radical scavenging power of the extracts of *A. eupatoria*, numerous studies have demonstrated its efficacy. A study conducted on the activity of extracts against the reactive species formed during inflammation associated the antioxidant capacity observed with the presence of PC. A hydroalcoholic extract and an EA fraction were able to react with DPPH, peroxy and hydroxyl radicals, as well as with other oxidizing species such as peroxynitrite, producing a significant radical scavenging activity and a powerful antioxidant activity. The tests carried out with polyphenol-rich extracts were the ones that showed the most efficacy (up to 97.5% in the DPPH reaction, and 79.5% in the ABTS (2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid)) reaction), which suggests a close relationship between the content in these compounds and the antioxidant capacity and radical scavenger (Al-snafi, 2015), as also other authors affirm that they observed a high correlation between the antioxidant function of hydroalcoholic extracts of *A. eupatoria* and their content in Pp (Muruzović, Mladenović, Stefanović, Vasić, & Čomić, 2016). Some of the PC associated with this capacity are certain phenolic acids,

Table 2
Traditional uses of species belonging to Rosaceae family.

Plant	Part Used	Mode of application	Properties	Infections Treated	Reference
<i>Agrimonia eupatoria</i>	AP	Infusion, decoction, dry extraction, wet extraction (using aqueous, oil or alcoholic solvents) and tincture	Astringent; Healing, hypotensive, anti-diarrhoea, antiemetic, analgesic, anti-inflammatory, anthelmintic	Mouthwash and gargling/ Cleaning varicose ulcers/ Renal, hepatic infections/ Controlling menstruation pain/ Detoxify blood/ Eliminating pimples and rashes	(Correia, González-Paramás, Amaral, Santos-Buelga, & Batista, 2006; de Santayana et al., 2014, 2018; Escudero, 1999; Febrer et al., 2001; Granica, Krupa, Klebowski, & Kiss, 2013; Muruzović, M. Ž. et al., 2016; Verde, 2008) (de Santayana et al., 2014; Escudero, 1999; Franco et al., 2013; Soutullo et al., 2015; Verde, 2008)
<i>Crataegus monogyna</i>	R, FL, L, FR	Tinctures, infusions, or liquid extractions	Cardiotonic, hypotensive, anti-diarrhoea, hepato-protective, anxiolytic, calming and sedative	Improving blood circulation/ Hepatitis processes/ Bronchitis and respiratory infections/ Insomnia	(Escudero, 1999; Franco et al., 2013; Rivera & Obón, 1998)
<i>Filipendula ulmaria</i>	FL	Infusions, decoction, dry or liquid extracts, tincture or syrup	Antiseptic, analgesic, anticoagulant, anti-inflammatory, antipyretic, anti-rheumatic, astringent, diuretic	Precursor of aspirin/ Cystitis/ Rheumatic pain	(Al-Snafi, 2019; Singh et al., 2019)
<i>Geum urbanum</i>	FL, R	Infusions or dried	Anti-inflammatory, haemostatic, astringent, anticholesterolemic, antihypertensive, sedative	Mouth, throat and gastrointestinal infections/ Haemorrhoids/ Genital discharges/ Skin infections	(Febrer et al., 2001; Tobyn et al., 2011)
<i>Potentilla erecta</i>	R	Infusion and decoction	Astringent, hypotensive, improve blood circulation	Intestinal colic/ Skin infections/ Urinary incontinence/ Menorrhagia/ Arthritic pain	(Blanco et al., 1999; Escudero, 1999; Verde, 2008)
<i>Rosa canina</i>	P, L, RH	Infusion, decoction, liquid extract, tincture, or syrup	Anti-diarrheic, astringent, healing, depurative, diuretic, venotonic, source of VC	Respiratory infections/ Anemia/ Control diarrhoea/ Collyrium/ Teas	

AP, aerial parts; FL, flowers; FR, fruits; L, leaves; P, petals; R, roots; RH, rose hips

Table 3
In vitro studies of *Agrimonia eupatoria*, *Crataegus monogyna*, *Filipendula ulmaria*, *Geum urbanum*, *Potentilla erecta*, *Rosa canina* as a source of bioactive compounds.

Effects	Study	Solvent	Compounds	Reference
<i>Agrimonia eupatoria</i>				
Antioxidant	DPPH, peroxyl, peroxy and hydroxyl radicals' neutralization. Protective effect against DNA damage.	HA, EA	PC, FL, T, PA, K	(Heilerová, Bučková, Tarapčík, Šilhár, & Labuda, 2018; Muruzović, M. Ž. et al., 2016; Santos et al., 2017) (Ghaima, 2013) (Khare, 2017)
Antibacterial & antibiofilm	<i>S. aureus</i> , <i>P. aeruginosa</i> , <i>E. coli</i> , <i>B. subtilis</i> inhibition	W, Et	Carb, G, T (Et) Te, PC (EO)	(Muruzović, M. Ž. et al., 2016)
	18 bacteria and 6 fungi strains	Ac	PC, F, T, PA	(Muruzović, M. Ž. et al., 2016)
<i>P. aeruginosa</i> and <i>P. mirabilis</i>	Total inhibition of blood coagulation cascade	H	Pp, Ps	(Tsirigotis-Maniecka, M., Pawlaczyk-Graja et al., 2019)
Anticoagulant	Pro-inflammatory cytokine and nitric oxide inhibition on cell cultures	W	PC	(Bae et al., 2010; Santos et al., 2017)
Anti-inflammatory	Human rhabdomyosarcoma, human cervical cancer and mouse embryo fibroblast	W, M	F, T	(Ad'hiyah et al., 2013)
Cytotoxic	Oxidative impact prevention on HT22 cells of the hippocampus	M	F	(Lee et al., 2010)
Neuroprotective				
<i>Crataegus monogyna</i>				
Antioxidant and photoprotection UV	DPPH radical scavenging effect, reducing power, inhibition of β -carotene bleaching and inhibition of lipid peroxidation of brain cells	M	PC, F	(Barros et al., 2011)
	Free radical formation limitation	CO ₂ s	N.A.	(Shortle et al., 2013)
	Photoprotection at cellular and mitochondrial level	Ch / AE	PC	(Jarzycka et al., 2013; Pawlaczyk, 2018)
	Inhibition of HepG2, NCI-H460, Hela and MCF-7	80% M	PC, F	(Rodrigues et al., 2012)
	Prolong the activated partial thromboplastin time test and the prothrombin time test	AE	Pp, Ps	(Pawlaczyk-Graja, 2018)
	<i>Lysteria monocytogenes</i> , <i>Micrococcus flavus</i> and <i>Bacillus subtilis</i> inhibition	70% Et	PC, F	(Tadić et al., 2008)
	DPPH and reducing power	W	F, T	(Samardžić et al., 2018)
	Decrease in the biosynthesis of 12-HETE and PGE2	M	PC	(Katanić et al., 2016)
	Inhibition of the COX-1 and COX-2 enzymes	W	F, AA	(Lima et al., 2014)
	Inhibition of NCI-H460, melanoma A375-C5 and MCF-7 cell lines	EA	T	(Nitta et al., 2013)
	Histidine activity decarboxylase inhibition	W, E	Pp	(Neagu et al., 2015)
	Inhibitory effect of acetylcholinesterase and tyrosinase			
<i>Geum urbanum</i>				
Antioxidant	DPPH, FRAP and linoleic acid peroxidation test	HA, EA, n-B	PC, T	(Owczarek et al., 2015).
	Inhibit prostaglandin biosynthesis and platelet activating factor-induced exocytosis	W	PC, F, PA	(Radu, Paun, Neagu, & Albu, 2015)
Anti-inflammatory	PPAR- α and PPAR- γ activation, NF- κ B inhibition	W	N.A.	(Tunón, Olavsdotter, & Bohlin, 1995)
	Inhibit prostaglandin biosynthesis and platelet-activating factor: induced exocytosis	DCIM	N.A.	(Vogl et al., 2013)
Cytotoxic	Against α -Synuclein fibrillation. Treatments against Parkinson's disease.	DCIM	T	(Granica et al., 2016)
Neuroprotective	Inhibition of acetylcholinesterase and tyrosinase	Et	N.A.	(Lobbens et al., 2016)
	Inhibition of Gram + bacteria from the genus <i>Staphylococcus</i>	Et	PC, F, PA	(Radu et al., 2015)
Antimicrobial	Inhibition of several Gram + and Gram - bacteria	EA	PC	(Dimitrova et al., 2017)
<i>Potentilla erecta</i>				
Antioxidant	Reduction of oxygen radicals cell-free-oxidant- generating systems and inflamed human colorectal biopsies	M	N.A.	(Jahantighi, Kiiani, Moghaddam, & Ghahari, 2016)
	Inhibition of UVB-induced inflammation	W	N.A.	(Langmead et al., 2002)
Anti-inflammatory	Inhibition of COX enzyme	-	T Agrimonin	(Hoffmann et al., 2016)
	Reduction of the production of IL-6 and PGE2 and NF- κ B activation in irradiated or TNF- α stimulated HaCaT keratinocytes	W	N.A.	(Tunón et al., 1995)
	Collagen gel contraction assay with VSMCs	Et	N.A.	(Wölfe et al., 2017)
Vasoconstrictive	In vitro cytotoxic activities against SMMC-7221 and HL-60 cells	Et	N.A.	(Spiridonov, Konovalov, & Arkhipov, 2005)
Cytotoxic	Inhibition of <i>B. subtilis</i> , <i>S. aureus</i> , <i>H. anomala</i> , <i>C.lipolitica</i>	Et	T, Te, Sa	(Synowicz et al., 2014)
Antimicrobial	Inhibition of <i>Herpes</i> virus types I and II, <i>Cowpox</i> and the <i>Influenza</i> virus	W	Ca	(Synowicz et al., 2014)
	Reduction of ROS in HepG2 and SH-SY5Y cells	-	T	(Tomczyk & Latté, 2009)
<i>Rosa canina</i>				
Antioxidant	DPPH, TEAC and FRAP	Et	PC	(Fetni et al., 2017)
	Inhibition of COX-1, COX-2 and of 5-LOX-mediated leukotriene B4 formation	EA, M	PC	(Ouerghemmi et al., 2016)
Anti-inflammatory	Inhibition of WiDr cells by induction of cell phase arrest and apoptosis	H, DCIM, M	Lipophilic compounds	(Wenzig et al., 2008)
Anti-tumor		DMSO	PC, VC	(Turan et al., 2018)

Ac, Acetone; AE, alkaline extraction; CE, commercial extract; Ch, chloroform; CO₂s: CO₂ supercritical; DMSO, dimethylsulfoxide; EA, ethyl acetate; Et, ether; FL, hexane HA; hydroalcoholic; M, methanol; W, water. AA, ascorbic acid; Ca, Catechins; EO, essential oil; Carb, carbohydrates; E, Flavonoids; FL, flavones; G, glycosides; K, kaempferol; NA, not analysed; PA, proanthocyanidin; PC, Phenolic compounds; Pp, polyphenols; Ps, Polysaccharides; Sa, saponins; T, tannins; Te, terpenoids; VC, vitamin C;

procyanidins, F, flavan-3-ols, T, proanthocyanidins (PA), Ca, quercetin, apigenin, kaempferol (K) and their O-glucosidic derivatives, having been reported to be up to 220.31 mg of the total PC per gram of extract (Al-snafi, 2015; Muruzović et al., 2016).

Regarding the anti-inflammatory capacity, it was possible to inhibit the production of pro-inflammatory cytokine and nitric oxide of microglial cells *in vivo* by applying the aqueous extract of *A. eupatoria* (Bae et al., 2010; Yoon et al., 2012). This extract also achieved a reduction in the increase in serum aminotransferase activity, as well as in the production of tumor necrosis factor α (TNF- α) in rats to which 10, 30, 100 and 300 mg/kg/day of aqueous extract had been orally administered, in addition to imposing a chronic consumption of alcohol. It was also observed that the extract produced a protective effect against chronic liver damage caused by Et (Al-snafi, 2015). Luteolin 7-glucuronide is an active molecule that is present in the extract and is associated with the decreases in the inflammatory process, an effect probably caused by the inhibition of oxidative stress and inflammation signaling mediated by TLRs (Yoon et al., 2012). On the other hand, the ethyl extract of *A. eupatoria* managed to reduce peripheral edema in *in vivo* models by 43.2% (when applying a single dose) and 52.2% (when a second dose was applied). Likewise, in this study the analgesic capacity of the ethyl extract was tested, resulting in the fact that it did not produce effects of general anesthesia, but was verified by the formalin and the writhing test, which produced an inhibitory effect of 43.5% in a single dose and 49.8% when applying a second dose (Al-snafi, 2015).

The cytotoxic effects of the aqueous and methanolic extracts of *A. eupatoria* were tested *in vitro* tests on human RD, HeLa and mouse embryo fibroblast cell lines, using different concentrations (6, 12, 24, 48 and 96 $\mu\text{g/mL}$) of the extracts, and throughout three rounds. In the results, it was observed that all the concentrations analyzed achieved the antitumor effect, although proportionally to the concentration, and that the methanolic extract achieved better results in the RD and HeLa cell lines than the aqueous extract, since the percentage growth inhibition was greater (Al-snafi, 2015; Tsirigotis-Maniecka et al., 2019). Some authors associate the antitumor activity observed in the RD and HeLa tumor cell lines with the presence of F and T in the extracts obtained from *A. eupatoria* (Ad'hiah, Al-Bederi, & Al-Sammarae, 2013).

Immunomodulatory capacity was demonstrated using a hydroalcoholic extract in the peritoneal cavity of mice, which resulted in immunostimulatory activity through the activation of phagocytes, lysozyme and peroxidase (Al-snafi, 2015).

In addition, the methanolic extract of *A. eupatoria* was shown to have neuroprotective activity by decreasing the damaging oxidative impact on HT22 cells of the mouse hippocampus produced by glutamate (Al-snafi, 2015; Lee et al., 2010). This effect was associated with various compounds present in the extract with antioxidant capacity. Among these compounds, a new flavonoid that was not previously characterized as standing out, was identified as kaempferol 3-O- β -D-(2'-O-acetyl-6'- E)-p-coumaroyl)-glucopyranoside. Other nine R known as F were found in the methanolic extract (Lee et al., 2010).

The antibacterial properties of the aqueous and ethanolic extracts of *A. eupatoria* were tested against several clinically important strains, such as *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Escherichia coli*. In this case, the ethanol (Et) extract was the one that showed the greatest inhibitory efficacy, specifically against *E. coli*, followed by *S. aureus*, with a 15 mm inhibition halo. *P. aeruginosa* was the most resistant strain to the extract. The effect of the EO of *A. eupatoria* was also tested, reported to be active against *Bacillus subtilis* (Al-snafi, 2015). In another study, 24 species of microorganisms were tested, both bacteria, for which ampicillin and tetracycline were used as control, as fungi, whose control substances were itraconazole and amphotericin B. The extract that showed the greatest antimicrobial potential was acetone (Ac) one (Minimal Inhibitory Concentration of 4469.5 $\mu\text{g/mL}$ for *P. aeruginosa* and 4315 $\mu\text{g/mL}$ for *P. mirabilis*), followed by aqueous, Et and, finally, diethyl ether. Additionally, Ac extract also produced a significant inhibition of biofilm formation from *P. aeruginosa* and

Proteus mirabilis (Muruzović et al., 2016).

In addition to all these bioactivities, antidiabetic properties are also associated with *A. eupatoria*, since the aqueous extract of this plant (1 mg/mL) achieves an antihyperglycemic effect in mice, producing insulin released by the B cells of the pancreas and the stimulation of glucose transporters. Furthermore, the extract reduced symptoms of this disorder characterized as weight loss, hyperphagia and polydipsia (Al-snafi, 2015).

Other activities related to the compounds present in this plant are healing, analyzed in comparison with fucidin, which achieved complete healing in rats in 14 days, while the aqueous extract achieved it in 12 days, and the ethanolic in just 10 days; antihypertensive effect, since it manages to lower blood pressure by 40% by administering the extract intravenously in cats; uricolytic capacity, by providing infusions or decoctions at 15% w/v in male rats in doses of 20 mL/kg (Al-snafi, 2015); anticoagulant function, tested *in vitro* using an extract of the AP of *A. eupatoria* rich in Pp and Ps that managed to totally inhibit the blood coagulation cascade and, therefore, the formation of clot plasma, at a concentration of 500 $\mu\text{g/mL}$ (Tsirigotis-Maniecka et al., 2019); or a lipid lowering activity, observed in healthy patients who were supplied with *A. eupatoria* tea, which produced an increase in high density lipoprotein (HDL) cholesterol, also achieving an improvement in lipid profiles (Ivanova, Vankova, & Nashar, 2013).

The European Medicines Agency (EMA) considered the properties of the herbal preparations (comminuted, tinctures or liquid extracts) and its pharmaceutical forms (comminuted for oral use or as infusion or decoction preparations for oromucosal and cutaneous uses). The uses included in the EMA report are the relief of mild diarrhoea (oral administration), mild inflammation of the mouth and throat (used for gargles), and minor skin inflammation and small superficial wounds (topic administration). It has been recommended to applied in adults and children over 12 years (EMA/HMPC/680595/2013, 2015).

2.2. *Crataegus monogyna*

2.2.1. Traditional importance

C. monogyna is widely distributed. It is considered native to Europe, but not in the northeastern part and the Mediterranean coast of North Africa. Roots (R), flowers (FL), leaves (L) and fruits (FR) from *C. monogyna* have different traditional uses in Iberian Peninsula (Garmendia & Castroviejo, 1998). Tinctures, infusions, or liquid extractions are the most typical ways of application (Font Quer, 2001). Even though L and FR are not considered to possess a high nutritional value, they have been also used as food (de Santayana et al., 2014). L and FL from *C. monogyna* have been found to have circulatory properties, improving blood circulation and being cardiogenic and hypotensive (Font Quer, 2001). FR and FL have been used at digestive levels to prevent diarrhea and used to protect the liver in hepatitis processes. FR and R were applied in bronchitis and respiratory infections. For nervous system infections, all plant parts were mostly utilized, usually to treat insomnia, and as anxiolytic, to calm and sedate (de Santayana et al., 2014; Escudero, 1999; Franco et al., 2013; Soutullo, Muñoz, Pazó, Alonso, & Boente, 2015; Verde, 2008).

2.2.2. Scientific studies

There are many studies that have been conducted of the bioactivities of *C. monogyna*, in which its antioxidant, cardioprotective, anticancer, anticoagulant and antimicrobial capabilities are proven, among others (Table 3, Table 4). The content of PC in this species is variable depending in the analyzed tissue, ranging from 55 in FR up to 377 μg GAE/g in stems (Luís, Domingues, & Duarte, 2011). Thus, this plant is rich in PC, such as phenolic acids, F, PAs and Ca, as well as tocopherols, ascorbic acid (AA), β -carotene and both saturated fatty acids (hexadecanoic and tricosanoic acid) and polyunsaturated ((9Z, 12Z)-octadeca-9,12-dienoic and (9Z, 12Z, 15Z)-octadeca-9,12,15-trienoic); as well as inorganic compounds such as Cu, Fe, Mg, Mn, Zn, etc. (Barros,

Table 4

In vivo studies of *Agrimonia eupatoria*, *Crataegus monogyna*, *Filipendula ulmaria*, *Geum urbanum*, *Potentilla erecta*, *Rosa canina* as a source of bioactive compounds.

Effects	Study	Solvent	Compounds	Reference
<i>Agrimonia eupatoria</i>				
Anti-inflammatory	Carrageenan-induced rat paw edema test in mice	W	PC	(Santos et al., 2017)
Hepatoprotective	Inhibition of serum aminotransferase activity and production of TNF- α on mice liver	W	N.A.	(Yoon et al., 2012)
Antidiabetic	Insulin release by the B cells of the pancreas and glucose transporters stimulation in mice	W	N.A.	(Gray & Flatt, 1998)
Antihypertensive	Blood pressure reduction by 40% when administered intravenously in cats	W, Et, Er, Ch, M, Ac, DCIE	N.A.	(Petkov, 1979)
Healing	Complete healing in 12 days in rats	W	Carb, G, T. In Etl also	(Ghaima, 2013)
	Complete healing in 10 days in rats	Et	Te and PC	
Lipid lowering	Increase in high density lipoprotein (HDL) cholesterol in humans	W	N.A.	(Ivanova et al., 2013)
<i>Crataegus monogyna</i>				
Anti-inflammatory	72.4% inflammation decrease in rats	70% Et	Te	(Tadić et al., 2008)
Antithrombotic	Inhibition of platelet aggregation in mice	50% Et	PC, F	(Arslan et al., 2015)
Cardioprotective and antiarrhythmic	Protective effect of myocardial dysfunction and infarction occurrence in rats	CO ₂ s	PC, F	(Shortle et al., 2013)
	Vasodilator effect and an inotropic effect in guinea pigs	A.E	PC, F	(Pawlaczyk-Graja, 2018)
Analgesic	Depression of the central nervous system. Analgesic effects mediated by endogenous opioid system	80% Et	N.A.	(Can, Özkay, Öztrk, & Öztürk, 2010)
<i>Filipendula ulmaria</i>				
Antioxidant and hepatoprotective	Hepatic oxidative stress reduction in Wistar rats	M	Pp	(Katanić et al., 2017)
Anti-inflammatory	Reduction carrageenan-induced paw edema in rats	M	Pp	(Katanić et al., 2016)
Cytotoxic	Inhibitory effect of colorectal carcinogenesis in rats	W	PC	(Bespalov et al., 2018)
	Cervix and vagina carcinoma reduction in mice	W	F, AA	(Spiridonov et al., 2005)
Gastroprotective	Acute stomach damage caused by Et reduction	W	F, T	(Samardžić et al., 2018)
Immunomodulatory	Complementary immune system activation	W	W-soluble Ps	(Olennikov et al., 2017)
<i>Geum urbanum</i>				
Hypotensive	Intravenous injection, has been reported to produce a reduction in blood pressure in cats	W	N.A.	(Petkov, 1979)
<i>Potentilla erecta</i>				
Vasoconstrictive	Occlusive patch test in humans	Et	N. A.	(Wölfle et al., 2017)
Anti-diarrhoea	Reduction of diarrhea caused by rotavirus in children	CE	T	(Subbotina et al., 2003)
Anti-ulcerogenic	Inhibition of ulcerative colitis in humans	CE	T	(Huber et al., 2007)
Antihyperglycemic	Improvement of diabetes symptoms in mice	M	Tomentoside	(Tomczyk & Latté, 2009)
<i>Rosa canina</i>				
Antioxidant	Has the power to reduce pain and stiffness on human patients with osteoarthritis	-	N.A.	(Nadpal et al., 2016; Turan et al., 2018)
Anti-inflammatory	Carrageenan-induced and PGE1-induced hind paw edema models, as well as on acetic acid induced increase in a capillary permeability model	W, Et	PC and VC	(Deliorman Orhan et al., 2007; Ouerghemmi et al., 2016)
Antinociceptive	ρ -Benzoquinone-induced abdominal constriction test	W, Et	PC and VC	(Deliorman Orhan et al., 2007))
Anti-tumor	Inhibited melanogenesis in mouse melanoma cells and guinea pig skin, and could be useful as a skin-whitening agent when taken orally	Et	PA	(Fujii et al., 2011)
Anti-obese	Inhibitory effect on the body weight gain and in the visceral fat gain in mice	Ac	T-t	(Ninomiya et al., 2007)

Ac, Acetone; AE: alkaline extraction; CE, commercial extract; Ch, chloroform, CO₂ s: CO₂ supercritical; DMSO, dimethylsulfoxide; EA, ethyl acetate; Er, ether; Et, ethanol; H, hexane; M, methanol; W, water.

AA, ascorbic acid; Ca, Catechins; Carb, carbohydrates; EO, essential oil; F, Flavonoids; Fl, flavones; G, glycosides; K, kaempferol; NA, not analysed; PA, proanthocyanidin; PC, Phenolic compounds; Pp, polyphenols; Ps, Polysaccharides; Sa, saponins; T, tannins; Te, terpenoids; To, tomentoside; T-t, Trans-tiltoside; VC, vitamin C.

Carvalho, & Ferreira, 2011; Jarzycka, Lewińska, Gancarz, & Wilk, 2013).

The antioxidant activity of *C. monogyna* extracts has been demonstrated *in vitro* using four methodologies applied to animal cell lines: DPPH radical scavenging effect, reducing power, inhibition of β -carotene bleaching and inhibition of lipid peroxidation of brain cells. All the analyzed extracts showed antioxidant capacity, being especially apparent when using the extracts of unripe FR (EC₅₀ between 5.42 and 20.83 μ g/mL), which coincides with the high amount of PC and F that contain these extracts. (Barros et al., 2011). Interestingly, in an essay on the effect of the inclusion of extracts in W-soluble gels with the intention of incorporating them into cosmetic and/or pharmaceutical preparations, the antioxidant capacity was verified, using the same methodologies mentioned above. The results showed a decrease of this property, but not significant regarding the use of isolated extracts (Barreira, Rodrigues, Carvalho, & Ferreira, 2013).

The antioxidant compounds contained in *C. monogyna* also have a

high potential for protection against UV radiation, both UVB and UVA, experimentally tested *in vitro* at the cellular and mitochondrial level, as well as high photostability (Jarzycka et al., 2013; Pawlaczyk-Graja, 2018). In addition, these compounds are associated with a cardioprotective effect, thanks to the limitation of free radical formation, reducing possible damage to the heart and the deposition of cholesterol in the arteries, as it has been demonstrated *in vitro*. In rat models, a protective effect of myocardial dysfunction and the occurrence of infarction was observed (Shortle, Kerry, Furey, & Gilroy, 2013). Moreover, the extracts achieved a vasodilator effect and an inotropic effect that affected sodium-potassium (Na⁺/K⁺)-ATPase and that favored the Ca²⁺ ions transport in cardiomyocytes. In ventricular myocytes of guinea pigs, a decrease in arrhythmia was observed through a mechanism of actions similar to that of class III antiarrhythmics (Pawlaczyk-Graja, 2018), as well as a negative chronotropic effect *in vitro*, reducing the contraction of cardiomyocytes through the activation of muscarinic receptors. Certain F isolated from *C. monogyna* such as

luteolin-7-glucoside, hyperoside, and rutin produced a cardioprotective effect in the hearts of isolated guinea pigs, favoring coronary flow (186%, 66%, and 66%, respectively) and velocity of relaxation (104%, 62%, and 73%, respectively) in doses of 0.5 mmol/L (Nabavi et al., 2015).

Going deeper into the content of the extracts, (Pawlaczyk-Graja, 2018) polyphenol-polysaccharide conjugates were isolated and characterized by using various spectrophotometric methods such as GLC-MS, FT-IR and NMR techniques. These were Pp matrices, including some F and poorly esterified Ps rich in galacturonic acid, capable of increasing blood clotting time with very low concentrations (31.25 µg/mL), as verified *in vitro*, although the FL extract was the one that showed an action with greater selectivity (Pawlaczyk-Graja, 2018). The antithrombotic effect of the methanolic extract was tested in mouse tail models injected with a carrageenan, using heparin as the control substance. Extract doses of between 200 and 300 mg/kg were the most effective, achieving an inhibition of platelet aggregation with a high significance (Arslan, Bektas, Bor, & Sener, 2015; Nabavi et al., 2015).

The oral administration of *C. monogyna* extracts to rats with edemas demonstrated the anti-inflammatory effect of this plant, even surpassing what was produced by indomethacin, a drug used as a positive control, since a 72.4% decrease in inflammation was reported when the extract was used, compared to 50% achieved by the control. This activity is associated with the presence of triterpenes, especially cycloartenol (Nabavi et al., 2015).

The antitumor capacity of *C. monogyna* extracts, highly related to the antiproliferative and the apoptotic effect of PC, was tested in four human tumor lines: NCI-H460, HepG2, HeLa and MCF-7, resulting in extracts obtained from FL being the most effective. HPLC-DAD-ESI/MS analyzes of the extracts revealed a predominance of F (Rodrigues et al., 2012). An interesting fact is that, in addition to not producing toxicity in non-tumor cells, the aqueous extract of *C. monogyna* is capable of exerting a protective effect against the toxicity caused by doxorubicin, a broad-spectrum chemotherapeutic drug whose use has been restricted (Shalizar Jalali & Hasanzadeh, 2013).

The antibacterial activity produced by the extracts of *C. monogyna* is moderate, especially against Gram-positive bacteria, such as *Listeria monocytogenes*, *Micrococcus flavus* and *B. subtilis* (Nabavi et al., 2015).

Another capacity that deserves to be highlighted is the regulation of the central nervous system, since it depresses its functionality as observed in mice, whose locomotor activity decreased after the administration of *C. monogyna*. An analgesic effect associated with the activation of opioid receptors was also observed; as well as a gastro-protective function comparable to that of ranitidine in rat models (Nabavi et al., 2015).

The EMA report for *Crataegus monogyna* offers a wide range of optional herbal preparations and pharmaceutical forms (comminuted/powdered, dry and liquid extracts using different extraction ratios and solvents, such as ethanol, methanol or sweet wine, juices, and tinctures). Leaf and flower preparations have been associated with the relief of nervous-based temporary heart complaints, such as palpitations, but it requires the previous exclusion of medical major and serious conditions. Same preparations may also relieve mild symptoms of mental stress and to aid sleep. Their use is contemplated for adults and children over 12 years (EMA/HMPC/159076/2014, 2016).

2.3. *Filipendula ulmaria*

2.3.1. Traditional uses

F. ulmaria is a common and widespread plant, native to Europe and Western Asia, while it has also been introduced but naturalized to North America, including some states from the United States (US) and Canada, where it has become naturalized. Flowered apexes from *F. ulmaria*, preferably collected when flowering is beginning, are consumed as an infusion, decoction, dry or liquid extract, tincture or syrup (Escudero, 1999). Antiseptic, analgesic, anticoagulant, anti-inflammatory,

antipyretic, anti-rheumatic, astringent, diaphoretic and diuretic properties have been associated with this species (Escudero, 1999; Franco et al., 2013; Rivera & Obón, 1998). Pharmaceutical industries utilized this plant as a precursor to aspirin since it contains salicylic acid. In addition, it has been applied for the treatment of cystitis and rheumatic pain (Font Quer, 2001). L and FL of *F. ulmaria* have been also used for aromatizing soups or wine and beers, respectively. It is a useful resource of other colors and Eos in cosmetics (Escudero, 1999).

2.3.2. Scientific studies

F. ulmaria is known to possess many biological activities, among which antioxidant, anti-inflammatory, antimicrobial and anti-proliferative capacities stand out (Katanić et al., 2017) (Tables 3 and 4). Another interesting property that has been associated with this MP is a healing function. PC are the constituents of *F. ulmaria* that mostly related to bioactive potentials, and can be divided into three main groups: phenolic acids (salicylic acid, gallic acid, ellagic acid, salicylaldehyde, etc.), F and their G (K, rutoside, epicatechin, Ca, quercetin, spiraeoside, apigenin, hyperoside, astragalin, etc.), and T (rugosin A, B1, B2, D, E1 and E2, tellimagrandin I and II, etc.) (Katanić et al., 2016).

The antioxidant properties of *F. ulmaria* extracts have been reported and demonstrated by many studies (Neagu, Paun, Albu, & Radu, 2015; Olennikov, Kashchenko, & Chirikova, 2017). A recent investigation analysed the radical scavenging activity of lyophilized FL infusions, in which the potential for *in vitro* neutralization of DPPH and OH radicals was evaluated, as well as the reducing power through the FRAP assay, obtaining IC₅₀ values of 8.45 µg/mL for DPPH and a FRAP value of 4.46 mmol Fe²⁺/g. However, the results suggest that this extract has no potential to neutralize OH radicals (Samardžić et al., 2018). Taking into account the aerial part and the root of *F. ulmaria*, the methanolic extracts of these parts achieved potent antioxidant effects demonstrated *in vivo* Wistar rats, since the administration of such extracts for 10 consecutive days reduced the hepatic oxidative stress induced by cisplatin, a drug used in antitumor therapies and that of which is related to numerous adverse effects. An increase in the activities of catalase and superoxide dismutase, and a reduction of lipid peroxidation and suppression of DNA damage in normal cells was observed (Katanić et al., 2017).

Regarding the anti-inflammatory activity of the extracts of *F. ulmaria*, results obtained both *in vitro* and *in vivo* show an important potential. FL extracts achieved a moderate effect of the activation of PPARs (Katanić et al., 2016). The infusion of lyophilized FL studied *ex vivo* in human platelets achieved a decrease in the biosynthesis of 12-HETE and PGE2 with IC₅₀ values of 3415 and 6768 µg/mL, respectively (Samardžić et al., 2018). The inhibition of NF-κB achieved by extracts from other plant parts of *F. ulmaria* was potentially significant (Katanić et al., 2017). Likewise, it was demonstrated that the extracts of *F. ulmaria* managed to reduce the production of cytokines *in vitro*, IL-1β, IL-6 and TNF-α, in addition to the inhibition of the activity of 15-LOX. The tests carried out with the extracts (50 µg/mL) of the AP and R of *F. ulmaria* achieved a significant inhibition of COX-1 and COX-2, being the predominant inhibitory effect of the extract of the AP (62.84% COX-1 inhibition and 46.43% COX-2 inhibition), and, therefore, comparable to the inhibitory values achieved using the control substances, indomethacin and NS-398 at concentrations of 1.25 µM and 5 µM, respectively (Katanić et al., 2016). In *in vivo* trials in rats, in which carrageenan had been injected, causing tail edema, it was observed that the administration of 100 mg of AP extract and 200 mg of R extract per kg of body weight, was capable of significantly reducing inflammation, thereby achieving the maximum peak of action at 24 h (Katanić et al., 2016).

Immunomodulatory properties of *F. ulmaria* have also been demonstrated. W-soluble Ps obtained from *F. ulmaria* FL have the ability to activate the complementary immune system in a dose-dependent manner. This system is related to the beginning of inflammatory processes, leukocyte activation and degranulation of basophils and mast

cells. (Olennikov et al., 2017).

Similarly, the hepatoprotective and neuroprotective properties of *F. ulmaria* have been investigated. The hepatoprotective capacity of an Et extract was tested *in vivo*, resulting in the aforementioned beneficial action, which was closely related to the antioxidant properties of the plant, as well as the ability to stabilize membranes. The characterization by LC-DAD-MSn of the extracts of AP and R allowed the broad phenolic composition present in this plant to be elucidated, among which the quercetin stands out, being identified in both plant parts, and related to strong antioxidant properties (Katanić et al., 2017). The neuroprotective capacity was tested through the inhibitory effect of the AChE and tyrosinase enzymes, using aqueous and ethanolic extracts. At concentrations of 3 mg/mL, ethanolic extract achieved a greater inhibition of AChE, 98.3% versus an inhibition of 77.03% produced by the aqueous extract. In the case of tyrosinase, the inhibitions achieved showed fewer differences: 90.65% for ethanolic extract and 87.88% for W extract. The PC associated with this bioactivity were Pp such as quercetin and rutin, and chlorogenic acid (Neagu et al., 2015).

On the other hand, the anti-tumor effects of *F. ulmaria* extracts, such as those produced by FL extracts, which inhibit the proliferation of NCI-H460 tumor cells and cause an increase in p21 levels (at G1-checkpoint cyclin-dependent kinase inhibitor) have also been reported. In addition, concentrations between 10 and 50 µg/mL of the ethanolic extracts of *F. ulmaria* produced cytotoxicity in Burkitt lymphoma cell lines (Lima et al., 2014). Additionally, the inhibitory activity of a decoction of *F. ulmaria* in colorectal tumor processes was evaluated *in vivo* in female rats to which carcinogenesis had been induced using methylnitrosourea. A potent inhibitory effect of colorectal carcinogenesis was observed, probably related to the anti-inflammatory potential of *F. ulmaria* extracts in certain stomach and intestinal pathologies (Bespalov et al., 2018). Another study that corroborates the antitumor action of the extracts of *F. ulmaria* was performed in mice with cervical dysplasia, to which extracts obtained by decoction of FL were locally administered. The extracts were able to reduce the frequency of occurrence of carcinoma of the cervix and vagina, induced by 712-dimethyl-benz (a) anthracene (Lima et al., 2014).

Other effects related to *F. ulmaria* extracts are gastroprotective capacity, demonstrated in animal models with acute stomach damage caused by ethanol, to which oral infusions of lyophilized FL (100–300 mg/kg) were orally administered (Samardžić et al., 2018). Likewise, some PC belonging to the group of T, found in EA extracts from P of *F. ulmaria*, such as rugosin D, rugosin A, rugosin A methyl ester and tellimagrandin II, were able to significantly inhibit histidine activity decarboxylase, an enzyme related to the symptoms of many allergies and stomach ulcers (Nitta et al., 2013). *F. ulmaria* tea intake is also associated with an inhibitory effect of the enzymes amylase and α-glucosidase, as well as the formation of advanced glycation end products (Olennikov et al., 2017).

The EMA report for *Filipendula ulmaria* includes herbal and pharmaceutical forms (comminuted for flowers and herbs, and powdered or tinctures with ethanol for the herb). They are recommended for the relief of cough and cold symptoms, as well as for pain and inflammation. Due to the lack of enough clinical safety data, the use of these herbal/pharmaceutical preparations is not recommended during pregnancy and lactation and in children and adolescents under 18 years of age (EMA/HMPC/434892/2010, 2013).

2.4. *Geum urbanum*

2.4.1. Traditional use

Geum urbanum is widely extended throughout Europe, where it can reach elevations up to 2100 m over sea level. It is also considered to be native to North Africa (Algeria, Morocco, Tunisia), and temperate Asia from Anatolia, Syria and Caucasian areas up to further occidental and central zones. References about Iberian traditional uses of *Geum urbanum* are not easily found. Nevertheless, scientific references from

other geographical areas point out that all parts of the plant have anti-inflammatory properties. Mostly, infusions of dried or fresh FL and R are valued since they were demonstrated to be astringent, anticholesterolemic, antihypertensive, sedative and haemostatic. Mouth, throat and gastrointestinal tract affections, such as loss of appetite or diarrhea have been treated using infusions. In addition, these infusions can be externally applied to wash hemorrhoids, genital discharges and skin infections (Al-Snafi, 2019; Singh et al., 2019).

2.4.2. Scientific studies

G. urbanum has been used for a long time in folk medicine due to its astringent and antiseptic properties. More recent studies have shown that its traditional use makes sense, considering *G. urbanum* is rich in PC like gallic, caffeic and chlorogenic acids, and also eugenol, carotenoids, F, T and vicianose sugar, which is a very rare disaccharide that can be isolated from its R (Al-Snafi, 2019; Owczarek, Gudej, & Olszewska, 2015) (Table 3, Table 4). Other studies also found ellagitannins and procyanidins located in the R of *G. urbanum* and the EO extracted from the AP have also been studied (Al-Snafi, 2019; Dimitrova et al., 2017).

Total phenolic acids, F and PA content were evaluated in aqueous extract and ethanolic *G. urbanum* extracts. In the first, total phenolic acids were 768.2 GAE/l, total F were 14.7 mg RE/l and PA were 37.5 mg CE/L, whereas in the second, total phenolic acid was 1261.5 mg GAE/l, a total flavonoid, 49.9 mg RE/l and PA, 60.1 mg CE/L (Al-Snafi, 2019). Ellagitannins and gallotannins are also known constituents of *G. urbanum*, and ellagic acid is also present. Ellagic acid is a potent antioxidant properties with beneficial properties to human health, such as hepatoprotective, cardioprotective, and chemopreventive effects (Owczarek, Olszewska, & Gudej, 2014). Another potent antioxidant detected in high amounts is gallic acid, which also showed antiviral activity (Owczarek et al., 2014).

The antioxidant activity of different *G. urbanum* extracts has been demonstrated in several studies. (Mantle, Eddeb, & Pickering, 2000) analysed a hydroalcoholic extract of *G. urbanum* L, demonstrating a powerful antioxidant effect through the ABTS assay, with values of 1.67 mmol Trolox/equivalent/mg of dry weigh. Hydromethanolic extracts of the underground and AP were evaluated through FRAP, DPPH, and linoleic acid oxidation assays. The assays had similar antioxidant activities, which the author correlates with the presence of PC (Owczarek et al., 2015). Aqueous and Et extract also showed a remarkable scavenging activity against 2, 2-diphenyl-1-picrylhydrazyl with values of IC₅₀ 7.8 µg/mL and IC₅₀ 1.3 µg/mL, respectively (Al-Snafi, 2019).

Regarding anti-inflammatory properties, *Geum urbanum* was able to inhibit prostaglandin biosynthesis and platelet-activating factor-induced exocytosis *in vitro* tests. Different extracts, in a concentration of 10 mg/mL, were tested on PPAR-α and PPAR-γ activation as well as on NF-κB inhibition. The best results were obtained with the dichloromethane extraction, which possessed moderate to strong effects (Al-Snafi, 2019).

G. urbanum was also tested through R infusions and decoctions. These extractions have been used externally for reducing gingivitis (bleeding and inflammation of gums), and other membranes mucous membranes (Granica et al., 2016). This action is attributed to its high content in Gemin A. This compound significantly affects the functions of stimulated neutrophils by reducing the surface expression of CD11b, and inhibiting the release of reactive oxygen species, and proteases (elastase, matrix metalloproteinase-9), chemokines and cytokines (interleukins IL-8, IL-1β), besides it also stimulates the release of TNF-α, which may be one of the stimulators of the apoptosis of neutrophil cells (Granica et al., 2016).

G. urbanum extracts were also tested for their *in vitro* neuroprotective effect via the inhibition of AChE and tyrosinase. The aqueous extract had an AChE inhibitory activity of 27.03%; 36.48% and 79.11% at concentration of 0.75 mg/mL, 1.5 mg/mL and 3 mg/mL respectively. The IC₅₀ was 2.3 mg/mL for this extract. For the ethanolic extracts,

AChE inhibitory activities were 54.74; 73.53 and 86.77 for the same concentrations. The IC₅₀ was 0.5 mg/mL. Even though all concentrations of the aqueous extracts inhibited tyrosinase in more than 50%, ethanolic extracts were more potent than the aqueous ones (Al-Snafi, 2019).

Furthermore (Dimitrova et al., 2017), also studied the antimicrobial potential of extracts from *G. urbanum* against Gram-positive. EA fractions from the AP and R had a MIC of 78 µg/mL and 156 µg/mL, respectively, and inhibited the visible growth of the bacteria tested, suppressing their respiratory activity up to 26.2–28.9%. The minimal bactericidal concentration calculated was 625 µg/mL (Dimitrova et al., 2017).

The presence of Lewy bodies and Lewy neuritis is thought to be a major pathological hallmark of Parkinson's disease and is also linked to the disease's development. These aggregations of protein primarily consist of fibrillated α-Synuclein; and for now, there is no treatment available targeting stabilization of α-Synuclein in its native state. In the study of (Lobbens et al., 2016), an ethanolic extract of *G. urbanum* was found to inhibit α-Synuclein fibrillation in a concentration-dependent way and to partly disintegrate preformed α-Synuclein fibrils.

Finally, *in vivo* studies performed in cats have shown that a 20% aqueous decoction of *G. urbanum*, administered by intravenous injection, produced a reduction in their blood pressure (Al-Snafi, 2019).

2.5. *Potentilla erecta*

2.5.1. Traditional use

P. erecta is a Euroasiatic plant widely spread throughout Europe, Caucasus, Anatolia and Western Siberia, up to 2,300 m above sea level. It can be also found in the northwest parts of Africa, Azores and Madeira and, as *F. erecta*, it has been naturalized in North America (Guillén & Rico, 1998). R from *P. erecta* are usually applied orally as an infusion to treat diarrhea and intestinal colic, due to its astringent properties. The decoction of the same part of plant can be topically used for skin infections and also hemorrhoids (Font Quer, 2001). *P. erecta* is also considered to be hypotensive, improve blood circulation and to control urinary incontinence, menorrhagia, and arthritic pain (Febrer et al., 2001; Tobyn, Denham, & Whitelegg, 2011).

2.5.2. Scientific studies

P. erecta is an interesting plant with different confirmed bioactivities, such as antioxidant, antimicrobial, anti-inflammatory and anti-diarrhea properties (Table 3, Table 4). These properties are mostly associated with the presence of different molecules like F, triterpenoids and phenolic acids, which are usually extracted from the R and rhizomes of this plant, being the most abundant T, with 15 to 25% of dry weight. The rhizome of this plant is rich in PA and hydrolysable T, mostly agrimoniin, a dimeric ellagitannin that represents about 2% of the dry mass (Wölfle, Hoffmann, Haarhaus, Rao Mittapalli, & Schempp, 2017). These compounds are thought to be beneficial for human health due to their antioxidant properties and because PA are capable of complex macromolecules and metal ions. In addition, they have been proven to play an important role in the anti-inflammatory activity of several diseases, probably due to their scavenging and antioxidant activities (Mari, Eletto, Pizza, Montoro, & Piacente, 2013).

A *P. erecta* rhizome extract showed antioxidant effects in cell-free oxidant-generating systems and in inflamed human colorectal cells, reducing the production of oxygen radicals (Tomczyk & Latté, 2009).

A study from (Wölfle et al., 2017) determined the anti-inflammatory activity of *P. erecta* in irradiated HaCaT keratinocytes, by measuring the formation of IL-6 and PGE₂. They also analysed the effect on TNF-α induced NF-κB activation, both tests having demonstrated great results in the reduction of inflammation. In this study, *P. erecta* also showed a blanching effect comparable to hydrocortisone, that it is thought to be partly attributable to a scavenging effect of NO and inhibition of eNOS. However, compared to glucocorticoids, *P. erecta* was not able to cause

nuclear translocation of the glucocorticoid receptor in HaCaT cells.

In a different study, extracts from *P. erecta* combined with an hydrolysable tannin Agrimonnin, known to be a potent radical scavenger, were able to reduce the inflammation caused by UVB-rays in HaCaT keratinocytes through the inhibition of cyclooxygenase-2 in a dose-dependent manner (Hoffmann et al., 2016; Wölfle et al., 2017). A further experiment demonstrated that the extracts produced a dose-dependent inhibition of UVB-induced inflammation in a *in vivo* model (Hoffmann et al., 2016).

The cytotoxic properties of *P. erecta* have been evaluated. (Tomczyk & Latté, 2009) studied the crude extract of *P. erecta* rhizome in 40% Et and determined that concentrations of 10 and 50 µg/mL were able to inhibit lymphoma cell growth. They also studied the effect of the extract on the herpes virus types I and II *in vitro* and it revealed a moderate antiviral effect, as well as its cytotoxic activity against Cowpox and the influenza virus type A2 (Tomczyk & Latté, 2009).

(Synowiec, Gniewosz, Bączek, & Przybył, 2014) evaluated the antibacterial properties of an aqueous extract of *P. erecta* rhizome and it displayed an inhibiting effect against Gram-positive bacteria, like *B. subtilis* ATCC 6633 and *S. aureus* ATCC 25923, and also against yeast-like *Hansenula anomala* R 26 and *Candida lipolitica* KKP 322.

As mentioned before, *P. erecta* presents anti-diarrhoea properties, which have been corroborated by different studies. The oligomeric and polymeric flavan-3-ols (also known as condensed T or PA) have shown therapeutic properties in the treatment of diarrhoea, by complexing secretory compounds such as cholera toxin. Moreover, they also created an unspecific complexation of mucosal proteins in the gut, forming a protective layer and inhibiting the intestinal motility associated with the secretory compounds (Tomczyk & Latté, 2009). This action was tested with a rhizome dry extract and the results showed a high efficacy in the therapy of children's diarrhoea due to Rotavirus. The treatment ensured a shorter duration of the diarrhoea (three days) compared with the placebo group (five days). In the treated group, 40% of the children were diarrhoea free 48 h after admission to the hospital. In the placebo group this only happened to 5% of the children, which leads the authors to assume that there is an interaction between the T and the Rotavirus proteins that contributes to the high efficacy of this extract, but they also admit that another unknown ingredient might also be involved (Tomczyk & Latté, 2009).

Another use for the *P. erecta* extract is in the treatment of active ulcerative colitis. Sixteen patients were treated with the extract for three weeks with a dosage of 2,400 mg/day, resulting in a significant decline in the colitis activity index (Tomczyk & Latté, 2009).

Besides, *P. erecta* even decreased the biochemical indices of lipid metabolism in the blood, specifically of malonaldehyde and endogenous lipids, after three administrations of a spirituous tincture from the rhizome, on albino rats under normal physiological conditions. The tincture was administered intragastrically for 14 days at 0.05 and 0.1 mL per 100 g of the animals' body weight (Tomczyk & Latté, 2009).

2.6. *Rosa canina*

2.6.1. Traditional uses

R. canina belongs to Europe, Caucasus, central area of Asia (Pakistan, Afghanistan, Iran, Iraq, Anatolia, Syria, Lebanon, and Palestine), and the northwest of Africa, Canary Islands and Azores Islands. Like the last three species, it was naturalized in North America, as well as in Chile, and in Southern Australia. Different parts of *Rosa canina* are used and are recommended to be collected during diverse seasons. Petals (P) should be collected before the flower blooms, rose hips (RH) by the end of summer or autumn, and L during spring. RH are the most used part of *Rosa canina* and can be infused, decocted, extracted as a liquid, tinctured or consumed as syrup. Anti-diarrheic, astringent, healing, depurative, diuretic, venotonic, and vitamin properties are associated with this species. Respiratory infections, such as a cough, cold or sore throat, have been treated with RH infusions, which

can be mixed with other plants (*Origanum* spp., *Laurus nobilis*, *Ficus carica* and *Citrus limon*). RH are well-known to be a vitamin C (VC) source, so infusions are used for treating anemia and to control diarrhea. Jams using RH and P are valued for their high vitamin content. In gastronomy, P are also a decoration element in salads. L have been used as tea substitute, L and dried hip roses have been added to beverages to provide aroma, color and taste. Rose water, which has been described as a collyrium, has been widely applied in cosmetics for providing aroma or color as well as for its astringent properties. Indirect uses of this plant can be applied, one example is the honey obtained from rose P (Blanco, Macia, & Morales, 1999; Escudero, 1999; Verde, 2008)

2.6.2. Scientific studies

Numerous studies have tried to determine the mechanisms of the action of *R. canina* of the different antioxidant, anti-inflammatory and diuretic properties (Tables 3 and 4). The anti-inflammatory activities observed *in vitro* were related to the inhibition of pro-inflammatory enzymes such as cyclooxygenases, lipoxygenases, cytokines, chemokines, metalloproteases, nitric oxide synthase, and related stress factors such as NF- κ B or C-reactive protein levels (Gruenwald, Uebelhack, & Moré, 2019). Among its different constituents, PC and VC stand out, due to their interesting properties (Cosmulescu, Trandafir, & Nour, 2017; Gruenwald et al., 2019). The nutritional content of *R. canina* has been described and characterized as follows: 643.38 ± 138.4 of ascorbic acid, 28.78 ± 2.07 of carotenes, 13.28 ± 0.85 of total sugars and 0.36 ± 0.03 of protein (all the values are expressed in mg per 100 g of fresh weigh) (Rosu et al., 2011). Regarding antioxidant properties, several compounds with high antioxidant proprieties have been identified in *R. canina* extracts: PC, such as quercitrin, gallic, and protocatechuic acids, and also VC. These compounds are known for their potent antioxidant radical scavenging effects and their capacity for neutralizing reactive oxygen and nitrogen species (Gruenwald et al., 2019). EA extracts of this plant have a total phenolic content of 197 μ g GAE/mg of dry content. Methanolic extracts have shown a wide variety of results depending on the pre-treatment of the sample. Fresh biomass displayed a lower phenolic amount, 7.27 μ g GAE/mg, than when using dry biomass which concentration was estimated in 255 μ g GAE/mg (Cosmulescu et al., 2017; Ouerghemmi et al., 2016). The antioxidant properties of the more concentrated extracts, both the EA and the methanolic, were tested by using different methods. The methanolic extract showed better results for the TEAC, FRAP and DPPH assays, while the EA showed good results for the ORAC assay (Ouerghemmi et al., 2016). This activity has also been studied *in vitro*, on HepG2 and SH-SY5Y cells against intracellular reactive oxygen species (Fetni, Bertella, Ouahab, Martinez Zapater, & De Pascual-Teresa Fernandez, 2017).

(Gruenwald et al., 2019) also determined that *R. canina* has different anti-inflammatory effects: reduces the pro-inflammatory molecules and C-reactive protein levels, diminishes NF- κ B signalling, inhibits pro-inflammatory enzymes, including COX1 and 2, 5-LOX and iNOS, reduces the chemotaxis and chemoluminescence of polymorphonuclear cells, and inhibits pro-inflammatory metalloproteases. The anti-inflammatory properties of a *R. canina* ethanolic extract were demonstrated *in vivo* models. The extract was able to significantly inhibit the inflammation in the studied models: carrageenan-induced and PGE1-induced hind paw edema models, on an acetic acid-induced increase in a capillary permeability model and on a pain-model based on the inhibition of p-benzoquinone-induced in writhing mice (Deliorman Orhan, Hartevioğlu, Küpeli, & Yesilada, 2007). Besides, it has been reported that in humans, *R. canina* has the power to reduce pain and stiffness in patients with osteoarthritis (Ouerghemmi et al., 2016).

Numerous studies have reported good cytotoxic activity against HeLa, MRC-5, human colon (HT-29) and breast (MCF-7) cancer cells as well anti-proliferative effects on human colon cancer (Caco-2) cells by increasing the number of apoptotic cells and cell cycle arrest at the S phase (Nadpal et al., 2016; Turan et al., 2018). Through the inhibition

of tyrosinase, ethanolic extract of *R. canina* has cytotoxic effect on mouse melanoma. This extract also has a selective cytotoxic effect on WiDr cells compared with normal colon cells, inducing cell cycle arrest at the S phase and apoptosis via reduced matrix metalloproteinases. In addition, an inhibition in the expression of telomerase was observed in WiDR cells at 48 and 72 h of treatment (Turan et al., 2018).

(Fuji, Ikeda, & Saito, 2011) suggested that PA present in orally administered *R. canina* extract were the compounds responsible for the skin pigmentation's inhibition of tyrosinase activity on the skin of brown guinea pigs.

Finally, different works point to *R. canina* as a source of biomolecules which may improve nutritional pathologies. A study conducted on mice demonstrated the anti-obese properties of *R. canina*. Acetone extracts of FR (with a concentration of 50 mg/kg/dw) and seeds (with a concentration of 12.5 and 25 mg/kg/dw) were tested on mice. These extracts were found to have a substantial inhibitory effect on weight gain and in visceral fat gain without affecting the food intake in mice for two weeks after the administration of the extracts. The study attributed these proprieties to *trans*-tiliroside (0.1–10 mg/kg/d) present in the *R. canina* extract, and this compound also significantly reduced blood glucose levels after glucose loading (1 g/kg) in mice (Ninomiya et al., 2007). W fractions of *R. canina*, mostly containing vitamins, carotenoids, flavonoids, tannins, sachharides and pectins, were demonstrated to have hypoglycaemic activity in induced diabetic rats. The administration of these extracts triggered an increment in the number of active islets of pancreas while reducing necrotic ones islets. Another functions related with the consumption of *R. canina* were the prevention of ulcer formation and anti-diarrheal. The antioxidant effect of extracts obtained from this plant has been also displayed to reduce the peroxidation of lipids or unsaturated fatty acids which action in hepatic and renal tissues have demonstrated to attenuate their associated diseases (Ayati et al., 2018; Mármol, Sánchez-de-Diego, Jiménez-Moreno, Ancín-Azpilicueta, & Rodríguez-Yoldi, 2017).

3. Conclusion

Although modern medicine has almost replaced the traditional use of plants, they are still a source of healing. The numerous studies compiled in this review confirm the knowledge obtained from the trial and error of thousands of years: the six plants selected possess compounds with many health benefits that justify their use in traditional medicine. Understanding the knowledge that has been revealed to us through popular wisdom is important, since this information may be the key to the development of different drugs, foods and other products in the future. Thus, conducting more studies about the phytochemical composition and the mechanisms of the actions of the compounds is essential.

CRedit authorship contribution statement

P. Garcia-Oliveira: Conceptualization, Methodology, Formal analysis, Investigation, Visualization, Supervision. **M. Fraga-Corral:** Conceptualization, Methodology, Formal analysis, Investigation, Visualization, Supervision. **A.G. Pereira:** Conceptualization, Methodology, Formal analysis, Investigation, Visualization, Supervision. **C. Lourenço-Lopes:** Conceptualization, Methodology, Formal analysis, Investigation, Visualization, Supervision. **C. Jimenez-Lopez:** Conceptualization, Methodology, Formal analysis, Investigation, Visualization, Supervision. **M.A. Prieto:** Conceptualization, Methodology, Formal analysis, Investigation, Visualization, Supervision. **J. Simal-Gandara:** Conceptualization, Methodology, Formal analysis, Investigation, Visualization, Supervision.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The research leading to these results was funded by FEDER under the European Union programme Interreg V Spain-Portugal (POCTEP, ref. 0377-Iberphenol-6-E; POCTEP, ref. 0181-Nanoeaters-01-E); by European Union Bio-Based Industries (Up4Health project) granted by H2020-BBI-JTI-2019; by MICINN supporting the Ramón&Cajal grant for Miguel Ángel Prieto Lage (RYC-2017-22891); by the Xunta de Galicia and University of Vigo supporting the post-doctoral grant of María Fraga Corral (ED481B-2019/096), the pre-doctoral grants for Antia González Pereira (ED481A-2019/0228) and Paula García Oliveira (ED481A-2019/295); and by Axudas Conecta Peme (Xunta de Galicia) supporting the IN852A 2018/58 NeuroFood Project.

References

- Aburjai, T., & Natsheh, F. M. (2003). Plants Used in Cosmetics. *Phytotherapy Research*, 17(9), 987–1000. <https://doi.org/10.1002/ptr.1363>.
- Ad'hiah, A. H., Al-Bederi, O. N. H., & Al-Sammarae, K. W. (2013). Cytotoxic effect of *Agrimonia eupatoria* L. against cancer cell lines *in vitro*. *Journal of the Association of Arab Universities for Basic and Applied Sciences*, 14(1), 87–92. <https://doi.org/10.1016/j.jaubas.2013.01.003>.
- Akram, M., Tahir, I. M., Shah, S. M. A., Mahmood, Z., Altaf, A., Ahmad, K., ... Mehboob, H. (2018). Antiviral potential of medicinal plants against HIV, HSV, influenza, hepatitis, and coxsackievirus: A systematic review. *Phytotherapy Research*, 32, 811–822. <https://doi.org/10.1002/ptr.6024>.
- Al-snafi, A. E. (2015). The Pharmacological and therapeutic importance of *Agrimonia eupatoria* - A review. *Asian Journal of Pharmaceutical Science & Technology*, 5(2), 112–117.
- Al-Snafi, A. E. (2019). Constituents and pharmacology of *Geum urbanum*-A review. Retrieved from www.iosrphr.org. *IOSR Journal of Pharmacy*, 9(5), 28–33.
- Arslan, R., Bektas, N., Bor, Z., & Sener, E. (2015). Evaluation of the antithrombotic effects of *Crataegus monogyna* and *Crataegus davisi* in the carrageenan-induced tail thrombosis model. *Pharmaceutical Biology*, 53(2), 275–279. <https://doi.org/10.3109/13880209.2014.914957>.
- Asioli, D., Aschemann-Witzel, J., Caputo, V., Vecchio, R., Annunziata, A., Næs, T., & Varela, P. (2017). Making sense of the “clean label” trends: A review of consumer food choice behavior and discussion of industry implications. *Food Research International*, 99, 58–71. <https://doi.org/10.1016/j.foodres.2017.07.022>.
- Ayati, Z., Amiri, M. S., Ramezani, M., Delshad, E., Sahebkar, A., & Emami, S. A. (2018). Phytochemistry, traditional uses and pharmacological profile of rose hip: A review. *Current Pharmaceutical Design*, 24(35), 4101–4124.
- Bae, H., Kim, H.-J., Shin, M., Lee, H., Yin, C. S., Ra, J., & Kim, J. (2010). Inhibitory effect of *Agrimoniae Herba* on lipopolysaccharide-induced nitric oxide and proinflammatory cytokine production in BV2 microglial cells. *Neurological Research*, 32(sup1), 53–57. <https://doi.org/10.1179/016164109x12537002794002>.
- Barreira, J. C. M., Rodrigues, S., Carvalho, A. M., & Ferreira, I. C. F. R. (2013). Development of hydrosoluble gels with *Crataegus monogyna* extracts for topical application: Evaluation of antioxidant activity of the final formulations. *Industrial Crops and Products*, 42(1), 175–180. <https://doi.org/10.1016/j.indcrop.2012.05.034>.
- Barros, L., Carvalho, A. M., & Ferreira, I. C. F. R. (2011). Comparing the composition and bioactivity of *Crataegus monogyna* flowers and fruits used in folk medicine. *Phytochemical Analysis*, 22(2), 181–188. <https://doi.org/10.1002/pca.1267>.
- Benarba, B., & Pandiella, A. (2018). Colorectal cancer and medicinal plants: Principle findings from recent studies. *Biomedicine and Pharmacotherapy*, 107, 408–423. <https://doi.org/10.1016/j.biopha.2018.08.006>.
- Bespalov, V. G., Alexandrov, V. A., Semenov, A. L., Vysochina, G. I., Kostikova, V. A., & Baranenko, D. A. (2018). The inhibitory effect of *Filipendula ulmaria* (L.) Maxim. on colorectal carcinogenesis induced in rats by methylnitrosourea. *Journal of Ethnopharmacology*, 227, 1–7. <https://doi.org/10.1016/j.jep.2018.08.013>.
- Blanco, E., Macia, M. J., & Morales, R. (1999). Medicinal and veterinary plants of El Caurel (Galicia, northwest Spain). *Journal of Ethnopharmacology*, 65, 113–124.
- Borneo, R., León, A. E., Aguirre, A., Ribotta, P., & Cantero, J. J. (2009). Antioxidant capacity of medicinal plants from the Province of Córdoba (Argentina) and their *in vitro* testing in a model food system. *Food Chemistry*, 112(3), 664–670. <https://doi.org/10.1016/j.foodchem.2008.06.027>.
- Cardoso, R. V. C., Fernandes, Á., González-Paramás, A. M., Barros, L., & Ferreira, I. C. F. R. (2019). Flour fortification for nutritional and health improvement: A review. *Food Research International*, 125(May), 108576. <https://doi.org/10.1016/j.foodres.2019.108576>.
- Castroviejo, S. (1986). *Flora iberica 1–8, 10–15, 17–18, 21*. CSIC, Madrid: Real Jardín Botánico 2012.
- Corbo, M. R., Bevilacqua, A., Petrucci, L., Casanova, F. P., & Sinigaglia, M. (2014). Functional Beverages: The Emerging Side of Functional Foods. *Comprehensive Reviews in Food Science and Food Safety*, 13(6), 1192–1206. <https://doi.org/10.1111/1541-4337.12109>.
- Cosmulescu, S., Trandafir, I., & Nour, V. (2017). Phenolic acids and flavonoids profiles of extracts from edible wild fruits and their antioxidant properties. *International Journal of Food Properties*, 20(12), 3124–3134. <https://doi.org/10.1080/10942912.2016.1274906>.
- Day, L. (2013). Proteins from land plants – Potential resources for human nutrition and food security. *Trends in Food Science & Technology*, 32(1), 25–42. <https://doi.org/10.1016/j.tifs.2013.05.005>.
- de Santayana, M. P., Morales, R., Tardío, J., Aceituno, L., & Molina, M. (2014). *Inventario Español de los Conocimientos Tradicionales relativos a la Biodiversidad*. <https://doi.org/10.1017/CBO9781107415324.004>.
- de Santayana, M. P., Morales, R., Tardío, J., Aceituno, L., & Molina, M. (2018). *Inventario español de los conocimientos tradicionales relativos a la biodiversidad: Segunda fase (Tomo II): Fichas. Alimentación y Medio Ambiente: Ministerio de Agricultura*.
- Deliorman Orhan, D., Hartevioğlu, A., Küpeli, E., & Yesilada, E. (2007). *In vivo* anti-inflammatory and antinociceptive activity of the crude extract and fractions from *Rosa canina* L. fruits. *Journal of Ethnopharmacology*, 112(2), 394–400. <https://doi.org/10.1016/j.jep.2007.03.029>.
- Dimitrova, L., Zaharieva, M. M., Popova, M., Kostadinova, N., Tsvetkova, I., Bankova, V., & Najdenski, H. (2017). Antimicrobial and antioxidant potential of different solvent extracts of the medicinal plant *Geum urbanum* L. *Chemistry Central Journal*, 11(1), 1–11. <https://doi.org/10.1186/s13065-017-0343-8>.
- El Haouari, M., & Rosado, J. A. (2016). Medicinal Plants with Antiplatelet Activity. *Phytotherapy Research*, 30(12), 1059–1071. <https://doi.org/10.1002/ptr.5619>.
- EMA/HMPC/159076/2014. (2016). Assessment report on *Crataegus* spp., folium cum flore.
- EMA/HMPC/434892/2010. (2013). Assessment report on *Filipendula ulmaria* (L.) Maxim., herba and *Filipendula ulmaria* (L.) Maxim., flos.
- EMA/HMPC/680595/2013. (2015). Assessment report on *Agrimonia eupatoria* L., herba.
- Escudero, F. A. M. (1999). Las plantas de extractos. Bases Para Un Plan de Desarrollo Del Sector [Medicinal and Aromatic Plants. Bases for a Development Plan of the Sector]. Mundi-Prensa, Madrid.
- Febrer, J. L. F., Blanquer, G., & Pí, A. S. (2001). Inventario de las plantas medicinales de uso popular en la ciudad de Valencia. *Mediana y Ciencias Sociales*, 13.
- Fetni, S., Bertella, N., Ouahab, A., Martínez Zapater, J. M., & De Pascual-Teresa Fernandez, S. (2017). Composition and biological activity of the Algerian plant *Rosa canina* L. by HPLC-UV-MS. *Arabian Journal of Chemistry*. <https://doi.org/10.1016/j.arabjc.2017.09.013>.
- Flood, J. (2010). The importance of plant health to food security. *Food Security*, 2(3), 215–231. <https://doi.org/10.1007/s12571-010-0072-5>.
- Font Quer, P. (2001). Plantas medicinales. El Dioscórides renovado (Third edit). Barcelona (Spain).
- Franco, R. R., Guitián, M. R., & Resúa, Á. (2013). Plantas utilizadas en medicina humana y veterinaria en el municipio de Triacastela, Lugo (NW España). *Recursos Rurais: Revista Oficial Do Instituto de Biodiversidade Agraria e Desenvolvimento Rural (IBADER)*, 35–43.
- Fujii, T., Ikeda, K., & Saito, M. (2011). Inhibitory effect of rose hip (*Rosa canina* L.) on melanogenesis in mouse melanoma cells and on pigmentation in brown guinea pigs. *Bioscience, Biotechnology and Biochemistry*, 75(3), 489–495. <https://doi.org/10.1271/bbb.100702>.
- Ganesan, P., & Choi, D. K. (2016). Current application of phytochemical-based nanocosmetics for beauty and skin therapy. *International Journal of Nanomedicine*, 11, 1987–2007. <https://doi.org/10.2147/IJN.S104701>.
- Garmendia, F. M., & Castroviejo, S. (1998). *Crataegus L. Flora Iberica*, 6, 404–414.
- González-Hernández, M. P., Romero, R., Rodríguez-Guitián, M., & Rigueiro, A. (2004). Medicinal use of some plants in Galicia (NW Spain). *Acta Horticulturae*, 629, 63–75.
- Granica, S., Kłębowska, A., Kosiński, M., Piwowarski, J. P., Dudek, M. K., Kazmierski, S., & Kiss, A. K. (2016). Effects of *Geum urbanum* L. root extracts and its constituents on polymorphonuclear leucocytes functions. Significance in periodontal diseases. *Journal of Ethnopharmacology*, 188, 1–12. <https://doi.org/10.1016/j.jep.2016.04.030>.
- Gruenewald, J., Uebelhack, R., & Moré, M. I. (2019). *Rosa canina* – Rose hip pharmacological ingredients and molecular mechanics counteracting osteoarthritis – A systematic review. *Phytotherapy Research*, 33(12), 152958. <https://doi.org/10.1016/j.phymed.2019.152958>.
- Guillén, A., & Rico, E. (1998). *Potentilla L. Flora Iberica*, 6, 96–140.
- Gulati, O. P., Ottaway, P. B., Jennings, S., Coppens, P., & Gulati, N. (2019). Botanical nutraceuticals (food supplements and fortified and functional foods) and novel foods in the EU, with a main focus on legislative controls on safety aspects. In *Nutraceutical and Functional Food Regulations in the United States and around the World* (Third Edit, pp. 277–321). Doi: 10.1016/b978-0-12-816467-9.00020-4.
- Hoffmann, J., Casetti, F., Bullerker, U., Haarhaus, B., Vagedes, J., Schemp, C. M., & Wölfl, U. (2016). Anti-inflammatory effects of agrimoniin-enriched fractions of *Potentilla erecta*. *Molecules*, 21(6), <https://doi.org/10.3390/molecules21060792>.
- Ivanova, D., Vankova, D., & Nashar, M. (2013). *Agrimonia eupatoria* tea consumption in relation to markers of inflammation, oxidative status and lipid metabolism in healthy subjects. *Archives of Physiology and Biochemistry*, 119(1), 32–37. <https://doi.org/10.3109/13813455.2012.729844>.
- Jaberian, H., Piri, K., & Nazari, J. (2013). Phytochemical composition and *in vitro* antimicrobial and antioxidant activities of some medicinal plants. *Food Chemistry*, 136(1), 237–244. <https://doi.org/10.1016/j.foodchem.2012.07.084>.
- Jarzycka, A., Lewińska, A., Ganczar, R., & Wilk, K. A. (2013). Assessment of extracts of *Helichrysum arenarium*, *Crataegus monogyna*, *Sambucus nigra* in photoprotective UVA and UVB; Photostability in cosmetic emulsions. *Journal of Photochemistry and*

- Photobiology B: Biology, 128, 50–57. <https://doi.org/10.1016/j.jphotobiol.2013.07.029>.
- Katanić, J., Boroja, T., Mihailović, V., Nikles, S., Pan, S. P., Rosić, G., ... Bauer, R. (2016). In vitro and in vivo assessment of meadowsweet (*Filipendula ulmaria*) as anti-inflammatory agent. *Journal of Ethnopharmacology*, 193, 627–636. <https://doi.org/10.1016/j.jep.2016.10.015>.
- Katanić, J., Matić, S., Pferschy-Wenzig, E. M., Kretschmer, N., Boroja, T., Mihailović, V., ... Bauer, R. (2017). *Filipendula ulmaria* extracts attenuate cisplatin-induced liver and kidney oxidative stress in rats: In vivo investigation and LC-MS analysis. *Food and Chemical Toxicology*, 99, 86–102. <https://doi.org/10.1016/j.fct.2016.11.018>.
- Laila, U., Akram, M., Shariati, M. A., Hashmi, A. M., Akhtar, N., Tahir, I. M., ... Ahmad, S. (2019). Role of medicinal plants in HIV/AIDS therapy. *Clinical and Experimental Pharmacology and Physiology*, (May), 1–11. Doi: 10.1111/1440-1681.13151.
- Lee, K. Y., Hwang, L., Jeong, E. J., Kim, S. H., Kim, Y. C., & Sung, S. H. (2010). Effect of Neuroprotective Flavonoids of *Agrimonia eupatoria* on Glutamate-Induced Oxidative Injury to HT22 Hippocampal Cells. *Bioscience, Biotechnology and Biochemistry*, 74(8), 1704–1706. <https://doi.org/10.1271/bbb.100200>.
- Lenhain, B. (2014). *The Practical Herbal Medicine Handbook*. Berkeley (California): Althea Press.
- Lima, M. J., Sousa, D., Lima, R. T., Carvalho, A. M., Ferreira, I. C. F. R., & Vasconcelos, M. H. (2014). Flower extracts of *Filipendula ulmaria* (L.) Maxim inhibit the proliferation of the NCI-H460 tumour cell line. *Industrial Crops and Products*, 59, 149–153. <https://doi.org/10.1016/j.indcrop.2014.05.009>.
- Lobbens, E. S., Breydo, L., Skamris, T., Vestergaard, B., Jäger, A. K., Jorgensen, L., ... van de Weert, M. (2016). Mechanistic study of the inhibitory activity of *Geum urbanum* extract against α -Synuclein fibrillation. *Biochimica et Biophysica Acta - Proteins and Proteomics*, 1864(9), 1160–1169. <https://doi.org/10.1016/j.bbapap.2016.06.009>.
- López, C. J., Caleja, C., Prieto, M. A., Barreiro, M. F., Barros, L., & Ferreira, I. C. F. R. (2018). Optimization and comparison of heat and ultrasound assisted extraction techniques to obtain anthocyanin compounds from *Arbutus unedo* L Fruits. *Food Chemistry*. <https://doi.org/10.1016/j.foodchem.2018.04.103>.
- Lubbe, A., & Verpoorte, R. (2011). Cultivation of medicinal and aromatic plants for specialty industrial materials. *Industrial Crops and Products*, 34(1), 785–801. <https://doi.org/10.1016/j.indcrop.2011.01.019>.
- Luís, A., Domingues, F., & Duarte, A. P. (2011). Bioactive Compounds, RP-HPLC Analysis of Phenolics, and Antioxidant Activity of Some Portuguese Shrub Species Extracts. *Natural Product Communications*, 6(12), 1934578X1100601219. Doi: 10.1177/1934578X1100601219.
- Mantle, D., Eddeb, F., & Pickering, A. T. (2000). Comparison of relative antioxidant activities of British medicinal plant species in vitro. *Journal of Ethnopharmacology*, 72(1–2), 47–51. [https://doi.org/10.1016/S0378-8741\(00\)00199-9](https://doi.org/10.1016/S0378-8741(00)00199-9).
- Mari, A., Eletto, D., Pizza, C., Montoro, P., & Piacente, S. (2013). Integrated mass spectrometry approach to profile proanthocyanidins occurring in food supplements: Analysis of *Potentilla erecta* L. rhizomes. *Food Chemistry*, 141(4), 4171–4178. <https://doi.org/10.1016/j.foodchem.2013.06.101>.
- Mármol, I., Sánchez-de-Diego, C., Jiménez-Moreno, N., Ancín-Azpilicueta, C., & Rodríguez-Yoldi, M. J. (2017). Therapeutic Applications of Rose Hips from Different Rosa Species. *International Journal of Molecular Sciences*, 18(6), <https://doi.org/10.3390/ijms18061137>.
- Martín Ortega, A. M., & Segura Campos, M. R. (2019). Medicinal Plants and Their Bioactive Metabolites in Cancer Prevention and Treatment. In *Bioactive Compounds* (pp. 85–109). Doi: 10.1016/b978-0-12-814774-0.00005-0.
- Martins, J., & Brijesh, S. (2018). Phytochemistry and pharmacology of anti-depressant medicinal plants: A review. *Biomedicine and Pharmacotherapy*, 104(May), 343–365. <https://doi.org/10.1016/j.biopha.2018.05.044>.
- Ministerio De Agricultura Alimentación y Medio Ambiente. (2019). *Anthos*. Sistema de información de las plantas de España.
- Mosquera Paans, M. (2013). *A nosa botica. Plantas medicinais* (First; F. Villegas Belmonte, Ed.). Pontevedra: Edicións do Cumio.
- Muruzović, M. Ž., Mladenović, K. G., Stefanović, O. D., Vasić, S. M., & Čović, L. R. (2016). Extracts of *Agrimonia eupatoria* as sources of biologically active compounds and evaluation of their antioxidant, antimicrobial, and antibiologic activities. *Journal of Food and Drug Analysis*, 24, 539–547.
- Nabavi, S. F., Habtemariam, S., Ahmed, T., Sureda, A., Daglia, M., Sobarzo-Sánchez, E., & Nabavi, S. M. (2015). Polyphenolic composition of *Crataegus monogyna* Jacq.: From chemistry to medical applications. *Nutrients*, 7(9), 7708–7728. <https://doi.org/10.3390/nu7095361>.
- Nadpal, J. D., Lesjak, M. M., Šibul, F. S., Anačkov, G. T., Četojević-Simin, D. D., Mimica-Dukić, N. M., & Beara, I. N. (2016). Comparative study of biological activities and phytochemical composition of two rose hips and their preserves: *Rosa canina* L. and *Rosa arvensis* Huds. *Food Chemistry*, 192, 907–914. <https://doi.org/10.1016/j.foodchem.2015.07.089>.
- Neagu, E., Paun, G., Albu, C., & Radu, G. L. (2015). Assessment of acetylcholinesterase and tyrosinase inhibitory and antioxidant activity of *Alchemilla vulgaris* and *Filipendula ulmaria* extracts. *Journal of the Taiwan Institute of Chemical Engineers*, 52, 1–6. <https://doi.org/10.1016/j.jtice.2015.01.026>.
- Ninomiya, K., Matsuda, H., Kubo, M., Morikawa, T., Nishida, N., & Yoshikawa, M. (2007). Potent anti-obese principle from *Rosa canina*: Structural requirements and mode of action of trans-tiliroside. *Bioorganic and Medicinal Chemistry Letters*, 17(11), 3059–3064. <https://doi.org/10.1016/j.bmcl.2007.03.051>.
- Nitta, Y., Kikuzaki, H., Azuma, T., Ye, Y., Sakae, M., Higuchi, Y., ... Ueno, H. (2013). Inhibitory activity of *Filipendula ulmaria* constituents on recombinant human histidine decarboxylase. *Food Chemistry*, 138(2–3), 1551–1556. <https://doi.org/10.1016/j.foodchem.2012.10.074>.
- Olennikov, D. N., Kashchenko, N. I., & Chirikova, N. K. (2017). Meadowsweet teas as new functional beverages: Comparative analysis of nutrients, phytochemicals and biological effects of four *Filipendula* species. *Molecules*, 22(1), 1–23. <https://doi.org/10.3390/molecules22010016>.
- Ortega-Ramirez, L. A., Rodriguez-Garcia, I., Leyva, J. M., Cruz-Valenzuela, M. R., Silva-Espinoza, B. A., Gonzalez-Aguilar, G. A., ... Ayala-Zavala, J. F. (2014). Potential of medicinal plants as antimicrobial and antioxidant agents in food industry: A hypothesis. *Journal of Food Science*, 79(2), <https://doi.org/10.1111/1750-3841.12341>.
- Ouerghemmi, S., Sebei, H., Siracusa, L., Ruberto, G., Saija, A., Cimino, F., & Cristani, M. (2016). Comparative study of phenolic composition and antioxidant activity of leaf extracts from three wild *Rosa* species grown in different Tunisia regions: *Rosa canina* L., *Rosa moschata* Herrm. and *Rosa sempervirens* L. *Industrial Crops and Products*, 94, 167–177. <https://doi.org/10.1016/j.indcrop.2016.08.019>.
- Owczarek, A., Gudej, J., & Olszewska, M. A. (2015). Antioxidant activity of *Geum rivale* L. and *Geum urbanum* L. *Acta Poloniae Pharmaceutica - Drug Research*, 72(6), 1239–1244.
- Owczarek, A., Olszewska, M. A., & Gudej, J. (2014). Quantitative determination of ellagic acid and gallic acid in *Geum rivale* L. and *G. urbanum* L. *Acta Biologica Cracoviensia Series Botanica*, 56(2), 74–78. <https://doi.org/10.2478/abscb-2014-0021>.
- Oyenihi, A. B., & Smith, C. (2019). Are polyphenol antioxidants at the root of medicinal plant anti-cancer success? *Journal of Ethnopharmacology*, 229(May), 54–72. <https://doi.org/10.1016/j.jep.2018.09.037>.
- Pahlow, M. (1994). *El gran libro de las plantas medicinales* (8th Editio; E. Everest, Ed.).
- Pandey, A., & Agnihotri, V. (2015). Antimicrobials from medicinal plants: Research initiatives, challenges, and the future prospects. In V. K. Gupta & M. G. Tuohy (Eds.), *Biotechnology of Bioactive Compounds: Sources and Applications* (pp. 123–150). <https://doi.org/10.1002/9781118733103.ch5>.
- Pawlaczky-Graja, I. (2018). Polyphenolic-polysaccharide conjugates from flowers and fruits of single-seeded hawthorn (*Crataegus monogyna* Jacq.): Chemical profiles and mechanisms of anticoagulant activity. *International Journal of Biological Macromolecules*, 116, 869–879. <https://doi.org/10.1016/j.ijbiomac.2018.05.101>.
- Pérez, I. C. (1998). El uso de las plantas medicinales. *Medicina Clínica*, 110(10), 399.
- Pinela, J., Prieto, M. A., Carvalho, A. M., Barreiro, M. F., Oliveira, M. B. P. P., Barros, L., & Ferreira, I. C. F. R. (2016). Microwave-assisted extraction of phenolic acids and flavonoids and production of antioxidant ingredients from tomato: A nutraceutical-oriented optimization study. *Separation and Purification Technology*, 164, 114–124. <https://doi.org/10.1016/j.seppur.2016.03.030>.
- Reverter, M., Tapissier-Bontemps, N., Sasal, P., & Saulnier, D. (2017). Use of Medicinal Plants in Aquaculture. In B. Austin & A. Newaj-Fyzul (Eds.), *Diagnosis and Control of Diseases of Fish and Shellfish* (First edit, pp. 223–261). Doi: 10.1002/9781119152125.
- Rivera, D., & Obón, C. (1998). *Guía de teoría y prácticas de Etnobotánica*. Colección Textos Docentes. Murcia.
- Rodrigues, S., Calhella, R. C., Barreira, J. C. M., Dueñas, M., Carvalho, A. M., Abreu, R. M. V., ... Ferreira, I. C. F. R. (2012). *Crataegus monogyna* buds and fruits phenolic extracts: Growth inhibitory activity on human tumor cell lines and chemical characterization by HPLC-DAD-ESI/MS. *Food Research International*, 49(1), 516–523. <https://doi.org/10.1016/j.foodres.2012.07.046>.
- Rodríguez González, A. (2008). *Plantas medicinais de Galiza* (A Nosa ter). Vigo.
- Roleira, F. M. F., Tavares-Da-Silva, E. J., Varela, C. L., Costa, S. C., Silva, T., Garrido, J., & Borges, F. (2015). Plant derived and dietary phenolic antioxidants: Anticancer properties. *Food Chemistry*, 183, 235–258. <https://doi.org/10.1016/j.foodchem.2015.03.039>.
- Romero, M. (2008). *Catálogo da flora de Galicia. Monografías Do Ibadar*, (Instituto de Biodiversidade Agraria e Desenvolvimento Rural, Da USC), 1, 177.
- Romero Martín, T. (1998). *Agrimonia L. Flora Iberica*, 6, 369–374.
- Roriz, C. L., Barros, L., Prieto, M. A., Morales, P., & Ferreira, I. C. F. R. (2017). Floral parts of *Gomphrena globosa* L. as a novel alternative source of betacyanins: Optimization of the extraction using response surface methodology. *Food Chemistry*, 229, 223–234.
- Rosu, C. M., Manzu, C., Olteanu, Z., Oprica, L., Oprea, A., Giornea, E., & Zamfirache, M. M. (2011). Several fruit characteristics of *Rosa* sp. genotypes from the Northeastern region of Romania. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 39(2), 203–208.
- Samaniego-Vaesken, M. L., Alonso-Aperte, E., & Varela-Moreiras, G. (2016). Voluntary fortification with folic acid in Spain: An updated food composition database. *Food Chemistry*, 193, 148–153. <https://doi.org/10.1016/j.foodchem.2014.06.046>.
- Samardžić, S., Arsenijević, J., Božić, D., Milenković, M., Tešević, V., & Maksimović, Z. (2018). Antioxidant, anti-inflammatory and gastroprotective activity of *Filipendula ulmaria* (L.) Maxim. and *Filipendula vulgaris* Moench. *Journal of Ethnopharmacology*, 213, 132–137. <https://doi.org/10.1016/j.jep.2017.11.013>.
- Sengupta, G., Gaurav, A., & Tiwari, S. (2018). Substituting medicinal plants through drug synthesis. *Synthesis of Medicinal Agents from Plants*. <https://doi.org/10.1016/b978-0-08-102071-5.00003-9>.
- Shalazar Jalali, A., & Hasanzadeh, S. (2013). *Crataegus monogyna* fruit aqueous extract as a protective agent against doxorubicin-induced reproductive toxicity in male rats. *Avicenna Journal of Phytomedicine*, 3(2), 159–15970.
- Shortle, E., Kerry, J., Furey, A., & Gilroy, D. (2013). Optimisation of process variables for antioxidant components from *Crataegus monogyna* by supercritical fluid extraction (CO₂) using Box-Behnken experimental design. *Journal of Supercritical Fluids*, 81, 112–118. <https://doi.org/10.1016/j.supflu.2013.05.007>.
- Singh, B., Singh, S., Gulati, M., Singh, H., Arora, R., & Arora, S. (2019). Herbal Products in Hypertension: Paradox or Paragon. In *Herbal Medicine: Back to the Future* (pp. 76–124). Doi: 10.2174/9789811403743119020004.
- Soutullo, E. C., Muñoz, C. F., Pazo, R. M., Alonso, R. M., & Boente, M. J. R. (2015). Hierbas medicinales: uso en la cultura gallega. Revbigio.
- Srivastava, A. K. (2018). Significance of medicinal plants in human life. In *Synthesis of Medicinal Agents from Plants* (pp. 1–24). Doi: 10.1016/B978-0-08-102071-5/00001-5.
- Stanković, M. S., Radić, Z. S., Blanco-Salas, J., Vázquez-Pardo, F. M., & Ruiz-Téllez, T. (2017). Screening of selected species from Spanish flora as a source of bioactive

- substances. *Industrial Crops and Products*, 95, 493–501. <https://doi.org/10.1016/j.indcrop.2016.09.070>.
- Synowiec, A., Gniewosz, M., Bączek, K., & Przybył, J. L. (2014). Antimicrobial effect of an aqueous extract of *Potentilla erecta* rhizome. *Herba Polonica*, 60(2), 18–28. <https://doi.org/10.2478/hepo-2014-0007>.
- Tobyn, G., Denham, A., & Whitelegg, M. (2011). *Potentilla erecta*, tormentil. Doi: 10.1016/B978-0-443-10344-5.00029-X.
- Tolve, R., Condelli, N., Caruso, M. C., Genovese, F., Di Renzo, G. C., Mauriello, G., & Galgano, F. (2019). Preparation and characterization of microencapsulated phytochemicals for the formulation of functional foods: Scale up from laboratory to semi-technical production. *Food Research International*, 116(2018), 1274–1281.
- Tomczyk, M., & Latté, K. P. (2009). *Potentilla*-A review of its phytochemical and pharmacological profile. *Journal of Ethnopharmacology*, 122(2), 184–204. <https://doi.org/10.1016/j.jep.2008.12.022>.
- Tsirigotis-Maniecka, M., Pawlaczyk-Graja, I., Ziewiecki, R., Balicki, S., Matulová, M., Capek, P., ... Gancarz, R. (2019). The polyphenolic-polysaccharide complex of *Agrimonia eupatoria* L. as an indirect thrombin inhibitor- isolation and chemical characterization. *International Journal of Biological Macromolecules*, 125, 124–132.
- Turan, I., Demir, S., Kilinc, K., Yaman, S. O., Misir, S., Kara, H., ... Deger, O. (2018). Cytotoxic effect of *Rosa canina* extract on human colon cancer cells through repression of telomerase expression. *Journal of Pharmaceutical Analysis*, 8(6), 394–399. <https://doi.org/10.1016/j.jpha.2017.12.005>.
- Van Hai, N. (2015). The use of medicinal plants as immunostimulants in aquaculture: A review. *Aquaculture*, 446, 88–96. <https://doi.org/10.1016/j.aquaculture.2015.03.014>.
- Verde, A. (2008). Guía de las plantas medicinales de Castilla-La Mancha.
- Verde, A., Rivera, D., Fajardo, J., Obón, C., & Cebrián, F. (2008). Guía de las plantas medicinales de Castilla La Mancha (y otros recursos de uso tradicional) (E. Altabán, Ed.).
- Vieira, V., Prieto, M. A., Barros, L., Coutinho, J. A. P., Ferreira, O., & Ferreira, I. C. F. R. (2017). Optimization and comparison of maceration and microwave extraction systems for the production of phenolic compounds from *Juglans regia* L. for the valorization of walnut leaves. *Industrial Crops and Products*, 107, 341–352.
- Wölfle, U., Hoffmann, J., Haarhaus, B., Rao Mittapalli, V., & Schempp, C. M. (2017). Anti-inflammatory and vasoconstrictive properties of *Potentilla erecta* – A traditional medicinal plant from the northern hemisphere. *Journal of Ethnopharmacology*, 204(April), 86–94. <https://doi.org/10.1016/j.jep.2017.03.058>.
- Yoon, S. J., Koh, E. J., Kim, C. S., Zee, O. P., Kwak, J. H., Jeong, W. J., ... Lee, S. M. (2012). *Agrimonia eupatoria* protects against chronic ethanol-induced liver injury in rats. *Food and Chemical Toxicology*, 50(7), 2335–2341. <https://doi.org/10.1016/j.fct.2012.04.005>.