

Title	Interface Vanishing for Solutions to Maxwell and Stokes Systems (Nonlinear Wave Phenomena and Applications)
Author(s)	Kobayashi, Takayuki; Suzuki, Takashi; Watanabe, Kazuo
Citation	数理解析研究所講究録 (2004), 1368: 65-71
Issue Date	2004-04
URL	http://hdl.handle.net/2433/25408
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Interface Vanishing for Solutions to Maxwell and Stokes Systems

Takayuki KOBAYASHI,* Takashi SUZUKI,** Kazuo WATANABE***
 (小林 孝行) (鈴木 貴) (渡邊 一雄)

*Department of Mathematics
 Faculty of Science and Engineering, Saga University
 **Division of Mathematical Science
 Graduate School of Engineering Science, Osaka University
 ***Department of Mathematics
 Gakushuin University

In the previous work [1], we studied interface regularity of three dimensional Maxwell system when the interface is C^2 , and that of Stokes system when it is flat. In this article, we continue the study and show refined interface vanishing theorems for general interface.

Geometric situation which we are concerned in is described as follows. Namely, $\Omega \subset \mathbf{R}^3$ denotes a bounded domain with Lipschitz boundary $\partial\Omega$, and $\mathcal{M} \subset \mathbf{R}^3$ is a Lipschitz hypersurface cutting Ω transversally. Thus, it holds that

$$\begin{aligned} \mathcal{M} \cap \Omega &\neq \phi \\ \Omega &= \Omega_+ \cup (\Omega \cap \mathcal{M}) \cup \Omega_- \quad (\text{disjoint union}) \end{aligned} \tag{0.1}$$

with the open subsets Ω_{\pm} of Ω . First, we consider the Maxwell system in magnetostatics,

$$\left. \begin{aligned} \nabla \times B &= J \\ \nabla \cdot B &= 0 \end{aligned} \right\} \quad \text{in } \Omega_{\pm}, \tag{0.2}$$

where $B = {}^t(B^1(x), B^2(x), B^3(x))$ and $J = {}^t(J^1(x), J^2(x), J^3(x))$ stand for three dimensional vector fields, indicating magnetic field and total current

density, respectively. Furthermore, $\nabla = {}^t(\partial_1, \partial_2, \partial_3)$ denotes the gradient operator and \times and \cdot are outer and inner products in \mathbf{R}^3 , so that $\nabla \times$ and $\nabla \cdot$ are the operations of rotation and divergence, respectively.

In the context of magnetoencephalography, Suzuki, Watanabe, and Shimogawara [3] introduced an interface vanishing theorem when the interface is given by the boundary ∂D of a smooth bounded domain $D \subset \mathbf{R}^3$ in use of the layer potential. More precisely, if J is continuous on $\overline{\Omega_{\pm}}$ and system (0.2) has a solution $B \in C(\mathbf{R}^3)^3 \cap C^1(\mathbf{R}^3 \setminus \partial D)^3$ for $\Omega_- = D$ and $\Omega_+ = \mathbf{R}^3 \setminus D$, then

$$[\nabla(n \cdot B)]_{\pm}^{\pm} = 0 \quad \text{on } \partial D$$

follows, regardless with the continuity of J across ∂D . Here, n denotes the outer unit normal vector to ∂D , $[A]_{\pm}^{\pm} = A_+ - A_-$, and

$$A_+(\xi) = \lim_{x \rightarrow \xi, x \in \mathbf{R}^3 \setminus D} A(x), \quad A_-(\xi) = \lim_{x \rightarrow \xi, x \in D} A(x)$$

for $\xi \in \partial D$. Then, Kobayashi, Suzuki and Watanabe [1] studied local version, the case where the bounded domain Ω is given with the interface $\mathcal{M} \cap \Omega$ as in (0.1), and showed that even if $n \times J$ has an interface on $\mathcal{M} \cap \Omega$, the normal component $n \cdot B$ of B gains the regularity in one rank. In this article, we refine the argument and reduce smoothness of the interface. This refinement is very useful to study similar problems for the Stokes system as will be described later.

To state our result for (0.2), we take some preliminaries on function spaces from Girault and Raviart [2]. Namely, if $D \subset \mathbf{R}^3$ is a bounded domain with Lipschitz boundary ∂D and n denotes the unit normal vector to ∂D , then for $p \in [1, \infty]$, $L^p(D)$ denotes the standard L^p space on D provided with the norm $\|\cdot\|_{L^p(D)}$, and the Sobolev space $W^{m,p}(D)$ is defined by

$$W^{m,p}(D) = \{u \in L^p(D) \mid \partial^{\alpha} u \in L^p(D) \quad \text{for } |\alpha| \leq m\}$$

for a positive integer m , where $\partial^{\alpha} = \partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} \partial_{x_3}^{\alpha_3}$ for the multi-index $\alpha = (\alpha_1, \alpha_2, \alpha_3)$. Put $H^m(D) = W^{m,2}(D)$. Given $\sigma \in (0, 1)$, we say that $u \in H^{m+\sigma}(D)$ if $u \in H^m(D)$ and

$$\int_D \int_D \frac{|\partial^{\alpha} u(x) - \partial^{\alpha} u(y)|^2}{|x - y|^{n+2\sigma}} dx dy < +\infty$$

for any α in $|\alpha| = m$ and $n = 3$. The space $H^s(\Gamma)$ is defined similarly with $n = 2$ through the local chart of Γ , where $s \in [0, 1]$ and $\Gamma \subset \partial D$ is a relatively

open connected set. Then, we set $H^{-s}(\Gamma) = H_0^s(\Gamma)'$, where $H_0^s(\Gamma)$ denotes the closure in $H^s(\Gamma)$ of the space composed of Lipschitz continuous functions on Γ with compact supports. Thus, we have $H_0^s(\Gamma) = H^s(\Gamma)$ if $\Gamma \subset \partial D$ is a closed surface, and in particular, it holds that $H^{1/2}(\partial D) = H_0^{1/2}(\partial D)$. In this context, let us remember the standard trace theorem that $H^1(D)|_{\partial D} \cong H^{1/2}(\partial D)$. We also put

$$H(\operatorname{div}, D) = \{u \in L^2(D)^3 \mid \nabla \cdot u \in L^2(D)\}$$

and

$$H(\operatorname{rot}, D) = \{u \in L^2(D)^3 \mid \nabla \times u \in L^2(D)^3\}.$$

Here and henceforth, $(\cdot, \cdot)_D$ and $((\cdot, \cdot))_D$ denote $L^2(D)$ and $L^2(D)^3$ inner products, respectively, and $\langle \cdot, \cdot \rangle_{\partial D}$ and $\langle\langle \cdot, \cdot \rangle\rangle_{\partial D}$ the duality pairing between $H^{-1/2}(\partial D)$ and $H^{1/2}(\partial D) = H_0^{1/2}(\partial D)$, and $H^{-1/2}(\partial D)^3$ and $H^{1/2}(\partial D)^3$, respectively. Then we have the following.

Proposition 0.1 *Each $v \in H(\operatorname{div}, D)$ admits the trace*

$$n \cdot v|_{\partial D} \in H^{-1/2}(\partial D),$$

and Green's formula

$$((v, \nabla \varphi))_D + (\nabla \cdot v, \varphi)_D = \langle n \cdot v, \varphi \rangle_{\partial D}$$

holds for $\varphi \in H^1(D)$.

Proposition 0.2 *Each $v \in H(\operatorname{rot}, D)$ admits the trace*

$$n \times v|_{\partial D} \in H^{-1/2}(\partial D)^3,$$

and the Stokes formula

$$((\nabla \times v, w))_D - ((v, \nabla \times w))_D = \langle\langle n \times v, w \rangle\rangle_{\partial D}$$

holds for $w \in H^1(D)^3$.

To discuss the interface regularity of the solution B to the Maxwell system (0.2), we take that

$$\Gamma_{\pm} = \partial\Omega_{\pm} \cap \mathcal{M}$$

with $\partial\Omega_{\pm}$ being the boundary of Ω_{\pm} . This means that Γ_+ and Γ_- coincide as sets, but are regarded as parts of the boundaries of Ω_+ and Ω_- , respectively. Henceforth, n denotes the outer unit normal vector to Γ_- so that $-n$ is the outer unit normal vector to Γ_+ . Unless otherwise stated, if \mathcal{M} is $C^{k,1}$, then $C^{k,1} \cap W_{loc}^{k+1,\infty}$ extension of the normal vector n defined on $\Gamma = \mathcal{M} \cap \Omega$ is always taken to Ω henceforth, where k is a non-negative integer. Furthermore, given a function $A(x)$ on Ω_{\pm} , we set

$$[A]_{\pm}^{\pm} = A_+ - A_- \quad \text{on} \quad \Gamma,$$

where $A_{\pm}(\xi) = \lim_{x \rightarrow \xi, x \in \Omega_{\pm}} A(x)$ for $\xi \in \Gamma$ are always taken in the sense of traces to Γ_{\pm} .

Suppose that B and J are in $L^2(\Omega_{\pm})^3$ and satisfy (0.2). This means that those relations hold piecewisely in Ω_{\pm} in the sense of distributions $\mathcal{D}'(\Omega_{\pm})$, that is,

$$\int_{\Omega_{\pm}} B \cdot \nabla \times C = \int_{\Omega_{\pm}} J \cdot C \quad \text{and} \quad \int_{\Omega_{\pm}} B \cdot \nabla \varphi = 0$$

for any $C \in C_0^{\infty}(\Omega_{\pm})^3$ and $\varphi \in C_0^{\infty}(\Omega_{\pm})$. Unless otherwise stated, those vector fields $B \in L^2(\Omega_{\pm})^3$ and $J \in L^2(\Omega_{\pm})^3$ are identified with the elements in $L^2(\Omega)^3$.

For the moment, we assume that \mathcal{M} is Lipschitz continuous and relation (0.2) holds for $B \in L^2(\Omega_{\pm})^3$ and $J \in L^2(\Omega_{\pm})^3$. This implies that $B \in H(\text{rot}, \Omega_{\pm}) \cap H(\text{div}, \Omega_{\pm})$, which assures the well-definedness of

$$n \times B|_{\Gamma_{\pm}} \in H^{-1/2}(\Gamma_{\pm})^3 \quad \text{and} \quad n \cdot B|_{\Gamma_{\pm}} \in H^{-1/2}(\Gamma_{\pm}),$$

and hence $B|_{\Gamma_{\pm}} \in H^{-1/2}(\Gamma_{\pm})^3$ follows. Furthermore,

$$[n \times B]_{-}^{+} = 0 \quad \text{and} \quad [n \cdot B]_{-}^{+} = 0 \quad (0.3)$$

if and only if

$$\nabla \times B = J \in L^2(\Omega)^3 \quad \text{and} \quad \nabla \cdot B = 0 \in L^2(\Omega) \quad (0.4)$$

as distributions in Ω , respectively. If both relations of (0.4) are satisfied, then $B \in H^1(\Omega)^3$ follows, because $B \in H^1(\Omega)^3$ is equivalent to $[B]_{-}^{+} = 0$ on Γ for $B \in H^1(\Omega_{\pm})^3$. A slightly weaker fact $B \in H_{loc}^1(\Omega)^3$ is also obtained by Corollary I.2.10 of [2],

$$H(\text{rot}, \Omega) \cap H(\text{div}, \Omega) \subset H_{loc}^1(\Omega)^3, \quad (0.5)$$

as (0.4) implies $B \in H_{loc}^1(\Omega)^3$.

Our first result is stated as follows. There, \mathcal{M} is supposed to be $C^{1,1}$ and hence $C^{1,1} \cap W_{loc}^{2,\infty}(\Omega)$ extension is taken to n .

Theorem 0.1 *If \mathcal{M} is $C^{1,1}$, and $B \in H^1(\Omega)^3$ and $J \in H(\text{rot}, \Omega_{\pm})$ satisfy (0.2), then it holds that $n \cdot B \in H_{loc}^2(\Omega)$.*

Above theorem is a slight improvement of a theorem of [1], but new argument for the proof is presented. Similarly to that case, $B \in H^1(\Omega)^3$ solves (0.2) in Ω as a distribution, so that

$$\int_{\Omega} B \cdot \nabla \times C = \int_{\Omega} J \cdot C \quad \text{and} \quad \int_{\Omega} B \cdot \nabla \varphi = 0$$

hold for any $C \in C_0^{\infty}(\Omega)^3$ and $\varphi \in C_0^{\infty}(\Omega)$. We note that $J \in H(\text{rot}, \Omega_{\pm})$ belongs to $J \in H(\text{rot}, \Omega)$ if and only if $[n \times J]_{-}^{+} = 0$ on Γ , and if this condition is satisfied furthermore, then we have

$$-\Delta B = \nabla \times J \in L^2(\Omega)^3$$

(as distributions in Ω), because $\nabla \times B = J \in H(\text{rot}, \Omega)$ and $\nabla \cdot B = 0 \in L^2(\Omega)^3$ are valid similarly in Ω . Then, $B \in H_{loc}^2(\Omega)^3$ is obtained from the standard elliptic regularity. Theorem 0.1 says, in contrast, that even if $n \times J$ has an interface on $\Gamma = \mathcal{M} \cap \Omega$, the normal component $n \cdot B$ of B gains the regularity in one rank. It is not difficult to suspect that the solenoidal condition $\nabla \cdot B = 0$ plays an essential role in such a regularity. In this connection, it must be noted that in Theorem 0.1, interface to $n \cdot J$ is not permitted. In fact, the first equation of (0.2) holds in Ω , and therefore,

$$\nabla \cdot J = \nabla \cdot (\nabla \times B) = 0$$

follows there. This implies $J \in H(\text{div}, \Omega)$, and hence $[n \cdot J]_{-}^{+} = 0$ holds on Γ .

The second theme of this article is the stationary Stokes system;

$$\left. \begin{aligned} -\Delta v &= \nabla p + f \\ \nabla \cdot v &= 0 \end{aligned} \right\} \quad \text{in} \quad \Omega_{\pm} \quad (0.6)$$

where $v = {}^t(v^1(x), v^2(x), v^3(x))$ denotes the velocity of fluid, $p = p(x)$ the pressure, and $f(x) = {}^t(f^1(x), f^2(x), f^3(x))$ the external force. The following theorem is proven by Theorem 0.1 and the vorticity formulation. There, \mathcal{M} is supposed to be $C^{2,1}$ so that $C^{2,1} \cap W_{loc}^{3,\infty}(\Omega)$ extension is taken to n . Actually, [1] showed the theorem when \mathcal{M} is flat.

Theorem 0.2 *If \mathcal{M} is $C^{2,1}$, $v \in H^2(\Omega)^3$, $p \in H^1(\Omega)$, and $f \in H^1(\Omega_{\pm})^3$ satisfy (0.6), and $[n \cdot \nabla p]_{\pm}^{\pm} = 0$ holds on Γ , then the condition $(n \cdot \nabla)^2(n \times v) \in H^1(\Omega)^3$ implies that $v \in H_{loc}^3(\Omega)^3$ and $p \in H_{loc}^2(\Omega)$.*

Standard regularity associated with the above theorem is obvious, so that $v \in H^2(\Omega)^3$, $p \in H^1(\Omega)$, and $f \in H^1(\Omega)^3$ imply $v \in H_{loc}^3(\Omega)$ and $p \in H_{loc}^2(\Omega)$ in (0.6). On the other hand, $f \in H(\text{div}, \Omega)$ can take place of the assumption

$$[n \cdot \nabla p]_{\pm}^{\pm} = \left[\frac{\partial p}{\partial n} \right]_{\pm}^{\pm} = 0 \quad \text{on} \quad \Gamma \quad (0.7)$$

in Theorem 0.2, because (0.6) holds in Ω and therefore, $f \in H(\text{div}, \Omega)$ gives that

$$-\Delta p = -\nabla \cdot \nabla p = \nabla \cdot f \in L^2(\Omega),$$

in Ω , or $\nabla p \in H(\text{div}, \Omega)$. This implies (0.7) and also $p \in H_{loc}^2(\Omega)$ from the standard elliptic regularity. In other words, if (0.6) holds in a natural L^2 setting in Ω , then $f \in H^1(\Omega_{\pm})^3 \cap H(\text{div}, \Omega)$ implies H^2 interface vanishing of the pressure and H^3 interface of the velocity only in the second normal derivative of the tangential component. Namely, we have the following.

Theorem 0.3 *If \mathcal{M} is $C^{2,1}$ and $v \in H^2(\Omega)^3$, $p \in H^1(\Omega)$, and $f \in L^2(\Omega)^3$ satisfy (0.6), then $f \in H(\text{div}, \Omega)$ implies $p \in H_{loc}^2(\Omega)$, and $f \in H^1(\Omega_{\pm})^3$ with $(n \cdot \nabla)^2(n \times v) \in H^1(\Omega)^3$, furthermore, gives that $v \in H_{loc}^3(\Omega)^3$.*

Those interface vanishing theorems are optimal in the sense that there is $v \in H^2(\Omega)^3$, $p \in H^2(\Omega)$, and $f \in H^1(\Omega_{\pm})^3 \cap H(\text{div}, \Omega)$ satisfying (0.6), (0.7), and $[(n \cdot \nabla)^2(n \times v)]_{\pm}^{\pm} \neq 0$ on Γ . Among them is the case that

$$\mathcal{M} = \{x = (x_1, x_2, x_3) \mid x_3 = 0\}$$

with $n = {}^t(0, 0, 1)$, $\Omega = \{x = (x_1, x_2, x_3) \mid x_1^2 + x_2^2 + x_3^2 < 1\}$,

$$v = \begin{pmatrix} \chi(x_1 - x_2)x_3|x_3| \\ \chi(x_1 - x_2)x_3|x_3| \\ 0 \end{pmatrix} \in H^2(\Omega)^3,$$

and $p = x_3|x_3| \in H^2(\Omega)$, where χ is a smooth function on \mathbf{R} with the support containing 0. In fact, we have

$$f = - \begin{pmatrix} 2\chi(x_1 - x_2)H(x_3) + \chi''(x_1 - x_2)x_3|x_3| \\ 2\chi(x_1 - x_2)H(x_3) + \chi''(x_1 - x_2)x_3|x_3| \\ 2|x_3| \end{pmatrix},$$

where $H = H(s)$ is the Heaviside function:

$$H(s) = \begin{cases} 1 & (x_3 > 0) \\ -1 & (x_3 < 0) \end{cases}$$

and this f is in $H^2(\Omega_{\pm})^3 \cap H(\text{div}, \Omega)$. Thus, here actual interface arises in the second normal derivative of the tangential component of the velocity, in spite that any other assumption in Theorems 0.2 and 0.3 is satisfied.

References

- [1] T. Kobayashi, T. Suzuki, and K. Watanabe, Interface regularity Maxwell and Stokes systems, to appear in Osaka J. Math..
- [2] V. Girault and P-A. Raviart "Finite element methods for Navier-Stokes equations." Springer-Verlag, Berlin, 1986.
- [3] T. Suzuki, K. Watanabe, and M. Shimogawara, *Current state and mathematical analysis of magnetoencephalography* (in Japanese), Osaka Univ. Research Reports in Math. no.1 (2000)