

Title	Remarks on a New Existence Theorem for Generalized Vector Equilibrium Problems and its Applications (Nonlinear Analysis and Convex Analysis)
Author(s)	Kalmoun, El Mostafa; Riahi, Hassan; Tanaka, Tamaki
Citation	数理解析研究所講究録 (2002), 1246: 165-173
Issue Date	2002-01
URL	http://hdl.handle.net/2433/41725
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Remarks on a New Existence Theorem for Generalized Vector Equilibrium Problems and its Applications

El Mostafa Kalmoun*, Hassan Riahi

Cadi Ayyad University
Faculty of Science Semlalia
Department of Mathematics
B.P. 2390, Marrakech-40000, Morocco

Tamaki Tanaka

Graduate School of Science and Technology
Niigata University
Niigata 950-2181, Japan

We consider a generalized vector equilibrium problem, which is the following set-valued vector version of Ky Fan's minimax inequality:

Find $\bar{x} \in C$ such as to satisfy $\varphi(\bar{x}, y) \not\subseteq K(\bar{x})$ for all $y \in C$, (GVEP)

where

- X and E are topological vector spaces,
- C is a nonempty closed convex subset of X ,
- $\varphi : C \times C \rightarrow 2^E$ is a set-valued map, and
- K is a set-valued map from C to E .

*The research of the first author is supported by the Matsumae International Foundation during his stay at Niigata University.

Using a particular case of the extended version of Fan-KKM theorem [6, Theorem 2.1], we can formulate the following general existence theorem for (GVEP) in topological vector spaces.

First, we need to recall the following definitions. Let $\psi : C \times C \rightarrow 2^E$ and $L : C \rightarrow 2^E$ be two other set-valued maps, and denote by $\mathcal{F}(C)$ the set of all finite subsets of C .

Definition 1. We say that ψ is diagonally quasi convex in its first argument relatively to L , in short L -diagonally quasi convex in x , if for any A in $\mathcal{F}(C)$ and any y in $co(A)$, we have $\psi(A, y) \not\subseteq L(y)$.

Definition 2. We say that φ is K -transfer semicontinuous in y if for any $(x, y) \in C \times X$ with $\varphi(x, y) \subset K(y)$, there exist an element $x' \in C$ and an open $V \subset X$ containing y such that $\varphi(x', v) \subset K(v)$ for all $v \in V$.

Theorem 1. ([7, Theorem 2.1]) Suppose that

(A0) $\psi(x, y) \not\subseteq L(y) \implies \varphi(x, y) \not\subseteq K(y) \forall x, y \in C$;

(A1) ψ is L -diagonally quasi-convex in x ;

(A2) φ is K -transfer semicontinuous in y ;

(A3) there is a nonempty compact subset B in X such that for each $A \in \mathcal{F}(C)$ there is a compact convex $B_A \subset X$ containing A such that, for every $y \in B_A \setminus B$, there exists $x \in B_A \cap C$ with

$$y \in \text{int}_X \{v \in X : \psi(x, v) \subseteq L(v)\}.$$

Then there exists $\bar{y} \in B$ such that $\varphi(x, \bar{y}) \not\subseteq K(\bar{y})$ for all $x \in C$.

Theorem 1 generalizes [2, Theorem 2.1], which is proved by means of a Fan-Browder fixed point theorem - an immediate consequence of the Fan-KKM theorem. As we will mention in the 'Assumptions analysis' subsection, our hypotheses are more general than those used in [2]. Besides, the scalar version of this result extends [10, Theorem 4] (we take $C_A = co(A \cup R) \cap X$ where R is the convex compact which contains C in [10, Theorem 4, (4iii)]). Other particular cases are [1, Theorem 2], [12, Theorem 2.1], [13, Theorem 2.11], [11, Theorem 1], [8, Corollary 2.4], [9, Lemma 2.1] and [3, Theorem 2]. The origin of this kind of results goes back to Ky Fan [5]. His classical

minimax inequality can be deduced from our result by setting $E = \mathbb{R}$, $K(x) = \mathbb{R}_+^*$ and $\varphi(x, y) = \psi(x, y) = f(x, y) - \sup_{x \in C} f(x, x)$ for all $x, y \in C$.

Let us turn to Theorem 1 and analyze its requirements by presenting different situations where assumptions (A0)-(A3) hold true. Let $(P(y))_{y \in C}$ a family of proper convex closed cones on E with $\text{int } P(y) \neq \emptyset$ for all $y \in C$.

- Pseudomonotonicity

Remark 1. (A0) holds provided one of the following statements is satisfied.

(a) $\varphi = \psi$ and $K = L$.

(b) $X = C$, $K(y) = -L(y) = -\text{int } P(y)$, $\psi(x, y) = \varphi(y, x)$ for all $x, y \in C$, and φ is P_x -pseudomonotone, that is,

$$\varphi(x, y) \not\subseteq \text{int } P(x) \implies \varphi(y, x) \not\subseteq -\text{int } P(x) \quad \forall x, y \in C.$$

- Convexity.

Remark 2. (A1) holds provided that, for every $y \in C$, one has either

(a) $\psi(y, y) \not\subseteq L(y)$, and

(b) the set $\{x \in C : \psi(x, y) \subseteq L(y)\}$ is convex,

or

(i) $L(y) = -\text{int } P(y)$ and $P(y)$ is w -pointed¹,

(ii) $\psi(y, y) \subseteq P(y)$, and

(iii) ψ is left P_y -quasiconvex, that is, for all $x_1, x_2, y \in C$ and all $\lambda \in [0, 1]$, one has either

$$\psi(x_1, y) \subseteq \psi(\lambda x_1 + (1 - \lambda)x_2, y) + P(y)$$

or

$$\psi(x_2, y) \subseteq \psi(\lambda x_1 + (1 - \lambda)x_2, y) + P(y).$$

- Continuity

¹A cone P is w -pointed if $P \cap -\text{int } P = \emptyset$.

Remark 3. (A2) holds provided that one of the following statements is satisfied.

- (a) φ is (transfer) u.s.c in y with compact values and if K has an open graph.
- (b) φ is (transfer) u.s.c in y and $K(x) = O$ for all $x \in C$, where O is an open subset of E .
- (c) For each $x \in C$, the set $\{y \in X : \varphi(x, y) \not\subseteq K(y)\}$ is closed in C .

- **Coercivity.**

Remark 4. (A3) holds if one of the following statements is satisfied.

- (a) C is compact.
- (b) There is $x_0 \in C$ such that $\psi(x_0, \cdot)$ is K -compact.
- (c) There is a nonempty compact subset B in C such that for each $y \in C \setminus B$ there exists $x \in B \cap C$ such that $\psi(x, y) \subseteq L(y)$.
- (d) There is a nonempty compact subset B of C and a compact convex subset $B' \in C$ such that for each $y \in C \setminus B$ there exists $x \in B' \cap C$ with

$$y \in \text{int} \{v \in X : \psi(x, v) \subseteq L(v)\}.$$

Besides, when the classical assumption (c) of Remark 3 is satisfied, (A3) holds provided that

- (e) there is a nonempty compact subset B in X such that for each $A \in \mathcal{F}(C)$ there is a compact convex $B_A \subset X$ containing A such that, for every $y \in C \setminus B$, there exists $x \in B_A \cap C$ with $\varphi(x, y) \subseteq K(y)$.

Applications

a) **Generalized vector variational like-inequalities**

Let us consider a set-valued operator T from C into $L(X, E)$, and a bifunction η from C to itself. We write for $\Pi \subset L(X, E)$ and $x \in C$, $\langle \Pi, x \rangle =$

$\{\langle \pi, x \rangle : \pi \in \Pi\}$, where $\langle \pi, x \rangle$ denotes the evaluation of the linear mapping π at x which is supposed to be continuous on $L(X, E) \times X^2$.

The generalized vector variational inequality problem (*GVVLIP*) takes the following form:

$$\text{Find } \bar{x} \in C \text{ such that } \langle T\bar{x}, \eta(y, \bar{x}) \rangle \not\subseteq -\text{int } P(\bar{x}) \quad \forall y \in C.$$

Thus (*GVVLIP*) is a particular case of (*GVEP*) if we take

$$\varphi(x, y) = \{\langle t, \eta(y, x) \rangle : t \in Tx\}.$$

For the reader's convenience, we recall the following definitions.

Definition 3. 1) T is said to be η -pseudomonotone if, for all $x, y \in C$,

$$\langle Tx, \eta(y, x) \rangle \not\subseteq -\text{int } P(x) \Rightarrow \langle Ty, \eta(y, x) \rangle \not\subseteq -\text{int } P(x).$$

2) T is said to be V -hemicontinuous if for any $x, y \in C$ and $t \in]0, 1[$ $T(tx + (1-t)y) \rightarrow T(y)$ as $t \rightarrow 0_+$ (i.e. for any $z_t \in T(tx + (1-t)y)$ there exists $z \in Ty$ such that for any $a \in C$, $\langle z_t, a \rangle \rightarrow \langle z, a \rangle$ as $t \rightarrow 0_+$).

It has to be observed that when T is single-valued, we recover the hemicontinuity used in [4]. if $\eta(x, y) = x - y$ for all $x, y \in C$, η is dropped from the definition of pseudomonotonicity.

The linearization lemma plays a significant role in variational inequalities. Chen [4] extended this lemma to the single-valued vector case. For our need in this subsection, we state it in the set-valued case by using standard Minty's argument. Consider the following problem, which may be seen as a dual problem of (*GVVLIP*),

$$\text{Find } \bar{x} \in C \text{ such that } \langle Ty, \eta(y, \bar{x}) \rangle \not\subseteq -\text{int } P(\bar{x}) \quad \forall y \in C. \quad (\text{GVVLIP}^*)$$

Lemma 1. Suppose that $\eta(\cdot, x)$ is affine and $\eta(x, x) = 0$ for each $x \in C$. If T is η -pseudomonotone and V -hemicontinuous then (*GVVLIP*) and (*GVVLIP**) are equivalent.

As an application of Theorem 1, we are now in position to formulate the following existence result for (*GVVLIP*).

²A typical situation when X is a reflexive Banach and E is a Banach

Theorem 2. *Suppose that*

- (i) *the mapping $\text{int } P(\cdot)$ has an open graph in $C \times L(X, E)$;*
- (ii) *for each $x \in C$, $\eta(\cdot, x)$ is affine, $\eta(x, \cdot)$ is continuous and $\eta(x, x) = 0$;*
- (iii) *T is compact valued, η -pseudomonotone and V -hemicontinuous;*
- (iv) *there is a nonempty compact subset B in C such that for each $A \in \mathcal{F}(C)$ there is a compact convex $B_A \subset C$ containing A such that, for every $y \in B_A \setminus B$, there exists $x \in B_A \cap C$ with*

$$y \in \text{int}_C \{v \in C : \langle Tv, \eta(x, v) \rangle \subseteq -\text{int } P(v)\}.$$

Then (GVVLIP) has at least one solution, which is in B .

Proof. Set $\varphi(x, y) = \langle Tx, \eta(x, y) \rangle$, $\psi(x, y) = \langle Ty, \eta(x, y) \rangle$ and $K(x) = -\text{int } P(x)$ for all $x, y \in C$. We can show that the assumptions of Theorem 1 are satisfied; see the proof of Theorem 4.1 in [7]. Therefore, from Theorem 1, there exists $\bar{x} \in B$ such that

$$\langle Ty, \eta(y, \bar{x}) \rangle \not\subseteq -\text{int } P(\bar{x}) \quad \forall y \in C.$$

Hence (GVVLIP*) has a solution in B , which completes the proof of the theorem according to Lemma 1. ■

b) Vector complementarity problems

A natural extension of the classical nonlinear complementarity problem, (CP) for short, is considered as follows. Let T be a single-valued operator from C , which is supposed to be a convex closed cone, to $L(X, E)$. The vector complementarity problem considered in this subsequent, (VCP) for short, is to find $\bar{x} \in C$ such that

$$\langle T(\bar{x}), \bar{x} \rangle \notin \text{int } P(\bar{x}), \text{ and } \langle T(\bar{x}), y \rangle \notin -\text{int } P(\bar{x}) \text{ for all } y \in C.$$

This problem collapses to (CP) when $E = \mathbb{R}$ and $P(x) = \mathbb{R}_+$ for all $x \in C$.

By means of vector variational inequalities, we can formulate the following existence theorem for (VCP).

Theorem 3. *Suppose that*

- (i) the set-valued map $\text{int } P(\cdot)$ has an open graph in $C \times L(X, E)$;
- (ii) T is pseudomonotone and hemicontinuous;
- (iv) there is a nonempty compact subset B in C such that for each $A \in \mathcal{F}(C)$ there is a compact convex $B_A \subset C$ containing A such that, for every $y \in B_A \setminus B$, there exists $x \in B_A \cap C$ with

$$y \in \text{int}_C \{v \in C : \langle Tv, x - v \rangle \in -\text{int } P(v)\}.$$

Then (VCP) has at least one solution, which is in B .

Proof. It is clear that all the assumptions of Theorem 2 are satisfied with $\eta(x, y) = x - y$ for all $x, y \in C$. Therefore there exists $\bar{x} \in B$ such that

$$\langle T\bar{x}, z - \bar{x} \rangle \notin -\text{int } P(\bar{x}) \quad \forall z \in C. \quad (1)$$

Since C is a convex cone, then setting in (1), $z = 0$ and $z = y + \bar{x}$ for an arbitrary $y \in C$, we get respectively

$$\langle T\bar{x}, \bar{x} \rangle \notin \text{int } P(\bar{x}) \text{ and } \langle T\bar{x}, y \rangle \notin -\text{int } P(\bar{x}).$$

Hence we conclude that \bar{x} is also a solution to (VCP). ■

c) Vector optimization

Here, to convey an idea about the use of vector variational-like inequalities in vector optimization which involves smooth vector mappings, we prove the existence of solutions to weak vector optimization problems, (WVOP) for short, by considering the concept of invexity. Let us state the problem as follows.

$$\text{Find } \bar{x} \in C \text{ such that } \phi(y) - \phi(\bar{x}) \notin -\text{int } P \text{ for all } y \in C, \quad (\text{WVOP})$$

where $\phi : C \rightarrow E$ is a given vector-valued function and P is a given convex cone in E .

Let $\eta : C \times C \rightarrow X$ be a given function, and denote by $\nabla\phi$ the Fréchet derivative of ϕ once the latter is assumed to be Fréchet differentiable.

Theorem 4. Suppose that P is a convex cone in E with $\text{int } P \neq \emptyset$, and let $\phi : C \rightarrow E$ be a Fréchet differentiable mapping. Assume that

(i) $\langle \nabla \phi(x), \eta(y, x) \rangle \notin -\text{int } P$ implies $\langle \nabla \phi(y), \eta(y, x) \rangle \notin -\text{int } P$ for all $x, y \in C$;

(ii) ϕ is P -invex with respect to η , that is,

$$\phi(y) - \phi(x) - \langle \nabla \phi(x), \eta(y, x) \rangle \in P \quad \forall x, y \in C.$$

(iii) $\nabla \phi$ is hemicontinuous;

(iv) for each $x \in C$, $\eta(\cdot, x)$ is affine, $\eta(x, \cdot)$ is continuous and $\eta(x, x) = 0$;

(v) there is a compact subset B in C such that for every finite subset A in C there is a compact convex $C_A \subset X$ containing A such as to satisfy, for every $y \in C \setminus B$, there exists $x \in C_A \cap C$ with $\langle \nabla \phi(x), \eta(x, y) \rangle \in -\text{int } P$.

Then (WVOP) has at least one solution.

Proof. First, by virtue of Theorem 2 with $T := \nabla \phi$, we get

$$\langle \nabla \phi(\bar{x}), \eta(y, \bar{x}) \rangle \notin -\text{int } P \quad \forall y \in C.$$

Then the P -invexity of ϕ allows us to conclude. ■

References

- [1] G. Allen, Variational inequalities, complementarity problems, and duality theorems, *Journal of Mathematical Analysis and Applications* 58 (1977), 1-10.
- [2] Q. H. Ansari and J.-C. Yao, An Existence Result for the Generalized Vector Equilibrium Problem, *Applied Mathematics Letters* 12 (1999), 53-56.
- [3] H. Ben-El-Mechaiekh, P. Deguire, and A. Granas, Une alternative non linéaire en analyse convexe at applications, *Compte Rendu de l'Académie des Sciences Paris Série I* 295 (1982), 257-259.

- [4] G. Y. Chen, Existence of Solutions for a Vector Variational Inequality: An Extension of Hartman-Stampacchia Theorem, *Journal of Optimization Theory and Applications* **74** (1992), 445-456.
- [5] Ky Fan, A minimax inequality and application, in "Inequalities" (Edited by O. Shisha) Vol. 3, pp. 103-113, Academic Press, New York, 1972.
- [6] E. M. Kalmoun and H. Riahi, Topological KKM Theorems and Generalized Vector Equilibria on G-Convex Spaces with Applications, *Proceedings of the American Mathematical Society*, **129** (2001), 1335-1348.
- [7] E. M. Kalmoun, H. Riahi and T. Tamaki, On Vector Equilibrium Problems: Remarks on a General Existence Theorem and Applications, submitted to *Journal of Nonlinear Analysis and Convex Analysis*, 2001.
- [8] K.-K Tan, J. Yu and X.-Z. Yuan, Some new minimax inequalities and applications to existence of equilibria in H -spaces, *Nonlinear Analysis* **24** (1995), 1457-1470.
- [9] E. Tarafdar and X.-Z Yuan, Generalized variational inequalities and its applications, *Nonlinear Analysis Theory Methods and Applications* **30** (1997), 4171-4181.
- [10] G. Tian, Generalizations of the FKKM theorem and the Ky Fan minimax inequality with applications to maximal elements, price equilibrium, and complementarity, *Journal of Mathematical Analysis and Applications* **170** (1992), 457-471.
- [11] C. L. Yen, A minimax inequality and its applications to variational inequalities, *Pacific Journal of Mathematics* **97** (1981), 477-481.
- [12] X. Z. Yuan, The study of minimax inequalities and applications to economies and variational inequalities, *Memoirs of the American Mathematical Society* **132** (1998), N. 625.
- [13] J. X. Zhou and G. Chen, Diagonal convexity conditions for problems in convex analysis and quasi-variational inequalities, *Journal of Mathematical Analysis and Applications* **132** (1988), 213-225.