

Title	HIGH ENERGY AND SMOOTHNESS ASYMPTOTIC EXPANSION OF THE SCATTERING AMPLITUDE : MAIN IDEAS OF THE APPROACH (Spectral and Scattering Theory and Related Topics)
Author(s)	Yafaev, Dimitri
Citation	数理解析研究所講究録 (2002), 1255: 55-75
Issue Date	2002-04
URL	<a href="http://hdl.handle.net/2433/41900">http://hdl.handle.net/2433/41900</a>
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

HIGH ENERGY AND SMOOTHNESS ASYMPTOTIC EXPANSION OF  
THE SCATTERING AMPLITUDE  
(MAIN IDEAS OF THE APPROACH)

D.Yafaev

Department of Mathematics, University Rennes-1, Campus Beaulieu, 35042,  
Rennes, France (e-mail : yafaev@univ-rennes1.fr)

Abstract

We find an explicit expression for the kernel of the scattering matrix for the Schrödinger operator containing at high energies all terms of power order. It turns out that the same expression gives a complete description of the diagonal singularities of the kernel in the angular variables. The formula obtained is in some sense universal since it applies both to short- and long-range electric as well as magnetic potentials.

1. INTRODUCTION

1. High energy asymptotics of the scattering matrix  $S(\lambda) : L_2(\mathbb{S}^{d-1}) \rightarrow L_2(\mathbb{S}^{d-1})$  for the Schrödinger operator  $H = -\Delta + V$  in the space  $\mathcal{H} = L_2(\mathbb{R}^d)$ ,  $d \geq 2$ , with a real short-range potential (bounded and satisfying the condition  $V(x) = O(|x|^{-\rho})$ ,  $\rho > 1$ , as  $|x| \rightarrow \infty$ ) is given by the Born approximation. To describe it, let us introduce the operator  $\Gamma_0(\lambda)$ ,

$$(\Gamma_0(\lambda)f)(\omega) = 2^{-1/2}k^{(d-2)/2}\hat{f}(k\omega), \quad k = \lambda^{1/2} \in \mathbb{R}_+ = (0, \infty), \quad \omega \in \mathbb{S}^{d-1}, \quad (1.1)$$

of the restriction (up to the numerical factor) of the Fourier transform  $\hat{f}$  of a function  $f$  to the sphere of radius  $k$ . Set  $R_0(z) = (-\Delta - z)^{-1}$ ,  $R(z) = (H - z)^{-1}$ . By the Sobolev trace theorem and the limiting absorption principle the operators  $\Gamma_0(\lambda)\langle x \rangle^{-r} : \mathcal{H} \rightarrow L_2(\mathbb{S}^{d-1})$  and  $\langle x \rangle^{-r}R(\lambda + i0)\langle x \rangle^{-r} : \mathcal{H} \rightarrow \mathcal{H}$  are correctly defined as bounded operators for any  $r > 1/2$  and their norms are estimated by  $\lambda^{-1/4}$  and  $\lambda^{-1/2}$ , respectively. Therefore it is easy to deduce (see, e.g., [14, 24]) from the usual stationary representation

$$S(\lambda) = I - 2\pi i\Gamma_0(\lambda)(V - VR(\lambda + i0)V)\Gamma_0^*(\lambda) \quad (1.2)$$

for the scattering matrix (SM) and the resolvent identity that

$$S(\lambda) = I - 2\pi i \sum_{n=0}^N (-1)^n \Gamma_0(\lambda)V(R_0(\lambda + i0)V)^n \Gamma_0^*(\lambda) + \sigma_N(\lambda), \quad (1.3)$$

where  $\|\sigma_N(\lambda)\| = O(\lambda^{-(N+2)/2})$  as  $\lambda \rightarrow \infty$ . Moreover, the operators  $\sigma_N$  belong to suitable Schatten - von Neumann classes  $\mathfrak{S}_{\alpha(N)}$  and  $\alpha(N) \rightarrow 0$  as  $N \rightarrow \infty$ .

Nevertheless the Born expansion (1.3) has at least three drawbacks. First, the structure of the  $n^{\text{th}}$  term is extremely complicated already for relatively small  $n$ . Second, (1.3) definitely fails for long-range potentials, and, finally, it fails as  $\lambda \rightarrow \infty$  for a perturbation of the operator  $-\Delta$  by first order differential operators even with short-range coefficients (magnetic potentials).

2. In the particular case when  $A = 0$  and  $V$  belongs to the Schwartz class a convenient form of the high-energy expansion of the kernel of SM (called often the scattering amplitude) was obtained in [3] (see also the earlier paper [7]). The method of [3] relies on a preliminary study of the scattering solutions of the Schrödinger equation defined, for example, by the formula

$$\psi_{\pm}(\xi) = u_0(\xi) - R(|\xi|^2 \mp i0)Vu_0(\xi), \quad u_0(x, \xi) = \exp(i\langle x, \xi \rangle), \quad \xi = \hat{\xi}|\xi| \in \mathbb{R}^d.$$

It is shown in [3] that (at least on all compact sets of  $x$ ) the function  $\psi_{\pm}(x, \xi)$  has the asymptotic expansion  $\psi_{\pm}(x, \xi) = e^{i\langle x, \xi \rangle} b_{\pm}(x, \xi)$  where

$$b_{\pm}(x, \xi) = b_{\pm}^{(N)}(x, \xi) = \sum_{n=0}^N (2i|\xi|)^{-n} b_n^{(\pm)}(x, \xi), \quad b_0(x, \xi) = 1, \quad N \rightarrow \infty. \quad (1.4)$$

The function  $b_{\pm}(x, \xi)$  is determined by the transport equation (see subs. 2.3 below), and the coefficients  $b_n^{(\pm)}(x, \xi) = b_n^{(\pm)}(x, \hat{\xi})$  are quite explicit. Therefore it is easy to deduce from (1.2) that, for any  $N$ , the kernel of SM admits the asymptotic expansion

$$s(\omega, \omega'; \lambda) = \delta(\omega, \omega') - \pi i (2\pi)^{-d} k^{d-2} \times \sum_{n=0}^N (2ik)^{-n} \int_{\mathbb{R}^d} e^{ik(\omega' - \omega, x)} V(x) b_n^{(-)}(x, \omega') dx + O(k^{d-3-N}), \quad (1.5)$$

where  $\delta(\cdot)$  is of course the Dirac-function on the unit sphere. We emphasize that the functions  $b_n^{(-)}(x, \omega')$  are growing as  $|x| \rightarrow \infty$  in the direction of  $\omega'$  and the rate of growth increases as  $n$  increases. Thus, expansion (1.5) loses the sense (for sufficiently large  $N$ ) if  $V(x)$  decreases only as some power of  $|x|^{-1}$ .

The generalization of the results of [3] to the case of short-range potentials  $V$  satisfying the condition  $\partial^{\alpha} V(x) = O(|x|^{-\rho_v - |\alpha|})$  for some  $\rho_v > 1$  was suggested in [21] where the asymptotics of the scattering amplitude was also deduced from that of the scattering solutions. We note finally the paper [4] where the leading term of the high-energy asymptotics of the scattering amplitude was found for short-range magnetic potentials.

3. In the present paper we suggest a new method which allows us to find an explicit function  $s_0(\omega, \omega'; \lambda)$  which describes with arbitrary accuracy the kernel  $s(\omega, \omega'; \lambda)$  of the SM  $S(\lambda)$  at high energies (as  $\lambda \rightarrow \infty$ ), both for short- and long-range electric and magnetic potentials. It turns out that the same function  $s_0(\omega, \omega'; \lambda)$  gives also all diagonal singularities of the kernel  $s(\omega, \omega'; \lambda)$  in the angular variables  $\omega, \omega' \in \mathbb{S}^{d-1}$ . We emphasize that our approach allows us to avoid a study of solutions of the Schrödinger equation.

We consider the Schrödinger operator

$$H = (i\nabla + A(x))^2 + V(x) \quad (1.6)$$

in the space  $\mathcal{H}$  with electric  $V(x)$  and magnetic  $A(x) = (A_1(x), \dots, A_d(x))$  potentials satisfying the assumptions

$$\left. \begin{aligned} |\partial^\alpha V(x)| &\leq C_\alpha (1 + |x|)^{-\rho_v - |\alpha|}, & \rho_v > 0, \\ |\partial^\alpha A(x)| &\leq C_\alpha (1 + |x|)^{-\rho_a - |\alpha|}, & \rho_a > 0, \end{aligned} \right\} \quad (1.7)$$

for all multi-indices  $\alpha$ . We suppose that potentials are real, that is  $V(x) = \overline{V(x)}$  and  $A_j(x) = \overline{A_j(x)}$ ,  $j = 1, \dots, d$ . Set  $\rho = \min\{\rho_v, \rho_a\}$ , and

$$V_0(x) = V(x) + |A(x)|^2, \quad V_1(x) = V_0(x) + i \operatorname{div} A(x).$$

Then

$$H = -\Delta + 2i\langle A(x), \nabla \rangle + V_1(x).$$

We emphasize that the cases  $\rho > 1$  (short-range potentials) and  $\rho \in (0, 1]$  (long-range potentials) are treated in almost the same way.

Let us formulate our main result. The answer is given in terms of approximate solutions of the Schrödinger equation

$$-\Delta \psi(x, \xi) + 2i\langle A(x), \nabla \rangle \psi(x, \xi) + V_1(x) \psi(x, \xi) = |\xi|^2 \psi(x, \xi). \quad (1.8)$$

To be more precise, we denote by  $u_\pm(x, \xi) = u_\pm^{(N)}(x, \xi)$  explicit functions (see Section 2, for their construction)

$$u_\pm(x, \xi) = e^{i\Theta_\pm(x, \xi)} b_\pm(x, \xi) \quad (1.9)$$

such that

$$(-\Delta + 2i\langle A(x), \nabla \rangle + V_1(x) - |\xi|^2) u_\pm(x, \xi) = e^{i\Theta_\pm(x, \xi)} r_\pm(x, \xi) =: q_\pm(x, \xi) \quad (1.10)$$

and  $r_\pm(x, \xi) = r_\pm^{(N)}(x, \xi)$  tends to zero faster than  $|x|^{-p}$  as  $|x| \rightarrow \infty$  and  $|\xi|^{-q}$  as  $|\xi| \rightarrow \infty$  where  $p = p(N) \rightarrow \infty$  and  $q = q(N) \rightarrow \infty$  as  $N \rightarrow \infty$  off any conical neighborhood of the direction  $\hat{x} = \mp \hat{\xi}$ . Note that the phase  $\Theta_\pm(x, \xi) = \langle x, \xi \rangle$  if  $A(x) = 0$  and  $V(x)$  is a short-range function and  $\Theta_\pm(x, \xi)$  satisfies the eikonal equation in the general case. The function  $b_\pm(x, \xi)$  is obtained as a solution of the corresponding transport equation.

As is well known (see [1]), off the diagonal  $\omega = \omega'$ , the kernel  $s(\omega, \omega'; \lambda)$  is a  $C^\infty$ -function of  $\omega, \omega' \in \mathbb{S}^{d-1}$  where it tends to zero faster than any power of  $\lambda^{-1}$  as  $\lambda \rightarrow \infty$ . Thus, it suffices to describe the structure of  $s(\omega, \omega'; \lambda)$  in a neighborhood of the diagonal  $\omega = \omega'$ . Let  $\omega_0 \in \mathbb{S}^{d-1}$  be an arbitrary point,  $\Pi_{\omega_0}$  be the plane orthogonal to  $\omega_0$  and  $\Omega_\pm(\omega_0, \delta) \subset \mathbb{S}^{d-1}$  be determined by the condition  $\pm \langle \omega, \omega_0 \rangle > \delta > 0$ . Set

$$x = \omega_0 z + y, \quad y \in \Pi_{\omega_0}, \quad (1.11)$$

and

$$\begin{aligned} s_0(\omega, \omega'; \lambda) &= s_0^{(N)}(\omega, \omega'; \lambda) = \mp \pi i k^{d-2} (2\pi)^{-d} \\ &\times \left( \int_{\Pi_{\omega_0}} \overline{(u_+(y, k\omega))} (\partial_z u_-(y, k\omega')) - u_-(y, k\omega') \overline{(\partial_z u_+(y, k\omega))} dy \right. \\ &\quad \left. - 2i \int_{\Pi_{\omega_0}} \langle A(y), \omega_0 \rangle \overline{u_+(y, k\omega)} u_-(y, k\omega') dy \right) \end{aligned} \quad (1.12)$$

for  $\omega, \omega' \in \Omega_{\pm} = \Omega_{\pm}(\omega_0, \delta)$ . Then, for any  $p, q$  and sufficiently large  $N = N(p, q)$ , the kernel

$$\tilde{s}^{(N)}(\omega, \omega'; \lambda) = s(\omega, \omega'; \lambda) - s_0^{(N)}(\omega, \omega'; \lambda) \quad (1.13)$$

belongs to the class  $C^p(\Omega \times \Omega)$  where  $\Omega = \Omega_+ \cup \Omega_-$ , and its  $C^p$ -norm is  $O(\lambda^{-q})$  as  $\lambda \rightarrow \infty$ . Thus, all singularities of  $s(\omega, \omega'; \lambda)$  both for high energies and in smoothness are described by the explicit formula (1.12). Let  $S_0(\lambda)$  be integral operator with kernel  $s_0(\omega, \omega'; \lambda)$ . In view of representation (1.9), formula (1.12) shows that we actually consider the singular part  $S_0(\lambda)$  of the SM as a Fourier integral or, more precisely, a pseudo-differential operator (PDO) acting on the unit sphere and determined by its amplitude.

By our construction of functions (1.9),  $u_+(x, \xi) = \overline{u_-(x, -\xi)}$  if  $A(x) = 0$ . Therefore in the case  $A = 0$  the singular part  $s_0(\omega, \omega'; \lambda)$  satisfies the same symmetry relation (the time reversal invariance)

$$s(\omega, \omega'; \lambda) = s(-\omega', -\omega; \lambda)$$

as kernel of the SM itself. Kernel (1.12) is also gauge invariant. This means that, for a smooth function  $\varphi(x)$ , the integrand in (1.12) is not changed if the functions  $u_{\pm}$  are replaced by  $e^{-i\varphi}u_{\pm}$  and the magnetic potential  $A$  is replaced by  $A - \nabla\varphi$ . We emphasize however that throughout the paper we do not use any particular gauge.

Formula (1.12) gives the singular part of the scattering amplitude off any neighborhood of the hyperplane  $\Pi_{\omega_0}$ . Since  $\omega_0 \in \mathbb{S}^{d-1}$  is arbitrary, this determines the singular part of  $s(\omega, \omega'; \lambda)$  for all  $\omega, \omega' \in \mathbb{S}^{d-1}$ . We note that the leading diagonal singularity of  $s(\omega, \omega', \lambda)$  was found in [23] for  $\rho_v \in (1/2, 1]$  and  $A = 0$ .

4. Our approach to the proof of formula (1.12) relies (even in the short-range case) on the expression of the SM via modified wave operators

$$W_{\pm}(H, H_0; J_{\pm}) = s - \lim_{t \rightarrow \pm\infty} e^{iHt} J_{\pm} e^{-iH_0 t}, \quad (1.14)$$

where PDO  $J_{\pm}$  are constructed in terms of the functions  $u_{\pm}(x, \xi)$ . Following [9], we kill neighborhoods of “bad” directions  $\hat{x} = \mp\hat{\xi}$  by appropriate cut-off functions  $\zeta_{\pm}(x, \xi)$ . Let

$$T_{\pm} = HJ_{\pm} - J_{\pm}H_0 \quad (1.15)$$

be the “effective” perturbation. The SM  $S(\lambda)$  corresponding to wave operators (1.14) admits (see [10, 23, 24, 19]) the representation

$$S(\lambda) = S_1(\lambda) + S_2(\lambda), \quad (1.16)$$

where

$$S_1(\lambda) = -2\pi i \Gamma_0(\lambda) J_+^* T_- \Gamma_0^*(\lambda) \quad (1.17)$$

and

$$S_2(\lambda) = 2\pi i \Gamma_0(\lambda) T_+^* R(\lambda + i0) T_- \Gamma_0^*(\lambda). \quad (1.18)$$

Both these expressions are correctly defined which will be discussed in Sections 5 and 4, respectively.

With the help of the so called propagation estimates [17, 12, 11] we show in Section 4 that the operator  $S_2(\lambda)$  has smooth kernel rapidly decaying as  $\lambda \rightarrow \infty$ . Therefore we call

$S_2(\lambda)$  the regular part of the SM. The singular part  $S_1(\lambda)$  is given by explicit expression (1.17) not depending on the resolvent of the operator  $H$ . However it contains the cut-off functions  $\zeta_{\pm}$ . In Section 5 we get rid of these auxiliary functions and, neglecting  $C^\infty$ -kernels decaying faster than any power of  $\lambda^{-1}$ , transform the kernel of  $S_1(\lambda)$  to the invariant expression (1.12).

## 2. THE EIKONAL AND TRANSPORT EQUATIONS

In this section we give a standard construction of approximate but explicit solutions of the Schrödinger equation. This construction relies on a solution of the corresponding eikonal and transport equations by iterations.

1. Let us plug expression (1.9) into the Schrödinger equation (1.8). Then

$$\begin{aligned} & (-\Delta + 2i\langle A(x), \nabla \rangle + V_1(x) - |\xi|^2)(e^{i\Theta}b) \\ &= e^{i\Theta}(|\nabla\Theta|^2b - i(\Delta\Theta)b - 2i\langle \nabla\Theta, \nabla b \rangle - \Delta b \\ & - 2\langle A, \nabla\Theta \rangle b + 2i\langle A, \nabla b \rangle + V_1b - |\xi|^2b), \quad \nabla = \nabla_x. \end{aligned} \quad (2.1)$$

We require that the phase  $\Theta(x, \xi)$  and the amplitude  $b(x, \xi)$  be approximate solutions of the eikonal and transport equations, that is

$$|\nabla\Theta|^2 - 2\langle A, \nabla\Theta \rangle + V_0 - |\xi|^2 = q_0(x, \xi), \quad (2.2)$$

and

$$-2i\langle \nabla\Theta, \nabla b \rangle + 2i\langle A, \nabla b \rangle - \Delta b + (-i\Delta\Theta + i\operatorname{div} A + q_0)b = r(x, \xi). \quad (2.3)$$

It follows from (2.1) that, for such functions  $\Theta$  and  $b$ , equality (1.10) is satisfied with the same function  $r(x, \xi)$  as in (2.3). When considering (2.2), (2.3), we always remove either a conical neighborhood of the direction  $\hat{x} = -\hat{\xi}$  (for the sign “+”) or  $\hat{x} = \hat{\xi}$  (for the sign “-”). We choose  $\Theta(x, \xi) = \Theta_{\pm}(x, \xi)$  in such a way that  $q_0(x, \xi) = q_0^{(\pm)}(x, \xi)$  defined by (2.2) is a short-range function of  $x$ , and it tends to 0 as  $|\xi| \rightarrow \infty$ . Then we construct  $b(x, \xi) = b_{\pm}(x, \xi)$  so that  $r(x, \xi) = r_{\pm}(x, \xi)$  decays as  $|x| \rightarrow \infty$  as an arbitrary given power of  $|x|^{-1}$ . It turns out that  $r(x, \xi)$  has a similar decay also in the variable  $|\xi|^{-1}$ .

If  $V$  is short-range and  $A = 0$ , then we can set  $\Theta_{\pm}(x, \xi) = \langle x, \xi \rangle$  and consider the transport equation (2.3) only. However, the eikonal equation (2.2) is necessary for any non-trivial magnetic potential or (and) long-range electric potential  $V$ . The transport equation is always unavoidable because, as we shall see below, the function  $\Delta\Theta_{\pm}$  decays at infinity as  $|x|^{-1-\rho}$  only and hence, for example, the choice  $b_{\pm} = 1$  in (1.9) is not sufficient.

We seek  $\Theta_{\pm}(x, \xi)$  in the form

$$\Theta_{\pm}(x, \xi) = \langle x, \xi \rangle + \Phi_{\pm}(x, \xi), \quad (2.4)$$

where  $(\nabla\Phi_{\pm})(x, \xi)$  tends to zero as  $|x| \rightarrow \infty$  off any conical neighborhood of the direction  $\hat{x} = \mp\hat{\xi}$ . We construct a solution of equation (2.2) by iterations. Actually, we set

$$\Phi_{\pm}(x, \xi) = \Phi_{\pm}^{(N_0)}(x, \xi) = \sum_{n=0}^{N_0} (2|\xi|)^{-n} \phi_n^{(\pm)}(x, \hat{\xi}) \quad (2.5)$$

and plug expressions (2.4) and (2.5) into equation (2.2). Comparing coefficients at the same powers of  $(2|\xi|)^{-n}$ ,  $n = -1, 0, \dots, N_0 - 1$ , we obtain the equations

$$\langle \hat{\xi}, \nabla \phi_0 \rangle = \langle \hat{\xi}, A \rangle, \quad \langle \hat{\xi}, \nabla \phi_1 \rangle + |\nabla \phi_0|^2 - 2\langle A, \nabla \phi_0 \rangle + V_0 = 0, \quad (2.6)$$

$$\langle \hat{\xi}, \nabla \phi_{n+1} \rangle + \sum_{m=0}^n \langle \nabla \phi_m, \nabla \phi_{n-m} \rangle - 2\langle A, \nabla \phi_n \rangle = 0, \quad n \geq 1, \quad (2.7)$$

so that the “error term” equals

$$q_0(x, \xi) = \sum_{n+m \geq N_0} (2|\xi|)^{-n-m} \langle \nabla \phi_n, \nabla \phi_m \rangle - 2(2|\xi|)^{-N_0} \langle A, \nabla \phi_{N_0} \rangle.$$

All equations (2.6), (2.7) have the form

$$\langle \hat{\xi}, \nabla \phi(x, \hat{\xi}) \rangle + f(x, \hat{\xi}) = 0 \quad (2.8)$$

and can be explicitly solved. Let the domain  $\Gamma_{\pm}(\epsilon, R) \subset \mathbb{R}^d \times \mathbb{R}^d$  be distinguished by the condition:  $(x, \xi) \in \Gamma_{\pm}(\epsilon, R)$  if either  $|x| \leq R$  or  $\pm \langle \hat{x}, \hat{\xi} \rangle \geq -1 + \epsilon$  for some  $\epsilon > 0$ . Of course, all constants below depend on  $\epsilon$  and  $R$ . The following assertion is almost obvious (see [23], for details).

**Lemma 2.1** *Suppose that*

$$|\partial_x^\alpha \partial_\xi^\beta f(x, \hat{\xi})| \leq C_{\alpha, \beta} |\xi|^{-|\beta|} (1 + |x|)^{-\rho - |\alpha|} \quad (2.9)$$

for  $x \in \Gamma_{\pm}(\epsilon, R)$  and some  $\rho > 1$ . Then the function

$$\phi^{(\pm)}(x, \hat{\xi}) = \pm \int_0^\infty f(x \pm t\hat{\xi}, \hat{\xi}) dt \quad (2.10)$$

satisfies equation (2.8) and the estimates

$$|\partial_x^\alpha \partial_\xi^\beta \phi^{(\pm)}(x, \hat{\xi})| \leq C_{\alpha, \beta} |\xi|^{-|\beta|} (1 + |x|)^{1 - \rho - |\alpha|}, \quad x \in \Gamma_{\pm}(\epsilon, R). \quad (2.11)$$

If estimates (2.9) are fulfilled for some  $\rho \in (0, 1)$  only, then the function

$$\phi^{(\pm)}(x, \hat{\xi}) = \pm \int_0^\infty (f(x \pm t\hat{\xi}, \hat{\xi}) - f(\pm t\hat{\xi}, \hat{\xi})) dt \quad (2.12)$$

satisfies both equation (2.8) and estimates (2.11).

Proceeding by induction, we can solve by formulas (2.10) or (2.12) all equations (2.6) and (2.7). The case where  $V$  and  $A$  are both short-range is discussed specially in subs. 3. Here we focus on the long-range case. Let us formulate the corresponding result.

**Proposition 2.2** *Let assumption (1.7) hold for some  $\rho \in (0, 1)$ . Then estimates*

$$|\partial_x^\alpha \partial_\xi^\beta \phi_n^{(\pm)}(x, \hat{\xi})| \leq C_{\alpha, \beta} |\xi|^{-|\beta|} (1 + |x|)^{1 - n\rho - |\alpha|}, \quad n = 1, 2, \dots,$$

and

$$|\partial_x^\alpha \partial_\xi^\beta q_0^{(\pm)}(x, \xi)| \leq C_{\alpha, \beta} |\xi|^{-N_0 - |\beta|} (1 + |x|)^{-N_0\rho - |\alpha|}.$$

are fulfilled on the set  $\Gamma_{\pm}(\epsilon, R)$  for all multi-indices  $\alpha$  and  $\beta$ . The function  $\phi_0^{(\pm)}(x, \hat{\xi})$  satisfies the same estimate as  $\phi_1^{(\pm)}(x, \hat{\xi})$ .

**Corollary 2.3** *The function (2.5) satisfies the estimates*

$$|\partial_x^\alpha \partial_\xi^\beta \Phi_\pm(x, \xi)| \leq C_{\alpha, \beta} |\xi|^{-|\beta|} (1 + |x|)^{1-\rho-|\alpha|}, \quad x \in \Gamma_\pm(\epsilon, R). \quad (2.13)$$

Below the number  $N_0$  in (2.4) is subject to the only restriction  $N_0 \rho \geq 2$ .

Of course, in particular cases the procedure above can be simplified. For example, if  $A = 0$  and  $V$  is long-range but  $\rho_v > 1/2$ , then

$$\Phi^{(\pm)}(x, \xi) = (2|\xi|)^{-1} \phi_1^{(\pm)}(x, \hat{\xi}) = \pm 2^{-1} \int_0^\infty (V(x \pm t\xi) - V(\pm t\xi)) dt.$$

2. An approximate solution of the transport equation (2.3) can be constructed by a procedure similar to the one given above. Using (2.4), we rewrite this equation as

$$-2i\langle \xi, \nabla b \rangle + 2i\langle A - \nabla \Phi, \nabla b \rangle - \Delta b + (-i\Delta \Phi + i \operatorname{div} A + q_0)b = r. \quad (2.14)$$

We look for the function  $b_\pm(x, \xi)$  in the form (1.4) with bounded in  $\xi$  coefficients  $b_n^{(\pm)}(x, \xi)$ . Plugging this expression into (2.14), we obtain the following recurrent equations

$$\langle \hat{\xi}, \nabla b_{n+1} \rangle = 2i\langle A - \nabla \Phi, \nabla b_n \rangle - \Delta b_n + (-i\Delta \Phi + i \operatorname{div} A + q_0)b_n, \quad n = 0, 1, \dots, N-1. \quad (2.15)$$

Then

$$r(x, \xi) = r^{(N)}(x, \xi) = (2i|\xi|)^{-N} \langle \hat{\xi}, \nabla b_{N+1} \rangle$$

All these equations have the form (cf. (2.8))

$$\langle \hat{\xi}, \nabla b_{n+1}(x, \xi) \rangle + f_n(x, \xi) = 0,$$

where a short-range function  $f_n$  depends on  $b_1, \dots, b_n$ . Therefore they can be solved by one of the formulas (2.10). Thus, using again Lemma 2.1, we obtain

**Proposition 2.4** *Let assumption (1.7) hold, let  $\rho_1 = \min\{1, \rho\}$  and let  $(x, \xi) \in \Gamma_\pm(\epsilon, R)$ . Then functions  $b_n^{(\pm)}$ ,  $n \geq 1$ , satisfy the estimates*

$$|\partial_x^\alpha \partial_\xi^\beta b_n^{(\pm)}(x, \xi)| \leq C_{\alpha, \beta} |\xi|^{-|\beta|} (1 + |x|)^{-\rho_1 n - |\alpha|}.$$

The right-hand side of equation (2.3) satisfies

$$|\partial_x^\alpha \partial_\xi^\beta r_\pm^{(N)}(x, \xi)| \leq C_{\alpha, \beta} |\xi|^{-N-|\beta|} (1 + |x|)^{-1-\rho_1(N+1)-|\alpha|}. \quad (2.16)$$

**Corollary 2.5** *The function (1.4) satisfies the estimates*

$$|\partial_x^\alpha \partial_\xi^\beta b_\pm(x, \xi)| \leq C_{\alpha, \beta} |\xi|^{-|\beta|} (1 + |x|)^{-|\alpha|}. \quad (2.17)$$

Combining Propositions 2.2 and 2.4, we get the final result.

**Theorem 2.6** *For the functions  $\Theta_\pm^{(N_0)}(x, \xi)$  and  $b_\pm^{(N)}(x, \xi)$  constructed in Propositions 2.2 and 2.4, respectively, and for the functions  $u_\pm(x, \xi) = u_\pm^{(N)}(x, \xi)$  defined by (1.9), equality (1.10) holds with the remainder  $r_\pm^{(N)}(x, \xi)$  satisfying estimates (2.16) in the region*



We emphasize that in contrast to the parameter  $N_0$  which is fixed, we need  $N \rightarrow \infty$ .

3. Of course, the functions  $b_n^{(\pm)}(x, \xi)$  contain different powers of  $|\xi|^{-1}$ . However, in the short-range case  $b_n^{(\pm)}$  depend on  $x$  and  $\hat{\xi}$  only. Suppose first that  $A = 0$ . Then  $\Phi = 0$  and equation (2.15) reduces to

$$\langle \hat{\xi}, \nabla b_{n+1} \rangle = -\Delta b_n + V b_n.$$

Thus, we obtain the following assertion.

**Proposition 2.7** *Let  $A = 0$  and let  $V$  satisfy assumption (1.7) with  $\rho_v > 1$ . Let  $u_{\pm}(x, \xi) = e^{i(x, \xi)} b_{\pm}(x, \xi)$  where  $b_{\pm}$  is the sum (1.4) and the functions  $b_n^{(\pm)}(x, \hat{\xi})$  are defined by recurrent formulas  $b_0^{(\pm)} = 1$  and*

$$b_{n+1}^{(\pm)}(x, \hat{\xi}) = \mp \int_0^{\infty} \left( -\Delta b_n^{(\pm)}(x \pm t\hat{\xi}, \hat{\xi}) + V(x \pm t\hat{\xi}) b_n^{(\pm)}(x \pm t\hat{\xi}, \hat{\xi}) \right) dt.$$

Then for  $(x, \xi) \in \Gamma_{\pm}(\epsilon, R)$  and  $\rho_2 = \min\{2, \rho_v\}$

$$|\partial_x^{\alpha} \partial_{\hat{\xi}}^{\beta} b_n^{(\pm)}(x, \hat{\xi})| \leq C_{\alpha, \beta} |\xi|^{-|\beta|} (1 + |x|)^{-(\rho_2 - 1)n - |\alpha|} \quad (2.18)$$

and the remainder (1.10) satisfies the estimates

$$|\partial_x^{\alpha} \partial_{\hat{\xi}}^{\beta} r_{\pm}^{(N)}(x, \xi)| \leq C_{\alpha, \beta} |\xi|^{-N - |\beta|} (1 + |x|)^{-(\rho_2 - 1)(N + 1) - |\alpha|}. \quad (2.19)$$

Let us write down explicit expressions for the first two functions  $b_n$ :

$$\begin{aligned} b_1^{(\pm)}(x, \hat{\xi}) &= \mp \int_0^{\infty} V(x \pm t\hat{\xi}) dt, \\ b_2^{(\pm)}(x, \hat{\xi}) &= - \int_0^{\infty} t(\Delta V)(x \pm t\hat{\xi}) dt + \frac{1}{2} \left( \int_0^{\infty} V(x \pm t\hat{\xi}) dt \right)^2. \end{aligned}$$

If a magnetic potential is non-trivial, then

$$\Phi_{\pm}(x, \hat{\xi}) = \phi_0^{(\pm)}(x, \hat{\xi}) = \mp \int_0^{\infty} \langle \hat{\xi}, A(x \pm t\hat{\xi}) \rangle dt \quad (2.20)$$

and

$$q_0^{(\pm)} = |\nabla \Phi_{\pm}|^2 - 2\langle A, \nabla \Phi_{\pm} \rangle + V_0.$$

Hence it follows from (2.15) that the coefficients  $b_n^{(\pm)}(x, \hat{\xi})$  are determined by formulas  $b_0^{(\pm)} = 1$  and

$$b_{n+1}^{(\pm)}(x, \hat{\xi}) = \mp \int_0^{\infty} f_n^{(\pm)}(x \pm t\hat{\xi}, \hat{\xi}) dt, \quad (2.21)$$

where

$$\begin{aligned} f_n^{(\pm)} &= 2i\langle A - \nabla \Phi_{\pm}, \nabla b_n^{(\pm)} \rangle - \Delta b_n^{(\pm)} \\ &+ (|\nabla \Phi_{\pm}|^2 - 2\langle A, \nabla \Phi_{\pm} \rangle + V_1 - i\Delta \Phi_{\pm}) b_n^{(\pm)}. \end{aligned} \quad (2.22)$$

Let us formulate the result obtained.

**Proposition 2.8** *Let  $A$  and  $V$  satisfy assumption (1.7) with  $\rho > 1$ , and let  $\rho_2 = \min\{2, \rho\}$ . Define  $\Theta(x, \xi)$  by formulas (2.4) and (2.20). Let the functions  $b_n^{(\pm)}$  be constructed by recurrent formulas (2.21), (2.22) and let  $b_{\pm}$  be the sum (1.4). Then estimates (2.18) on  $b_n^{(\pm)}$  and (2.19) on the remainder (1.10) hold.*

### 3. WAVE OPERATORS AND THE SCATTERING MATRIX

1. Let us recall briefly some basic facts about PDO (see, e.g., [6] or [20]). Let

$$(Af)(x) = (2\pi)^{-d/2} \int_{\mathbb{R}^d} e^{i\langle x, \xi \rangle} a(x, \xi) \hat{f}(\xi) d\xi,$$

where  $\hat{f} = \mathcal{F}f$  is the Fourier transform of  $f$  from, say, the Schwartz space  $\mathcal{S}(\mathbb{R}^d)$  and the symbol  $a \in C^\infty(\mathbb{R}^d \times \mathbb{R}^d)$ . Sometimes it is more convenient to consider more general PDO determined by their amplitudes. We define such operators in terms of the corresponding sesquilinear forms

$$(\mathbf{A}f, g) = (2\pi)^{-d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} e^{i\langle x, \xi' - \xi \rangle} \mathbf{a}(x, \xi, \xi') \hat{f}(\xi') \overline{\hat{g}(\xi)} d\xi d\xi' dx, \quad (3.1)$$

where the amplitude  $\mathbf{a}(x, \xi, \xi')$  is also a  $C^\infty$ -function of all its variables.

It is standard to assume that  $a$  and  $\mathbf{a}$  belong to Hörmander classes. Set  $\langle x \rangle = (1 + |x|^2)^{1/2}$ ,  $\langle \xi \rangle = (1 + |\xi|^2)^{1/2}$ . By definition, the symbol  $a$  (or the corresponding operator  $A$ ) belongs to the class  $\mathcal{S}^{n,m}(\rho, \delta)$ ,  $\rho > 0$ ,  $\delta < 1$ , if for all multi-indices  $\alpha$  and  $\beta$

$$|(\partial_x^\alpha \partial_\xi^\beta a)(x, \xi)| \leq C_{\alpha, \beta} \langle x \rangle^{n - |\alpha| \rho + |\beta| \delta} \langle \xi \rangle^{m - |\beta|}.$$

The operators  $A$  from these classes send the Schwartz space  $\mathcal{S}(\mathbb{R}^d)$  into itself. For the amplitudes  $\mathbf{a}$  we do not have to keep track of the dependence on  $\xi$  and  $\xi'$ . Thus,  $\mathbf{a} \in \mathcal{S}^n(\rho, \delta)$  if for all multi-indices  $\alpha, \beta, \beta'$ , any compact set  $K \subset \mathbb{R}^d$  and  $\xi, \xi' \in K$

$$|(\partial_x^\alpha \partial_\xi^\beta \partial_{\xi'}^{\beta'} \mathbf{a})(x, \xi, \xi')| \leq C_{\alpha, \beta, \beta'}(K) \langle x \rangle^{n - |\alpha| \rho + (|\beta| + |\beta'|) \delta}.$$

Under this assumption the form (3.1) is well-defined as an oscillating integral for  $\hat{f}, \hat{g} \in C_0^\infty(\mathbb{R}^d)$ . We omit in notation  $\rho$  and  $\delta$  if  $\rho = 1$  and  $\delta = 0$ .

Actually, we need a more special class of PDO with oscillating symbols

$$a(x, \xi) = e^{i\Phi(x, \xi)} \alpha(x, \xi), \quad (3.2)$$

where  $\Phi \in \mathcal{S}^{1-\rho, 0}$ ,  $\rho \in (0, 1)$ , and  $\alpha \in \mathcal{S}^{n,m}$ . We denote by  $\mathcal{C}^{n,m}(\Phi)$  the class of symbols or operators obeying the conditions above. The definition of the class  $\mathcal{C}^n(\Phi)$  in the case of oscillating amplitudes is quite similar. Since  $\mathcal{C}^{n,m}(\Phi) \subset \mathcal{S}^{n,m}(\rho, 1 - \rho)$ , the standard PDO calculus works in the classes  $\mathcal{C}^{n,m}(\Phi)$  if  $\rho > 1/2$ . In the general case the oscillating factors  $\exp(i\Phi(x, \xi))$  or  $\exp(i\Phi(x, \xi, \xi'))$  should be explicitly taken into account.

The proof of the following assertion can be found either in [13] or [25]. We often use the notation  $\langle x \rangle$  and  $\langle \xi \rangle$  for the operators of multiplication by these functions in the coordinate and momentum representations, respectively.

**Proposition 3.1** *Let  $a \in \mathcal{C}^{n,m}(\Phi)$ ,  $n \leq 0$  and  $m \leq 0$ . Then the operator  $A \langle x \rangle^{-n}$  is bounded in the space  $L_2(\mathbb{R}^d)$ .*

This result extends naturally to PDO defined by (3.1).

We need also a class  $\Xi_{\pm}$  of symbols such that  $a(x, \xi) = 0$  if  $\mp \langle \hat{x}, \hat{\xi} \rangle \leq \varepsilon$  for some  $\varepsilon > 0$ . Moreover, we assume that  $a(x, \xi) = 0$  if  $|x| \leq \varepsilon$  or  $|\xi| \leq \varepsilon$  for symbols from this class. Then we set

$$\mathcal{S}_{\pm}^{n,m}(\rho, \delta) = \mathcal{S}^{n,m}(\rho, \delta) \cap \Xi_{\pm}, \quad \mathcal{C}_{\pm}^{n,m}(\Phi) = \mathcal{C}^{n,m}(\Phi) \cap \Xi_{\pm}.$$

2. Let  $H_0 = -\Delta$  and the operator  $H$  defined by (1.6) act in the space  $\mathcal{H} = L_2(\mathbb{R}^d)$ . Denote by  $E_0$  and  $E$  their spectral projections. Note that, as shown in [8, 22] where the proof of [18] was extended to magnetic potentials, the operator  $H$  does not have positive eigenvalues. In the long-range case the wave operators (1.14) exist only for a special choice of identifications  $J_{\pm}$ . We construct  $J_{\pm}$  as PDO.

Let  $\sigma \in C^{\infty}(-\gamma, \gamma)$ ,  $\gamma > 1$ , be such that  $\sigma(\tau) = 1$  if  $\tau \in [-1, 1 - 2\varepsilon]$  for some  $\varepsilon \in (0, 1/2)$  and  $\sigma(\tau) = 0$  if  $\tau \in [1 - \varepsilon, 1]$ . Let  $\eta \in C^{\infty}(\mathbb{R}^d)$  be such that  $\eta(x) = 0$  in a neighborhood of zero and  $\eta(x) = 1$  for large  $|x|$ . We denote by  $\vartheta$  a  $C^{\infty}(\mathbb{R}_+)$ -function which equals to zero in a neighborhood of 0 and  $\vartheta(\lambda) = 1$  for, say,  $\lambda \geq \lambda_0$  (for some  $\lambda_0 > 0$ ). Set  $\zeta_{\pm}(x, \xi) = \sigma(\mp \eta(x) \langle \hat{\xi}, \hat{x} \rangle) \vartheta(|\xi|^2)$ .

Let  $u_{\pm}(x, \xi)$  be the function (it depends on  $N_0$  and  $N$ ) defined in the previous section (see Theorem 2.6). Following [9], we construct  $J_{\pm}$  by the formula

$$(J_{\pm}f)(x) = (2\pi)^{-d/2} \int_{\mathbb{R}^d} u_{\pm}(x, \xi) \zeta_{\pm}(x, \xi) \hat{f}(\xi) d\xi. \quad (3.3)$$

Thus,  $J_{\pm}$  is a PDO with symbol (3.2) where  $\Phi = \Phi_{\pm}$  and  $\alpha_{\pm}(x, \xi) = b_{\pm}(x, \xi) \zeta_{\pm}(x, \xi)$ . We emphasize however that in contrast to [9] the symbol  $a_{\pm}(x, \xi)$  of the operator  $J_{\pm}$  is quite an explicit function. This is essential for construction of the asymptotic expansion of the SM. Due to the cut-off functions  $\zeta_{\pm}(x, \xi)$  and estimates (2.17) on  $b_{\pm}(x, \xi)$ , we have that  $\alpha_{\pm} \in \mathcal{S}^{0,0}$ . The function  $\Phi_{\pm}(x, \xi)$  is of course singular on the set  $\hat{x} = \mp \hat{\xi}$  but satisfies the estimates of the class  $\mathcal{S}^{1-\rho,0}$  on the support of  $\zeta_{\pm}$ . Abusing somewhat terminology, we write  $J_{\pm} \in \mathcal{C}^{0,0}(\Phi_{\pm})$ . By Proposition 3.1, the operator  $J_{\pm}$  is bounded.

It is shown in [9, 23, 19] that the wave operators (1.14) exist which implies the intertwining property  $W_{\pm}(H, H_0; J_{\pm})H_0 = HW_{\pm}(H, H_0; J_{\pm})$ . Moreover, they are isometric on the subspace  $E_0(\lambda_0, \infty)\mathcal{H}$  and are complete, that is

$$\text{Ran}(W_{\pm}(H, H_0; J_{\pm})E_0(\lambda_0, \infty)) = E(\lambda_0, \infty)\mathcal{H}.$$

In the short-range case

$$s - \lim_{t \rightarrow \pm\infty} (J_{\pm} - \vartheta(H_0))e^{-iH_0t} = 0,$$

so that the wave operators  $W_{\pm}(H, H_0; J_{\pm})$  coincide with the usual wave operators  $W_{\pm}(H, H_0)$  (times  $\vartheta(H_0)$ ). The scattering operator is defined by the standard relation

$$\mathbf{S} = \mathbf{S}(H, H_0; J_+, J_-) = W_+^*(H, H_0; J_+)W_-(H, H_0; J_-).$$

It commutes with the operator  $H_0$  and is unitary on the space  $E_0(\lambda_0, \infty)\mathcal{H}$ .

3. Let us calculate the perturbation (1.15). According to (1.10), we have that

$$\begin{aligned} g_{\pm}(x, \xi) &:= (-\Delta + 2i\langle A(x), \nabla \rangle + V_1(x) - |\xi|^2)(u_{\pm}(x, \xi) \zeta_{\pm}(x, \xi)) \\ &= q_{\pm}(x, \xi) \zeta_{\pm}(x, \xi) - 2\langle \nabla u_{\pm}(x, \xi), \nabla \zeta_{\pm}(x, \xi) \rangle \\ &\quad - u_{\pm}(x, \xi) (\Delta \zeta_{\pm})(x, \xi) + 2i u_{\pm}(x, \xi) \langle A(x), \nabla \zeta_{\pm}(x, \xi) \rangle. \end{aligned} \quad (3.4)$$

Now it follows from (3.3) that

$$\begin{aligned}
(T_{\pm}f)(x) &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} g_{\pm}(x, \xi) \hat{f}(\xi) d\xi \\
&= (2\pi)^{-d/2} \int_{\mathbb{R}^d} e^{i(x, \xi)} (t_{\pm}^{(r)}(x, \xi) + t_{\pm}^{(s)}(x, \xi)) \hat{f}(\xi) d\xi \\
&= : (T_{\pm}^{(r)}f)(x) + (T_{\pm}^{(s)}f)(x), \tag{3.5}
\end{aligned}$$

where  $t_{\pm}^{(r)} = \exp(i\Phi_{\pm})\tau_{\pm}^{(r)}$ ,  $t_{\pm}^{(s)} = \exp(i\Phi_{\pm})\tau_{\pm}^{(s)}$  and

$$\tau_{\pm}^{(r)} = r_{\pm}\zeta_{\pm}, \quad \tau_{\pm}^{(s)} = -2ib_{\pm}\langle \xi + \nabla\Phi_{\pm} - A, \nabla\zeta_{\pm} \rangle - 2\langle \nabla b_{\pm}, \nabla\zeta_{\pm} \rangle - b_{\pm}\Delta\zeta_{\pm}.$$

Due to the cut-off functions  $\zeta_{\pm}$ ,  $\nabla\zeta_{\pm}$  and  $\Delta\zeta_{\pm}$ , the next result follows directly from Propositions 2.2 and 2.4.

**Proposition 3.2** *Let assumption (1.7) hold and let  $\rho_1 = \min\{1, \rho\}$ . Then*

$$t_{\pm}^{(r)} \in C^{-1-\rho_1(N+1), -N}(\Phi_{\pm}) \quad \text{and} \quad t_{\pm}^{(s)} \in C_{\pm}^{-1,1}(\Phi_{\pm}).$$

4. Let  $\mathfrak{H} = L_2(\mathbb{S}^{d-1})$ , let the operator  $\Gamma_0(\lambda) : \mathcal{S}(\mathbb{R}^d) \rightarrow \mathfrak{H}$  be defined by formula (1.1) and let  $(Uf)(\lambda) = \Gamma_0(\lambda)f$ . Then  $U : \mathcal{H} \rightarrow \mathcal{H} = L_2(\mathbb{R}_+; \mathfrak{H})$  extends by continuity to a unitary operator and  $UH_0U^*$  acts in the space  $\tilde{\mathcal{H}}$  as multiplication by the independent variable  $\lambda$ . Since  $SH_0 = H_0S$ , the operator  $USU^*$  acts in the space  $\mathcal{H}$  as multiplication by the operator-function  $S(\lambda) : \mathfrak{H} \rightarrow \mathfrak{H}$  known as the SM.

We need a stationary formula (see [10, 23, 24, 19]) for the SM  $S(\lambda)$  in the case where identifications  $J_+$  and  $J_-$  for  $t \rightarrow +\infty$  and  $t \rightarrow -\infty$  are different. Since auxiliary wave operators

$$s - \lim_{t \rightarrow \pm\infty} e^{iH_0t} J_{\pm}^* J_{\mp} e^{-iH_0t} = 0,$$

we have the following result.

**Proposition 3.3** *Let assumption (1.7) hold. Then the SM admits the representation (1.16) where  $S_1(\lambda)$  and  $S_2(\lambda)$  are given by formulas (1.17) and (1.18), respectively.*

Let us discuss here the precise meaning of the expression  $A^b(\lambda) := \Gamma_0(\lambda)A\Gamma_0^*(\lambda)$  where  $A$  is an operator acting on functions defined on  $\mathbb{R}^d$ . Put

$$\delta_{\varepsilon}(|\xi|^2 - \lambda) = \varepsilon\pi^{-1} \left( (|\xi|^2 - \lambda)^2 + \varepsilon^2 \right)^{-1},$$

and let  $\gamma_j \in C_0^{\infty}(\mathbb{R}_+)$  be an arbitrary function such that  $\gamma_j(k) = 1$ . Taking into account (1.1), we define (see, e.g., [24]) the sesquilinear form  $(A^b(\lambda)w_1, w_2)$  for  $w_j \in C^{\infty}(\mathbb{S}^{d-1})$  by the relation

$$(A^b(\lambda)w_1, w_2) = 2k^{-d+2} \lim_{\varepsilon \rightarrow 0} (A\mathcal{F}^*\delta_{\varepsilon}(|\xi|^2 - \lambda)\hat{\psi}_1, \mathcal{F}^*\delta_{\varepsilon}(|\xi|^2 - \lambda)\hat{\psi}_2), \tag{3.6}$$

where  $k = \lambda^{1/2}$ ,

$$\hat{\psi}_j(\xi) = w_j(\hat{\xi})\gamma_j(|\xi|), \quad j = 1, 2,$$

provided the limit in the right-hand side exists. The form  $(A^b(\lambda)w_1, w_2)$  is well defined if the limit (3.6) exists for all  $w_j \in C^{\infty}(\mathbb{S}^{d-1})$ . This is, of course, true if  $G = \mathcal{F}A\mathcal{F}^*$  is an

integral operator with kernel  $G(\xi, \xi')$  which is continuous near the surface  $|\xi| = |\xi'| = k$ . In this case  $A^b(\lambda)$  is also an integral operator on  $\mathbb{S}^{d-1}$  with kernel

$$g(\omega, \omega'; \lambda) = 2^{-1} k^{d-2} G(k\omega, k\omega'). \quad (3.7)$$

Furthermore, by the Sobolev trace theorem, limit (3.6) exists and hence the operator  $A^b(\lambda)$  is well-defined (and is bounded in the space  $L_2(\mathbb{S}^{d-1})$ ) if

$$A = \langle x \rangle^{-r} B \langle x \rangle^{-r} \quad (3.8)$$

for a bounded operator  $B$  in  $L_2(\mathbb{R}^d)$  and  $r > 1/2$ . This means that the operators  $A^b(\lambda)$  are also well-defined for PDO  $A$  of order  $n < -1$ .

We note that the stationary representation of the SM is determined exactly by the limits as the one in the right-hand side of (3.6).

To estimate in the next section the regular part  $S_2(\lambda)$  of the SM, we need the following obvious remark.

**Lemma 3.4** *Suppose that (3.8) is satisfied for  $r > d/2$ . Set  $u_0(x, \omega, \lambda) = \exp(i\lambda^{1/2}\langle \omega, x \rangle)$ . Then the operator  $A^b(\lambda)$  has continuous kernel*

$$g(\omega, \omega'; \lambda) = 2^{-1} k^{d-2} (2\pi)^{-d} (B \langle x \rangle^{-r} u_0(\omega', \lambda), \langle x \rangle^{-r} u_0(\omega, \lambda)).$$

Moreover, this function belongs to the class  $C^p(\mathbb{S}^{d-1} \times \mathbb{S}^{d-1})$  for  $p < r - d/2$  and its  $C^p$ -norm is bounded by  $Ck^{d-2+p}$ .

To treat the singular part  $S_1(\lambda)$ , we apply definition (3.6) to the PDO  $A = J_+^* T_-$  determined by its amplitude  $\mathbf{a}(x, \xi, \xi')$ . In this case, by Proposition 3.2,  $\mathbf{a}$  is of order  $-1$ , and hence the operators  $A^b(\lambda)$  are defined only under special assumptions on  $\mathbf{a}$ . According to (3.4), (3.5), up to an integral operator with smooth kernel,  $A$  has the amplitude which, due to the functions  $\nabla \zeta_-(x, \xi')$  and  $\Delta \zeta_-(x, \xi')$ , equals zero if  $\langle \hat{x}, \hat{\xi}' \rangle$  is close to 1 or  $-1$  (in a neighborhood of the conormal bundle of each sphere  $|\xi'| = k$ ). In this case, as shown in [25], the operators  $A^b(\lambda)$  are correctly defined by formula (3.6) in a space of functions on  $\mathbb{S}^{d-1}$  (the case of PDO determined by their symbols was considered earlier in [15]). Moreover, they are also PDO, and an explicit expression for their amplitudes was given in [25]. However, our construction of the singular part of the scattering matrix in Section 5 is, at least formally, independent of the results of [25]. It is important that this construction allows us to get rid of the cut-off functions  $\zeta_{\pm}$  and to obtain an arbitrary close approximation to the SM.

#### 4. THE REGULAR PART

In this section we show that the regular part (1.18) of the SM is negligible.

1. Recall that the functions  $u_{\pm} = u_{\pm}^{(N)}$  were constructed in Theorem 2.6 and that the corresponding operators  $J_{\pm} = J_{\pm}^{(N)}$  and  $T_{\pm} = T_{\pm}^{(N)}$  were defined by equations (3.3) and (3.5), respectively. Our main analytical result here is the following

**Proposition 4.1** For any  $p$  and  $q$  there exists  $N$  such that for  $T_{\pm} = T_{\pm}^{(N)}$  the operators

$$B_{p,q,N}(\lambda) = \langle x \rangle^p \langle \xi \rangle^q T_{+}^{*} R(\lambda + i0) T_{-} \langle \xi \rangle^q \langle x \rangle^p$$

are bounded uniformly in  $\lambda \geq \lambda_0 > 0$ .

This result will be checked in the following subsections. Let us first of all show that it implies regularity of the operator  $S_2(\lambda)$ .

**Theorem 4.2** For any  $p$  and  $q$  there exists  $N$  such that for  $T_{\pm} = T_{\pm}^{(N)}$ , the operator (1.18) has kernel  $s_2(\omega, \omega'; \lambda)$  which belongs to the class  $C^p(\mathbb{S}^{d-1} \times \mathbb{S}^{d-1})$  and the  $C^p$ -norm of this kernel is  $O(\lambda^{-q})$  as  $\lambda \rightarrow \infty$ .

Remark that  $\Gamma_0(\lambda) \langle \xi \rangle^{-q_0} = (1 + \lambda)^{-q_0/2} \Gamma_0(\lambda)$  and hence

$$S_2(\lambda) = 2\pi i (1 + \lambda)^{-q_0} \Gamma_0(\lambda) \langle x \rangle^{-p_0} B_{p_0, q_0, N}(\lambda) \langle x \rangle^{-p_0} \Gamma_0^{*}(\lambda).$$

Let  $p_0 > d/2 + p$  and  $q_0 \geq q - 1 + (d + p)/2$ . We suppose here that  $N = N(p_0, q_0)$  is the same as in Proposition 4.1, so that the operators  $B_{p_0, q_0, N}(\lambda)$  are bounded uniformly in  $\lambda \geq \lambda_0$ . Then, as shown in Lemma 3.4, the kernel of the operator  $S_2(\lambda)$  belongs to the class  $C^p(\mathbb{S}^{d-1} \times \mathbb{S}^{d-1})$ , and its  $C^p$ -norm is bounded by  $Ck^{d-2+p-2q_0}$  which is estimated by  $Ck^{-2q}$ . This concludes the proof of Theorem 4.2.

In the following subsections we shall give an idea of the proof of Proposition 4.1.

2. We need some results on the boundedness of combinations of PDO  $T$  with symbols  $t \in C_{\pm}^{n,m}(\Phi)$  (see subs. 1 of Section 3) where  $\Phi \in \mathcal{S}^{1-\rho,0}$  with functions of the generator of dilations

$$\mathbb{A} = \frac{1}{2} \sum_{j=1}^d (x_j D_j + D_j x_j).$$

We denote by  $\mathbb{P}_{\pm} = E_{\mathbb{A}}(\mathbb{R}_{\pm})$  the spectral projection of the operator  $\mathbb{A}$ .

First we formulate a strengthening of a result of [11].

**Proposition 4.3** Let  $t \in C_{\pm}^{n,m}(\Phi)$  for one of the signs and some  $n, m$ . Then there exists  $k$  such that the operator  $\langle \mathbb{A} \rangle^{-k} T$  is bounded.

Of course, this result is of interest only if at least one of the indices  $n$  or  $m$  is positive.

The following assertion is also motivated by the results of [11].

**Proposition 4.4** Let  $t \in \mathcal{S}_{\pm}^{n,m}(\rho, \delta)$  for some  $n, m$  and  $\rho > 0, \delta < 1$ . Then the operator  $\langle \mathbb{A} \rangle^k \mathbb{P}_{\pm} T \langle \xi \rangle^q \langle x \rangle^p$  is bounded for arbitrary  $p, q$  and  $k$ .

The following resolvent estimates were deduced in [17, 12, 11] from the famous Mourre estimate [16]. To obtain estimates at high energies, we use additionally the dilation transformation.

**Proposition 4.5** Let assumption (1.7) hold. Then for  $\operatorname{Re} z > 0, \operatorname{Im} z \geq 0$  the operator-functions

$$\langle \mathbb{A} \rangle^{-p} R(z) \langle \mathbb{A} \rangle^{-p}, \quad p > 1/2, \quad (4.1)$$

$$\langle \mathbb{A} \rangle^{-1+p_2} \mathbb{P}_{-} R(z) \langle \mathbb{A} \rangle^{-p_1}, \quad \langle \mathbb{A} \rangle^{-p_1} R(z) \mathbb{P}_{+} \langle \mathbb{A} \rangle^{-1+p_2} \quad (4.2)$$

for each  $p_1 > 1/2$ ,  $p_2 < p_1$  and

$$\langle \mathbf{A} \rangle^p \mathbf{P}_- R(z) \mathbf{P}_+ \langle \mathbf{A} \rangle^p \quad (4.3)$$

for arbitrary  $p$  are continuous in norm with respect to  $z$ . Moreover, the norms of the operators (4.1) – (4.3) at  $z = \lambda + i0$  are bounded by  $C\lambda^{-1}$  as  $\lambda \rightarrow \infty$ .

3. Now we are able to check Proposition 4.1. Let us first show that the operators

$$\langle x \rangle^p \langle \xi \rangle^q (T_+^{(r)})^* R(\lambda + i0) T_-^{(r)} \langle \xi \rangle^q \langle x \rangle^p$$

are uniformly bounded provided  $N$  is large enough. Note that the operators  $\langle x \rangle^\sigma T_\pm^{(r)} \langle \xi \rangle^q \langle x \rangle^p$  are bounded by Propositions 3.1 and 3.2 if  $(N+1)\rho_1 \geq \sigma + p - 1$  and  $N \geq q$ . Thus, it suffices to use that

$$\| \langle x \rangle^{-\sigma} R(\lambda + i0) \langle x \rangle^{-\sigma} \| = O(\lambda^{-1/2}), \quad \sigma > 1/2,$$

which follows, for example, from the result of Proposition 4.5 about operator (4.1).

Let us further consider the singular part  $T_\pm^{(s)}$  of  $T_\pm$ . Recall that, according to Proposition 3.2,  $T_\pm^{(s)} \in \mathcal{C}_\pm^{-1,1}(\Phi_\pm)$ . We need to prove the uniform boundedness of four operators

$$\langle x \rangle^p \langle \xi \rangle^q (T_+^{(s)})^* \mathbf{P}_- R(\lambda + i0) \mathbf{P}_+ T_-^{(s)} \langle \xi \rangle^q \langle x \rangle^p, \quad (4.4)$$

$$\langle x \rangle^p \langle \xi \rangle^q (T_+^{(s)})^* \mathbf{P}_+ R(\lambda + i0) \mathbf{P}_- T_-^{(s)} \langle \xi \rangle^q \langle x \rangle^p \quad (4.5)$$

and

$$\langle x \rangle^p \langle \xi \rangle^q (T_+^{(s)})^* \mathbf{P}_\pm R(\lambda + i0) \mathbf{P}_\pm T_-^{(s)} \langle \xi \rangle^q \langle x \rangle^p. \quad (4.6)$$

The operator (4.4) can be factorized into a product of three operators

$$\langle x \rangle^p \langle \xi \rangle^q (T_+^{(s)})^* \langle \mathbf{A} \rangle^{-k}, \langle \mathbf{A} \rangle^k \mathbf{P}_- R(\lambda + i0) \mathbf{P}_+ \langle \mathbf{A} \rangle^k \quad \text{and} \quad \langle \mathbf{A} \rangle^{-k} T_-^{(s)} \langle \xi \rangle^q \langle x \rangle^p.$$

The first and the third factors are bounded for sufficiently large  $k$  by Proposition 4.3 while the second operator has the form (4.3), and hence it is bounded by  $C\lambda^{-1}$  by Proposition 4.5.

The operator (4.5) can be factorized into a product of three operators

$$\langle x \rangle^p \langle \xi \rangle^q (T_+^{(s)})^* \mathbf{P}_+ \langle \mathbf{A} \rangle^\sigma, \langle \mathbf{A} \rangle^{-\sigma} R(\lambda + i0) \langle \mathbf{A} \rangle^{-\sigma} \quad \text{and} \quad \langle \mathbf{A} \rangle^\sigma \mathbf{P}_- T_-^{(s)} \langle \xi \rangle^q \langle x \rangle^p.$$

The first and the third factors are bounded for each  $\sigma$  by Proposition 4.4 while the second operator has the form (4.1), and hence it is bounded for any  $\sigma > 1/2$  by  $C\lambda^{-1}$  by Proposition 4.5.

Finally, we factorize the operator (4.6) (for the sign “+”, for example) into a product of three operators  $\langle x \rangle^p \langle \xi \rangle^q (T_+^{(s)})^* \mathbf{P}_+ \langle \mathbf{A} \rangle^\sigma$ ,  $\langle \mathbf{A} \rangle^{-\sigma} R(\lambda + i0) \mathbf{P}_+ \langle \mathbf{A} \rangle^{-1+\sigma-\varepsilon}$ ,  $\varepsilon > 0$ , and  $\langle \mathbf{A} \rangle^{1-\sigma+\varepsilon} T_-^{(s)} \langle \xi \rangle^q \langle x \rangle^p$ . The first factor is bounded for any  $\sigma$  by Proposition 4.4. The second operator has the form (4.2), and hence it is bounded for any  $\sigma > 1/2$  by  $C\lambda^{-1}$  by Proposition 4.5. The last factor is bounded by Proposition 4.3 if  $\sigma$  is sufficiently large.

The cross-terms containing  $T_+^{(r)}$  and  $T_-^{(s)}$  can be considered quite similarly. We need to prove the uniform boundedness of two operators  $\langle x \rangle^p \langle \xi \rangle^q (T_+^{(r)})^* R(\lambda + i0) \mathbf{P}_\tau T_-^{(s)} \langle \xi \rangle^q \langle x \rangle^p$ ,

where  $\tau = "+"$  or  $\tau = "-"$ . First, using Proposition 3.2, for any  $l$  we can choose  $N$  such that the operator  $\langle x \rangle^p \langle \xi \rangle^q (T_+^{(\tau)})^* \langle \mathbb{A} \rangle^l$  is bounded and hence it suffices to consider the operators  $\langle \mathbb{A} \rangle^{-l} R(\lambda + i0) \mathbb{P}_\tau T_-^{(s)} \langle \xi \rangle^q \langle x \rangle^p$ . If  $\tau = "-"$ , then these operators are uniformly bounded for any  $l > 1/2$  according to Proposition 4.4 and the estimate of Proposition 4.5 on the operator (4.1). If  $\tau = "+"$ , then according to Proposition 4.3 the operator  $\langle \mathbb{A} \rangle^{-k} T_-^{(s)} \langle \xi \rangle^q \langle x \rangle^p$  is bounded for sufficiently large  $k$ . So it remains to use that the operator  $\langle \mathbb{A} \rangle^{-l} R(\lambda + i0) \mathbb{P}_+ \langle \mathbb{A} \rangle^k$  has the form (4.2), and hence it is bounded by  $C\lambda^{-1}$  for  $l > k + 1$  by Proposition 4.5.

This concludes our sketch of the proof of Proposition 4.1 and hence of Theorem 4.2.

## 5. THE SINGULAR PART

1. Let us discuss the precise meaning of the formula (1.12). Recall that  $\omega_0 \in \mathbb{S}^{d-1}$  is an arbitrary point,  $\Pi = \Pi_{\omega_0}$  is the hyperplane orthogonal to  $\omega_0$  and  $\Omega_\pm = \Omega_\pm(\omega_0, \delta) \subset \mathbb{S}^{d-1}$  is determined by the condition  $\pm \langle \omega, \omega_0 \rangle > \delta > 0$ . The coordinates  $(z, y)$  in  $\mathbb{R}^d$  are defined by equation (1.11). Set

$$h_\pm(x, \xi) = e^{i\Phi_\pm(x, \xi)} b_\pm(x, \xi), \quad (5.1)$$

so that

$$u_\pm(x, \xi) = e^{i(x, \xi)} h_\pm(x, \xi).$$

Then (1.12) can be rewritten as

$$s_0(\omega, \omega'; \lambda) = (2\pi)^{-d+1} \int_{\Pi} e^{ik(y, \omega' - \omega)} \mathbf{a}_0(y, \omega, \omega'; \lambda) dy, \quad (5.2)$$

where  $\omega, \omega' \in \Omega_\pm$  and

$$\begin{aligned} \mathbf{a}_0(y, \omega, \omega'; \lambda) = & \pm 2^{-1} k^{d-2} \left( k \langle \omega + \omega', \omega_0 \rangle \overline{h_+(y, k\omega)} h_-(y, k\omega') \right. \\ & \left. + i h_-(y, k\omega') \overline{(\partial_z h_+)(y, k\omega)} - i \overline{h_+(y, k\omega)} (\partial_z h_-)(y, k\omega') - 2 \langle A(y), \omega_0 \rangle \overline{h_+(y, k\omega)} h_-(y, k\omega') \right) \end{aligned} \quad (5.3)$$

Formula (5.2) shows that  $S_0(\lambda)$  is, actually, regarded as a PDO with amplitude  $\mathbf{a}_0(y, \omega, \omega'; \lambda)$ . It is convenient to define the operator  $S_0(\lambda)$  via its sesquilinear form. Indeed, suppose, for example, that  $\omega \in \Omega = \Omega_+$  and denote by  $\Sigma$  and  $\zeta$  the orthogonal projections of  $\Omega$  and of a point  $\omega \in \Omega$  on the hyperplane  $\Pi$  which we identify with  $\mathbb{R}^{d-1}$ . We also identify below points  $\omega \in \Omega$  and  $\zeta \in \Sigma$  and functions

$$w(\omega) = \tilde{w}(\zeta) \quad (5.4)$$

on  $\Omega$  and  $\Sigma$ . Set

$$\tilde{\mathbf{a}}_0(y, \zeta, \zeta'; \lambda) = (1 - |\zeta|^2)^{-1/2} (1 - |\zeta'|^2)^{-1/2} \mathbf{a}_0(y, \omega, \omega'; \lambda).$$

Then it follows from (5.2) that for arbitrary  $w_j \in C_0^\infty(\Omega)$ ,  $j = 1, 2$ ,

$$(S_0(\lambda)w_1, w_2) = (2\pi)^{-d+1} \int_{\Pi} \int_{\Pi} \int_{\Pi} e^{ik(y, \zeta' - \zeta)} \tilde{\mathbf{a}}_0(y, \zeta, \zeta'; \lambda) \tilde{w}_1(\zeta') \overline{\tilde{w}_2(\zeta)} d\zeta d\zeta' dy. \quad (5.5)$$



Since  $\tilde{\mathbf{a}}_0 \in \mathcal{S}^0(\rho, 1 - \rho)$ , the right-hand side of the last equation is well-defined as an oscillating integral which gives the precise sense to its left-hand side. Of course, we can make the change of variables  $y \mapsto k^{-1}y$  in (5.5) transforming PDO  $S_0(\lambda)$  to the standard form, but this operation is not really necessary. It follows from (5.1) that amplitude (5.3) contains an oscillating factor  $\exp(i\Xi)$  where

$$\Xi(y, \omega, \omega'; k) = \Phi_-(y, k\omega') - \Phi_+(y, k\omega), \quad (5.6)$$

and hence the operator  $S_0(\lambda)$  is bounded according to Proposition 3.1.

2. It follows from Theorem 4.2 that the operator (1.17) contains all power terms of the high-energy expansion of the SM as well as of its diagonal singularity. However, the obvious drawback of the expression (1.17) is that it depends on the cut-off functions  $\zeta_{\pm}$ . Our final goal is to show that, up to negligible terms, it can be transformed to the invariant expression (1.12).

We proceed from relation (3.6) where  $\mathbf{A} = J_+^* T_-$ . Recall that  $J_+$  and  $T_-$  are PDO defined by formulas (3.3) and (3.4), (3.5), respectively. Therefore for all  $f_1, f_2 \in \mathcal{S}$

$$(T_- f_1, J_+ f_2) = (2\pi)^{-d} \int_{\mathbf{R}^d} \left( \int_{\mathbf{R}^d} \int_{\mathbf{R}^d} e^{i(x, \xi' - \xi)} \mathbf{a}(x, \xi, \xi') \hat{f}_1(\xi') \overline{\hat{f}_2(\xi)} d\xi d\xi' \right) dx, \quad (5.7)$$

where

$$\mathbf{a}(x, \xi, \xi') = \overline{j_+(x, \xi)} t_-(x, \xi') \quad (5.8)$$

and  $j_+, t_-$  are the symbols of the operators  $J_+, T_-$ , respectively. According to Propositions 2.2, 2.4 and 3.2, the amplitude  $\mathbf{a}(x, \xi, \xi')$  belongs to the Hörmander class  $\mathcal{S}^{-1}(\rho, 1 - \rho)$ . To obtain a convenient representation for (5.7), we have to change the order of integrations over  $x$  and  $\xi, \xi'$  in (5.7) and then calculate the integral over  $x$ . Below we do not go into details of standard manipulations with oscillating integrals. Note only that, strictly speaking, we have to introduce into (5.7) a function  $\varphi(\epsilon x)$  such that  $\varphi \in C_0^\infty(\mathbf{R}^d)$ ,  $\varphi(0) = 1$ , and pass to the limit  $\epsilon \rightarrow 0$  at the very end of our calculations. Denote

$$G(\xi, \xi') = \int_{\mathbf{R}^d} e^{i(x, \xi' - \xi)} \mathbf{a}(x, \xi, \xi') dx \quad (5.9)$$

and let  $G$  be integral operator with kernel  $G(\xi, \xi')$ . Then, at least formally,  $G = (2\pi)^d \mathcal{F} J_+^* T_- \mathcal{F}^*$ . We set  $\zeta = \zeta_-$ , then  $\zeta_+(x, \xi) = \zeta(x, -\xi)$ . It follows from (3.3), (3.5) and (5.8), (5.9) that

$$G(\xi, \xi') = \int_{\mathbf{R}^d} \overline{u_+(x, \xi)} \zeta(x, -\xi) g_-(x, \xi') dx. \quad (5.10)$$

Standard arguments show that, off the diagonal,  $G(\xi, \xi')$  is a smooth function, and it rapidly tends to zero as  $|\xi| \rightarrow \infty$  and  $|\xi'| \rightarrow \infty$ . Applying (3.6) to functions  $w_1$  and  $w_2$  with disjoint supports, it is easy to show that off the diagonal  $\omega = \omega'$  the kernel  $s_1(\omega, \omega', \lambda)$  of the operator  $S_1(\lambda)$  satisfies the relation (cf. (3.7))

$$s_1(\omega, \omega'; \lambda) = -\pi i k^{d-2} G(k\omega, k\omega'), \quad \omega \neq \omega'. \quad (5.11)$$

Combining these results with Theorem 4.2, we obtain

**Theorem 5.1** *Let assumption (1.7) hold, and let  $\omega \in \Omega$ ,  $\omega' \in \Omega'$  for some open sets  $\Omega, \Omega' \subset \mathbb{S}^{d-1}$  such that  $\text{dist}(\Omega, \Omega') > 0$ . Then for any  $p$  and  $q$  the kernel  $s(\omega, \omega', \lambda)$  of the SM belongs to the space  $C^p(\Omega \times \Omega')$  and its  $C^p$ -norm is bounded by  $C\lambda^{-q}$  as  $\lambda \rightarrow \infty$ .*

3. Our study of the function (5.10) in a neighborhood of the diagonal  $\xi = \xi'$  relies on integration by parts. Let us plug (3.4) into (5.10) and denote by  $G_j(\xi, \xi')$ ,  $j = 1, 2, 3, 4$ , the integrals corresponding to the four functions in the right-hand side of (3.4):

$$\begin{aligned} G_1(\xi, \xi') &= \int_{\mathbb{R}^d} \overline{u_+(x, \xi)} \zeta(x, -\xi) q_-(x, \xi') \zeta(x, \xi') dx, \\ G_2(\xi, \xi') &= -2 \int_{\mathbb{R}^d} \overline{u_+(x, \xi)} \zeta(x, -\xi) \langle \nabla u_-(x, \xi'), \nabla \zeta(x, \xi') \rangle dx, \\ G_3(\xi, \xi') &= - \int_{\mathbb{R}^d} \overline{u_+(x, \xi)} \zeta(x, -\xi) u_-(x, \xi') \Delta \zeta(x, \xi') dx, \\ G_4(\xi, \xi') &= 2i \int_{\mathbb{R}^d} \overline{u_+(x, \xi)} \zeta(x, -\xi) u_-(x, \xi') \langle A(x), \nabla \zeta(x, \xi') \rangle dx. \end{aligned}$$

Let us consider first the function  $G_1$  where  $q_- = e^{i\Theta} r_-$ . By virtue of Theorem 2.6, the function  $\overline{u_+(x, \xi)} \zeta(x, -\xi)$  satisfies estimates (2.17) for all  $x, \xi \in \mathbb{R}^d$  and the function  $r_-(x, \xi') \zeta(x, \xi')$  satisfies estimates (2.16) for all  $x, \xi' \in \mathbb{R}^d$ . Hence the integrand in  $G_1(\xi, \xi')$  is estimated by  $C|\xi|^{-N}(1+|x|)^{-1-\rho_1(N+1)}$ , where  $N$  can be chosen arbitrary large. Using also the estimates on derivatives of these functions and estimates (2.13) on the phase functions  $\Phi_{\pm}$ , we see that  $G_1(\xi, \xi')$  is a smooth function of  $\xi, \xi'$  rapidly decreasing as  $|\xi| = |\xi'| \rightarrow \infty$ .

Let  $\omega$  and  $\omega'$  belong to some conical neighborhood of a point  $\omega_1 \in \mathbb{S}^{d-1}$  where, for example,  $\langle \omega_1, \omega_0 \rangle > 0$ . Then  $\zeta(x, -\xi)(\nabla \zeta)(x, \xi') = (\nabla \zeta)(x, \xi')$  so that the function  $\zeta(x, -\xi)$  in the integrals  $G_j(\xi, \xi')$ ,  $j = 2, 3, 4$ , can be omitted. All these integrals will be transformed by integration by parts. Integrating in  $G_3(\xi, \xi')$  by parts, we find that

$$\begin{aligned} &G_2(\xi, \xi') + G_3(\xi, \xi') = \\ &+ \int_{\mathbb{R}^d} \langle u_-(x, \xi') (\overline{\nabla u_+}(x, \xi) - \overline{u_+(x, \xi)} (\nabla u_-)(x, \xi')), \nabla \zeta(x, \xi') \rangle dx. \end{aligned} \quad (5.12)$$

Due to the function  $\nabla \zeta(x, \xi')$ , the integrals (5.12) as well as  $G_4(\xi, \xi')$  are actually taken over the half-space  $z \geq 0$  only. Therefore integrating once more by parts and taking into account the equality  $\zeta(y, \xi') = 1$ , we obtain that

$$\begin{aligned} &G_2(\xi, \xi') + G_3(\xi, \xi') = \\ &+ \int_{z \geq 0} (\overline{u_+(x, \xi)} (\Delta u_-)(x, \xi') - u_-(x, \xi') (\overline{\Delta u_+}(x, \xi))) \zeta(x, \xi') dx \\ &+ \int_{\Pi} (\overline{u_+(y, \xi)} (\partial_z u_-)(y, \xi') - u_-(y, \xi') (\overline{\partial_z u_+}(y, \xi))) dy \end{aligned} \quad (5.13)$$

and

$$\begin{aligned} G_4(\xi, \xi') &= -2i \int_{z \geq 0} \text{div} (A(x) \overline{u_+(x, \xi)} u_-(x, \xi')) \zeta(x, \xi') dx \\ &\quad - 2i \int_{\Pi} \langle A(y), \omega_0 \rangle \overline{u_+(y, \xi)} u_-(y, \xi') dy. \end{aligned} \quad (5.14)$$

It is now convenient to formulate an intermediary result.

**Proposition 5.2** *The function (5.10) is the sum*

$$G = G_1 + G_2 + G_3 + G_4.$$

Here  $G_1(\xi, \xi')$  is a smooth function of  $\xi, \xi'$  rapidly decreasing as  $|\xi| = |\xi'| \rightarrow \infty$ . The functions  $G_2 + G_3$  and  $G_4$  satisfy equalities (5.13) and (5.14), respectively.

4. In the following we need to calculate the operators  $\mathbf{A}^b(\lambda)$  for two special classes of integral operators  $G = \mathcal{F}\mathbf{A}\mathcal{F}^*$  acting on functions of  $\xi \in \mathbb{R}^d$ . For the operators from the first class the passage to the limit (3.6) is quite direct (cf. (3.7)).

**Proposition 5.3** *Let an operator  $G$  be defined by its kernel*

$$G(\xi, \xi') = \int_{\Pi} e^{i(y, \xi' - \xi)} \mathbf{a}(y, \xi, \xi') dy,$$

where  $\mathbf{a} \in \mathcal{S}^p(\rho, \delta)$  for some  $p$  and  $\rho > 0$ ,  $\delta < 1$ . Then the operator  $\mathbf{A}^b(\lambda)$  exists for all  $\lambda > 0$  and is the integral operator on the unit sphere with kernel

$$g(\omega, \omega'; \lambda) = 2^{-1} k^{d-2} \int_{\Pi} e^{ik(y, \omega' - \omega)} \mathbf{a}(y, k\omega, k\omega') dy, \quad \omega, \omega' \in \Omega_{\pm}.$$

Kernels of the operators from the second class are defined in terms of integrals over a half-space.

**Proposition 5.4** *Let an operator  $G$  have kernel*

$$G(\xi, \xi') = (|\xi|^2 - |\xi'|^2) \int_{z \geq 0} e^{i(x, \xi' - \xi)} \mathbf{a}(x, \xi, \xi') dx, \quad (5.15)$$

where  $\mathbf{a} \in \mathcal{S}^p(\rho, \delta)$  for some  $p$  and  $\rho > 0$ ,  $\delta < 1$ . Assume moreover that

$$\mathbf{a}(x, \xi, \xi') = 0 \quad \text{if} \quad \langle \xi + \xi', x \rangle \geq c_0 |\xi + \xi'| |x| \quad (5.16)$$

for some  $c_0 \in (0, 1)$ . Then  $\mathbf{A}^b(\lambda) = 0$  for all  $\lambda > 0$ .

The proof relies on condition (5.16). Let  $\mathbf{A}_1 = \mathcal{F}^* G_1 \mathcal{F}$  where

$$G_1(\xi, \xi') = \int_{z \geq 0} e^{i(x, \xi' - \xi)} \mathbf{a}(x, \xi, \xi') dx.$$

Then the operator  $\mathbf{A}_1^b(\lambda)$  is well-defined (cf. [15, 25]) due to (5.16). Taking into account the factor  $|\xi|^2 - |\xi'|^2$  in (5.15), it is easy to show that  $\mathbf{A}^b(\lambda) = 0$ .

5. Now we are in a position to derive formula (1.12) for the singular part of the SM. To that end, we have to calculate the limit in the right-hand side of (3.6) for  $\mathbf{A} = J_+^* T_-$  and show that the expression obtained coincides, up to negligible terms, with the form  $-(2\pi i)^{-1} (S_0(\lambda) w_1, w_2)$ . Let us proceed from Proposition 5.2.

According to (3.7) the contribution of  $G_1$  to  $S_1(\lambda)$  is given by the expression  $-\pi i k^{d-2} \times G_1(k\omega, k\omega')$  which is a smooth function of  $\omega, \omega'$  and rapidly decays as  $k \rightarrow \infty$ . Hence this term can be neglected.

Let us further consider the integrals (5.13) and (5.14) over  $\Pi$ . By virtue of Proposition 5.3, the contribution of each integral to the kernel of  $S_1(\lambda)$  equals its value at  $\xi = k\omega$ ,  $\xi' = k\omega'$  times (compare with (5.11)) the numerical factor  $-\pi i k^{d-2} (2\pi)^{-d}$ . The sum of these expressions coincides with (1.12).

It remains to show that the sum of the integrals over the half-space  $z \geq 0$  in (5.13) and (5.14) is negligible. It follows from relation (1.10) that

$$\begin{aligned} & \overline{u_+(x, \xi)}(\Delta u_-)(x, \xi') - u_-(x, \xi')\overline{(\Delta u_+)(x, \xi)} - 2i \operatorname{div} \left( A(x) \overline{u_+(x, \xi)} u_-(x, \xi') \right) \\ &= \left( \overline{q_+(x, \xi)} u_-(x, \xi') - q_-(x, \xi') \overline{u_+(x, \xi)} \right) + (|\xi|^2 - |\xi'|^2) \overline{u_+(x, \xi)} u_-(x, \xi'). \end{aligned}$$

To consider the integral

$$\int_{z \geq 0} e^{i\Theta_-(x, \xi') - i\Theta_+(x, \xi)} \left( \overline{r_+(x, \xi)} b_-(x, \xi') - r_-(x, \xi') \overline{b_+(x, \xi)} \right) \zeta(x, \xi') dx, \quad (5.17)$$

we use again that, by Proposition 2.4 and Corollary 2.5, the functions  $r_-(x, \xi') \zeta(x, \xi')$  and  $b_-(x, \xi') \zeta(x, \xi')$  satisfy estimates (2.16) and (2.17), respectively, for all  $x, \xi' \in \mathbb{R}^d$ . The same result for the functions  $b_+(x, \xi)$  and  $r_+(x, \xi)$  holds true in the half-space  $z \geq 0$  which does not contain the “bad” direction  $\hat{x} = -\hat{\xi}$ . By Corollary 2.3, the function  $\Phi_-(x, \xi') - \Phi_+(x, \xi)$  satisfies estimates (2.13) for all  $z \geq 0$  off a conical neighborhood of the direction  $\hat{x} = \hat{\xi}'$  where  $\zeta(x, \xi') = 0$ . Therefore the integral (5.17) is a smooth function of  $\xi, \xi'$  rapidly decreasing as  $|\xi| = |\xi'| \rightarrow \infty$ . Hence, similarly to the function  $G_1(\xi, \xi')$ , this integral does not contribute to  $S_0(\lambda)$ .

Let us, finally, consider the kernel

$$G_0(\xi, \xi') = (|\xi|^2 - |\xi'|^2) \int_{z \geq 0} e^{i(x, \xi' - \xi)} \overline{h_+(x, \xi)} h_-(x, \xi') \zeta(x, \xi') dx,$$

where the functions  $h_{\pm}(x, \xi)$  are defined by formula (5.1). Due to the factor  $\zeta(x, \xi')$ , the function  $G_0(\xi, \xi')$  satisfies the conditions of Proposition 5.4 and hence  $(\mathcal{F}^* G_0 \mathcal{F})^b(\lambda) = 0$  for all  $\lambda > 0$ .

Now we can formulate our main result on the asymptotics of the kernel  $s(\omega, \omega'; \lambda)$  of the SM.

**Theorem 5.5** *Let assumption (1.7) hold, let  $p, q$  be arbitrary numbers and  $N = N(p, q)$  be sufficiently large. Let functions  $\Theta_{\pm}^{(N_0)}(x, \xi)$  and  $b_{\pm}^{(N)}(x, \xi)$  be constructed in Propositions 2.2 and 2.4, respectively, and let  $u_{\pm}^{(N)}(x, \xi)$  be defined by formula (1.9). Define, for  $\omega, \omega' \in \Omega_{\pm}$ , the kernel  $s_0^{(N)}(\omega, \omega'; \lambda)$  by formula (1.12). Then the remainder (1.13) belongs to the class  $C^p(\Omega \times \Omega)$  and the  $C^p$ -norm of this kernel is  $O(\lambda^{-q})$  as  $\lambda \rightarrow \infty$ .*

This result gives simultaneously the high-energy and smoothness expansion of the kernel of the SM. As was already mentioned, we actually formulate the result in terms of the corresponding amplitude  $\mathbf{a}_0(y, \omega, \omega'; \lambda)$  related to the kernel of the SM by formula (5.2). Indeed, it follows from (5.1), (5.3) and (5.6) that

$$\mathbf{a}_0(y, \omega, \omega'; \lambda) = \pm 2^{-1} k^{d-1} \exp(i\Xi(y, \omega, \omega'; k)) \sum_{n=0}^N (2ik)^{-n} \sigma_n(y, \omega, \omega'),$$

$$\Xi(y, \omega, \omega'; k) = \sum_{n=0}^{N_0} (2k)^{-n} \theta_n(y, \omega, \omega'), \quad \theta_n(y, \omega, \omega') = \phi_n^{(-)}(y, \omega') - \phi_n^{(+)}(y, \omega)$$

and the functions  $\phi_n^{(\pm)}$  are constructed in Proposition 2.2. Note that  $\theta_0 \in \mathcal{S}^{1-\rho_a}$  and  $\theta_n \in \mathcal{S}^{1-n\rho}$  for  $n \geq 1$ . The coefficients  $\sigma_n(y, \omega, \omega')$  are expressed in terms of functions  $\phi_n^{(\pm)}$  and  $b_n^{(\pm)}$  constructed in Proposition 2.4. It is easy to see that  $\sigma_n \in \mathcal{S}^{-n\rho_1}$  for  $n \geq 0$ . In particular,  $S_0(\lambda) \in C^0(\Xi)$ .

## REFERENCES

- [1] S. Agmon, Some new results in spectral and scattering theory of differential operators in  $\mathbb{R}^n$ , *Seminaire Goulaouic Schwartz, Ecole Polytechnique*, 1978.
- [2] S. Arians, Geometric approach to inverse scattering for the Schrödinger equation with magnetic and electric potentials, *J. Math. Phys.* **38** (1997), 2761-2773.
- [3] V. S. Buslaev, Trace formulas and certain asymptotic estimates of the resolvent kernel for the Schrödinger operator in three-dimensional space, in *Topics in Math. Phys.* **1**, Plenum Press, 1967.
- [4] V. S. Buslaev, Dispersion relations in nonrelativistic scattering theory with spin-orbit interaction, *Soviet Physics JETP* **12** (1960), 990-992.
- [5] V. Enss, R. Weder, The geometrical approach to multidimensional inverse scattering, *J. Math. Phys.* **36** (1995), 3902-3921.
- [6] L. Hörmander, *The Analysis of Linear Partial Differential Operators III*, Springer-Verlag, 1985.
- [7] W. Hunziker, Potential scattering at high energies, *Helv. Phys. Acta* **36** (1963), 838-856.
- [8] T. Ikebe, J. Uchiyama, On the asymptotic behavior of eigenfunctions of second-order elliptic operators, *J. Math. Kyoto Univ.* **11** (1971), 425-448.
- [9] H. Isozaki, H. Kitada, Modified wave operators with time-independent modifies, *J. Fac. Sci, Univ. Tokyo*, **32** (1985), 77-104.
- [10] H. Isozaki, H. Kitada, Scattering matrices for two-body Schrödinger operators, *Sci. Papers College Arts and Sci., Univ. Tokyo*, **35** (1985), 81-107.
- [11] A. Jensen, Propagation estimates for Schrödinger-type operators, *Trans. Amer. Math. Soc.* **291** (1985), 129-144.
- [12] A. Jensen, E. Mourre, P. Perry, Multiple commutator estimates and resolvent smoothness in quantum scattering theory, *Ann. Inst. Henri Poincaré, Phys. théor.* **41** (1984), 207-225.

- [13] H. Kitada, Micro-local resolvent estimates for 2-body Schrödinger operators, *J. Math. Soc. Japan* **39** (1987), 455-476.
- [14] L. D. Landau and E. M. Lifshitz, *Quantum mechanics*, Pergamon Press, 1965.
- [15] N. Lerner, D. Yafaev, Trace theorems for pseudo-differential operators, *J. Analyse Math.* **74** (1998), 113-164.
- [16] E. Mourre, Absence of singular continuous spectrum for certain self-adjoint operators, *Comm. Math. Phys.* **78** (1981), 391-408.
- [17] E. Mourre, Opérateurs conjugués et propriétés de propagation, *Comm. Math. Phys.* **91** (1983), 279-300.
- [18] M. Reed and B. Simon, *Methods of Modern Mathematical Physics IV*, Academic Press, 1978.
- [19] P. Roux and D. R. Yafaev, The scattering matrix for the Schrödinger equation with a long-range electromagnetic potential, Preprint IRMAR, Rennes (2002).
- [20] M. A. Shubin, *Pseudodifferential Operators and Spectral Theory*, Springer-Verlag, 1987.
- [21] M. M. Skriganov, Uniform coordinate and spectral asymptotics for solutions of the scattering problem for the Schrödinger equation, *J. Soviet Math.* **10**, no. 1 (1978), 120-141.
- [22] J. Uchiyama, Polynomial growth or decay of eigenfunctions of second-order elliptic operators, *Publ. RIMS, Kyoto Univ.*, **23** (1987), 975-1006.
- [23] D. R. Yafaev, The scattering amplitude for the Schrödinger equation with a long-range potential, *Comm. Math. Phys.* **191** (1998), 183-218.
- [24] D. R. Yafaev, Scattering theory: some old and new problems, *Lect. Notes Math.* **1735**, Springer-Verlag, 2000.
- [25] D. R. Yafaev, A class of pseudo-differential operators with oscillating symbols, *St. Petersburg Math. J.* **11** no. 2 (2000).