

|             |  |
|-------------|--|
| Title       | Variants of subtlety in $SP_{\kappa\lambda}$ : comparison of ideals defined by combinatorial principles (Studies in Relative Consistency Proofs with Particular Emphasis on Set Theoretic Methods) |
| Author(s)   | Abe, Yoshihiro   |
| Citation    | 数理解析研究所講究録 (2003), 1304: 47-66   |
| Issue Date  | 2003-02  |
| URL         | <a href="http://hdl.handle.net/2433/42767">http://hdl.handle.net/2433/42767</a>  |
| Right       |  |
| Type        | Departmental Bulletin Paper  |
| Textversion | publisher  |

# Variants of subtlety in $P_\kappa\lambda$

— comparison of ideals defined by combinatorial principles —

2002. 10. 10

神奈川大学工学部 阿部 吉弘 (Yoshihiro Abe)<sup>1</sup>

Dept. of Math., Faculty of Engineering,

Kanagawa University

## 概要

A type of subtlety for  $P_\kappa\lambda$  called “ $A$ -subtle” is presented and compared with Menas’ notion “ $M$ -subtle”. Using it we prove almost ineffability is consistencywise stronger than Shelah property although  $P_\kappa\lambda$  is  $A$ -subtle for every  $\lambda \geq \kappa$  if  $\kappa$  is subtle. The following are also shown (i) “ $\{x \in P_\kappa\lambda : x \cap \kappa < |x|\}$  is  $A$ -subtle” has rather strong consequences. (ii) The subtle ideals are not  $\lambda$ -saturated, and completely ineffable ideal is not precipitous. (iii)  $NAIn_{\kappa,\lambda} = NIn_{\kappa,\lambda}$  and  $I_{\kappa,\lambda}$  does not have the partition property if  $\lambda^{<\kappa} = 2^\lambda$ . (iv) We can not prove “ $\kappa$  is  $\lambda^{<\kappa}$ -ineffable whenever  $\kappa$  is  $\lambda$ -ineffable”.

## 1 Notations and basic facts

Throughout this paper  $\kappa$  denotes a regular uncountable cardinal and  $\lambda$  a cardinal  $\geq \kappa$ . Let  $A$  be a set and  $a$  a set of ordinals with  $|a| \leq |A|$ . For any such pair  $(a, A)$ ,  $P_a A$  denotes the set  $\{x \subset A : |x| < |a|\}$ . Thus  $P_\kappa\lambda$  denotes the set  $\{x \subset \lambda : |x| < \kappa\}$  and, for  $x \in P_\kappa\lambda$ ,  $P_{x \cap \kappa} x = \{s \subset x : |s| < |x \cap \kappa|\}$ .

Combinatorial properties originated for regular uncountable cardinals have been translated into  $P_\kappa\lambda$  in [10], [6] and [7]. For  $x \in P_\kappa\lambda$ ,  $\hat{x}$  denotes the set  $\{y \in$

---

<sup>1</sup>2000 *Mathematical Subject Classification*: Primary 03E. Reserach partially supported by “Grant-in-Aid for Scientific research (C), The Ministry of Education, Science, Sports and Culture of Japan 09640299, and Japan Society for the Promotion of Science 14540142”

$P_\kappa\lambda : x \subset y$ . We say  $X \subset P_\kappa\lambda$  is *unbounded* if  $X \cap \hat{x} \neq \emptyset$  for all  $x \in P_\kappa\lambda$ . Let  $I_{\kappa\lambda} = \{X \subset P_\kappa\lambda : X \text{ is not unbounded}\}$ .

We say  $I$  is an ideal on  $P_\kappa\lambda$  if the following hold:

1.  $I \subset PP_\kappa\lambda$ ,
2.  $\emptyset \in I$  and  $P_\kappa\lambda \notin I$ ,
3.  $I_{\kappa,\lambda} \subset I$ ,
4.  $I$  is  $\kappa$ -complete;  $\cup X \in I$  for any  $X \subset I$  with  $|X| < \kappa$ .

Thus  $I_{\kappa,\lambda}$  is the minimal ideal on  $P_\kappa\lambda$ .

We say  $X \subset P_\kappa\lambda$  is *closed* if  $\cup D \in X$  for any  $\subset$ -chain  $D \subset X$  with  $|D| < \kappa$ .  $X$  is called *club* if it is closed and unbounded.

**Fact 1.1.** *Let  $X \subset P_\kappa\lambda$ . Then,  $X$  is club if and only if there exists  $f : \lambda \times \lambda \rightarrow \lambda$  such that  $C_f := \{x \in P_\kappa\lambda : f''(x \times x) \subset x \text{ and } x \cap \kappa \in \kappa\} \subset X$ .*

We say  $X$  is *stationary* if  $X \cap C \neq \emptyset$  for any club  $C$ . Let  $NS_{\kappa,\lambda} = \{X \subset P_\kappa\lambda : X \text{ is nonstationary}\}$ . Let  $I^+ = PP_\kappa\lambda \setminus I$ . For  $X \in I^+$   $I \upharpoonright X$  denotes the set  $\{Y \subset P_\kappa\lambda : Y \cap X \in I\}$ , which is an ideal on  $P_\kappa\lambda$  extending  $I$ . We say an ideal  $I$  is *normal* if  $I$  is closed under diagonal unions;  $\nabla X := \{x \in P_\kappa\lambda : x \in \cup\{X_\alpha : \alpha \in x\}\} \in I$  for any  $X = \{X_\alpha : \alpha < \lambda\} \subset I$ . Note that  $I$  is normal if and only if every regressive function on  $X \in I^+$  is constant on some  $Y \in P(X) \cap I^+$ , where a function  $f$  is said to be *regressive* if  $f(x) \in x$  for any  $x$  in  $\text{dom}(f) \setminus \{\emptyset\}$ .

The relation  $\prec$  is defined by  $y \prec z$  if  $y \in P_{z \cap \kappa} z$ . An ideal  $I$  is *strongly normal* if for any  $X \in I^+$  and  $f : X \rightarrow P_\kappa\lambda$  such that  $f(x) \prec x$  for all  $x \in X$  there exists  $Y \in P(X) \cap I^+$  such that  $f \upharpoonright Y$  is constant. This is equivalent to the following: for any  $\{X_s : s \in P_\kappa\lambda\} \subset I$ ,  $\nabla_{\prec} X_s := \{x : x \in \cup\{X_s : s \prec x\}\} \in I$ . Clearly every strongly normal ideal is normal. A filter  $F$  on  $P_\kappa\lambda$  and an ideal  $I$  on  $P_\kappa\lambda$  are *dual* to each other if the following holds:

$$X \in F \text{ if and only if } P_\kappa\lambda \setminus X \in I \text{ for every } X \subset P_\kappa\lambda.$$

The dual filter of  $I$  will be denoted by  $I^*$ .

For  $f : P_\kappa\lambda \rightarrow P_\kappa\lambda$ , let  $C_f = \{x \in P_\kappa\lambda : f''P_{x \cap \kappa} x \subset P_{x \cap \kappa} x\}$ . We define an ideal  $WNS_{\kappa,\lambda}$  by:

$$X \in WNS_{\kappa,\lambda} \text{ if and only if } X \subset P_\kappa\lambda \text{ and } X \cap C_f = \emptyset \text{ for some } f : P_\kappa\lambda \rightarrow P_\kappa\lambda.$$

The following are well-known [5], [8] and [1].

- Fact 1.2.** (1)  $NS_{\kappa,\lambda}$  is the minimal normal ideal on  $P_\kappa\lambda$ .  
(2)  $WNS_{\kappa,\lambda}$  is the minimal strongly normal ideal on  $P_\kappa\lambda$  and  $NS_{\kappa,\lambda} \subsetneq WNS_{\kappa,\lambda}$ .  
(3)  $WNS_{\kappa,\lambda}$  is proper if and only if  $\kappa$  is Mahlo or  $\kappa = \nu^+$  with  $\nu^{<\nu} = \nu$ .  
(4) If  $\kappa$  is Mahlo, then  $\{x \in P_\kappa\lambda : x \cap \kappa = |x \cap \kappa|\}$  is inaccessible  $\in WNS_{\kappa,\lambda}^*$ .  
(5) If  $h : P_\kappa\lambda \rightarrow \lambda$  is a bijection, then  $WNS_{\kappa,\lambda} = NS_{\kappa,\lambda} \upharpoonright \{x : h \text{``} P_{x \cap \kappa} x = x\}$ .  
(6) If  $\{s_\alpha : \alpha < \lambda^{<\kappa}\}$  is an enumeration of  $P_\kappa\lambda$  and  $f : P_\kappa\lambda \rightarrow P_\kappa\lambda^{<\kappa}$  is defined by  $f(x) = \{\alpha : s_\alpha \prec x\}$ , then  $f_*(WNS_{\kappa,\lambda}) := \{X \subset P_\kappa\lambda^{<\kappa} : f^{-1}(X) \in WNS_{\kappa,\lambda}\} = WNS_{\kappa,\lambda^{<\kappa}}$  and  $\{x \in P_\kappa\lambda : f(x) \cap \lambda = x\} \in WNS_{\kappa,\lambda}^*$ .

All the notions defined above for  $P_\kappa\lambda$  can be naturally translated into  $P_{x \cap \kappa} x$  if  $x \cap \kappa$  is regular uncountable. For instance,  $X \subset P_{x \cap \kappa} x$  is unbounded if for any  $y \in P_{x \cap \kappa} x$  there is  $z \in X$  such that  $y \subset z$ , and  $I_{x \cap \kappa, x}$  denotes the set  $\{X \subset P_{x \cap \kappa} x : X \text{ is not unbounded in } P_{x \cap \kappa} x\}$  which is a  $x \cap \kappa$ -complete ideal on  $P_{x \cap \kappa} x$ .

First we observe a type of subtlety for  $P_\kappa\lambda$ :  $X \subset P_\kappa\lambda$  is *A-subtle* if for any  $\{S_x \subset P_{x \cap \kappa} x : x \in P_\kappa\lambda\}$  and  $C \in WNS_{\kappa,\lambda}^*$ , there exist  $y \prec z$  both in  $C \cap X$  such that  $S_y = S_z \cap P_{y \cap \kappa} y$ .

The following are shown in §2:

- Theorem 1.3.** (1)  $\kappa$  is subtle if and only if  $P_\kappa\lambda$  is *A-subtle* for every  $\lambda \geq \kappa$ .  
(2) If  $X \subset P_\kappa\lambda$  is *A-subtle*, there exists  $\{S_x \subset P_{x \cap \kappa} x : x \in X\}$  such that  $\{x \in X : S_x = S \cap x\} \in WNS_{\kappa,\lambda}^+$  for every  $S \subset P_\kappa\lambda$ .  
(3) If  $\kappa$  is weakly Mahlo and  $2^{\alpha^{<\kappa}} \leq \lambda$  for every  $\alpha < \lambda$ , there exists  $\{S_x \subset P_{x \cap \kappa} x : x \in P_\kappa\lambda \text{ and } x \cap \kappa \text{ is regular}\}$  such that  $S_x$  is a club in  $P_{x \cap \kappa} x$  and  $\{x : S_x \not\subset C\} \in NS_{\kappa,\lambda}$  for any club  $C \subset P_\kappa\lambda$ .

The last statement is false for  $\lambda = \kappa$ .

In the next section we study large cardinal aspects of *A-subtlety* to prove an analogue of Baumgartner's theorem [4] for regular uncountable cardinals:

- Theorem 1.4.** If  $X \subset P_\kappa\lambda$  is *A-subtle*, then  $\{x \in X : X \cap P_{x \cap \kappa} x \text{ is } \Pi_n^m\text{-indescribable}\}$  is *A-subtle* for every  $m, n < \omega$ .

Thus our subtlety takes an appropriate place in  $P_\kappa\lambda$  combinatorics. The next follows immediately.

**Corollary 1.5.** *If  $cf(\lambda) \geq \kappa$  and  $X \subset P_\kappa \lambda$  is almost ineffable, then  $\{x \in X : X \cap P_{x \cap \kappa} x \text{ is Shelah}\}$  is almost ineffable.*

Hence “ $\kappa$  is almost  $\lambda$ -ineffable” is an essentially stronger hypothesis than “ $\kappa$  is  $\lambda$ -Shelah”. Kamo [12] already proved:

**Fact 1.6.** (Kamo) *If  $\kappa$  is  $\lambda$ -ineffable, then  $\{x \in P_\kappa \lambda : P_{x \cap \kappa} x \text{ is not almost ineffable}\}$  is not ineffable.*

Thus we have the same hierarchy of combinatorial properties for  $P_\kappa \lambda$  as for regular uncountable cardinals. Our proof is applicable for Kamo’s theorem and more simple than his.

Another corollary is:

**Corollary 1.7.** *If  $\{x \in P_\kappa \lambda : x \cap \kappa < |x|\}$  is  $A$ -subtle, then  $V \neq L[U]$*

Note that  $L \models$  “ $\kappa$  is subtle” if  $\kappa$  is subtle.

In §4 we turn to saturation of subtle ideals and show:

**Theorem 1.8.** (1) *The ideals of non-subtle sets are not  $\lambda$ -saturated.*

(2) *The ideal of non-completely ineffable sets is not precipitous*

The last section is devoted to almost ineffability and ineffability. Our results might be surprising comparing with Kamo’s theorem:

**Theorem 1.9.** *Suppose that  $\lambda^{<\kappa} = 2^\lambda$ . Then,  $X$  is almost ineffable if and only if  $X$  is ineffable.*

As a corollary we get:

**Corollary 1.10.** (1) *We can not prove in ZFC that  $\kappa$  is  $\lambda^{<\kappa}$ -ineffable whenever  $\kappa$  is  $\lambda$ -ineffable.*

(2) *If  $\lambda^{<\kappa} = 2^\lambda$ ,  $I_{\kappa,\lambda}$  does not have the partition property*

## 2 The Subtle ideals on $P_\kappa \lambda$ .

Menas [16] tried to introduce subtlety into  $P_\kappa \lambda$  as follows (we call  $M$ -subtle in this paper):

**Definition 2.1.** Let  $X \subset P_\kappa\lambda$ .  $X$  is  $M$ -subtle if for any  $\{S_x \subset x : x \in P_\kappa\lambda\}$  and a club  $C \subset P_\kappa\lambda$  there exist  $y \subsetneq z$  both in  $C \cap X$  such that  $S_y = S_z \cap y$ .

This is not an essential generalization as proved in the same paper.

**Fact 2.2.** For every  $\lambda \geq \kappa$ ,  $P_\kappa\lambda$  is  $M$ -subtle if and only if  $\kappa$  is subtle.

We present a new definition of subtlety for  $P_\kappa\lambda$  and use the word “ $A$ -subtle” for it. In place of the filter of club sets we use  $WNS_{\kappa,\lambda}$ .

**Definition 2.3.** For  $X \subset P_\kappa\lambda$ ,  $X$  is  $A$ -subtle if for any  $\{S_x \subset P_{x \cap \kappa}x : x \in P_\kappa\lambda\}$  and  $C \in WNS_{\kappa,\lambda}^*$ , there are  $y \prec z$  both in  $C \cap X$  such that  $S_y = S_z \cap P_{y \cap \kappa}y$ .

Set  $I_M = \{X \subset P_\kappa\lambda : X \text{ is not } M\text{-subtle}\}$  and  $I_A = \{X \subset P_\kappa\lambda : X \text{ is not } A\text{-subtle}\}$ .

**Remark 2.4.** If  $\lambda^{<\kappa} = \lambda$ , then  $\{x \in P_\kappa\lambda : h''P_{x \cap \kappa}x \subset x\} \in WNS_{\kappa,\lambda}^*$  for any bijection  $h : P_\kappa\lambda \rightarrow \lambda$ . Thus, in this case,  $X \subset P_\kappa\lambda$  is  $A$ -subtle if and only if for any  $\{S_x \subset x : x \in P_\kappa\lambda\}$  and  $C \in WNS_{\kappa,\lambda}^*$  there are  $y \prec z$  both in  $C \cap X$  such that  $S_y = S_z \cap y$ .

We say  $\kappa$  is  $\lambda$ -subtle if  $P_\kappa\lambda$  is  $A$ -subtle (in  $P_\kappa\lambda$ ). If  $P_\kappa\lambda$  is  $A$ -subtle, then it is  $M$ -subtle. So  $\kappa$  is assumed to be subtle in the rest of this section.

We collect several facts for subtle ideals:

**Proposition 2.5.** (1)  $I_M \subset I_A$ .

(2)  $I_M$  is a normal ideal on  $P_\kappa\lambda$ .

(3)  $I_A$  is a strongly normal ideal on  $P_\kappa\lambda$

(4)  $\{x \in P_\kappa\lambda : x \cap \kappa \text{ is Mahlo}\} \in I_M^*$ .

(5) If  $\kappa \leq \delta < \lambda$  and  $X \subset P_\kappa\lambda$  is  $A$ -subtle, then  $X \upharpoonright \delta := \{x \cap \delta : x \in X\} \subset P_\kappa\delta$  is  $A$ -subtle.

(6) If  $\kappa$  is Mahlo and  $\{x \in P_\kappa\lambda : X \cap P_{x \cap \kappa}x \text{ is } A\text{-subtle}\} \in WNS_{\kappa,\lambda}^+$ , then  $X$  is  $A$ -subtle.

(7) Let  $\{s_\alpha : \alpha < \delta\}$  be an enumeration of  $P_\kappa\lambda$  and  $f(x) = \{\alpha : s_\alpha \prec x\}$  for  $x \in P_\kappa\lambda$ . Then,  $f''X \subset P_\kappa\delta$  is  $A$ -subtle if and only if  $X \subset P_\kappa\lambda$  is  $A$ -subtle.

(8) If  $\lambda$  is regular,  $X \subset P_\kappa\lambda$  and  $\{\alpha < \lambda : X \cap P_\kappa\alpha \text{ is } M\text{-subtle}\} \in NS_\lambda^+$ , then  $X$  is  $M$ -subtle.

*Proof.* We only show (3). Let  $X \subset P_\kappa\lambda$  be subtle and  $f : X \rightarrow P_\kappa\lambda$  such that  $f(x) \prec x$  for every  $x \in X$ . Suppose to the contradiction that  $f^{-1}(\{a\})$  is not subtle for any  $a \in P_\kappa\lambda$ . For  $a \in P_\kappa\lambda$ , we fix  $\{S_x^a \subset P_{x \cap \kappa} : x \in P_\kappa\lambda\}$  and  $D_a \in WNS_{\kappa,\lambda}^*$  such that  $S_y^a \neq S_z^a \cap P_{y \cap \kappa}y$  for any  $y \prec z$  both in  $D_a \cap f^{-1}(\{a\})$ .

Let  $h : P_\kappa\lambda \times P_\kappa\lambda \rightarrow P_\kappa\lambda$  be a bijection and set  $T_x = h''(\{f(x)\} \times S_x^{f(x)}) \cap P_{x \cap \kappa}x$ . Note that  $C = \{x \in P_\kappa\lambda : h''(P_{x \cap \kappa}x \times P_{x \cap \kappa}x) \subset P_{x \cap \kappa}x\} \in WNS_{\kappa,\lambda}^*$ . Since  $X$  is  $A$ -subtle and  $E = C \cap \Delta_{\prec} D_a \in WNS_{\kappa,\lambda}^*$ , there exist  $y \prec z$  both in  $E \cap X$  such that  $T_y = T_z \cap P_{y \cap \kappa}y$ . Then,  $f(y) = f(z)$  and  $S_y^{f(y)} = S_z^{f(z)} \cap P_{y \cap \kappa}y$ . Set  $a = f(y) = f(z)$ . We have  $y \prec z$  are both in  $D_a \cap f^{-1}(\{a\})$  and  $S_y^a = S_z^a$ , which contradicts our assumption.  $\square$

A natural question arises:

**Question 2.6.** *Can it be proved that  $I_M \subsetneq I_A$ ?*

It turns out that “ $A$ -subtle” is neither an essential generalization.

**Theorem 2.7.** *If  $\kappa$  is subtle, then  $P_\kappa\lambda$  is  $A$ -subtle.*

*Proof.* Let  $S_x \subset P_{x \cap \kappa}x$  for  $x \in P_\kappa\lambda$  and  $D \in WNS_{\kappa,\lambda}^*$ . Since  $\kappa^{<\kappa} = \kappa$ ,  $WNS_{\kappa^+, \lambda}$  is proper.

We first show  $\{x \in P_{\kappa^+ \lambda} : D \cap P_\kappa x \in WNS_{\kappa,x}^*\} \in WNS_{\kappa^+, \lambda}^*$ . Let  $f : P_\kappa\lambda \rightarrow P_\kappa\lambda$  such that  $C_f \subset D$ . If  $\{x \in P_{\kappa^+ \lambda} : f''P_\kappa x \subset P_\kappa x\} \notin WNS_{\kappa^+, \lambda}^*$ ,  $X := \{x \in P_{\kappa^+ \lambda} : \kappa \subset x \wedge \exists y_x \in P_\kappa x (f(y) \notin P_\kappa x)\} \in WNS_{\kappa^+, \lambda}^+$ . Note that  $\kappa = |x \cap \kappa|$  for every  $x \in X$ . By strong normality we have  $y \in P_\kappa\lambda$  such that  $Y := \{x \in X : y_x = y\} \in WNS_{\kappa^+, \lambda}^+$ .  $Y \cap \widehat{f(y)} = \emptyset$ . Contradiction. Thus  $Z := \{x \in P_{\kappa^+ \lambda} : \kappa \subset x \wedge f \upharpoonright P_\kappa x : P_\kappa x \rightarrow P_\kappa x\} \in WNS_{\kappa^+, \lambda}^*$ . For  $x \in Z$   $D_x := \{s \in P_\kappa x : f''P_{s \cap \kappa}x \subset P_{s \cap \kappa}x\} \in WNS_{\kappa,x}^*$ . For every  $x \in Z$   $D \cap P_\kappa x \in WNS_{\kappa,x}^*$  since  $D_x \subset C_f \cap P_\kappa x \subset D$ .

Note that  $\kappa$  is subtle if and only if  $P_\kappa\kappa$  is  $A$ -subtle. Thus  $P_\kappa y$  is  $A$ -subtle for every  $y \in Z$ . Now we consider  $\{S_x : x \in P_\kappa y\}$  and  $D \cap P_\kappa y$ . There exist  $x_1 \prec x_2$  both in  $D \cap P_\kappa y$  such that  $S_{x_1} = S_{x_2} \cap P_{x_1 \cap \kappa}x_1$ .  $\square$

The following observations suggest two ideals may be the same. The first appears in [16].

**Fact 2.8.** *If  $X \subset P_\kappa\lambda$  is  $M$ -subtle and  $S_x \subset x$  for each  $x \in X$ , then for any club  $C \subset P_\kappa\lambda$  there exist  $x, y$  both in  $C \cap X$  such that  $x \subset y$ ,  $x \cap \kappa < y \cap \kappa$ , and  $S_x = S_y \cap x$ .*

**Proposition 2.9.** *If  $\kappa$  is subtle, then  $X = \{x \in P_\kappa\lambda : x \cap \kappa = |x|\}$  is  $A$ -subtle.*

*Proof.* Note that  $X \notin WNS_{\kappa,\lambda}$  ([1]). Let  $f : P_\kappa\lambda \rightarrow P_\kappa\lambda$  and  $S_x \subset P_{x \cap \kappa}x$  for all  $x \in P_\kappa\lambda$ . We build a chain  $\langle x_\alpha : \alpha < \kappa \rangle$  as follows:

Choose  $x_0 \in X \cap C_f$  arbitrarily and  $x_{\alpha+1} \in X \cap C_f$  so that  $x_\alpha \prec x_{\alpha+1}$ . For limit  $\alpha$  let  $x_\alpha = \bigcup \{x_\beta : \beta < \alpha\}$ .

Set  $x = \bigcup \{x_\alpha : \alpha < \kappa\}$ . Then,  $x \cap \kappa = |x| = \kappa$ ,  $P_{x \cap \kappa}x = \bigcup \{P_{x_\alpha \cap \kappa}x_\alpha : \alpha < \kappa\}$ , and there exists a club  $D \subset \kappa$  such that for every  $\alpha \in D$   $x_\alpha \cap \kappa = \alpha = |x_\alpha|$ . Note that  $x_\alpha \in C_f$  if  $\alpha$  is regular. Let  $g : P_{x \cap \kappa}x \rightarrow \kappa$  be any bijection. Then, we have a club  $E \subset \{\alpha \in D : g \text{``} P_\alpha x_\alpha \subset \alpha \text{'}$ . Since  $E$  is subtle in  $\kappa$ , there exist regular  $\beta < \gamma$  both in  $E$  such that  $g \text{``} S_{x_\beta} = g \text{``} S_{x_\gamma} \cap \beta$ . Since  $\beta$  and  $\gamma$  are regular,  $x_\beta$  and  $x_\gamma$  belong to  $C_f$ .  $\square$

S. Baldwin[3] and others observed the consistency strength of the stationarity of  $\{x \in P_\kappa\lambda : x \cap \kappa < |x|\}$ , the complement of the set we mentioned now.

**Fact 2.10.** ([3],[14],[1])

- (1) *If  $\kappa$  is weakly inaccessible and  $\lambda$  Ramsey  $> \kappa$ , then  $\{x \in P_\kappa\lambda : x \cap \kappa < |x|\}$  is stationary.*
- (2) *If  $\{x \in P_\kappa\kappa^+ : x \cap \kappa < |x|\}$  is stationary, then  $0^\dagger$  exists.*
- (3) *If  $P_\kappa\lambda$  is Shelah, then  $\{x \in P_\kappa\lambda : x \cap \kappa < |x|\} \in NSh_{\kappa,\lambda}^*$ .*

**Corollary 2.11.** *Suppose that  $0^\dagger$  does not exist and  $S = \{x \in P_\kappa\lambda : h \text{``} P_{x \cap \kappa}x = x\}$  where  $h : P_\kappa\lambda \rightarrow \lambda$  is a bijection. Then,  $I_M \upharpoonright S = I_A$ . Thus  $I_M = I_A$  if  $I_M$  is strongly normal.*

The same relation holds between subtlety and its another weakening, which relates to “ethereal” introduced by Kunen:  $X \subset \kappa$  is *ethereal* if for any  $\langle S_\alpha \in [\alpha]^\alpha : \alpha < \kappa \rangle$  and a club  $C \subset \kappa$ , there are  $\beta < \gamma$  both in  $C \cap X$  such that  $|S_\beta \cap S_\gamma| = |\beta|$ .

**Definition 2.12.** *Let  $X \subset P_\kappa\lambda$ . We say  $X$  is weakly subtle if for any  $\{S_x \in I_{x \cap \kappa, x}^+ : x \in P_\kappa\lambda\}$  and a club  $C \subset P_\kappa\lambda$ , there are  $y \prec z$  both in  $C \cap X$  such that  $S_y \cap S_z \in I_{y \cap \kappa, y}^+$ . Let  $I_W = \{X \subset P_\kappa\lambda : X \text{ is not weakly subtle}\}$ .*

We have the following:

- Proposition 2.13.** (1)  $I_W$  is a normal ideal.  
 (2) If  $P_\kappa\lambda$  is weakly subtle, then  $\kappa$  is ethereal.  
 (3)  $I_W \upharpoonright S = I_A$  where  $S = \{x \in P_\kappa\lambda : h \text{``} P_{x \cap \kappa}x = x\}$  for a bijection  $h : P_\kappa\lambda \rightarrow \lambda$ .



Three subtle ideals are interesting from the view of the diamond principles for  $P_\kappa\lambda$ . The following two cardinal version of diamond principle by Jech is wellknown.

**Definition 2.14.** Let  $X \subset P_\kappa\lambda$ . Then,

$\diamond_0(X)$ : there exist  $\{S_x \subset x : x \in X\}$  such that  $\{x \in X : S_x = S \cap x\}$  is stationary for any  $S \subset \lambda$ . We simply write  $\diamond_0$  for  $\diamond_0(P_\kappa\lambda)$ . Let  $J_0 = \{X \subset P_\kappa\lambda : \diamond_0(X) \text{ does not hold}\}$ .

The following are known (see [9], [10]):

- Fact 2.15.** (1)  $J_0$  is a normal ideal on  $P_\kappa\lambda$ .  
 (2)  $L \models$  " $\diamond_0(X)$  for any  $X \subset P_{\omega_1}\lambda$ ".  
 (3) If  $2^{<\kappa} < \lambda$ , then  $J_0$  is proper.  
 (4) If  $\diamond_0$  holds, then  $NS_{\kappa,\lambda}$  is not  $2^\lambda$  saturated.

In the context of  $I_A$  and  $I_W$  another version of diamond arises.

**Definition 2.16.** Let  $X \subset P_\kappa\lambda$ . Then,

$\diamond_1(X)$ : there exists  $\{S_x \subset P_{x \cap \kappa} : x \in X\}$  such that  $\{x \in X : S_x = S \cap P_{x \cap \kappa}\}$  is stationary for any  $S \subset P_\kappa\lambda$ .

$\diamond_2(X)$ : there exists  $\{S_x \subset P_{x \cap \kappa} : x \in X\}$  such that  $\{x \in X : S_x = S \cap P_{x \cap \kappa}\} \in WNS_{\kappa,\lambda}^+$  for any  $S \subset P_\kappa\lambda$ .

$J_1$  and  $J_2$  are similarly defined as  $J_0$ .

Of course  $J_0 \subset J_1 \subset J_2$  and it is easily seen:

**Lemma 2.17.** (1)  $J_1$  is a normal ideal on  $P_\kappa\lambda$ .

(2)  $J_2$  is a strongly normal ideal on  $P_\kappa\lambda$

(3) If  $\lambda^{<\kappa} = \lambda$  and  $h : P_\kappa\lambda \rightarrow \lambda$  is a bijection, then  $J_2 = J_0 \upharpoonright S$  with  $S = \{x : h''P_{x \cap \kappa} = x\}$ .

(4)  $L \models$  " $J_2 = WNS_{\omega_1,\lambda}$ ".

(5) If  $J_2$  is proper, then any ideal  $\subset WNS_{\kappa,\lambda}$  is not  $2^{\lambda^{<\kappa}}$  saturated.

**Theorem 2.18.** If  $\kappa$  is subtle, then  $J_2$  is proper.

*Proof.* We show  $\diamond_2(X)$  holds for every  $A$ -subtle  $X \subset P_\kappa\lambda$ .

By induction on  $\prec$  we define  $S_x \subset P_{x \cap \kappa}$  for  $x \in X$  as well as  $C_x \subset P_{x \cap \kappa}$ .

If  $x$  is a  $\prec$  minimal element of  $X$ , then  $S_x = C_x = \emptyset$ .

Suppose  $S_y$  and  $C_y$  is defined for every  $y \in X \cap P_{x \cap \kappa} x$ . If there exist  $S \subset P_{x \cap \kappa} x$  and  $C \in WNS_{x \cap \kappa, x}^*$  such that  $S_y \neq S \cap P_{y \cap \kappa} y$  for any  $y \in C$ , then let  $S_x$  and  $C_x$  be any such  $S$  and  $C$ . We say this is the substantial case. Otherwise let  $S_x = C_x = P_{x \cap \kappa} x$ . To show  $\{S_x : x \in X\}$  is a witness of  $\diamond_2(X)$ , let  $S \subset P_\kappa \lambda$ ,  $D \in WNS_{\kappa, \lambda}^*$ , and  $S_x \neq S \cap P_{x \cap \kappa} x$  for any  $x \in X \cap D$ . Since  $X \cap D$  is  $A$ -subtle, we may assume that  $D \cap P_{x \cap \kappa} x \in WNS_{x \cap \kappa, x}^*$  for every  $x \in X \cap D$ . Thus  $S \cap P_{x \cap \kappa} x$  and  $D \cap P_{x \cap \kappa} x$  witness the substantial case occurs for every  $x \in X \cap D$ . However  $X \cap D$  is subtle hence there exist  $y \prec z$  both in  $X \cap D$  such that  $S_y = S_z \cap P_{y \cap \kappa} y$ . In particular  $y \in D \cap P_{z \cap \kappa} z$ . Contradiction.  $\square$

Note that  $\diamond_\kappa$  holds if  $\kappa$  is ethereal and  $\kappa^{<\kappa} = \kappa$  [13].

**Question 2.19.** (1) Does  $\diamond_1$  hold if  $P_\kappa \lambda$  is weakly subtle?  
 (2) If  $\kappa$  is ethereal, then  $P_\kappa \lambda$  is weakly subtle?

Let  $\text{Reg} = \{x \in P_\kappa \lambda : x \cap \kappa \text{ is regular}\}$ . We conclude this section by:

**Proposition 2.20.** Suppose  $\kappa$  is weakly Mahlo and  $2^{\alpha^{<\kappa}} \leq \lambda$  for every  $\alpha < \lambda$ . Then, there exists  $\{S_x : x \in P_\kappa \lambda, x \cap \kappa \text{ is regular}\}$  such that

1.  $S_x \subset P_{x \cap \kappa} x$  club,
2. for every club  $C \subset P_\kappa \lambda$   $\{x : S_x \not\subset C\} \in NS_{\kappa, \lambda} \upharpoonright \text{Reg}$ .

*Proof.* Let  $\{C_\alpha : \kappa^+ \leq \alpha < \lambda\} = \{X : \exists \beta < \lambda \text{ } X \text{ is a club of } P_\kappa \beta\}$  be an enumeration and  $C_\alpha$  a club of  $P_\kappa \beta(\alpha)$ . Set  $B = \{x : \forall \alpha \in x \beta(\alpha) \in x\}$ . Then,  $B \in NS_{\kappa, \lambda}^*$ . Let  $S_x = \{y \in P_{x \cap \kappa} x : \forall \alpha \in x y \cap \beta(\alpha) \in C_\alpha\}$ .

For every  $\alpha$   $\{x \in P_\kappa \lambda : C_\alpha \cap P_{x \cap \kappa}(x \cap \beta(\alpha)) \text{ is a club of } P_{x \cap \kappa}(x \cap \beta(\alpha))\} \in (NS_{\kappa, \lambda} \upharpoonright \text{Reg})^*$ . Thus,  $\{x \in P_\kappa \lambda : \{z \in P_{x \cap \kappa} x : z \cap \beta(\alpha) \in C_\alpha\} \text{ is a club of } P_{x \cap \kappa} x\} \in (NS_{\kappa, \lambda} \upharpoonright \text{Reg})^*$ . Hence  $A = \{x \in P_\kappa \lambda : S_x \text{ is a club of } P_{x \cap \kappa} x\} \in (NS_{\kappa, \lambda} \upharpoonright \text{Reg})^*$ .

Pick any club  $C \subset P_\kappa \lambda$ . We have  $f : \lambda \times \lambda \rightarrow \lambda$  with  $C_f \subset C$ . Define  $g$  by  $C_f \upharpoonright \beta = C_{g(\beta)}$  for  $\beta < \lambda$ . Note that  $\beta = \beta(g(\beta))$ .

Let  $x \in A \cap B \cap C_f$ . For every  $\beta \in x$   $g(\beta) \in x$ . Hence for every  $y \in S_x$  and  $\beta \in x$  we have  $y \cap \beta \in C_{g(\beta)} = C_f \upharpoonright \beta$ . If  $\{\xi, \zeta\} \subset y$ , then  $\{\xi, \zeta, f(\xi, \zeta)\} \subset \beta$  for some  $\beta \in x$ . Choose  $z \in C_f$  such that  $y \cap \beta = z \cap \beta$ . Then,  $f(\xi, \zeta) \in z \cap \beta = y \cap \beta$ . Hence  $y \in C_f$ . We have shown that  $S_x \subset C_f \subset C$  for every  $x \in A \cap B \cap C_f$ .  $\square$

**Remark 2.21.** This is false for  $\lambda = \kappa$ .

### 3 Subtlety and large cardinals

Recall that  $X \subset \kappa$  is  $\Pi_n^m$ -*indescribable* if for any  $R \subset V_\kappa$  and  $\Pi_n^m$  sentence  $\varphi$  such that  $(V_\kappa, \in, R) \models \varphi$ , there exists  $\alpha \in X$  such that  $(V_\alpha, \in, R \cap V_\alpha) \models \varphi$ .

**Fact 3.1.** (1) *Suppose that  $\lambda$  is weakly compact. Then,  $X \subset P_\kappa \lambda$  is  $M$ -subtle if and only if  $\{\alpha < \lambda : X \cap P_\kappa \alpha \text{ is not } M\text{-subtle}\}$  is not  $\Pi_1^1$ -indescribable.*

(2)  *$I_M$  is a normal ideal on  $P_\kappa \lambda$  such that  $\{x \in P_\kappa \lambda : x \cap \kappa \text{ is not } \Pi_n^m\text{-indescribable}\} \in I_M$  for every  $m, n < \omega$ .*

Carr[7] defined  $P_\kappa \lambda$ -version of indescribability.

**Definition 3.2.** *A sequence  $\langle V_\alpha(\kappa, \lambda) : \alpha \leq \kappa \rangle$  is recursively defined as follows:*

$$\begin{aligned} V_0(\kappa, \lambda) &= \lambda \\ V_{\alpha+1}(\kappa, \lambda) &= P_\kappa(V_\alpha(\kappa, \lambda)) \cup V_\alpha(\kappa, \lambda) \\ V_\alpha(\kappa, \lambda) &= \bigcup \{V_\beta(\kappa, \lambda) : \beta < \alpha\} \quad \text{if } \alpha \text{ is a limit ordinal} \end{aligned}$$

This definition can be carried out for  $x \in P_\kappa \lambda$  if  $x \cap \kappa$  is inaccessible. For such  $x$  we consider the structure  $(V_{x \cap \kappa}(x \cap \kappa, x), \in)$  in the same way as  $(V_\kappa(\kappa, \lambda), \in)$ .

**Definition 3.3.** *We say  $X \subset P_\kappa \lambda$  is  $\Pi_n^m$ -indescribable if for any  $R \subset V_\kappa(\kappa, \lambda)$  and  $\Pi_n^m$  sentence  $\varphi$  such that  $(V_\kappa(\kappa, \lambda), \in, R) \models \varphi$ , there exists  $x \in X$  such that  $x \cap \kappa = |x \cap \kappa|$  and  $(V_{x \cap \kappa}(x \cap \kappa, x), \in, R \cap V_{x \cap \kappa}(x \cap \kappa, x)) \models \varphi$ .*

**Lemma 3.4.** *If  $X \subset P_\kappa \lambda$  is  $A$ -subtle and  $S_x \subset P_{x \cap \kappa} x$  for  $x \in P_\kappa \lambda$ , then  $\{x \in X : \{y \in X \cap P_{x \cap \kappa} x : S_y = S_x \cap P_{y \cap \kappa} y\} \text{ is not } \Pi_n^m\text{-indescribable for some } m, n\}$  is not  $A$ -subtle.*

*Proof.* Otherwise, by  $\kappa$ -completeness of  $I_A$ ,  $Y := \{x \in X : \{y \in X \cap P_{x \cap \kappa} x : S_y = S_x \cap P_{y \cap \kappa} y\} \text{ is not } \Pi_n^m\text{-indescribable}\}$  is subtle for some  $m, n < \omega$ . We may assume that  $x \cap \kappa$  is inaccessible for all  $x \in Y$ . For  $x \in Y$  there exist  $R_x \subset V_{x \cap \kappa}(x \cap \kappa, x)$  and a  $\Pi_n^m$  sentence  $\varphi_x$  such that  $(V_{x \cap \kappa}(x \cap \kappa, x), \in, R_x) \models \varphi_x$  while  $(V_{y \cap \kappa}(y \cap \kappa, y), \in, R_x \cap V_{y \cap \kappa}(y \cap \kappa, y)) \models \neg \varphi_x$  for any  $y \in X \cap P_{x \cap \kappa} x$  with  $S_y = S_x \cap P_{y \cap \kappa} y$ . By  $\kappa$ -completeness again, we can assume for all  $x \in Y$   $\varphi_x = \varphi$  for some  $\varphi$ .

Since  $Y$  is subtle, there are  $y \prec z$  both in  $Y$  such that  $R_y = R_z \cap V_{y \cap \kappa}(y \cap \kappa, y)$  and  $S_y = S_z \cap P_{y \cap \kappa} y$ . Then,  $y \in X \cap P_{z \cap \kappa} z$ ,  $S_y = S_z \cap P_{y \cap \kappa} y$  and  $(V_{y \cap \kappa}(y \cap \kappa, y), \in, R_z \cap V_{y \cap \kappa}(y \cap \kappa, y)) \models \varphi_z$ , which is a contradiction.  $\square$

As a corollary we have:

**Theorem 3.5.**  $\{x \in P_\kappa\lambda : P_{x \cap \kappa}x \text{ is not } \Pi_n^m\text{-indescribable}\} \in I_A$  for every  $m, n < \omega$ .

This theorem derives strong facts.

**Lemma 3.6.** *If  $\{x \in P_\kappa\lambda : 2^{x \cap \kappa} \leq |x|\}$  is  $A$ -subtle, then  $\{x \in P_\kappa\lambda : o(x \cap \kappa) \geq 1\}$  is  $A$ -subtle.*

*Proof.* Note that  $\kappa$  is measurable if  $P_\kappa 2^\kappa$  is  $\Pi_1^1$ -indescribable ([6],[7]). Let  $X = \{x \in P_\kappa\lambda : P_{x \cap \kappa} 2^{x \cap \kappa} \text{ is } \Pi_1^1\text{-indescribable}\}$ . Then,  $X$  is  $A$ -subtle. For  $x \in X$   $x \cap \kappa$  is measurable and  $\{y \in P_{x \cap \kappa} 2^{x \cap \kappa} : y \cap \kappa \text{ is measurable}\}$  is  $\Pi_1^1$ -indescribable. Hence  $o(x \cap \kappa) \geq 1$ .  $\square$

Note that  $L[U] \models$  "there exist  $\kappa < \lambda$  such that  $\{x \in P_\kappa\lambda : x \cap \kappa < |x|\} \in NS_{\kappa,\lambda}^+$ ".

**Corollary 3.7.** (1)  $L[U] \models$  " $\{x \in P_\kappa\lambda : x \cap \kappa < |x|\}$  is not  $A$ -subtle".

(2)  $L[U] \models$  " $\neg \exists \kappa(\kappa \text{ is } \kappa^+\text{-Shelah})$ ".

Thus the existence of a cardinal  $\kappa$  such that  $\{x \in P_\kappa\lambda : x \cap \kappa < |x|\}$  is  $A$ -subtle is rather strong in consistency strength. On the other hand subtlety is a  $\Pi_1^1$  property of  $P_\kappa\lambda$ . The following proposition says the size of  $\kappa$  is not necessarily large.

**Proposition 3.8.** *The least cardinal  $\kappa$  such that  $\kappa$  is  $\kappa^+$ -subtle is not  $\kappa^+$ -Shelah.*

In the rest of this section we compare the almost ineffability and Shelah property using  $I_A$ , which reveals very useful in large cardinal hierarchy.

Carr [6] defined Shelah property as a  $P_\kappa\lambda$  generalization of weak compactness. We show: *if  $P_\kappa\lambda$  is subtle then there exist many  $x \in P_\kappa\lambda$  such that  $P_{x \cap \kappa}x$  is Shelah.*

**Definition 3.9.** *Let  $X \subset P_\kappa\lambda$ . We say  $X$  is Shelah if for any  $\{f_x \in {}^x x : x \in P_\kappa\lambda\}$  there is  $f : \lambda \rightarrow \lambda$  such that for every  $y \in P_\kappa\lambda$  the set  $\{x \in X \cap \hat{y} : f \upharpoonright y = f_x \upharpoonright y\} \in I_{\kappa,\lambda}^+$ .*

*We say  $X$  is almost ineffable (ineffable) if for any  $\{f_x \in {}^x x : x \in P_\kappa\lambda\}$  there is  $f : \lambda \rightarrow \lambda$  such that  $\{x \in X : f \upharpoonright x = f_x\} \in I_{\kappa,\lambda}^+$  ( $NS_{\kappa,\lambda}^+$ ).*

*Let  $NSh_{\kappa\lambda} = \{X \subset P_\kappa\lambda : X \text{ is not Shelah}\}$  and  $NAIn_{\kappa\lambda} = \{X \subset P_\kappa\lambda : X \text{ is not almost ineffable}\}$ .*

*We often say  $\kappa$  is  $\lambda$ -Shelah (almost  $\lambda$ -ineffable) if  $P_\kappa\lambda$  is Shelah (almost ineffable).*

Clearly  $X$  is Shelah if  $X$  is almost ineffable, and  $X$  is almost ineffable if  $X$  is ineffable. It is known that  $NSh_{\kappa\lambda}$  and  $NAIn_{\kappa\lambda}$  are strongly normal ideals if  $cf(\lambda) \geq \kappa$ . Moreover, Kamo [12] proved the following:

**Fact 3.10.** (Kamo) *If  $X \subset P_\kappa\lambda$  is ineffable and  $cf(\lambda) \geq \kappa$ , then  $\{x \in X : X \cap P_{x \cap \kappa}x$  is almost ineffable $\}$  is ineffable.*

The following follows immediately from definition and the remark after Definition 2.1 with strong normality of  $NAIn_{\kappa\lambda}$ .

**Proposition 3.11.** *If  $cf(\lambda) \geq \kappa$  and  $X \subset P_\kappa\lambda$  is almost ineffable, then  $X$  is  $A$ -subtle.*

Carr [7] proved the following:

**Fact 3.12.** *Let  $X \subset P_\kappa\lambda$  is  $\Pi_1^1$ -indescribable, then  $X$  is Shelah. The converse is also true if  $cf(\lambda) \geq \kappa$ .*

**Corollary 3.13.** *If  $X \subset P_\kappa\lambda$  is subtle, then  $Y = \{x \in X : X \cap P_{x \cap \kappa}x$  is not Shelah $\}$  is not  $A$ -subtle.*

**Corollary 3.14.** *Let  $cf(\lambda) \geq \kappa$ . If  $X \subset P_\kappa\lambda$  is almost ineffable, then  $\{x \in P_\kappa\lambda : X \cap P_{x \cap \kappa}x$  is Shelah $\}$  is almost ineffable. In particular,  $\{x \in P_\kappa\lambda : x \cap \kappa$  is  $x$ -Shelah $\} \in NAIn_{\kappa,\lambda}^*$  if  $\kappa$  is almost  $\lambda$ -ineffable.*

This corollary tells that almost ineffability is much stronger hypothesis than Shelah property. For instance, suppose that  $\kappa$  is almost  $\kappa^+$ -ineffable. Then  $\{x \in P_\kappa\lambda : o.t.(x) = (x \cap \kappa)^+\} \in NAIn_{\kappa,\lambda}^*$  hence  $\{x \in P_\kappa\lambda : x \cap \kappa$  is  $(x \cap \kappa)^+$ -Shelah $\} \in NAIn_{\kappa,\lambda}^*$ . Thus, below the least  $\alpha$  that is almost  $\alpha^+$ -ineffable, stationary many  $\beta$  which is  $\beta^+$ -Shelah exist.

## 4 Saturation of subtle ideals on $P_\kappa\lambda$

Now we turn to saturation of subtle ideals.

**Proposition 4.1.** *Let  $I$  be a normal  $\lambda$  saturated ideal on  $P_\kappa\lambda$  with  $\lambda$  regular. Then,  $\{x \in P_\kappa\lambda : o.t.(x)$  is regular $\} \in I^*$  hence  $\{x \in P_\kappa\lambda : x \cap \kappa < |x|\} \in I^*$ .*

*Proof.* Let  $G$  be  $P_I$  generic for  $V$ . By  $\lambda$  saturation the generic ultrapower  $Ult(V, G)$  is well-founded. Let  $j : V \rightarrow M \cong Ult(V, G)$  be an generic embedding. Since  $\lambda$  is regular in  $V[G]$ ,  $M \models \text{“}\lambda \text{ is regular”}$  and  $[(o.t.(x) | x \in P_\kappa \lambda)]_G = o.t.(j \text{“}\lambda) = \lambda$ . Clearly  $\{x \in P_\kappa \lambda : x \cap \kappa \in \kappa \wedge x \setminus \kappa \neq \emptyset\} \in I^*$ .  $\square$

**Theorem 4.2.** *Neither  $I_A$  nor  $I_M$  is  $\lambda$  saturated.*

*Proof.* First suppose that  $\lambda$  is regular. We know  $X = \{x \in P_\kappa \lambda : x \cap \kappa = |x| \text{ and } \kappa \in x\}$  is  $A$ -subtle. For  $x \in X$   $o.t.(x)$  is singular.

Second assume  $\lambda$  is singular and the subtle ideal on  $P_\kappa \lambda$ , say  $I$ , is  $\lambda$  saturated. In fact  $I$  is  $\delta$  saturated for some regular  $\delta < \lambda$ . Then  $I \upharpoonright \delta$  is  $\delta$  saturated and  $\{x \in P_\kappa \delta : x \cap \kappa = |x|\} \notin I \upharpoonright \delta$ . Contradiction.  $\square$

**Remark 4.3.** *By Cummings’ theorem  $I_A$  is not  $\lambda^+$ -saturated if  $cf(\lambda) < \kappa$ .*

By definition ineffability can be seen as a strengthening of Shelah property. One of the strongest version is complete ineffability defined as follows:

**Definition 4.4.** *An ideal  $I$  on  $P_\kappa \lambda$  is  $(\lambda, \lambda)$ -distributive if for any  $X \in I^+$  and  $\{W_\alpha : \alpha < \lambda\}$  which is an  $I$ -partition of  $X$  with  $|W_\alpha| \leq \lambda$  there exist  $Y \in P(X) \cap I^+$  and  $\{X_\alpha : \alpha < \lambda\}$  such that  $X_\alpha \in W_\alpha$  and  $Y \setminus X_\alpha \in I$  for every  $\alpha < \lambda$ .*

*We say  $\kappa$  is completely  $\lambda$  ineffable if there is a normal  $(\lambda, \lambda)$ -distributive ideal on  $P_\kappa \lambda$ . When  $\kappa$  is completely  $\lambda$  ineffable, the minimal normal  $(\lambda, \lambda)$ -distributive ideal on  $P_\kappa \lambda$ , say  $I$ , is called completely ineffable ideal and  $X \in I^+$  is said to be completely ineffable.*

If  $I$  is a normal  $(\lambda, \lambda)$ -distributive ideal on  $P_\kappa \lambda$  and  $X \in I^+$ , the following hold [11]:

1. For any  $\{f_x \in {}^x x : x \in X\}$  there is  $f : \lambda \rightarrow \lambda$  such that  $\{x \in X : f_x = f \upharpoonright x\} \in I^+$
2.  $I$  is strongly normal.

In the rest of this section we assume  $cf(\lambda) \geq \kappa$  and  $I$  is a normal  $(\lambda, \lambda)$ -distributive ideal on  $P_\kappa \lambda$ . Hence  $\kappa$  is Mahlo,  $\lambda^{<\kappa} = \lambda$ , and  $I$  is strongly normal. Let  $\mathbb{P}_I = (I, \subset)$ ,  $G \subset \mathbb{P}_I$  be  $V$  generic, and  $j : V \prec M \cong Ult(V, G)$  a generic elementary embedding defined in  $V[G]$ .

By normality of  $I$   $P(\lambda)^V \subset M$ . Moreover  $\{\langle P_{x \cap \kappa} x \mid x \in P_\kappa \lambda \rangle\}_G = j'' P_\kappa \lambda \in M$ ,  $j'' P_\kappa \lambda = P_\kappa j'' \lambda$ , and  $j(X) \cap j'' P_\kappa \lambda = j'' X \in M$  for every  $X \in P(P_\kappa \lambda)^V$  by strong normality. Conversely we have:

**Lemma 4.5.** *For every  $[f]_G \in j''^\lambda j'' V \cap M$  there exists  $g \in V$  such that  $[f]_G = j(g) \upharpoonright j'' \lambda = j'' g$ .*

*Proof.* Suppose  $f \in V$ ,  $X \in I^+$  with  $X \Vdash "[f]_G : j'' \lambda \rightarrow j'' V"$ . For  $Y := \{s \in X : \text{dom}(f(s)) = s\} \in G$ ,  $\alpha < \lambda$ , and  $x \in V$ , let  $X_{\alpha, x} = \{s \in Y \cap \hat{\alpha} : f(s)(\alpha) = x\}$ . Then  $W_\alpha := \{X_{\alpha, x} : x \in V\} \cap I^+$  is a disjoint  $I$ -partition of  $Y$  and  $|W_\alpha| \leq \lambda$ . By  $(\lambda, \lambda)$ -distributivity there is  $g : \lambda \rightarrow V$  such that for every  $\alpha < \lambda$   $Y \setminus X_{\alpha, g(\alpha)} \in I$ . Hence  $Z := \Delta_\alpha X_{\alpha, g(\alpha)} \in I^+ \cap P(Y)$  and for every  $s \in Z$  and  $\alpha \in s$   $f(s)(\alpha) = g(\alpha)$ , that is,  $Z \Vdash "[f]_G = j(g) \upharpoonright j'' \lambda"$ .  $\square$

**Remark 4.6.** *By  $(\lambda, \lambda)$ -distributivity of  $I$ ,  $\lambda$  remains a cardinal in  $V[G]$  hence in  $M$ .*

**Corollary 4.7.** (1) *If  $X \in M$ ,  $X \subset j'' V$ , and  $|X|^M \leq \lambda$ , then  $X = j'' Y$  for some  $Y \in V$ .*

(2)  $P_\kappa j'' \lambda = j''(P_\kappa \lambda)$  and  $P(P_\kappa j'' \lambda)^M = \{j'' X : X \in P(P_\kappa \lambda)\}$ .

*Proof.* (1) Note that the collapsing map  $\pi : j'' \lambda \rightarrow \lambda$  is a bijection in  $M$ . Choose any surjection  $f \in M$  from  $j'' \lambda$  to  $X$ . By the lemma  $f = j(g) \upharpoonright j'' \lambda$  for some  $g \in V$ . Then  $X = j(g)''(j'' \lambda) = \{j(g)(j(\alpha)) : \alpha < \lambda\} = \{j(g(\alpha)) : \alpha < \lambda\}$ . Hence  $X = j'' Y$  where  $Y = g'' \lambda$ . Now (2) is clear.  $\square$

Next we present another characterization of completely ineffable subsets of  $P_\kappa \lambda$ .

**Definition 4.8.** *Let  $A$  be a set of ordinals. We inductively define  $In_\alpha(\kappa, A) \subset P(P_\kappa A)$  as follows:*

(1)  $X \in In_0(\kappa, A)$  if  $X \in NS_{\kappa, A}^+$ , that is, for every  $f : A \times A \rightarrow P_\kappa A$  there exists  $x \in X$  such that  $f''(x \times x) \subset P(x)$ .

(2)  $X \in In_{\alpha+1}(\kappa, A)$  if for every  $f : P_\kappa A \rightarrow P_\kappa A$  such that  $f(x) \subset x$  for any  $x \in P_\kappa A$  there is  $S \subset A$  with  $\{x \in X : f(x) = x \cap S\} \in In_\alpha(\kappa, A)$ .

(3) If  $\alpha$  is a limit ordinal,  $In_\alpha(\kappa, A) = \bigcap_{\beta < \alpha} In_\beta(\kappa, A)$ .

Thus  $X \in In_1(\kappa, \lambda)$  if and only if  $X \subset P_\kappa \lambda$  is ineffable. Clearly  $In_\alpha(\kappa, A) \subset In_\beta(\kappa, A)$  if  $\beta < \alpha$ . Hence there is  $\alpha$  such that  $In_\alpha(\kappa, A) = In_{\alpha+1}(\kappa, A)$ . If

$In_\alpha(\kappa, A) = In_{\alpha+1}(\kappa, A) \neq \emptyset$ ,  $P(P_\kappa A) \setminus In_\alpha(\kappa, A)$  is the minimal normal  $(\lambda, \lambda)$ -distributive ideal, that is, completely ineffable ideal on  $P_\kappa A$ .

**Definition 4.9.** We say  $I$  is precipitous if  $\Vdash_{\mathbb{P}_I}$  “ $Ult(V, G)$  is well-founded”.

**Lemma 4.10.** Let  $cf(\lambda) \geq \kappa$  and  $I$  be a normal  $(\lambda, \lambda)$ -distributive precipitous ideal on  $P_\kappa \lambda$ . For any  $X \in P(P_\kappa \lambda)$  and  $\alpha$ ,  $X \in In_\alpha(\kappa, \lambda)$  if and only if  $M \models “j''X \in In_\alpha(\kappa, j''\lambda)”$ .

*Proof.* By induction on  $\alpha$ . Assume first  $X$  is stationary,  $f : j''\lambda \times j''\lambda \rightarrow P_\kappa j''\lambda$ , and  $f \in M$ . Since  $P_\kappa j''\lambda = j''(P_\kappa \lambda)$ ,  $f = j(g) \upharpoonright j''\lambda \times j''\lambda$  for some  $g \in V$  such that  $g : \lambda \times \lambda \rightarrow P_\kappa \lambda$ . There is  $x \in X$  such that  $g''(x \times x) \subset P(x)$ . Then  $M \models “j(g)(a) \subset j(x)$  for all  $a \in j(x) \times j(x)”$ . Since  $j(x) = j''x \subset j''\lambda$ ,  $M \models “f(a) \subset j(x)$  for all  $a \in j(x) \times j(x)”$ . Hence  $M \models “j''X \in NS_{\kappa, j''\lambda}^+”$ .

Assume conversely  $M \models “j''X \in NS_{\kappa, j''\lambda}^+”$ ,  $f \in V$ , and  $f : \lambda \times \lambda \rightarrow P_\kappa \lambda$ . Since  $M \models “j(f) \upharpoonright (j''\lambda \times j''\lambda) : j''\lambda \times j''\lambda \rightarrow P_\kappa j''\lambda”$ , there is  $y \in j''X$  such that  $j(f)''(y \times y) \subset P(y)$ . For  $x \in X$  with  $j(x) = y$ ,  $j(f)(j(a)) \subset j(x)$  for every  $a \in x \times x$ . Hence  $f''(x \times x) \subset P(x)$ , showing that  $X$  is stationary.

The case  $\alpha$  is a limit ordinal is clear by our definition of  $In_\alpha(\kappa, \lambda)$  and  $In_\alpha(\kappa, j''\lambda)$ . We prove for  $\alpha = \beta + 1$ . Suppose first  $X \in In_\alpha(\kappa, \lambda)$ ,  $M \models “f : P_\kappa j''\lambda \rightarrow P_\kappa j''\lambda$  and  $f(j(x)) \subset j(x)$  for every  $x \in P_\kappa \lambda”$ . We have  $g \in V$  with  $f = j(g) \upharpoonright j''P_\kappa \lambda$ . Since  $g(x) \subset x$  for every  $x \in P_\kappa \lambda$ , there is  $S \subset \lambda$  such that  $Y := \{x \in X : g(x) = x \cap S\} \in In_\beta(\kappa, \lambda)$ . By inductive hypothesis  $j''Y \in In_\beta(\kappa, j''\lambda)$ . For every  $x \in Y$   $j(g)(j(x)) = j(g(x)) = j(x \cap S) = j''(x \cap S) = j(x) \cap j''S$  and  $j''S \in M$ . Now  $f(y) = y \cap j''S$  for every  $y \in j''Y$  hence  $j''X \in In_\alpha(\kappa, j''\lambda)$ .

Suppose second  $M \models “j''X \in In_\alpha(\kappa, j''\lambda)”$ ,  $f \in V$ , and  $f(x) \subset x$  for every  $x \in P_\kappa \lambda$ . Set  $j(f) \upharpoonright j''P_\kappa \lambda = g$ . Since  $M \models “g(x) \subset x$  for all  $x \in j''P_\kappa \lambda”$ , there is  $T \subset j''\lambda$  such that  $S := \{y \in j''X : g(y) = y \cap T\} \in In_\beta(\kappa, j''\lambda)$ . By the fact  $S \in P(P_\kappa j''\lambda)$  we have  $Y \in P(P_\kappa \lambda)^V$  with  $S = j''Y$ . For every  $x \in Y$   $g(j(x)) = j(x) \cap T$ . By the same reason for  $S$ ,  $T = j''T_1$  for some  $T_1 \in V$ . For every  $x \in Y$   $j(f(x)) = g(j(x)) = j(x) \cap j''T_1 = j''(x \cap T_1) = j(x \cap T_1)$ . So  $f(x) = x \cap T_1$  for all  $x \in Y$ . The inductive hypothesis shows  $Y \in In_\beta(\kappa, \lambda)$ . Hence  $X \in In_\alpha(\kappa, \lambda)$ .  $\square$

**Theorem 4.11.** Suppose  $cf(\lambda) \geq \kappa$  and  $I$  is a normal  $(\lambda, \lambda)$ -distributive precipitous ideal on  $P_\kappa \lambda$ . Then  $\{x : x \cap \kappa \text{ is completely o.t.}(x)\text{-ineffable}\} \in I^*$ .



*Proof.* Since  $V \models \text{“}\kappa \text{ is completely } \lambda\text{-ineffable”}$ ,  $In_\alpha(\kappa, \lambda) = In_{\alpha+1}(\kappa, \lambda) \neq \emptyset$  for some  $\alpha$ .  $In_\alpha(\kappa, \lambda) \neq \emptyset$  implies  $In_\alpha(\kappa, j''\lambda) \neq \emptyset$ . To show  $In_\alpha(\kappa, j''\lambda) = In_{\alpha+1}(\kappa, j''\lambda)$ , assume  $X \in In_\alpha(\kappa, j''\lambda) - In_{\alpha+1}(\kappa, j''\lambda)$ . Since  $X \in P(P_\kappa j''\lambda)^M$ , there is  $Y \in P(P_\kappa \lambda)^V$  such that  $X = j''Y$ . Then  $Y \in In_\alpha(\kappa, \lambda) = In_{\alpha+1}(\kappa, \lambda)$  hence  $X = j''Y \in In_{\alpha+1}(\kappa, j''\lambda)$ . Contradiction.  $\square$

**Lemma 4.12.** *Let  $\kappa$  be Mahlo,  $X \subset P_\kappa \lambda$ , and  $\alpha$  an ordinal.*

- (1) *If  $\{x \in P_\kappa \lambda : X \cap P_{x \cap \kappa} x \in In_\alpha(x \cap \kappa, x)\} \in In_\alpha(\kappa, \lambda)$ , then  $X \in In_\alpha(\kappa, \lambda)$ .*
- (2) *If  $X \in In_\alpha(\kappa, \lambda)$ , then  $\{x \in P_\kappa \lambda : X \cap P_{x \cap \kappa} x \notin In_\alpha(x \cap \kappa, x)\} \in In_\alpha(\kappa, \lambda)$ .*
- (3) *If  $X \subset P_\kappa \lambda$  is completely  $\lambda$  ineffable, then  $\{x \in X : X \cap P_{x \cap \kappa} x \text{ is not completely ineffable}\}$  is completely ineffable.*

*Proof.* (1) By induction on  $\alpha$ . Suppose that  $\{x \in P_\kappa \lambda : X \cap P_{x \cap \kappa} x \in NS_{x \cap \kappa, x}^+\} \in NS_{\kappa, \lambda}^+$  and  $f : \lambda^2 \rightarrow P_\kappa \lambda$ . There is  $x$  such that  $X \cap P_{x \cap \kappa} x \in NS_{x \cap \kappa, x}^+$  and  $f \upharpoonright x \times x : x \times x \rightarrow P(x)$ . Then we can find  $y \in X \cap P_{x \cap \kappa} x$  such that  $f''(y \times y) \subset P(y)$ . Hence  $X$  is stationary.

Let  $\alpha$  be a limit ordinal and  $Y := \{x \in P_\kappa \lambda : X \cap P_{x \cap \kappa} x \in In_\alpha(x \cap \kappa, x)\} \in In_\alpha(\kappa, \lambda)$ . For any  $\beta < \alpha$   $Y \subset \{x \in P_\kappa \lambda : X \cap P_{x \cap \kappa} x \in In_\beta(x \cap \kappa, x)\} \in In_\alpha(\kappa, \lambda) \subset In_\beta(\kappa, \lambda)$ . By induction hypothesis  $X \in In_\beta(\kappa, \lambda)$ . Thus  $X \in In_\alpha(\kappa, \lambda)$ .

Let  $\alpha = \beta + 1$  and  $f(x) \subset x$  for all  $x \in P_\kappa \lambda$ . For each  $x \in Y$  there is  $S_x \subset x$  such that  $\{y \in X \cap P_{x \cap \kappa} x : f(y) = y \cap S_x\} \in In_\beta(x \cap \kappa, x)$ . Since  $Y \in In_{\beta+1}(\kappa, \lambda)$  we have  $S \subset \lambda$  such that  $Z := \{x \in Y : S_x = x \cap S\} \in In_\beta(\kappa, \lambda)$ . For  $x \in Z$  and  $y \in X \cap P_{x \cap \kappa} x$   $f(y) = y \cap S_x = y \cap x \cap S = y \cap S$ . So  $\{x \in P_\kappa \lambda : \{y \in X : f(y) = y \cap S\} \cap P_{x \cap \kappa} x \in In_\beta(x \cap \kappa, x)\} \in In_\beta(\kappa, \lambda)$ . By induction hypothesis  $\{y \in X : f(y) = y \cap S\} \in In_\beta(\kappa, \lambda)$  hence  $X \in In_{\beta+1}(\kappa, \lambda)$ .

(2) We prove this by induction on  $\kappa$ . We may assume  $A := \{x \in X : X \cap P_{x \cap \kappa} x \in In_\alpha(x \cap \kappa, x)\} \in In_\alpha(\kappa, \lambda)$ . For any  $x \in A$ , by inductive hypothesis,  $B_x := \{y \in P_{x \cap \kappa} x : X \cap P_{x \cap \kappa} x \cap P_{y \cap \kappa} y = X \cap P_{y \cap \kappa} y \notin In_\alpha(y \cap \kappa, y)\} \in In_\alpha(x \cap \kappa, x)$ . By (1) and the fact  $B_x = \{y \in P_\kappa \lambda : X \cap P_{y \cap \kappa} y \notin In_\alpha(y \cap \kappa, y)\} \cap P_{x \cap \kappa} x$ , the conclusion holds.

(3) Let  $\beta$  be the least ordinal such that  $In_{\beta+1}(\kappa, \lambda) = In_\beta(\kappa, \lambda)$  and  $X \in In_\beta(\kappa, \lambda)$ . We may assume  $W := \{x \in X : X \cap P_{x \cap \kappa} x \text{ is completely ineffable}\} \in In_\beta(\kappa, \lambda)$ . For  $x \in W$  let  $\beta_x$  be the least ordinal such that  $In_{\beta_x+1}(x \cap \kappa, x) = In_{\beta_x}(x \cap \kappa, x) \neq \emptyset$ . By (2), for each  $x \in W$   $T_x := \{y \in P_{x \cap \kappa} x : X \cap P_{x \cap \kappa} x \cap P_{y \cap \kappa} y = X \cap P_{y \cap \kappa} y \notin In_{\beta_x}(y \cap \kappa, y)\} \in In_{\beta_x}(x \cap \kappa, x)$ . Set  $\gamma = \max(\beta, \bigcup_{x \in W} \beta_x)$  and  $T = \{x \in P_\kappa \lambda :$

$X \cap P_{x \cap \kappa} x \notin \text{In}_\gamma(x \cap \kappa, x)$ . Then  $T_x \subset T \cap P_{x \cap \kappa} x$  for every  $x \in W$ . Hence  $T \in \text{In}_\gamma(\kappa, \lambda) = \text{In}_\beta(\kappa, \lambda)$   $\square$

Now we conclude by the following corollary:

**Corollary 4.13.** *If  $cf(\lambda) \geq \kappa$ , then the completely ineffable ideal on  $P_\kappa \lambda$  is not precipitous.*

## 5 Ineffability and almost ineffability

In this section we show  $\text{NAIn}_{\kappa, \lambda} = \text{NIn}_{\kappa, \lambda}$  if  $cf(\lambda) < \kappa$  and  $\lambda^{<\kappa} = 2^\lambda$ . Thus, two ideals are the same for “small” cofinality points if GCH holds.

Then we use it to prove that for  $\lambda$  with cofinality less than  $\kappa$ , “ $\kappa$  is  $\lambda$ -ineffable” does not always imply “ $\kappa$  is  $\lambda^{<\kappa}$ -ineffable”. This contrasts to the fact “ $\kappa$  is  $\lambda^{<\kappa}$ -(super)compact if  $\kappa$  is  $\lambda$ -(super)compact”.

**Fact 5.1.** (1)  $\text{WNS}_{\kappa, \lambda} \subsetneq \text{NAIn}_{\kappa, \lambda} \subset \text{NIn}_{\kappa, \lambda}$ .

(2) If  $P_\kappa \lambda \notin \text{NAIn}_{\kappa, \lambda}$  and  $cf(\lambda) \geq \kappa$ , then  $\lambda^{<\kappa} = \lambda$ .

(3) If  $cf(\lambda) \geq \kappa$ , then  $\{x \in P_\kappa \lambda : P_{x \cap \kappa} x \in \text{NAIn}_{x \cap \kappa, x}\} \in \text{NIn}_{\kappa, \lambda}$  and  $\text{NAIn}_{\kappa, \lambda} \subsetneq \text{NIn}_{\kappa, \lambda}$ .

We have known quite little when  $cf(\lambda) < \kappa$  while the following conjecture has seemed reasonable comparing with supercompactness:

**Conjecture 5.2.** *Suppose that  $cf(\lambda) < \kappa$  and  $\kappa$  is (almost)  $\lambda$ -ineffable. Then,  $\lambda^{<\kappa} = \lambda^+$  and  $\kappa$  is  $\lambda^+$ -ineffable.*

**Lemma 5.3.** *If  $\lambda^{<\kappa} = 2^\lambda$ , then there exists  $X \in \text{WNS}_{\kappa, \lambda}^*$  such that  $I_{\kappa, \lambda} \upharpoonright X = \text{NS}_{\kappa, \lambda} \upharpoonright X$ .*

*Proof.* Let  $\lambda^{\times \lambda} = \{f_s : s \in P_\kappa \lambda\}$  be an enumeration and set  $X = \{x \in P_\kappa \lambda : x \in \bigcap \{C_{f_s} : s \prec x\}\}$ . For every  $s \in P_\kappa \lambda$   $C_{f_s} \in \text{NS}_{\kappa, \lambda}^* \subset \text{WNS}_{\kappa, \lambda}^*$  hence  $X \in \text{WNS}_{\kappa, \lambda}^*$ . Let  $Y \notin I_{\kappa, \lambda} \upharpoonright X$ . For every club  $C \subset P_\kappa \lambda$  there exists  $s \in P_\kappa \lambda$  such that  $C_{f_s} \subset C$ . We have  $x \in Y \cap X$  with  $s \prec x$ . Then,  $x \in C_{f_s} \cap Y \cap X \subset C \cap Y \cap X$  hence  $Y \cap X \in \text{NS}_{\kappa, \lambda}^+$ .  $\square$

Since  $\text{WNS}_{\kappa, \lambda} \subset \text{NAIn}_{\kappa, \lambda}$  we have:

**Theorem 5.4.**  $NAIn_{\kappa,\lambda} = NIn_{\kappa,\lambda}$  if  $\lambda^{<\kappa} = 2^\lambda$ .

By lemma 4.12 we have the following:

**Lemma 5.5.** If  $\kappa$  is  $\lambda$ -ineffable, then  $\{x \in P_\kappa\lambda : P_{x \cap \kappa}x \in NIn_{x \cap \kappa, x}\} \notin NIn_{\kappa,\lambda}$ .

**Theorem 5.6.** Suppose that  $\lambda^{<\kappa} = 2^\lambda$  and  $\kappa$  is  $\lambda^+$ -ineffable. Then,  $\{x \in P_\kappa\lambda^+ : x \cap \kappa \text{ is } o.t.(x \cap \lambda)\text{-ineffable, not } o.t.(x)\text{-ineffable, and } o.t.(x) = o.t.(x \cap \lambda)^+\} \notin NIn_{\kappa,\lambda^+}$ .

*Proof.* We know  $A = \{x \in P_\kappa\lambda^+ : x \cap \kappa \text{ is almost } o.t.(x)\text{-ineffable, } o.t.(x) = o.t.(x \cap \lambda)^+, cf(o.t.(x \cap \lambda)) < x \cap \kappa, \text{ and } o.t.(x \cap \lambda)^{<x \cap \kappa} = 2^{o.t.(x \cap \lambda)}\} \in NIn_{\kappa,\lambda^+}^*$  ([12], [2]). Every  $x \in A$  is almost  $o.t.(x \cap \lambda)$ -ineffable hence  $o.t.(x \cap \lambda)$ -ineffable by the previous theorem. Now the conclusion follows by lemma 5.5.  $\square$

We conclude by the negation of  $I_{\kappa,\lambda}^+ \rightarrow (I_{\kappa,\lambda}^+)^2$  with  $\lambda^{<\kappa} = 2^\lambda$ .

**Definition 5.7.** For  $X \subset P_\kappa\lambda$  let  $[X]^2 = \{(x, y) \in X \times X : x \subsetneq y\}$ . We say  $I_{\kappa,\lambda}$  has the partition property if for every  $X \notin I_{\kappa,\lambda}$  and  $F : [X]^2 \rightarrow 2$  there exists  $H \notin I_{\kappa,\lambda}$  such that  $F \upharpoonright [H]^2$  is constant.

**Theorem 5.8.** If  $\lambda^{<\kappa} = 2^\lambda$ , then  $I_{\kappa,\lambda}$  does not have the partition property.

*Proof.* Suppose otherwise and let  $I = I_{\kappa,\lambda} \upharpoonright X = NS_{\kappa,\lambda} \upharpoonright X$  with  $X$  as in [?]. For every  $X \in I^+$  and  $F : [X]^2 \rightarrow 2$  there exists  $H \notin NS_{\kappa,\lambda}$  such that  $F \upharpoonright [H]^2$  is constant. Hence  $NIn_{\kappa,\lambda} \subset I$  ([15]). However  $I \subset WNS_{\kappa,\lambda} \subsetneq NIn_{\kappa,\lambda}$ . Contradiction.  $\square$

**Remark 5.9.** P. Matet proved that  $I_{\kappa,\kappa^+}$  does not have the partition property if  $2^\kappa = \kappa^+$ . While M. Shioya [17] constructed the model in which  $I_{\kappa,\lambda}$  has the partition property with  $\kappa$  supercompact.

## 参考文献

- [1] Y. Abe, *Saturation of fundamental ideals on  $P_\kappa\lambda$* , J. Math. Soc. Japan 48 (1996), 511-524.
- [2] Y. Abe, *Combinatorial characterization of  $\Pi_1^1$ -indescribability in  $P_\kappa\lambda$* , Arch. Math. Logic 37 (1998), 261-272.

- [3] S. Baldwin, *The consistency strength of certain stationary subsets of  $P_\kappa\lambda$* , Proc. Amer. Math. Soc. 92 (1984), 90-92.
- [4] J. Baumgartner, *Ineffability properties of cardinals 1*, Infinite and finite sets (P. Erdős 60th Birthday Colloquium, Keszthely, Hungary, 1973), Colloquia Mathematica Societatis János Bolyai, vol. 10, North-Holland, Amsterdam (1975), 109-130.
- [5] D. M. Carr, *The minimal normal filter on  $P_\kappa\lambda$* , Proc. Amer. Math. Soc. 86 (1982), 316-320.
- [6] D. M. Carr, *The structure of ineffability properties of  $P_\kappa\lambda$* , Acta Math. Hung. 47 (1986), 325-332.
- [7] D. M. Carr,  *$P_\kappa\lambda$ -generalizations of weak compactness*, Z. Math. Logik Grundlagen Math. 31 (1985), 393-401.
- [8] D. M. Carr, J. P. Levinski and D. H. Pelletier, *On the existence of strongly normal ideals on  $P_\kappa\lambda$* , Arch. Math. Logic 30 (1990), 59-72.
- [9] H. D. Donder and P. Matet, *Two cardinal versions of diamond*, Israel J. Math. 83 (1993), 1-43.
- [10] T. Jech, *Some combinatorial problems concerning uncountable cardinals*, Ann. Math. Logic 5 (1973), 165-198.
- [11] C. A. Johnson, *Some partition relations for ideals on  $P_\kappa\lambda$* , Acta Math. Hung. 56 (1990), 269-282.
- [12] S. Kamo, *Remarks on  $P_\kappa\lambda$ -combinatorics*, Fund. Math. 145 (1994), 141-151.
- [13] J. Ketonen, *Some combinatorial principles*, Trans. Amer. Math. Soc. 188 (1974), 387-394.
- [14] J. P. Levinski, *Instances of the conjecture of Chang*, Israel J. Math. 48 (1984), 225-243.
- [15] M. Magidor, *Combinatorial characterization of supercompact cardinals*, Proc. Amer. Math. Soc. 42 (1974), 279-285.

- [16] T. K. Menas, *On strong compactness and supercompactness*, Ann. Math. Logic 7 (1974), 327-359.
- [17] M. Shioya, *Partition properties of subsets of  $P_\kappa\lambda$* , Fund. Math. 161 (1999), 325-329.