

Title	Weak decays of He-4(Lambda)
Author(s)	Parker, JD; Athanas, MJ; Barnes, PD; Bart, S; Bassalleck, B; Chrien, R; Diebold, G; Franklin, GB; Johnston, K; Pile, P; Quinn, B; Rozon, F; Sawafta, R; Schumacher, RA; Stearns, R; Sukaton, I; Sutter, R; Szymanski, JJ; Zeps, VJ
Citation	PHYSICAL REVIEW C (2007), 76(3)
Issue Date	2007-09
URL	http://hdl.handle.net/2433/49820
Right	Copyright 2007 American Physical Society
Type	Journal Article
Textversion	publisher

Weak decays of ${}^4_{\Lambda}\text{He}$

J. D. Parker,^{1,*} M. J. Athanas,^{1,†} P. D. Barnes,^{1,‡} S. Bart,² B. Bassalleck,³ R. Chrien,² G. Diebold,¹ G. B. Franklin,¹ K. Johnston,^{4,§} P. Pile,² B. Quinn,¹ F. Rozon,¹ R. Sawafta,^{2,||} R. A. Schumacher,¹ R. Stearns,⁵ I. Sukaton,¹ R. Sutter,² J. J. Szymanski,^{6,‡} and V. J. Zeps^{1,¶}

¹*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*

²*Brookhaven National Laboratory, Upton, New York 11973, USA*

³*University of New Mexico, Albuquerque, New Mexico 87131, USA*

⁴*University of Houston, Houston, Texas 77004, USA*

⁵*Vassar College, Poughkeepsie, New York 12604, USA*

⁶*Indiana University Cyclotron Facility, Bloomington, Indiana 47408, USA*

(Received 21 May 2007; published 4 September 2007; corrected 12 September 2007)

We measured the lifetime and the mesonic and nonmesonic decay rates of the ${}^4_{\Lambda}\text{He}$ hypernucleus. The hypernuclei were created using a 750 MeV/c momentum K^- beam on a liquid ${}^4\text{He}$ target by the reaction ${}^4\text{He}(K^-, \pi^-){}^4_{\Lambda}\text{He}$. The ${}^4_{\Lambda}\text{He}$ lifetime was directly measured using protons from $\Lambda p \rightarrow np$ nonmesonic decay (also referred to as proton-stimulated decay) and was found to have a value of $\tau = 245 \pm 24$ ps. The mesonic decay rates were determined from the observed numbers of π^- 's and π^0 's as $\Gamma_{\pi^-}/\Gamma_{\text{tot}} = 0.270 \pm 0.024$ and $\Gamma_{\pi^0}/\Gamma_{\text{tot}} = 0.564 \pm 0.036$, respectively, and the values of the proton- and neutron-stimulated decay rates were extracted as $\Gamma_p/\Gamma_{\text{tot}} = 0.169 \pm 0.019$ and $\Gamma_n/\Gamma_{\text{tot}} \leq 0.032$ (95% CL), respectively. The effects of final-state interactions and possible three-body ΛNN decay contributions were studied in the context of a simple model of nucleon-stimulated decay. Nucleon-nucleon coincidence events were observed and were used in the determination of the nonmesonic branching fractions. The implications of the results of this analysis were considered for the empirical $\Delta I = \frac{1}{2}$ rule and the decay rates of the ${}^4_{\Lambda}\text{H}$ hypernucleus.

DOI: [10.1103/PhysRevC.76.035501](https://doi.org/10.1103/PhysRevC.76.035501)

PACS number(s): 21.80.+a, 13.75.Ev, 27.10.+h

I. INTRODUCTION

Studies of the hyperon-nucleon interaction investigate weak processes that are not well understood in multiple baryon systems. In particular, the study of nonmesonic decays of light hypernuclei is the only tractable method of investigating the $\Delta S = 1$ weak baryon-baryon interaction. By comprehensive measurements in s -shell hypernuclei, various ΛN initial states can be investigated. Few-body nuclear structure calculations can then relate the decay rates to the underlying weak ΛN interactions. These studies are complementary to investigations of the parity-violating contributions to the nucleon-nucleon interaction. However, the ΛN system can be prepared in a particle-stable state so that only weak processes contribute to the decay rate.

A particularly interesting aspect of $\Delta S = 1$ transitions is the empirical $\Delta I = \frac{1}{2}$ rule. We note that if strong interactions are

neglected, the $\Delta S = 1$ nonleptonic decays of hyperon systems would be expected to be dominated by a weak quark-level Hamiltonian based on one W -exchange of the form:

$$H_{\text{weak}} = \frac{G_f}{\sqrt{2}} \sin \theta_c \cos \theta_c [\bar{u}\gamma_{\mu}(1 - \gamma_5)s\bar{d}\gamma^{\mu}(1 - \gamma_5)u]. \quad (1)$$

The isospin character of this fundamental Hamiltonian does not favor $\Delta I = \frac{1}{2}$ weak transitions over the $\Delta I = \frac{3}{2}$ ones. However, it has long been recognized that the $\Delta I = \frac{1}{2}$ transitions dominate all observed nonleptonic decays of strange hadrons. Although it is generally believed that the presence of the strong interaction accounts for this empirical $\Delta I = \frac{1}{2}$ selection rule, the various explanations for the dominance of the $\Delta I = \frac{1}{2}$ amplitudes [1,2] have indicated that the mechanism may be specific to the given system and not a universal feature of the $\Delta S = 1$ weak Hamiltonian. It remains an open question whether all $\Delta S = 1$ nonleptonic decays obey the rule or it is specific to the pionic decay channels (the only tested class of nonleptonic decays). The study of hypernuclear decays provides an opportunity to answer this question by providing the only experimentally accessible nonpionic test of the $\Delta I = \frac{1}{2}$ rule.

The total decay width of a hypernucleus includes contributions from mesonic and nonmesonic decay modes. The mesonic decay modes, defined experimentally by the observation of the appropriate pion in the final state, are dominated by single-body processes analogous to the decay of the free Λ , $\Lambda \rightarrow \pi^- p$ and $\Lambda \rightarrow \pi^0 n$, along with additional multibody contributions such as $\Lambda p \rightarrow nn\pi^+$. The nonmesonic modes are expected to be dominated by the proton-stimulated reaction

*Present address: Department of Physics, Kyoto University, Kyoto 606-8502, Japan; jparker@scphys.kyoto-u.ac.jp

†Present address: VAST Scientific, Cambridge, Massachusetts 02139, USA.

‡Present address: Los Alamos National Laboratory, Los Alamos, New Mexico 87544, USA.

§Present address: Louisiana Tech University, Ruston, Louisiana 71272, USA.

||Present address: Quartek Corporation, Greensboro, NC 27410, USA.

¶Present address: Bluegrass Community and Technical College, Lexington, Kentucky 40506, USA.

$\Lambda p \rightarrow np$ and neutron-stimulated reaction $\Lambda n \rightarrow nn$ with rates Γ_p and Γ_n , respectively. Other multibody interactions such as $\Lambda NN \rightarrow NNN$ may also contribute with rate Γ_{mb} . Determination of the nucleon-stimulated decay rates is more complicated than the mesonic case and is ultimately model dependent. The total decay width can be written in terms of the widths of these mesonic and nonmesonic decay modes as:

$$\Gamma_{\text{total}} = \underbrace{\Gamma_{\text{mesonic}}}_{\Gamma_{\pi^-} + \Gamma_{\pi^0} + \Gamma_{\pi^+}} + \underbrace{\Gamma_{\text{nonmesonic}}}_{\Gamma_p + \Gamma_n + \Gamma_{mb}}, \quad (2)$$

where contributions from semileptonic and weak radiative Λ decays have been neglected as they contribute only about 0.3% to the total free Λ decay width [3].

The two-body nonmesonic decay modes, $\Lambda N \rightarrow NN$, are readily distinguishable from the mesonic decay modes, $\Lambda \rightarrow \pi N$, because of the large energy ($M_\Lambda - M_n = 176$ MeV) available for the final-state nucleons. These modes are sensitive to weak interaction couplings (such as $g_{\Lambda N \rho}$ or g_{NNK}) not available to the free hyperon decays. Also, it has been suggested that several of the weak $\Lambda N \rightarrow NN$ amplitudes are dominated by direct-quark processes [4–6] in which no intermediate meson is present in the interaction. Such direct processes may not adhere to the $\Delta I = \frac{1}{2}$ rule.

In nuclei the Λ -particle is not Pauli blocked, thus hypernuclei generally decay with the Λ -particle initially in the lowest $1s$ -state. To determine properties of the underlying $\Lambda N \rightarrow NN$ interaction, it is convenient to consider systems that confine the nucleons to the lowest orbitals. Figure 1 shows the spin-isospin character of the initial (ΛN) and final (nN) states for s -shell hypernuclei in a single-particle shell-model picture. The $I = 1$ final states are accessible to both proton- and neutron-stimulated decays, whereas the $I = 0$ final states are available only for proton-stimulated decay. Further complications arise as one increases A because the high probability of final-state interactions dilutes the correlation between nucleons observed and nucleons responsible for the decay. Table I shows that a complete set of measurements of the nonmesonic decay widths for the s -shell hypernuclei allows one to isolate specific ΛN initial states. Thus, one can determine the spin-isospin structure of the fundamental $\Lambda N \rightarrow nN$ weak interaction by combining measurements of these few-body hypernuclei with detailed finite nucleus calculations that take into account the differences in the initial- and final-state phase space. However, at this time the experimental data are incomplete and often have large estimated errors.




FIG. 1. Spin-isospin selection rules for the $\Lambda N \rightarrow nN$ process in $1s$ -shell hypernuclei.

TABLE I. Allowed initial states for $1s$ -shell hypernuclei.

Species	Λn	Λp
${}^5_\Lambda\text{He}$	${}^3S_1, {}^1S_0$	${}^3S_1, {}^1S_0$
${}^4_\Lambda\text{He}$	1S_0	${}^3S_1, {}^1S_0$
${}^4_\Lambda\text{H}$	${}^3S_1, {}^1S_0$	1S_0
${}^3_\Lambda\text{H}$	${}^3S_1, {}^1S_0$	${}^3S_1, {}^1S_0$

A phenomenological model first put forth by Block and Dalitz [7,8] can be used to relate the nonmesonic decay rates of the s -shell hypernuclei listed in Table I. Specifically, the current experimental results for ${}^4_\Lambda\text{He}$ (this work) and ${}^5_\Lambda\text{He}$ (Ota *et al.* [9]) can be extended to the ${}^4_\Lambda\text{H}$ hypernucleus, for which measurements are currently scarce. The model employs the mean nucleon density, ρ_A , at the location of the Λ and makes several assumptions concerning the ΛN interaction: (1) the nucleon-stimulated Λ decay is treated as incoherent, (2) final-state interactions are not included, and (3) three-body nucleon-stimulated decays are neglected. These assumptions are seen to be adequate at the present level of accuracy of the experimental quantities. Defining the quantity R_{NS} as the rate for an N -stimulated decay originating from ΛN relative spin state S , the nonmesonic decay rates for the s -shell hypernuclei ${}^4_\Lambda\text{H}$, ${}^4_\Lambda\text{He}$, and ${}^5_\Lambda\text{He}$ are derived as:

$$\Gamma_{nm}({}^4_\Lambda\text{H}) = \frac{\rho_A^4}{6}(R_{n0} + 3R_{n1} + 2R_{p0}), \quad (3)$$

$$\Gamma_{nm}({}^4_\Lambda\text{He}) = \frac{\rho_A^4}{6}(2R_{n0} + R_{p0} + 3R_{p1}), \quad (4)$$

$$\Gamma_{nm}({}^5_\Lambda\text{He}) = \frac{\rho_A^5}{8}(R_{n0} + 3R_{n1} + R_{p0} + 3R_{p1}), \quad (5)$$

where the mean nucleon density, ρ_A , is defined as:

$$\rho_A \equiv \int \rho_A(\vec{r}) |\psi(\vec{r})|^2 d\vec{r}, \quad (6)$$

for the nucleon density, $\rho_A(\vec{r})$, and the Λ wave function, $\psi(\vec{r})$. Also, it has been assumed that the mean nucleon density, ρ_A , has the same value for both ${}^4_\Lambda\text{H}$ and ${}^4_\Lambda\text{He}$.

Taking the ratio of the rates of neutron-stimulated ${}^4_\Lambda\text{He}$ decay to proton-stimulated ${}^4_\Lambda\text{H}$ decay yields:

$$\frac{\Gamma_n({}^4_\Lambda\text{He})}{\Gamma_p({}^4_\Lambda\text{H})} = \frac{R_{n0}}{R_{p0}} = 2, \quad (7)$$

where the value of 2 is the $\Delta I = \frac{1}{2}$ rule prediction for this ratio. Similarly, an expression for the ${}^4_\Lambda\text{H}$ neutron-stimulated rate can be found by considering the ratio of neutron-stimulated ${}^4_\Lambda\text{H}$ decay to proton-stimulated ${}^4_\Lambda\text{He}$ decay:

$$\frac{\Gamma_n({}^4_\Lambda\text{H})}{\Gamma_p({}^4_\Lambda\text{He})} = \frac{R_{n0} + 3R_{n1}}{R_{p0} + 3R_{p1}} = \frac{\Gamma_n({}^5_\Lambda\text{He})}{\Gamma_p({}^5_\Lambda\text{He})}, \quad (8)$$

independent of any assumptions about the $\Delta I = \frac{1}{2}$ rule.

In this article, we present the results of Brookhaven National Laboratory experiment E788, which measured the lifetime and partial widths of the ${}^4_\Lambda\text{He}$ hypernucleus. Section II presents the experimental apparatus used for these measurements,

and a discussion of the out-of-beam tracking and particle identification algorithm is given in Sec. III. The excitation energy spectra for the ${}^4\text{He}(K^-, \pi^-){}^4_{\Lambda}\text{He}$ production reaction are discussed in Sec. IV. The extraction of the lifetime of ${}^4_{\Lambda}\text{He}$ and particle-emission spectra are presented in Secs. V and VI, respectively. The determination of the ${}^4_{\Lambda}\text{He}$ mesonic and nonmesonic partial decay rates is discussed in Sec. VII. In the case of the nonmesonic proton- and neutron-stimulated decays, the rates are found from both the single-particle kinetic energy spectra of Sec. VI as well as from observed multiple-nucleon coincidence events, giving two essentially independent determinations. A comparison with the results of other experiments and theoretical calculations and the extension of these results to the ${}^4_{\Lambda}\text{H}$ hypernucleus is presented in Sec. VIII.

II. EXPERIMENTAL SETUP

The partial decay rates and lifetime of ${}^4_{\Lambda}\text{He}$ were measured using the LESB-II kaon beamline of the AGS at Brookhaven National Laboratory. The beamline parameters are given in Table II. The experimental layout is shown in Fig. 2. A beam of 750 MeV/c kaons incident on a liquid helium target produced ${}^4_{\Lambda}\text{He}$ through the ${}^4\text{He}(K^-, \pi^-){}^4_{\Lambda}\text{He}$ reaction, and pions from the production reaction were detected near 0° in the Moby Dick spectrometer. Detector packages located on each side of the target area were used to detect and identify pions, protons, neutrons, and γ 's from the hypernuclear decays.

The incident kaon trajectories were determined with drift chambers located just upstream of the target (ID2X through ID7U). This information was combined with data from the drift chamber ID1X, located upstream of the K^- spectrometer, to determine the incident kaon momentum. Although the kaon-to-pion ratio at the target was only 1:13, the kaons were cleanly separated using a combination of the time-of-flight (TOF) between scintillator IS1 (not shown) located upstream of the

TABLE II. LESB-II and Moby Dick spectrometer description.

Item	Description
LESB-II separated beamline/spectrometer	
Momentum	750 MeV/c
Momentum acceptance	5% FWHM
$\pi : K$ ratio at target	13 : 1
K^- flux	$2 \times 10^5 \text{s}^{-1}$
Momentum measurement	Drift chambers ID1X – ID7U and hodoscope MH
$\pi - K$ Particle ID	IS1–IT1, 2 TOF scintillators and CP Čerenkov π -veto
Moby Dick spectrometer	
Momentum	680 MeV/c
Momentum acceptance	5% FWHM
Momentum measurement	Drift chambers ID9Y – ID15Y
$\pi - K$ particle ID	IT1,2 – IS2,3 TOF scintillators

K^- spectrometer and scintillators IT1 and IT2 and a critical angle Čerenkov detector (CP). The outgoing pion momentum was measured with the Moby Dick spectrometer located at zero degrees downstream of the target, and the outgoing particle identification was accomplished with time-of-flight measurements.

The 30.5-cm-long liquid helium target vessel consisted of a 6.35-cm-diameter Dacron cylinder with a wall thickness of 0.020 cm. To minimize material between the decay vertex and the out-of-beam decay-particle detectors, the vacuum vessel containing the Dacron shell was constructed from a PVC foam cylinder of density 0.053 gm/cm^3 with inner and outer radii of 7.6 cm and 11.43 cm, respectively.

The separation of the decay protons and pions was accomplished by combining the crude range information from all




FIG. 2. Shown here is the E788 experimental apparatus used to carry out the ${}^4_{\Lambda}\text{He}$ hypernuclear decay measurement. Also shown are the focusing quadrupoles at the exit of the K^- spectrometer and at the entrance of the Moby Dick spectrometer.

TABLE III. Descriptions of the target area detectors utilized for the hypernuclear decay measurement. An overview of the layout of these detector elements is shown in Fig. 2.

Name	Function	Dimensions (cm ³)
Drift chambers		
ID	$K - \pi$ trajectories	$12.2 \times 12.2 \times 0.43$
OD	y and z track positions (1 pair each side)	$30.5 \times 30.5 \times 1.9$
Scintillation counters		
IT1, IT2	$K - \pi$ TOF and ${}^4_\Lambda\text{He}$ formation time	$15.0 \times 4.0 \times 1.27$
CP	critical angle pion Čerenkov counter	$15.0 \times 4.0 \times 0.6$
OT1	decay time and dE/dx (5 each side)	$22.0 \times 4.5 \times 0.64$
OT2	TOF, dE/dx , and range (10 each side)	$150.0 \times 12.0 \times 1.5$
NV	range and charged particle veto (9 each side)	$182.9 \times 22.9 \times 0.95$
NA	charged particle veto (2 each side)	$182.9 \times 22.9 \times 0.95$
ND	range and neutral particle detection (50 each side)	$182.9 \times 15.2 \times 5.1$

out-of-beam detectors with the measured rate of energy loss (dE/dx), total energy loss, and the measured TOF between the decay-timing and TOF scintillator layers (also referred to in this article as OT1 and OT2, respectively). The precision timing scintillators of the decay-timing layer also played an important part in the determination of the lifetime of the ${}^4_\Lambda\text{He}$ hypernucleus. With no magnetic field in the region of the out-of-beam detectors, the detector system of E788 was unable to distinguish a π^- from a π^+ , making it possible to measure only the sum of the two widths Γ_{π^-} and Γ_{π^+} . However, the width Γ_{π^+} has been seen to be small compared to the π^- decay width ($\Gamma_{\pi^+}/\Gamma_{\pi^-} = 0.043 \pm 0.017$ [10]) and was ignored in this analysis. Large-volume ($\approx 1.4\text{m}^3$) neutron detector arrays were placed near the target to directly measure the decay neutrons and were used to determine the neutron energies by TOF. The neutron detector arrays were also used to detect the γ 's from the decay of π^0 's originating from ${}^4_\Lambda\text{He}$ decay. Thus, this experiment directly measured Γ_{total} , Γ_{π^-} , Γ_{π^0} , Γ_p , and Γ_n , along with the kinetic energy spectra for the decay π^- 's, protons, and neutrons. Details of the detector dimensions are given in Table III.

III. OUT-OF-BEAM TRACK RECONSTRUCTION AND IDENTIFICATION

The task of the out-of-beam tracking was to sort through the scintillator hits recorded in the out-of-beam detector arrays and determine the trajectories of the charged and neutral particles that created them. A hit in the out-of-beam detector array was defined as a coincidence of signals in the photomultiplier tubes at each end of a scintillator element, requiring ADC and TDC information for both of the PMTs. Each hit was characterized by a hit time, energy deposition, and hit position. For a hit to be considered for tracking, the recorded time (measured relative to the event start-time as set by the mean time of in-beam scintillators IT1 and IT2) and measured energy deposition must have been greater than zero.

As a preliminary step, neutron detector hits that satisfied the rudimentary conditions described above were examined, and correlated hits were grouped into *clusters* based on relative hit

times and positions (i.e., groups of hits that were sufficiently close in time and space were considered to have arisen from a single particle). The hit clusters were classified as *charged* if the group included any hits in the OT2 or NV layers; otherwise, the cluster was considered to have originated from a neutral particle.

Charged tracks were indicated by a coincidence of hits in the OT1 and OT2 layers. The charged tracking began with a search for hits in the OT1 layer, and for each OT1 hit found, a search was made for corresponding hits in OT2. If the hit pairs were properly time ordered (i.e., the OT1 hit preceded the OT2 hit), they were considered to be part of a charged track and were then subjected to a more detailed tracking procedure. The final track was found from a linear least-squares fit to the OT1 and OT2 hits, the $K\pi$ reaction vertex, out-of-beam drift chamber hits, and any hits in the neutron veto layer or neutron detector array that belonged to the same cluster as the OT2 hit. Interesting quantities such as the particle's velocity between the OT1 and OT2 layers and the particle's range and total energy loss in the out-of-beam detector system were then determined.

The tracking information found above was used to distinguish protons from pions, a task that was complicated by the absence of a momentum-analyzing magnetic field. To determine the identity of the charged particles, the measured velocity, range, and energy loss of a charged track were compared with the expected behavior of the hypernuclear decay products via the quantity:

$$D_{\pi,p} = \left\{ \left[\frac{\Delta E_{\text{meas}} - \Delta E_{\pi,p}(\beta_{\text{fit}})}{\sigma_E} \right]^2 + \left(\frac{\beta_{\text{meas}} - \beta_{\text{fit}}}{\sigma_\beta} \right)^2 \right\}^{\frac{1}{2}}, \quad (9)$$

where ΔE_{meas} is the measured energy deposition in the OT2 scintillator layer, β_{meas} is the velocity fraction as measured in the time-of-flight region of the out-of-beam detector array, and $\sigma_E \sim 1/\sqrt{E}$ and $\sigma_\beta \sim \beta^2$ are the resolutions for the measured quantities. To find the quantity β_{fit} (and thus the theoretical energy deposition $\Delta E_{\pi,p}$), the above expression for $D_{\pi,p}$ was minimized by varying the value of β_{fit} within limits determined

by the measured range of the particle. The PID (particle identification) for a given track was then determined by testing the values of D_{π} (calculated assuming the track was created by a pion) and D_p (calculated assuming the track was created by a proton) against predetermined limits. The particle was then classified as either a proton, a pion, or an *unknown* (neither D_{π} nor D_p was within the appropriate limits). The unknown particles consisted mostly of electrons and were well separated from the pions and protons.

A neutral particle appeared as a single hit or small cluster of hits in the out-of-beam detector arrays, providing little information for tracking. A neutral *track* was taken as a straight line connecting the $K\pi$ reaction vertex with the out-of-beam neutral cluster, where the time and position of the reaction vertex were known from the in-beam tracking. For each of the neutral clusters found in the preliminary grouping, a cluster hit time, position, and energy were calculated. The hit time and position were determined by the member hit that was closest to the target, and the energy was taken as the sum of the energy deposition for all member hits. Additionally, each cluster was characterized by a cluster size defined simply as the total number of member hits. Once the neutral track was determined by the production vertex and cluster hit position, the velocity fraction of the neutral particle could be calculated.

The next task was to separate neutrons from γ 's and attempt to suppress the large accidental neutral background. To reduce accidentals, an energy deposition threshold of $E_{\text{dep}} > 5 \text{ MeVee}$ ($\text{MeV electron equivalent}$) was applied to the neutral clusters, and clusters with corresponding hits in the auxiliary veto counters NA were discarded. Good neutron candidate events were then selected by requiring the neutral cluster to have fewer than three member hits and a value for the neutron kinetic energy (as derived from the measured time-of-flight) from 50 to 200 MeV. The lower bound at 50 MeV avoids the energy region where accidentals dominate, and the upper bound at 200 MeV, which is well above the maximum expected kinetic energy for neutrons from hypernuclear decay ($\sim 170 \text{ MeV}$), removes γ 's from consideration. Good γ events were selected by requiring the measured β of the candidate track be in the range $0.666 < \beta < 2.0$, where the lower bound at 0.666 is well above the highest β expected for neutrons from hypernuclear decay. Even with the cuts described above, a sizable background still remained in the neutron and γ samples. The subtraction of this background will be discussed in subsequent sections. A cancellation effect inherent in the subtraction of the background reduces the sensitivity of the final results to the selection of the cut parameters given above.

IV. ${}^4_{\Lambda}\text{He}$ EXCITATION SPECTRA

Reconstruction of the K^- and π^- trajectories through their respective spectrometers allowed the calculation of the invariant mass of the unobserved strange system. Unlike p -shell hypernuclei, the ground state is also a substitutional state and thus the (K^-, π^-) reaction near zero degrees is ideal for creating ${}^4_{\Lambda}\text{He}$. The main source of background for the production reaction was in-flight kaon decays. Those decays




FIG. 3. The excitation energy spectrum for ${}^4\text{He}(K^-, \pi^-)X$ is shown for several cases: (a) unbiased by out-of-beam tags, (b) tagged by a proton in the out-of-beam detector package, (c) tagged by a pion, and (d) tagged by a neutron. Overlaid on each plot is the result of the fit to the ground-state peak (including appropriate backgrounds as described in the text).

that may satisfy the (K^-, π^-) trigger condition include (i) $K^- \rightarrow \pi^- \pi^0$, (ii) $K^- \rightarrow \mu^- \bar{\nu}_\mu$, and (iii) $K^- \rightarrow \mu^- \bar{\nu}_\mu \pi^0$. For kaons that decayed within the area of the target, the two-body decays were easily removed using the decay kinematics, whereas the three-body decays had to be modeled with a Monte Carlo simulation and fit to the data. Some kaons decayed within the beamline chambers or Moby Dick spectrometer or traversed the experimental area completely. These events were characterized by a small apparent $K\pi$ scattering angle and were removed by requiring an angle greater than 25 mrad. Events with production vertices outside the target region or poor beam track reconstruction were also rejected.

The resulting excitation energy spectrum for unbiased (K^-, π^-) events is shown in Fig. 3(a), where *unbiased* refers to the trigger condition in which only an incoming K^- and outgoing π^- were required (i.e., no information from the out-of-beam detectors was considered). Also shown are the excitation energy spectra for events with a coincident proton, pion, or neutron [Figs. 3(b), 3(c), and 3(d), respectively], where the coincident decay particles were identified as described in the previous section. (Note that the low number of counts in the unbiased spectrum as compared to the coincident spectra is a result of a hardware prescale in the unbiased trigger.) Because the coincident protons were required to have kinetic energies greater than 35 MeV, only events in which the Λ was bound to a nucleus and decayed through a nonmesonic channel contribute to the proton-tagged spectrum of Fig. 3(b).

A fit of a Gaussian shape to this peak determined an experimental resolution of 2.3-MeV root-mean-square and defined the zero of excitation energy. The unbiased spectrum, Fig. 3(a), was then determined to consist of three parts: (i) a broad background from $K^- \rightarrow \mu^- \bar{\nu}_\mu \pi^0$ contamination whose shape was determined by Monte Carlo simulation and whose magnitude was found from a fit to the lower part of the spectrum, (ii) events corresponding to the production of the bound ${}^4_\Lambda\text{He}$ ground state with mean and width determined from Fig. 3(b) and amplitude determined by a fit to the left half of the peak, and (iii) additional structure above zero excitation energy that we associate with the production of unbound Λ hyperons.

The excitation energy spectrum for coincident decay pions of Fig. 3(c) is shown with a two parameter fit to the left half of the ground-state peak. The fit determines the amplitude of the ground-state peak and includes a flat background term. The requirement of a decay-pion tag virtually eliminates the residual $K^- \rightarrow \mu^- \bar{\nu}_\mu \pi^0$ contamination but retains events from both bound and unbound Λ production as would be expected. For the neutron-tagged excitation energy spectrum of Fig. 3(d), the coincident neutrons consist of a mixture of neutrons from nonmesonic ${}^4_\Lambda\text{He}$ decay, accidental neutrons, and neutrons created when π^- 's from ${}^4_\Lambda\text{He}$ decay or quasifree Λ decay were absorbed in the material around the target area.

V. LIFETIME OF ${}^4_\Lambda\text{He}$

The lifetime of a ${}^4_\Lambda\text{He}$ hypernucleus was defined event-by-event as the time between the formation of the hypernucleus and its subsequent decay as determined from the velocity and trajectory of the incident kaon and the energy and trajectory of the charged hypernuclear decay product. The ${}^4_\Lambda\text{He}$ hypernucleus was assumed to be created at rest, and the position of the reaction vertex (and decay point) was inferred from the in-beam and out-of-beam tracks using a distance of closest approach (DCA) algorithm. The velocity of the incoming K^- was derived from its momentum as measured by the upstream K^- spectrometer, and its trajectory was determined by the six drift chambers located just upstream of the ${}^4\text{He}$ target. A time measurement was also made for the beam kaon as it passed the in-beam scintillators IT1 and IT2. This measurement set the start time for the event and was taken as the average of the two counters (producing a combined resolution of $\sigma_{\text{IT}} = 44.3$ ps). The time of hypernuclear formation was then simply found from the velocity of the K^- and its path length from the point midway between the IT1 and IT2 timing counters to the reaction point. (The velocity of the beam kaon was assumed to have remained constant as it traversed the experimental area.) For the out-of-beam charged track, the energy was determined from the TOF as measured between the decay-timing (OT1) and TOF (OT2) scintillator layers. The decay time at the vertex position was determined after making corrections for the unmeasured energy losses in the target region.

The events used in the ${}^4_\Lambda\text{He}$ mean lifetime determination were subject to the in-beam tracking cuts described in Sec. IV for the hypernuclear excitation energy spectra. Protons were chosen in the out-of-beam to avoid the contamination of quasifree Λ decay events present in the π^- -tagged sample.




FIG. 4. (a) The ${}^4_\Lambda\text{He}$ lifetime distribution for events with a coincident proton overlaid with the fit of the expected distribution. (b) The time-difference distribution for protons from the reaction ${}^4\text{He}(\pi^-, p)X$ used to determine the time resolution function.

Also, protons with kinetic energy below 70 MeV were not considered for the lifetime measurement because the uncertainty in the energy loss correction increases at lower energies. To ensure that the kaon and proton tracks were correlated with a hypernuclear formation event: (i) the excitation energy was required to be within $\pm 2\sigma$ of the mean, (ii) the K^- and proton tracks must have had a DCA of less than 2 cm, and (iii) the reaction vertex must have been within the liquid ${}^4\text{He}$ containment vessel. The resulting lifetime distribution for these proton-tagged events can be seen in Fig. 4(a).

The mean lifetime of ${}^4_\Lambda\text{He}$ was determined by a fit to the distribution of Fig. 4(a). The actual time distribution expected for this measurement has the form:

$$L_\tau(t) = \int_{-\infty}^{\infty} dt' R(t') P_\tau(t - t'), \quad (10)$$

where $P_\tau(t)$ is the decay probability distribution:

$$P_\tau(t) = \frac{1}{\tau} e^{-t/\tau} \theta(t), \quad (11)$$

and $R(t)$ is the resolution function as determined from the time distribution of prompt protons originating from the reaction $\pi^- + {}^4\text{He} \rightarrow p + X$ [shown in Fig. 4(b)] and $\theta(t)$ is the usual theta function. The measured distribution of Fig. 4(b) gives a resolution of $\sigma_\tau \simeq 150$ ps. The fit to the hypernuclear lifetime distribution of Fig. 4(a) was performed using a likelihood function derived assuming Poisson statistics from the function:

$$f(t) = C_1^2 + C_2 t + C_3^2 L_\tau(t), \quad (12)$$

which includes two background terms in addition to the expected lifetime distribution of Eq. (10). By




FIG. 5. Shown here are the kinetic energy spectra for (a) protons and (b) pions detected in coincidence with ${}^4_{\Lambda}\text{He}$ formation. These spectra have been corrected for PID inefficiencies, detector acceptance, and unmeasured energy losses in the target region.

maximizing the likelihood function on the four parameter space (C_1, C_2, C_3, τ), the value for the mean lifetime of ${}^4_{\Lambda}\text{He}$ was found to be $\tau = 245 \pm 24$ ps. [The values for the background terms C_1 and C_2 were found to be negligible compared to $C_3^2 L_{\tau}(t)$.]

VI. PARTICLE EMISSION SPECTRA

The kinetic energy spectra for the decay protons, pions, and neutrons are shown in Figs. 5(a), 5(b), and 6(b), respectively. These events were subject to the cuts described in Sec. IV for the ${}^4_{\Lambda}\text{He}$ excitation energy spectra. To reduce contamination from out-of-beam particles not originating from a hypernuclear ground-state event, a cut was also placed on the value of the excitation energy. In the case of proton- and neutron-tagged events, the excitation energy, EE , was required to be within $\pm 2\sigma$ of the mean of the ground-state peak, whereas in the case of pion-tagged events, a tighter cut of $\mu - 2\sigma < EE < \mu$ (referred to here as the “ -2σ cut”) was employed to reduce contamination from quasifree Λ decay events. (The quantities μ and σ are the previously determined mean and width of the ${}^4_{\Lambda}\text{He}$ ground-state peak.) For the proton and pion spectra, corrections for the unmeasured energy losses in the target region have been included.

To obtain the neutron kinetic energy spectrum of Fig. 6(b), a considerable background consisting of accidental neutrons and neutrons from π^- interactions had to be subtracted from




FIG. 6. (a) The raw neutron kinetic energy spectrum with background contributions from accidental neutrons (dashed line) and neutrons from π^- absorption (dotted line) overlaid. (b) The neutron spectrum after background subtraction and corrections for detector acceptance.

the raw neutron kinetic energy spectrum shown in Fig. 6(a). The expected shape of the accidental neutron spectrum was generated assuming a flat distribution in β^{-1} (i.e., a flat time distribution). The distribution for neutrons from π^- absorption was then extracted from the data by considering neutron-tagged events with excitation energy in the range $10 < EE < 30$ MeV. The neutrons from these events consist of a mixture of accidental neutrons and neutrons from π^- absorption only. The normalizations for the background shapes were found via a detailed comparison of the unbiased and neutron-tagged excitation spectra of Figs. 3(a) and 3(d). The background contributions thus determined are overlaid on the plot of Fig. 6(a).

The kinetic energy spectra of Figs. 5(a), 5(b), and 6(b) also include corrections for small PID inefficiencies and detector acceptance which were derived with the aid of a Monte Carlo simulation of the E788 out-of-beam detector system. Protons, neutrons, and π^- 's with realistic initial energy distributions were propagated through the target area and out-of-beam detector system. The response of the scintillator counters was modeled using an approximate form of the Bethe-Bloch equation of energy loss in the case of charged particles and the DEMONS software package [11] for neutron interactions. Unmeasured energy losses in the target materials and detector wrappings, pion decays and interactions, and corrections for the nonlinear response of the scintillator material at large energy deposition were also included. The tracking and particle

identification was performed with the same code as used for the real data, and the results were used to derive the necessary corrections.

VII. EXTRACTION OF PARTIAL RATES

The decay modes considered in this analysis were the π° , π^- , proton-stimulated, neutron-stimulated, and three-body nucleon-stimulated modes with branching fractions defined as:

$$B_a = \frac{\Gamma_a}{\Gamma_{\text{tot}}} = \frac{N_a}{N_{\text{HN}}^{\text{tot}}}, \quad (13)$$

where $a = \pi^\circ, \pi^-, p, n$, and mb ; N_a is the total number of hypernuclear events decaying to mode a ; and $N_{\text{HN}}^{\text{tot}} = 27800 \pm 1500$ is the total number of ${}^4_\Lambda\text{He}$ hypernuclear events found from the fit to the ground-state peak of the unbiased excitation spectrum of Fig. 3(a). This total also includes corrections for a hardware prescale in the unbiased trigger. The π° branching fraction was extracted by considering γ -tagged hypernuclear formation events, whereas the π^- branching fraction was determined from the observed π^- kinetic energy distribution. The nonmesonic decay branching fractions were extracted by two essentially independent methods utilizing (i) the single-particle kinetic energy spectra with final-state interactions and three-body ΛNN decay contributions considered within a simple model and (ii) multiple nucleon coincidence events. The final values for the ${}^4_\Lambda\text{He}$ branching fractions were then determined by a χ^2 minimization that considers the mesonic and nonmesonic rates simultaneously.

A. The π° decay mode

The π° decay rate was found by considering events for which one or more γ 's were detected in coincidence with ${}^4_\Lambda\text{He}$ hypernuclear formation. The γ 's were identified as described in Sec. III, and the candidate events were subject to the same cuts as the excitation spectra of Sec. IV. Three possible sources of γ rays were considered for the π° rate extraction: π° decay, π^- interactions, and accidental hits. Thus, the possible backgrounds include γ -tagged events arising from accidentals, π^- 's and π° 's from quasifree Λ decay, π^- 's from ${}^4_\Lambda\text{He}$ decay, and π° 's from the $K^- \rightarrow \mu^- \bar{\nu}_\mu \pi^\circ$ decay background. To find the π° branching fraction, these background events were subtracted as described below.

To determine the quantitative contributions of each type of background, the various sources of γ 's were isolated by considering γ -tagged events with particular excitation energies. First, events with excitation energy below -10 MeV, which consist almost exclusively of accidental γ 's and γ 's from the K^- decay background, were used to estimate the detection efficiencies for accidentals and γ 's from π° decay. Next, the γ 's from π^- interactions were studied using events with excitation energy in the range $10 < EE < 30$ MeV. These events consist of accidental γ 's and γ 's from the K^- decay background (as in the previous case) with additional background γ events originating from quasifree Λ decay. Using the accidental and π° decay γ detection probabilities determined above, the contribution from quasifree $\Lambda \rightarrow p\pi^-$ decay could be extracted, allowing the determination of the

detection probability for γ 's from π^- interactions. Finally, the ${}^4_\Lambda\text{He}$ π° decay events were selected by requiring the excitation energy to be within $\pm 2\sigma$ of the ground-state peak and subtracting the background contributions found above.

After properly accounting for the backgrounds and γ detection efficiency, the π° branching fraction was found to be $B_{\pi^\circ} = 0.552 \pm 0.076(\text{stat.}) \pm 0.061(\text{syst.})$. So, for a Λ embedded within the ${}^4_\Lambda\text{He}$ hypernucleus, the π° decay rate:

$$\frac{\Gamma_{\pi^\circ}}{\Gamma_\Lambda} = B_{\pi^\circ} \times \frac{\Gamma_{\text{tot}}}{\Gamma_\Lambda} = 0.59 \pm 0.10, \quad (14)$$

expressed here in units of the total decay rate of the free Λ , is enhanced by a factor of about 1.6 as compared to the free Λ decay partial width of $\Gamma_{\pi^\circ}^{\text{free}}/\Gamma_\Lambda = 0.358 \pm 0.005$ [3]. (The quantity $\Gamma_{\text{tot}}/\Gamma_\Lambda = 1.07 \pm 0.11$ was derived from the ${}^4_\Lambda\text{He}$ hypernuclear lifetime found in Sec. V).

B. The π^- decay mode

The π^- decay rate was determined by a fit to the observed π^- kinetic energy spectrum using a theoretical distribution provided by the model calculation of Kumagai-Fuse *et al.* [12]. Their predicted π^- spectrum [shown in Fig. 7(a)] was derived using the resonating group method and includes final-state interactions and pion distortion effects. The effects of detector acceptance and resolution unique to E788 were applied to the model by using the spectrum of Fig. 7(a) as input to the Monte Carlo simulation described in Sec. VI and reconstructing the initial kinetic energy distribution with the analysis code.

A fit of the model prediction to the PID-corrected π^- spectrum of Fig. 7(b) was then performed using the fit function:

$$N_\pi(E) = M_\pi^{2\sigma} h_\pi(E), \quad (15)$$

where $h_\pi(E)$ is a parametrization of the expected π^- kinetic energy distribution as reconstructed by the analysis code (normalized to the detector acceptance and in units of counts/bin per event) and $M_\pi^{2\sigma}$ is the fit parameter representing the acceptance-corrected number of pions within the -2σ excitation energy cut. The result of the fit is shown in Fig. 7(b), where the value of $M_\pi^{2\sigma}$ was found to be 3690 ± 190 . Correcting for the excitation energy cut yields a value for the total number of π^- 's:

$$M_\pi^{\text{total}} = g_\pi M_\pi^{2\sigma} = 7490 \pm 480, \quad (16)$$

where $g_\pi = 2.030 \pm 0.076$ is the ratio of the number of counts in the ground-state peak of the π^- -tagged excitation spectrum of Fig. 3(c) to the number of events within the -2σ cut.

The π^- -decay branching fraction for the ${}^4_\Lambda\text{He}$ hypernucleus was then found by Eq. (13) as $B_{\pi^-} = 0.269 \pm 0.022(\text{stat.}) \pm 0.014(\text{syst.})$. In the present case of ${}^4_\Lambda\text{He}$ decay, the observed π^- decay rate:

$$\frac{\Gamma_{\pi^-}}{\Gamma_\Lambda} = B_{\pi^-} \times \frac{\Gamma_{\text{tot}}}{\Gamma_\Lambda} = 0.289 \pm 0.037, \quad (17)$$

in units of the free Λ decay, is seen to be suppressed by a factor of about 2.2 as compared to the free Λ decay value of $\Gamma_{\pi^-}^{\text{free}}/\Gamma_\Lambda = 0.639 \pm 0.005$ [3]. This suppression is mainly due to the Pauli blocking of the final-state proton.




FIG. 7. (a) The theoretical π^- kinetic energy spectrum of Kumagai-Fuse *et al.* [12] and (b) a fit of this expected shape to the measured distribution after including the effects of detector acceptance and resolution. The measured spectrum of (b) was corrected for PID inefficiencies and unmeasured energy losses (but not detector acceptance).

C. Nonmesonic rates from single-particle KE spectra

The rates for proton- and neutron-stimulated decay were extracted simultaneously from the observed proton and neutron kinetic energy spectra using a simple model that included the effects of final-state interactions (FSI). In the model, two types of decay events were considered: ΛN decay with no FSI (referred to here as *no-scatter*) and decay events where one of the decay nucleons scatters from one of the spectator nucleons (*rescatter*). The expected no-scatter and rescatter nucleon kinetic energy distributions (shown in Fig. 8) were derived from a Monte Carlo simulation of the ΛN interaction. (Also shown is the distribution for nucleons from ΛNN decay, which will be discussed later in this section.) For the Monte Carlo events, the momenta of the four initial-state baryons were generated from a Gaussian distribution with a root-mean-square width of 125 MeV/ c for the nucleons and 50 MeV/ c for the Λ . The distributions were correlated such that the sum of the four momenta added to zero, and the energy of each spectator nucleon was determined by setting its mass to be on-shell. For no-scatter events, the energy distribution for the primary decay nucleons was determined by choosing a direction for the participating nucleon isotropically in the Λ -nucleon center-of-mass frame and boosting to the laboratory system. The rescatter events were simulated by choosing a direction for one of the nucleons isotropically in the decay-spectator nucleon system (*s-wave*) and boosting to the laboratory frame.




FIG. 8. Shown here are the predicted no-scatter and rescatter kinetic energy distributions for nucleons from ΛN decay (solid and dashed lines, respectively) along with the distribution for the primary nucleons from ΛNN decay (dotted line).

As in the π^- case above, the expected kinetic energy distributions were fit to the observed spectra, and the results were combined with the model to extract the nucleon-stimulated decay rates. To include the effects of the resolution and acceptance of the detector system, the expected no-scatter and rescatter distributions of Fig. 8 were used as input to the Monte Carlo simulation discussed in Sec. VI and then reconstructed using the analysis code. The resulting spectra were designated $h^0(E)$ for the nucleons that did not undergo FSI and $h^1(E)$ for the rescattered nucleons. (These distributions differ for protons and neutrons due to differing energy resolutions and detection acceptance.) The distributions are in units of counts per bin per event and are normalized to the detector acceptance. The fits were then performed using functions of the form:

$$N_a(E) = \underbrace{M_a^0 h_a^0(E)}_{\text{no-scatter}} + \underbrace{M_a^1 h_a^1(E)}_{\text{rescatter}}, \quad (18)$$

where $a = p, n$ and the fit parameters M_a^0 and M_a^1 represent the acceptance-corrected numbers of no-scatter and rescatter nucleons, respectively. The fit parameters M_a^0 and M_a^1 were related via our model as described below.

The effects of final-state interactions were incorporated by defining a rescatter probability, ϵ , as the probability for one of the primary-decay nucleons from ΛN decay to scatter off of one of the spectator nucleons. The value of ϵ was determined from the data as described below. The scattering processes were assumed to occur with equal probability (i.e., ϵ has the same value for all proton-neutron rescatter combinations). Because the effects of the FSI correction were seen to be small, this assumption was considered adequate. Assigning a probability to each possible final state in terms of the rescatter probability per spectator nucleon, ϵ , and the

nucleon-stimulated branching fractions, B_p and B_n , and then summing over no-scatter and rescatter nucleons yields:

$$M_p^0 = (1 - 2\epsilon)B_p N_{\text{HN}}^{\text{tot}}, \quad (19)$$

$$M_n^0 = (1 - 2\epsilon)B_p N_{\text{HN}}^{\text{tot}} + 2(1 - 2\epsilon)B_n N_{\text{HN}}^{\text{tot}}, \quad (20)$$

$$M_p^1 = 4\epsilon(B_p + B_n)N_{\text{HN}}^{\text{tot}}, \quad (21)$$

$$M_n^1 = 4\epsilon(B_p + B_n)N_{\text{HN}}^{\text{tot}}. \quad (22)$$

Using these expressions, the parameters M_p^1 and M_n^1 were eliminated from the fit functions:

$$N_p(E) = M_p^0 h_p^0(E) + \frac{2\epsilon}{1 - 2\epsilon} (M_p^0 + M_n^0) h_p^1(E), \quad (23)$$

$$N_n(E) = M_n^0 h_n^0(E) + \frac{2\epsilon}{1 - 2\epsilon} (M_p^0 + M_n^0) h_n^1(E). \quad (24)$$

The three parameters M_p^0 , M_n^0 , and ϵ were then determined by a fit to the data (shown in Fig. 9). Because the fit shapes $h^0(E)$ and $h^1(E)$ include the effects of the detector efficiency and resolution, the observed nucleon kinetic energy spectra used for the fits were not efficiency corrected. The fits were carried out simultaneously by varying the quantity ϵ to obtain a minimum combined χ^2 . Using the fit values, the nucleon-stimulated branching fractions were then calculated using Eqs. (19) and (20), and the results are listed in Table IV. The upper limits for the neutron-stimulated branching fraction and the ratio Γ_n/Γ_p were determined by the prescription of Feldman and Cousins [13], which defines the confidence levels




FIG. 9. Shown here are the fits of the expected no-scatter and rescatter distributions to the observed (a) proton and (b) neutron kinetic energy spectra. The no-scatter contribution is indicated by a dashed line, whereas the rescatter contribution is shown as a dotted line.

(CL) in a consistent way while imposing the physical boundary at zero. The upper limits are given at the 95% CL. Also listed are the results for the case of no FSI found by fixing the rescatter probability ϵ to zero.

The model was also extended to include possible contributions from three-body ΛNN decays. The expected kinetic energy distribution for these nucleons was generated using the same Monte Carlo code as for the ΛN interaction described above, and the resulting distribution is shown in Fig. 8. In this case, the momentum of the single on-shell spectator was initially selected, fixing the momentum and energy of the participating ΛNN system. Final-state momenta for these three baryons were then selected in the system's center-of-mass frame according to a phase-space distribution generated by the CERNLIB GENBOD code. The nucleon momenta were then boosted to the laboratory frame.

As in the case of FSI above, the contribution of ΛNN decay, $\Gamma_{mb}/\Gamma_{\text{tot}}$, was determined by a fit to the observed kinetic energy spectra of Fig. 9. Because the ΛNN decay nucleons and rescattered ΛN nucleons are indistinguishable at the level of statistics of the current measurement, the fits were performed with an $h^1(E)$ that consisted of a linear combination of the ΛNN and rescatter distributions. The results for the nucleon-stimulated branching fractions are shown in Table IV for the cases of (i) ΛNN decay with no FSI for the ΛN decays (i.e., $\epsilon = 0$) and (ii) a mixture of ΛNN decay and FSI with equal strengths (i.e., the quantity M_p^1 consisted of a 50/50 mixture of ΛNN decay events and rescattered ΛN decay events). From the case of FSI, the upper limit for the contribution of final-state interactions to the total ${}^4\text{He}$ decay rate was found as $\Gamma_{nm}^{\text{FSI}}/\Gamma_{\text{tot}} = 4\epsilon(\Gamma_{nm}/\Gamma_{\text{tot}}) \leq 0.11$ (95% CL). Also, the multibaryon decay branching fraction found for the case of ΛNN decay with no FSI gives an upper limit of $\Gamma_{mb}/\Gamma_{\text{tot}} \leq 0.097$ (95% CL) for the contribution of the ΛNN decay process.

D. Nonmesonic rates from multiple nucleon coincidence data

The ΛN interaction produces two energetic nucleons that may be detected in the out-of-beam detector arrays resulting in both proton-neutron (pn) and neutron-neutron (nn) coincidence events. The observation of such events gives another window onto the nucleon-stimulated decay process. In the present experiment, a total of 87 pn coincidence events and 19 nn coincidence events were observed. These events were subject to the same in-beam tracking cuts as the excitation spectra of Sec. IV along with a $\pm 2\sigma$ excitation energy cut. The protons and neutrons were identified in the out-of-beam detector arrays as described in Sec. III.

The cosine of the separation angle for the two final-state nucleons, $\cos\theta_{NN}$, is shown in Figs. 10(a) and 10(b) for pn and nn coincidence events, respectively. If the spectator nucleons are ignored, the two nucleons resulting from the ΛN interaction would be expected to emerge roughly back-to-back with a separation angle near 180° (from simple momentum conservation). The observed distribution for pn events is strongly peaked near the value of 180° with only about 10% of the pn events having a separation angle less than 140° .

TABLE IV. Results of the nonmesonic decay rate extraction considering FSI and ΛNN decay contributions. The decay rates are given in units of Γ_{tot} , and quoted errors are statistical only. The upper limits given for $\Gamma_n/\Gamma_{\text{tot}}$ and Γ_n/Γ_p (listed directly beneath the associated quantities) are at the 95% confidence level and include systematic errors.

	No FSI or ΛNN	FSI	ΛNN	FSI + ΛNN	Systematic error
$\chi^2/\text{d.o.f.}$	0.78	0.72	0.69	0.71	
ϵ		0.045 ± 0.035		$0.030^{+0.024}_{-0.021}$	
$\Gamma_{mb}/\Gamma_{\text{tot}}$			0.042 ± 0.028	$0.017^{+0.015}_{-0.012}$	
$\Gamma_p/\Gamma_{\text{tot}}$	0.158 ± 0.011	0.160 ± 0.015	0.134 ± 0.026	0.150 ± 0.016	± 0.0089
$\Gamma_n/\Gamma_{\text{tot}}$	-0.0066 ± 0.0060	-0.0089 ± 0.0075	$-0.0133^{+0.0089}_{-0.0071}$	$-0.0112^{+0.0084}_{-0.0077}$	± 0.0104
	≤ 0.018	≤ 0.017	≤ 0.015	≤ 0.017	
$\Gamma_{nm}/\Gamma_{\text{tot}}$	0.1510 ± 0.0099	0.151 ± 0.013	0.162 ± 0.020	0.156 ± 0.017	± 0.0096
Γ_n/Γ_p	-0.042 ± 0.037	$-0.056^{+0.046}_{-0.043}$	-0.100 ± 0.068	-0.075 ± 0.055	± 0.063
	≤ 0.10	≤ 0.10	≤ 0.097	≤ 0.098	

The latter events may be indicative of FSI or ΛNN decay contributions. The number of counts is low in the nn spectrum, but the back-to-back peak does not seem to be as pronounced for these events. The distributions for the sum of the kinetic energies of the coincident nucleons are shown in Fig. 11. For the ΛN interaction, the total energy available to the two decay nucleons is ~ 166 MeV (i.e., the difference in the Λ and n masses less the binding energy of the ${}^4_{\Lambda}\text{He}$ hypernucleus). The pn kinetic energy distribution of Fig. 11(a) peaks near 166 MeV as expected. The observed spread is largely consis-

tent with a spread due to the Fermi momentum of the baryons within the nucleus coupled with detector resolution, and the apparent tail out to lower energies may be the result of FSI or ΛNN decays. (The observed spectrum is in qualitative agreement with the results of a Monte Carlo simulation which includes such effects.) The nn kinetic energy distribution of Fig. 11(b), however, does not behave as expected and may be indicative of nn coincidence events originating from sources other than Λn decays (i.e., some mixture of background neutrons and neutrons from Λp or ΛNN decays).

The nucleon-stimulated branching fractions were extracted from the observed pn and nn coincidence events by relating




FIG. 10. The cosine of the nucleon-nucleon separation angle is shown for (a) pn and (b) nn coincidence events. Overlaid on these plots is the relative geometric acceptance for NN coincidence events normalized to the number of counts in each spectrum.




FIG. 11. The sum of the kinetic energy for the two observed nucleons is shown for (a) pn and (b) nn events. The line is drawn at the expected value of 166 MeV.

the number of each type of coincidence event to the branching fractions as:

$$N_{pn} = (P_{\text{acc}}^{(1)} P_p^{\Lambda p} + P_{pn}^{\Lambda p}) B_p g_2 N_{\text{HN}}^{\text{tot}}, \quad (25)$$

$$N_{nn} = (P_{\text{acc}}^{(1)} P_n^{\Lambda n} + P_{nn}^{\Lambda n}) B_n g_2 N_{\text{HN}}^{\text{tot}} + N_{nn}^{\text{bg}}, \quad (26)$$

where g_2 is the fraction of a Gaussian distribution contained within $\pm 2\sigma$ of the mean, $P_{\text{acc}}^{(1)}$ is the probability of detecting exactly one accidental neutron per event, $P_p^{\Lambda p}$ is the probability of detecting one proton from Λp decay, $P_{pn}^{\Lambda p}$ is the probability of detecting one proton and one neutron from Λp decay, and $P_n^{\Lambda n}$ and $P_{nn}^{\Lambda n}$ are the probabilities of detecting one and two neutrons from Λn decay, respectively. With the exception of $P_{\text{acc}}^{(1)}$, which was extracted from the data, the above detection probabilities had to be estimated using the Monte Carlo simulation described in the previous section. Final-state interactions were included in the Monte Carlo at the level determined for the case of FSI only.

The term N_{nn}^{bg} appearing in Eq. (26) is the expected number of nn background events due to accidental neutrons, neutrons from π^- absorption, and neutrons from proton-stimulated decays. As with the accidental detection probability above, the detection probabilities for neutrons from π^- interactions were determined from the data with no presumptions about the extent of FSI or ΛNN decays, whereas the detection probabilities for neutrons from Λp decay were again estimated using the Monte Carlo with FSI included. The number of expected background nn coincidence events was found to be $N_{nn}^{\text{bg}} = 18.3 \pm 6.1$ with the largest contribution arising from neutrons from π^- absorption (~ 13 events). Such a large background is consistent with the observed nn energy-sum spectrum of Fig. 11(b).

The resulting values for the branching fractions are listed in Table V along with the results from the analysis of the single-particle spectra of the previous section for the case of FSI only. Also shown is the weighted average of the two determinations.

In addition to the pn and nn events, a handful of pp coincidence events were also observed and were used to set an upper limit on the Λpp branching fraction. After the in-beam tracking cuts and $\pm 2\sigma$ excitation energy cut, a total of 8 pp coincidence events remained. Although these events may be

indicative of a Λpp three-body decay mode, they could also be Λp decay events for which the spectator proton has been detected. As in the pn and nn cases, the number of pp events of each type may be expressed in terms of the ${}^4_{\Lambda}\text{He}$ branching fractions as:

$$N_{pp}^{\Lambda p} = g_2 \varepsilon_q P_{pp}^{\Lambda p} B_p N_{\text{HN}}^{\text{tot}}, \quad (27)$$

$$N_{pp}^{\Lambda pp} = g_2 \varepsilon_q P_{pp}^{\Lambda pp} B_{pp} N_{\text{HN}}^{\text{tot}}, \quad (28)$$

where $\varepsilon_q = 0.986$ is the measured efficiency of the charged trigger, $P_{pp}^{\Lambda p}$ and $P_{pp}^{\Lambda pp}$ are the probabilities of detecting two protons from Λp and Λpp decay, respectively, and B_{pp} is the Λpp branching fraction. The probability of detecting two protons from Λp decay is negligibly small when interactions with the spectator nucleons are ignored, but when FSI is included at the level determined in Sec. VII C, it becomes significant. The probabilities $P_{pp}^{\Lambda p}$ and $P_{pp}^{\Lambda pp}$ were estimated using the Monte Carlo simulation discussed in Sec. VII C, including FSI for the Λp case. The expected number of background pp events, $N_{pp}^{\Lambda p}$, was then calculated from Eq. (27) as 6.40 ± 0.69 . Assuming Poisson statistics and using the prescription of Feldman and Cousins [13], the upper limit for the number of observed pp events originating from Λpp decay was found as $N_{pp}^{\Lambda pp} \leq 9.29$ (95% CL for a total number of eight observed counts and an expected background of six counts). Using Eq. (28), the upper limit for the Λpp branching fraction was then found as $B_{pp} \leq 0.026$.

As mentioned above, the contribution from Λp decay events becomes negligible when final-state interactions are ignored. In this case, all eight observed pp events are considered to have originated from Λpp decay, and the Λpp branching fraction is found to lie in the range $0.008 < B_{pp} < 0.041$ at the 95% CL.

E. Combined ${}^4_{\Lambda}\text{He}$ decay rates

The final values for the ${}^4_{\Lambda}\text{He}$ branching fractions were determined by considering a set of parameters (B_{π^-} , B_{π^0} , B_p , B_n , N) representing the π^- , π^0 , and proton- and neutron-stimulated branching fractions along with the total number of ${}^4_{\Lambda}\text{He}$ hypernuclei. The optimal values for this set of parameters were found by minimizing a χ^2 function of the

TABLE V. Nonmesonic decay rates as determined from the single-particle kinetic energy spectra and multiple nucleon coincidence data. The errors given consist of statistical plus systematic errors, and upper limits are given at the 95% CL. Also shown is the weighted average of the two determinations.

	Single-particle	Multiple coincidence	Average
$\Gamma_p / \Gamma_{\text{tot}}$	0.160 ± 0.018	0.165 ± 0.028	0.161 ± 0.015
$\Gamma_n / \Gamma_{\text{tot}}$	-0.0089 ± 0.0135 ≤ 0.018	0.0031 ± 0.0338 ≤ 0.070	-0.0073 ± 0.0125 ≤ 0.018
$\Gamma_{nm} / \Gamma_{\text{tot}}$	0.151 ± 0.017	0.168 ± 0.042	0.153 ± 0.016
Γ_n / Γ_p	-0.056 ± 0.082 ≤ 0.11	0.019 ± 0.205 ≤ 0.42	-0.046 ± 0.076 ≤ 0.11

TABLE VI. The ${}^4_\Lambda\text{He}$ decay rates as determined by the present analysis. The errors quoted include contributions from systematic error, and upper limits are given at the 95% CL.

Mesonic rates		Nonmesonic rates	
$\Gamma_{\pi^0}/\Gamma_{\text{tot}}$	0.564 ± 0.036	$\Gamma_p/\Gamma_{\text{tot}}$	0.169 ± 0.019
$\Gamma_{\pi^-}/\Gamma_{\text{tot}}$	0.270 ± 0.024	$\Gamma_n/\Gamma_{\text{tot}}$	-0.0032 ± 0.0183 ≤ 0.032
$\frac{\Gamma_{\pi^0} + \Gamma_{\pi^-}}{\Gamma_{\text{tot}}}$	0.835 ± 0.021	$\Gamma_{nm}/\Gamma_{\text{tot}}$	0.165 ± 0.021
$\Gamma_{\pi^0}/\Gamma_{\pi^-}$	2.09 ± 0.29	Γ_n/Γ_p	-0.019 ± 0.108 ≤ 0.19

form:

$$\chi^2 = \frac{(M_{\pi^-} - B_{\pi^-}N)^2}{\sigma_{\pi^-}^2} + \frac{(M_{\pi^0} - B_{\pi^0}N)^2}{\sigma_{\pi^0}^2} + \frac{(M_p - B_pN)^2}{\sigma_p^2} + \frac{[M_n - (B_p + 2B_n)N]^2}{\sigma_n^2} + \frac{(M_{pn} - B_pN)^2}{\sigma_{pn}^2} + \frac{(M_{nn} - B_nN)^2}{\sigma_{nn}^2} + \frac{(N_{\text{HN}}^{\text{tot}} - N)^2}{\sigma_N^2}, \quad (29)$$

where the values M_a are the acceptance-corrected numbers of event type a with associated error σ_a . Additionally, the parameters representing the branching fractions were subject to the constraint:

$$B_p + B_n + B_{\pi^-} + B_{\pi^0} = 1, \quad (30)$$

and any possible contributions from three-body ΛNN decay, which were previously seen to be small, were neglected. The minimization was carried out using the ROOT MINUIT minimization routine, and the results are listed in Table VI (in units of Γ_{tot}).

VIII. DISCUSSION

The ${}^4_\Lambda\text{He}$ decay rates determined in the present work are listed in Table VII (in units of Γ_Λ). Also listed are the results of another measurement of ${}^4_\Lambda\text{He}$ decay performed by Outa *et al.* [14] along with the results of some earlier bubble chamber experiments [8, 15, 16]. The values taken from Ref. [8] are the result of a reanalysis of various older results performed by Schumacher. The decay rates found in this work are seen to be in good agreement with past measurements.

The results reported for the experiment of Outa *et al.* were published in 1998, some 8 years after the running of our experiment. Their experiment created ${}^4_\Lambda\text{He}$ by the ${}^4\text{He}(K_{\text{stopped}}^-, \pi^-)$ reaction and detected the π^0 , π^- , and protons from the subsequent weak decay. Unlike our experiment, their apparatus was not able to detect neutrons, requiring them to determine the neutron-stimulated rate by subtraction. Our results for the neutron-stimulated rate and neutron to proton ratio, $\Gamma_n/\Gamma_\Lambda \leq 0.035$ and $\Gamma_n/\Gamma_p \leq 0.19$, provide stricter upper limits as compared to $\Gamma_n/\Gamma_\Lambda \leq 0.09$ and $\Gamma_n/\Gamma_p \leq 0.60$ for Outa *et al.* (all at the 95% CL). The upper limits for Outa *et al.* given above were derived from the results listed in Table VII using the same method as for our results. There

TABLE VII. Measured properties of the ${}^4_\Lambda\text{He}$ hypernucleus. Listed here are the results of the present analysis along with the results of another recent experiment on ${}^4_\Lambda\text{He}$ decay [14] and some older results from bubble chamber experiments [10, 15, 16]. The errors quoted for this analysis include contributions from systematic error.

	This work	Outa <i>et al.</i> [14]	Earlier results
$\Gamma_{\text{tot}}/\Gamma_\Lambda$	1.07 ± 0.11	$1.03^{+0.12}_{-0.10}$	1.15 ± 0.48 [8]
$\Gamma_{\pi^0}/\Gamma_\Lambda$	0.604 ± 0.073	0.53 ± 0.07	
$\Gamma_{\pi^-}/\Gamma_\Lambda$	0.289 ± 0.039	0.33 ± 0.05	
$\Gamma_{\pi^0}/\Gamma_{\pi^-}$	2.09 ± 0.29	1.59 ± 0.20	2.49 ± 0.34 [15] 2.20 ± 0.39 [16] 0.043 ± 0.017 [10]
$\Gamma_{\pi^+}/\Gamma_{\pi^-}$			
Γ_p/Γ_Λ	0.180 ± 0.028	0.16 ± 0.02	
Γ_n/Γ_Λ	≤ 0.035	$0.01^{+0.04}_{-0.01}$	
$\Gamma_{nm}/\Gamma_\Lambda$	0.177 ± 0.029	0.17 ± 0.05	
$\Gamma_{nm}/\Gamma_{\pi^-}$	0.61 ± 0.10	0.51 ± 0.16	0.56 ± 0.09 [8]
Γ_n/Γ_p	≤ 0.19	$0.06^{+0.28}_{-0.06}$	0.40 ± 0.15 [15] 0.29 ± 0.13 [16]

is also a slight difference in the mesonic decay rates, where our $\Gamma_{\pi^0}/\Gamma_{\pi^-}$ ratio is seen to be in better agreement with the earlier bubble chamber results.

Selected theoretical investigations of the ${}^4_\Lambda\text{He}$ nonmesonic partial decay rates are shown in Table VIII. These OPE (one pion exchange) models incorporate a quark-based model for short range interactions and are currently considered to be the most successful of the OPE-based models. The results of Jun [17] and Inoue *et al.* [5] are shown for OPE only and OPE + quark interaction mechanism. The work of Jun utilizes a phenomenological four-baryon point interaction model to describe the short-range quark interactions, whereas Inoue *et al.* uses a direct quark (DQ) interaction based on the effective weak Hamiltonian of Gilman and Wise [18]. The calculation of Sasaki *et al.* [6] employs the direct quark mechanism of Inoue *et al.* coupled with a $\pi + K + \sigma$ exchange potential. Unlike previous models, this model is able to reproduce the value of the proton asymmetry observed in polarized ${}^5_\Lambda\text{He}$ nonmesonic decay. All of the above models adequately reproduce the observed ${}^4_\Lambda\text{He}$ nonmesonic decay rates.

In regards to the $\Delta I = \frac{1}{2}$ rule, it is interesting to note that the direct quark models predict a significant contribution from $\Delta I = \frac{3}{2}$ transitions arising from the effective weak Hamiltonian used, whereas models based on the phenomeno-

TABLE VIII. Results of selected theoretical investigations of the ${}^4_\Lambda\text{He}$ partial decay rates. The partial rates listed here are given in units of the free Λ decay width Γ_Λ .

	Jun [17]		Inoue [5]		Sasaki [6]
	OPE	OPE+4BPI	OPE	OPE+DQ	DQ+
Γ_n/Γ_Λ	0.0008	0.0373	0.009	0.038	0.015
Γ_p/Γ_Λ	0.1478	0.1602	0.145	0.214	0.169
$\Gamma_{nm}/\Gamma_\Lambda$	0.1486	0.1975	0.154	0.253	0.184
Γ_n/Γ_p	0.005	0.23	0.061	0.178	0.091

logical four-baryon point interaction are able to reproduce the experimental results with little or no $\Delta I = \frac{3}{2}$ component. All of these models enforce the $\Delta I = \frac{1}{2}$ rule for the meson-exchange component.

A. FSI, three-body ΛNN decays, and interference effects

Using the simple model for nonmesonic decay introduced in Sec. VII C, the contributions of final-state interactions and three-body ΛNN decays were studied utilizing the single-particle proton and neutron kinetic energy spectra, and upper limits for each case were determined as:

$$\frac{\Gamma_{nm}^{\text{FSI}}}{\Gamma_{\text{tot}}} \leq 0.11, \quad \text{and} \quad \frac{\Gamma_{mb}}{\Gamma_{\text{tot}}} \leq 0.097, \quad (31)$$

for final-state interactions and ΛNN decays, respectively (95% CL). Within the model used here, the effects of FSI and ΛNN decays on pertinent quantities were seen to be small and, thus, have little effect on the final interpretation of the data.

In the analysis of the nonmesonic decay rates, the various decay modes have been assumed to add incoherently, neglecting possible quantum interference effects. It has been suggested [19,20] that such interference effects may be the origin of the so-called Γ_n/Γ_p puzzle, and Garbarino *et al.* [20] point out that any interference effects would be less pronounced for multiple coincidence measurements, providing a cleaner extraction of the nonmesonic decay rates. For the case of ${}^4_\Lambda\text{He}$ presented here, the results of the two determinations (listed in Table V) are consistent, suggesting that interference effects are not significant at this level of statistics. A detailed study of the angular correlations in the multiple nucleon coincidence data might shed more light on the contributions of interference effects as well as FSI and ΛNN decays, but the number of coincidence events in the present data set is too limited to provide anything conclusive.

B. Phenomenological analysis of s -shell hypernuclei

As mentioned in the introduction, measurements of the nonmesonic decay rates for the s -shell hypernucleus ${}^4_\Lambda\text{H}$ are currently quite limited. The most recent results from Outa *et al.* [14] give a value for the total nonmesonic rate of $\Gamma_{nm}/\Gamma_\Lambda({}^4_\Lambda\text{H}) = 0.17 \pm 0.11$, but no measurements of the proton- and neutron-stimulated partial rates have been made. Using the present results for ${}^4_\Lambda\text{He}$, these partial rates can

be calculated in the context of the previously described phenomenological model of Block and Dalitz.

From Eq. (7), the upper limit of $\Gamma_n/\Gamma_\Lambda({}^4_\Lambda\text{He}) \leq 0.035$ determined for the current measurement implies an upper limit of $\Gamma_p/\Gamma_\Lambda({}^4_\Lambda\text{H}) \leq 0.017$ for the ${}^4_\Lambda\text{H}$ proton-stimulated rate under the assumption of pure $\Delta I = \frac{1}{2}$ transitions. A measurement of the ${}^4_\Lambda\text{H}$ proton-stimulated rate greater than this value would indicate a violation of the $\Delta I = \frac{1}{2}$ rule for the ΛN weak interaction. The ${}^4_\Lambda\text{H}$ neutron-stimulated decay rate was found from Eq. (8) as $\Gamma_n/\Gamma_\Lambda({}^4_\Lambda\text{H}) = 0.081 \pm 0.023$. This calculation utilizes the result of Outa *et al.* for the ratio of the neutron- to proton-stimulated rates for ${}^5_\Lambda\text{He}$, $\Gamma_n/\Gamma_p({}^5_\Lambda\text{He}) = 0.45 \pm 0.11 \pm 0.03$ [9] but makes no assumptions regarding the $\Delta I = \frac{1}{2}$ rule. Using this result, an alternate determination of the ${}^4_\Lambda\text{H}$ proton-stimulated rate can be made from the result of Outa *et al.* for the total nonmesonic decay width of ${}^4_\Lambda\text{H}$:

$$\Gamma_p({}^4_\Lambda\text{H}) = \Gamma_{nm}({}^4_\Lambda\text{H}) - \Gamma_n({}^4_\Lambda\text{H}) = (0.09 \pm 0.11)\Gamma_\Lambda, \quad (32)$$

which, again, is independent of any assumptions about the isospin nature of the ΛN interaction. Because of the large error, this result is consistent with the upper limit found above. Thus, at the present level of statistics for the ${}^4_\Lambda\text{H}$ measurement, nothing conclusive can be said about the $\Delta I = \frac{1}{2}$ rule for the ΛN weak interaction.

The values of the ${}^4_\Lambda\text{H}$ nonmesonic decay rates derived from the above discussion represent the most current knowledge (although indirect) of the ${}^4_\Lambda\text{H}$ proton- and neutron-stimulated partial rates. The upper limit determined for the ${}^4_\Lambda\text{H}$ proton-stimulated decay rate suggests a direct measurement of this decay mode as an easy test of the $\Delta I = \frac{1}{2}$ rule for nucleon-stimulated decays (i.e., a measurement in excess of this upper limit would show a clear violation of the $\Delta I = \frac{1}{2}$ rule for the ΛN weak interaction). Measurements of the ${}^4_\Lambda\text{H}$ nonmesonic decay rates along with more precise determinations of the ${}^4_\Lambda\text{He}$ and ${}^5_\Lambda\text{He}$ nonmesonic decay rates would help to further determine the isospin structure of the ΛN weak interaction. Multiple nucleon coincidence studies can potentially provide the cleanest extraction of the nucleon-stimulated partial rates and would be a worthwhile avenue of future research.

ACKNOWLEDGMENTS

This work was partially supported by U.S. DOE grant numbers DE-FG02-87ER40315, DE-AC02-98CH10886, and DE-FG03-94ER40821.

- [1] L. B. Okun, *Leptons and Quarks* (North-Holland, Amsterdam, 1982).
- [2] N. Isgur, K. Maltman, J. Weinstein, and T. Barnes, *Phys. Rev. Lett.* **64**, 161 (1990).
- [3] W.-M. Yao *et al.*, *J. Phys. G* **33**, 1 (2006).
- [4] K. Maltman and M. Shmatikov, *Nucl. Phys.* **A585**, 343c (1995); T. Inoue, S. Takeuchi, and M. Oka, *ibid.* **A597**, 563 (1996).
- [5] T. Inoue, M. Oka, T. Motoba, and K. Itonaga, *Nucl. Phys.* **A633**, 312 (1998).

- [6] K. Sasaki, M. Izaki, and M. Oka, *Phys. Rev. C* **71**, 035502 (2005).
- [7] M. M. Block and R. H. Dalitz, *Phys. Rev. Lett.* **11**, 96 (1963); C. B. Dover, *Few-Body Syst. Suppl.* **2**, 77 (1987); J. Cohen, *Phys. Rev. C* **42**, 2724 (1990); R. A. Schumacher, in *Properties and Interactions of Hyperons*, Proceedings of the U.S.-Japan Seminar, Maui, Hawaii, edited by B. F. Gibson, P. D. Barnes, and K. Nakai (World Scientific, Singapore, 1994), p. 85.
- [8] R. A. Schumacher, *Nucl. Phys.* **A547**, 143c (1992).
- [9] H. Outa *et al.*, *Nucl. Phys.* **A754**, 157c (2005).

- [10] G. Keyes, J. Sacton, J. H. Wickens, and M. M. Block, *Nuovo Cimento A* **31**, 401 (1976).
- [11] R. C. Byrd, P. L. McGaughey, W. C. Sailor, R. C. Hammock, and Y. Yariv, *Nucl. Instrum. Methods A* **313**, 437 (1992).
- [12] I. Kumagai-Fuse, S. Okabe, and Y. Akaishi, *Phys. Rev. C* **54**, 2843 (1996).
- [13] G. J. Feldman and R. D. Cousins, *Phys. Rev. D* **57**, 3873 (1998).
- [14] H. Outa, M. Aoki, R. S. Hayano, T. Ishikawa, M. Iwasaki, A. Sakaguchi, E. Takada, H. Tamura, and T. Yamazaki, *Nucl. Phys. A* **639**, 251c (1998).
- [15] M. M. Block *et al.*, in *Proceedings of the International Conference on Hyperfragments, St. Cergue, Switzerland, 1963*, edited by W. O. Lock (CERN, Geneva, Switzerland, 1964), p. 63.
- [16] J. G. Fetkovich, J. McKenzie, B. R. Riley, I.-T. Wang, M. Derrick, T. Fields, L. G. Hyman, and G. Keyes, *Phys. Rev. D* **6**, 3069 (1972).
- [17] J. H. Jun, *Phys. Rev. C* **63**, 044012 (2001).
- [18] F. J. Gilman and M. B. Wise, *Phys. Rev. D* **20**, 2392 (1979).
- [19] M. Oka, *Nucl. Phys. A* **754**, 117 (2005).
- [20] G. Garbarino, A. Parreño, and A. Ramos, *Phys. Rev. C* **69**, 054603 (2004).