

Title	On Rogosinski theorem(Study on Calculus Operators in Univalent Function Theory)
Author(s)	Nunokawa, Mamoru
Citation	数理解析研究所講究録 (2007), 1538: 51-54
Issue Date	2007-02
URL	http://hdl.handle.net/2433/59046
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

On Rogosinski theorem

Mamoru Nunokawa

Emeritus Professor of University Gunma
Hoshikuki-cho 798-8, Chuo-ward, Chiba 260-0808, Japan
e-mail : mamoru_nuno@doctor.nifty.jp

1 Introduction

Let $F(z)$ be analytic and univalent in the unit disc $\mathbf{E} = \{z \mid |z| < 1\}$ and let $D = F(\mathbf{E})$ be the image of \mathbf{E} under the mapping $w = f(z)$. Let $f(z)$ be analytic in \mathbf{E} , but not necessarily univalent, and $f(\mathbf{E}) \subset D$. Then $f(z)$ is said to be subordinate to $F(z)$ in \mathbf{E} , denoted by $f(z) \prec F(z)$. It is well known that if $f(z) \prec F(z)$ in \mathbf{E} , then there exists a function $w(z)$, analytic in \mathbf{E} and with $|w(z)| < 1$, such that

$$f(z) = F(w(z)), \quad z \in \mathbf{E}.$$

If $f(0) = F(0)$, then $w(0) = 0$ and $|w(z)| \leq |z|$ in \mathbf{E} .

Rogosinski[1] proved the following theorem.

Theorem A. *Let $f(z) \prec F(z)$ in \mathbf{E} . Then*

$$\int_0^{2\pi} |f(re^{i\theta})|^p d\theta \leq \int_0^{2\pi} |F(re^{i\theta})|^p d\theta$$

where $0 < p$ and $0 \leq r < 1$.

2 Obtained results

Theorem 1. *Let $f(z) \prec F(z)$ in \mathbf{E} and $F(z) \neq 0$ in \mathbf{E} .*

Then

$$\int_0^{2\pi} \frac{1}{|f(re^{i\theta})|^p} d\theta \leq \int_0^{2\pi} \frac{1}{|F(re^{i\theta})|^p} d\theta$$

where $0 < p$ and $0 \leq r < 1$.

Proof. From the assumption of the Theorem, $f(z)^{-p}$ and $F(z)^{-p}$ are analytic in \mathbb{E} and so, from the Poisson integral form of harmonic function theory, we have

$$\begin{aligned} \frac{1}{f(z)^p} &= \frac{1}{F(w(z))^p} \\ &= \frac{1}{2\pi} \int_{|\zeta|=R} \frac{1}{F(\zeta)^p} \left(\operatorname{Re} \frac{\zeta + w(z)}{\zeta - w(z)} \right) d\zeta \end{aligned}$$

where $z = re^{i\theta}$, $\zeta = Re^{i\varphi}$, $|z| = r < |\zeta| = R < 1$, and $|w(z)| \leq |z|$.
Since

$$\operatorname{Re} \left(\frac{\zeta + w(z)}{\zeta - w(z)} \right) > 0 \text{ in } \mathbb{E},$$

it follows that

$$\begin{aligned} & \int_0^{2\pi} \frac{1}{|f(re^{i\theta})|^p} d\theta \\ & \leq \int_0^{2\pi} \frac{1}{2\pi} \int_{|\zeta|=R} \frac{1}{|F(\zeta)|^p} \left(\operatorname{Re} \frac{\zeta + w(z)}{\zeta - w(z)} \right) d\varphi d\theta \\ & = \frac{1}{2\pi} \int_{|\zeta|=R} \int_0^{2\pi} \frac{1}{|F(\zeta)|^p} \left(\operatorname{Re} \frac{\zeta + w(z)}{\zeta - w(z)} \right) d\theta d\varphi \\ & = \frac{1}{2\pi} \int_{|\zeta|=R} \left\{ \frac{1}{|F(Re^{i\varphi})|^p} \int_{|z|=r} \left(\operatorname{Re} \frac{\zeta + w(z)}{\zeta - w(z)} \right) \frac{dz}{iz} \right\} d\varphi \\ & = \int_0^{2\pi} \frac{1}{|F(Re^{i\varphi})|^p} d\varphi \end{aligned}$$

Putting $R \rightarrow r$, we have

$$\int_0^{2\pi} \frac{1}{|f(re^{i\theta})|^p} d\theta \leq \int_0^{2\pi} \frac{1}{|F(re^{i\theta})|^p} d\theta.$$

□

Prof. Owa (Kinki Univ.) pointed out another proof as the following : if $f(z) \prec F(z)$ in \mathbb{E} and $F(z) \neq 0$ in \mathbb{E} , then $\frac{1}{f(z)} \prec \frac{1}{F(z)}$ and applying Theorem A, we can obtain a proof of Theorem 1.

From Theorem A and Theorem 1, we obtain the following theorem.

Theorem 1'. Let $f(z) \prec F(z)$ in \mathbb{E} and $F(z) \neq 0$ in \mathbb{E} .

Then

$$\int_0^{2\pi} |f(re^{i\theta})|^p d\theta \leq \int_0^{2\pi} |F(re^{i\theta})|^p d\theta$$

where p is arbitrary real number and $0 \leq r < 1$.

Theorem 2. Let $f(z) \prec F(z) = z^m(a_m + a_{m+1}z + a_{m+2}z^2 + \dots)$ in \mathbb{E} and let z_k , $k = 1, 2, 3, \dots, n$, $0 < |z_1| \leq |z_2| \leq |z_3| \leq \dots \leq |z_n|$, are the zeros of $F(z)$ in \mathbb{E} which are to

be written with their multiplicities.

Then, if $F(z) \neq 0$ on certain circle $|z| = r < 1$, $z = re^{i\theta}$, we have

$$\int_0^{2\pi} \frac{1}{|f(re^{i\theta})|^p} d\theta \geq \frac{2\pi}{r^{m+n}} \prod_{k=1}^n |z_k|$$

where $0 < p$.

Proof. Without generalization, we can choose R , $0 < R < 1$ in such a manner that $F(z) \neq 0$ on the circle $|z| = R$. Let us construct a function $B(z)$ which has the same zeros with the same multiplicities in $|z| < R < 1$ as $F(z)$ has, and so, we choose

$$B(z) = \left(\frac{z}{R}\right)^m \prod_{k=1}^l \frac{R(z - z_k)}{R^2 - \bar{z}_k z}, \quad l \leq n.$$

Putting

$$g(z) = \left(\frac{B(z)}{F(z)}\right)^p, \quad 0 < p \quad \text{and} \quad z = re^{i\theta},$$

then $g(z)$ is analytic in $|z| < R$ and $g(z) \neq 0$ in $|z| < R$. From the Poisson integral form of harmonic functions, we have

$$g(z) = \frac{1}{2\pi} \int_{|\zeta|=R} g(\zeta) \operatorname{Re} \left(\frac{\zeta + z}{\zeta - z} \right) d\varphi$$

where $|z| = r < |\zeta| = R < 1$ and $\zeta = Re^{i\varphi}$.

Then, we have

$$\begin{aligned} \left(\frac{B(w(z))}{F(w(z))}\right)^p &= \left(\frac{B(w(z))}{f(z)}\right)^p \\ &= \frac{1}{2\pi} \int_{|\zeta|=R} \left(\frac{B(\zeta)}{F(\zeta)}\right)^p \operatorname{Re} \left(\frac{\zeta + w(z)}{\zeta - w(z)} \right) d\varphi. \end{aligned}$$

Here, we have

$$\operatorname{Re} \left(\frac{\zeta + w(z)}{\zeta - w(z)} \right) > 0 \quad \text{in} \quad |z| < R,$$

$$|B(w(z))| < 1 \quad \text{on} \quad |z| = r < R < 1,$$

and

$$|B(\zeta)| = 1 \quad \text{on} \quad |\zeta| = R.$$

Then, it follows that

$$\begin{aligned} \frac{1}{|f(re^{i\theta})|^p} &> \frac{|B(w(re^{i\theta}))|^p}{|f(re^{i\theta})|^p} \\ &= \left| \frac{1}{2\pi} \int_{|\zeta|=R} \left(\frac{B(\zeta)}{F(\zeta)}\right)^p \operatorname{Re} \left(\frac{\zeta + w(z)}{\zeta - w(z)} \right) d\varphi \right|. \end{aligned}$$

Therefore, we have

$$\begin{aligned}
& \int_0^{2\pi} \frac{1}{|f(re^{i\theta})|^p} d\theta \\
& > \int_0^{2\pi} \left| \frac{1}{2\pi} \int_{|\zeta|=R} \left(\frac{B(\zeta)}{F(\zeta)} \right)^p \operatorname{Re} \left(\frac{\zeta + w(z)}{\zeta - w(z)} \right) d\varphi \right| d\theta \\
& = \left| \frac{1}{2\pi} \int_{|\zeta|=R} \left(\frac{B(\zeta)}{F(\zeta)} \right)^p \int_0^{2\pi} \operatorname{Re} \left(\frac{\zeta + w(z)}{\zeta - w(z)} \right) d\theta d\varphi \right| \\
& = \left| \int_{|\zeta|=R} \left(\frac{B(\zeta)}{F(\zeta)} \right)^p d\varphi \right| \\
& = \left| \int_{|\zeta|=R} \left(\frac{B(\zeta)}{F(\zeta)} \right)^p \frac{d\zeta}{i\zeta} \right| \\
& = \left| 2\pi \left(\frac{B(0)}{F(0)} \right)^p \right| \\
& = 2\pi \frac{\prod_{k=1}^l |z_k|}{R^{m+l}} > 2\pi \frac{\prod_{k=1}^n |z_k|}{R^{m+n}}.
\end{aligned}$$

Putting $R \rightarrow r$, we have

$$\int_0^{2\pi} \frac{1}{|f(re^{i\theta})|^p} d\theta > 2\pi \frac{\prod_{k=1}^n |z_k|}{r^{m+n}}.$$

This completes the proof of Theorem 2. □

References

- [1] W. Rogosinski, *On the coefficients of subordinate functions*, Proc. London Math. Soc., (2), 48(1943), 48-82.