

Title	A continuous version of Gale's feasibility theorem(Discrete and Continuous Structures in Optimization)
Author(s)	Nozawa, Ryohei
Citation	数理解析研究所講究録 (1996), 945: 83-92
Issue Date	1996-04
URL	http://hdl.handle.net/2433/60221
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

A continuous version of Gale's feasibility theorem

Ryôhei Nozawa

札幌医科大学医学部 野澤 亮平

1. Introduction

There are several approaches to formulate flow problems on continuous networks. In this paper, using a formulation due to Iri (1979) and Strang (1983), we establish a continuous version of Gale's feasibility theorem [1].

The theorem is known as the "Supply - Demand Theorem" in a special case. By means of a cut capacity, this gives a necessary and sufficient condition for an existence of feasible flows.

Let us recall our formulation of continuous network and state a continuous version of the Supply - Demand Theorem. As for a discrete version, one can refer to Ford and Fulderson's book (1962). In this discussion, we assume that all functions and sets are sufficiently smooth. Let Ω be a bounded domain of n -dimensional Euclidean space R^n and $\partial\Omega$ be the boundary. Let A, B be disjoint subsets of $\partial\Omega$ which are regarded as a source and a sink. In our continuous network, every flow is represented by a vector field and every feasible flow σ satisfies the capacity constraint which is written as

$$\sigma(x) \in \Gamma(x) \text{ for all } x \in \Omega,$$

where Γ is a set-valued mapping from Ω to R^n . The flow value of σ is defined by $\sigma \cdot \nu$ on $\partial\Omega$. We call Ω with this capacity constraint a continuous network.

Furthermore, every cut is identified with a subset of Ω in our network. Let S be a cut and ν^S be the unit outer normal to S . Then the cut capacity $C(S)$ is defined by

$$C(S) = \int_{\Omega \cap \partial S} \beta(\nu^S(x), x) ds(x),$$

where

$$\beta(v, x) = \sup_{w \in \Gamma(x)} v \cdot w$$

for $v \in R^n$ and ds is the surface element. If the capacity constraint is isotropic, that is, $\Gamma(x) = \{w \in R^n \mid |w| \leq c(x)\}$ with some nonnegative function $c(x)$, then

$$C(S) = \int_{\Omega \cap \partial S} c(x) ds(x).$$

Let a, b be real-valued functions on A, B respectively and let ν be the unit outer normal to Ω . Then the problem of supply-demand in a simple case is stated as follows:

$$\begin{aligned}
 \text{(SD)} \quad & \text{Find } \sigma \text{ such that} \\
 & \sigma(x) \in \Gamma(x) \text{ for all } x \in \Omega, \\
 & \operatorname{div} \sigma = 0 \text{ on } \Omega, \quad -\sigma \cdot \nu = 0 \text{ on } \partial\Omega - (A \cap B), \\
 & -\sigma \cdot \nu \leq a \text{ on } A, \quad \sigma \cdot \nu \geq b \text{ on } B.
 \end{aligned}$$

The Supply-Demand theorem assures that (SD) has a solution if and only if

$$\text{(G)} \quad C(S) \geq \int_{B \cap \partial S} b ds - \int_{A \cap \partial S} a ds \quad \text{for each cut } S.$$

This can be proved by the aid of a continuous version of max-flow min-cut theorem under some assumptions. However, we can not apply the same method to a variant of (SD), which is called a symmetric type by Ford and Fulkerson.

On the other hand, Neumann [5] and Oettli and Yamasaki [8] investigated a problem of feasibility of flows and proved similar results in their own network formulations. Their method is based on a generalized Hahn-Banach Theorem and is applicable even for a symmetric supply-demand problem. In the next section, we give a concrete formulation of our problem in a more general form than (SD), and give a corresponding condition which is equivalent with an existence of solutions for the problem under suitable assumptions. Finally in §3, we consider (SD) as a special case and examine the assumptions.

2. Problem setting and a main theorem

Let Ω be a bounded domain in n -dimensional Euclidean space R^n with Lipschitz boundary $\partial\Omega$. One can consider $n - 1$ -dimensional surface measure on $\partial\Omega$ which is equal to $n - 1$ -dimensional Hausdorff measure H_{n-1} on $\partial\Omega$. We note that the unit outer normal ν to Ω is defined and essentially bounded measurable on $\partial\Omega$ with respect to H_{n-1} . Let Γ be a set-valued mapping from Ω to R^n which satisfies the following two conditions:

- (H1) $\Gamma(x)$ is a compact convex set containing 0 for all $x \in \Omega$.
 (H2) Let $\varepsilon > 0$ and Ω_0 be a compact subset of Ω .

Then there is $\delta > 0$ such that

$$\Gamma(x) \subset \Gamma(y) + B(0, \varepsilon) \text{ if } x, y \in \Omega_0 \text{ and } |x - y| < \delta.$$

In what follows, we assume that each feasible flow is represented by an essentially bounded vector field σ on Ω satisfying the following capacity constraints:

$$\sigma(x) \in \Gamma(x) \quad \text{for a.e. } x \in \Omega.$$

Furthermore if $\operatorname{div} \sigma \in L^n(\Omega)$, then $\sigma \cdot \nu$ can be defined as a function in $L^\infty(\partial\Omega)$ in a weak sense by Kohn and Temam [2]. Let $F \in L^n(\Omega)$ and $\lambda, \mu \in L^\infty(\partial\Omega)$ with $\lambda \leq \mu$. Then for the quintuple $(\Omega, \Gamma, F, \mu, \lambda)$, our problem is stated as follows:

- (P) Find $\sigma \in L^\infty(\Omega; R^n)$ such that $\sigma(x) \in \Gamma(x)$ for a.e. $x \in \Omega$,
 $\operatorname{div} \sigma = F$ a.e. on Ω and $\lambda \leq \sigma \cdot \nu \leq \mu$ H_{n-1} -a.e. on $\partial\Omega$

Problem (SD) considered in §1 can be written in this form with $F = 0$.

To specify the class of cuts, we consider the space $BV(\Omega)$ of functions of bounded variation on Ω :

$$BV(\Omega) = \{u \in L^1(\Omega) \mid \nabla u \text{ is a Radon measure of bounded variation on } \Omega\},$$

where $\nabla u = (\partial u / \partial x_1, \dots, \partial u / \partial x_n)$ is understood in the sense of distribution. We denote the characteristic function of a subset S of Ω by χ_S and set

$$Q = \{S \subset \Omega \mid \chi_S \in BV(\Omega)\}.$$

Let $S \in Q$. Then the reduced boundary $\partial^* S$ of S is the set of all $x \in \partial S$ where Federer's normal $\nu = \nu(x)$ to S exists. It is known that $\partial^* S$ is a measurable set with respect to both the measure of total variation of $|\nabla \chi_S|$ and H_{n-1} , $|\nabla \chi_S|(R^n - \partial^* S) = 0$ and $|\nabla \chi_S|(E) = H_{n-1}(E)$ for each $|\nabla \chi_S|$ -measurable subset E of $\partial^* S$. Furthermore let $\gamma u \in L^1(\partial\Omega)$ be the trace of $u \in BV(\Omega)$. Then [4; Theorem 6.6.2] implies that $\gamma \chi_S = \chi_{\partial^* S \cap \partial\Omega}$ H_{n-1} -a.e. on $\partial\Omega$. Accordingly, replacing ds by H_{n-1} and ∂S by $\partial^* S$, we can define the cut capacity as follows:

$$C(S) = \int_{\Omega \cap \partial^* S} \beta(\nu^S(x), x) dH_{n-1},$$

where $\beta(\cdot, x)$ is the support functional of $\Gamma(x)$ as defined in §1. Let $\nabla u / |\nabla u|$ be the Radon-Nikodym derivative of ∇u with respect to $|\nabla u|$ and set

$$\psi(u) = \int_{\Omega} \beta(\nabla u / |\nabla u|, x) d|\nabla u|(x)$$

for $u \in BV(\Omega)$. Then $C(S) = \psi(\chi_S)$. Since β is continuous and nonnegative by (H1) and (H2), $C(S)$ is finite. We set

$$\lambda(S) = \int_{\partial\Omega \cap \partial^* S} \lambda dH_{n-1}, \quad \mu(S) = \int_{\partial\Omega \cap \partial^* S} \mu dH_{n-1}, \quad F(S) = \int_S F dx.$$

for convenience sake, and consider the condition

$$(C) \quad C(S) \geq \lambda(S) - F(S) \text{ and } C(S) \geq -\mu(\Omega - S) + F(\Omega - S) \\ \text{hold for all } S \in Q.$$

Now we can state a continuous version of Gale's feasibility theorem.

THEOREM 2.1. *Assume that (H1) and (H2) hold. If (P) has a solution, then condition (C) holds. Conversely if $\cup_{x \in \Omega} \Gamma(x)$ is bounded and condition (C) holds, then (P) has a solution.*

To prove this theorem, we need some lemmas. First applying an isoperimetric inequality due to [4] we have

LEMMA 2.2. *There is $\sigma_0 \in L^\infty(\Omega; R^n)$ such that $\operatorname{div} \sigma_0 = F$ a.e. on Ω .*

PROOF: First assume that $\int_\Omega F dx = 0$. We use a max-flow min-cut theorem of Strang's type (1983):

$$\sup\{t \geq 0 \mid \operatorname{div} \sigma = -tF \text{ a.e. on } \Omega, \sigma \cdot \nu = 0 \text{ } H_{n-1}\text{-a.e. on } \partial\Omega \\ \text{for some } \sigma \in L^\infty(\Omega; R^n) \text{ with } \|\sigma\|_\infty \leq 1\} \\ = \inf\{H_{n-1}(\Omega \cap \partial^* S) / \int_S F dx \mid \int_S F dx > 0, S \subset \Omega, \chi_S \in BV(\Omega)\}.$$

(The proof is in [6].) To prove the existence of σ_0 , it is sufficient to show that the supremum is positive. We can prove that the infimum is positive as follows. According to [4; p.303] there is a positive constant k such that $\min(m_n(S), m_n(\Omega - S)) \leq kH_{n-1}(\Omega \cap \partial^* S)^{n/(n-1)}$, where m_n denotes the Lebesgue measure on R^n . Since

$$\int_S F dx \leq \left(\int_S 1 dx\right)^{(n-1)/n} \cdot \left(\int_S |F|^n dx\right)^{1/n} \leq \|F\|_n (m_n(S))^{(n-1)/n}$$

and

$$\int_S F dx = \int_{\Omega - S} -F dx \leq \left(\int_{\Omega - S} 1 dx\right)^{(n-1)/n} \cdot \left(\int_{\Omega - S} |F|^n dx\right)^{1/n} \\ \leq \|F\|_n (m_n(\Omega - S))^{(n-1)/n},$$

we can conclude that

$$\int_S F dx \leq k_1 H_{n-1}(\Omega \cap \partial^* S)$$

with $k_1 = \|F\|_n k^{(n-1)/n}$ for all $S \in Q$. It follows that the infimum is not less than $1/k_1$.

Finally in case of $\int_\Omega F dx \neq 0$, consider σ_1 such that $\operatorname{div} \sigma_1$ equals constantly $\int_\Omega F dx$, σ_2 such that $\operatorname{div} \sigma_2 = F - \int_\Omega F dx$ and set $\sigma_0 = \sigma_1 + \sigma_2$. Then $\operatorname{div} \sigma_0 = F$. This completes the proof.

From now on we fix σ_0 in Lemma 2.2. For $\sigma \in L^\infty(\Omega; R^n)$ such that $\operatorname{div} \sigma \in L^n(\Omega)$ and $u \in BV(\Omega)$, according to [2] we can define the distribution $(\sigma \nabla u)$ by

$$(\sigma \nabla u)(\varphi) = - \int_\Omega u \nabla \varphi \cdot \sigma dx - \int_\Omega u \varphi \operatorname{div} \sigma dx$$

for $\varphi \in C_0^\infty(\Omega)$. Since $BV(\Omega) \subset L^{n/(n-1)}(\Omega)$, each integral in the definition is finite. Furthermore it is known that $(\sigma \nabla u)$ is regarded as a bounded measure and that

$$(\sigma \nabla u)(\Omega) + \int_\Omega u \operatorname{div} \sigma dx = \int_{\partial\Omega} \gamma u \sigma \cdot \nu dH_{n-1}$$

holds. This is Green's formula due to Kohn and Temam [2; Proposition 1.1]. (See also [6; Theorem 2.3].) Using this formula, we can prove

LEMMA 2.3. *If (P) has a solution, then (C) holds.*

PROOF: Let σ be a solution of (P). Then by Green's formula stated above,

$$\begin{aligned} C(S) &\geq (\sigma \nabla \chi_S)(\Omega) = \int_{\partial\Omega \cap \partial^* S} \sigma \cdot \nu dH_{n-1} - \int_S \operatorname{div} \sigma dx \\ &\geq \lambda(S) - F(S). \end{aligned}$$

Another inequality in (C) can be similarly proved.

To prove the converse, we follow the idea in [5] and [8]. Let us consider the Sobolev space

$$W^{1,1}(\Omega) = \{u \in L^1(\Omega) \mid \nabla u \in L^1(\Omega; R^n)\},$$

which is a linear subspace of $BV(\Omega)$. We set

$$U = L^1(\Omega; R^n) \times L^1(\partial\Omega) \text{ and } V = \{(\nabla u, \gamma u) \mid u \in W^{1,1}(\Omega)\}.$$

Since $\gamma u \in L^1(\partial\Omega)$ for $u \in W^{1,1}(\Omega)$, V is a linear subspace of U . Let $u^+ = \max(u, 0)$ and $u^- = -\min(u, 0)$. Note that $u^+, u^- \in W^{1,1}(\Omega)$. We define a functional Φ on V by

$$\begin{aligned} \Phi(\nabla u, \gamma u) &= \int_{\Omega} \sigma_0 \cdot \nabla u dx - \int_{\partial\Omega} \sigma_0 \cdot \nu \gamma u dH_{n-1} \\ &\quad + \int_{\partial\Omega} \lambda \gamma u^+ dH_{n-1} - \int_{\partial\Omega} \mu \gamma u^- dH_{n-1} \end{aligned}$$

and set

$$K = \{\sigma \in L^\infty(\Omega; R^n) \mid \sigma(x) \in \Gamma(x) \text{ for a.e. } x \in \Omega\}.$$

For $v \in L^1(\Omega; R^n)$, we define a functional ρ on U by

$$\rho(v, \alpha) = \int_{\Omega} \beta(v(x), x) dx = \sup_{\phi \in K} \int_{\Omega} v \cdot \phi dx$$

for $(v, \alpha) \in U$. The last equality follows from a measurable selection theorem. (Cf. Castaing and Valadier (1977).) Since $\rho(v, \alpha)$ is independent of α , it is sometimes denoted by $\rho(v)$. We note that $\psi(u) = \rho(\nabla u)$ for all $u \in W^{1,1}(\Omega)$. The inequality $\lambda \leq \mu$ implies the next lemma.

LEMMA 2.4. *Φ is superlinear on V , that is, concave and positively homogeneous, and ρ is sublinear on U , that is, $-\rho$ is superlinear. Furthermore ρ is continuous at the origin of U if $\cup_{x \in \Omega} \Gamma(x)$ is bounded.*

Condition (C) can be replaced by an inequality with Φ and ρ .

LEMMA 2.5. *If (C) holds, then $\Phi \leq \rho$ on V .*

PROOF: We use equalities of coarea formula type which are stated in [6]: Let $u \in W^{1,1}(\Omega)$. Set $N_t = \{x \in \Omega \mid u(x) \geq t\}$ and $M_t = \Omega - N_t$ for any real number t . Then $N_t, M_t \in Q$ for a.e. t and

$$\psi(u) = \int_{-\infty}^{\infty} \psi(\chi_{N_t}) dt.$$

Furthermore by [6; Lemma 4.6]

$$\begin{aligned} \int_{\Omega} F u dx &= \int_0^{\infty} \left(\int_{\Omega} F \chi_{N_t} dx - \int_{\Omega} F \chi_{M_t} dx \right) dt, \\ \int_{\partial\Omega} \lambda \gamma u^+ dH_{n-1} &= \int_0^{\infty} \int_{\partial\Omega} \lambda \gamma \chi_{N_t} dH_{n-1} dt, \\ \int_{\partial\Omega} \mu \gamma u^- dH_{n-1} &= \int_0^{\infty} \int_{\partial\Omega} \mu \gamma \chi_{M_t} dH_{n-1} dt. \end{aligned}$$

It follows from these equalities and (C) that

$$\begin{aligned}
\rho(\nabla u) &= \psi(u) = \int_{-\infty}^{\infty} \psi(\chi_{N_t}) dt = \int_0^{\infty} \psi(\chi_{N_t}) dt + \int_0^{\infty} \psi(\chi_{\Omega - M_{-t}}) dt \\
&= \int_0^{\infty} C(N_t) dt + \int_0^{\infty} C(\Omega - M_{-t}) dt \\
&= \int_0^{\infty} (\lambda(N_t) - F(N_t)) dt + \int_0^{\infty} (-\mu(M_{-t}) + F(M_{-t})) dt \\
&\geq \int_0^{\infty} \left(\int_{\partial\Omega} \lambda \gamma \chi_{N_t} dH_{n-1} - \int_{\Omega} F \chi_{N_t} dx \right) dt \\
&\quad + \int_0^{\infty} \left(- \int_{\partial\Omega} \mu \gamma \chi_{M_{-t}} dH_{n-1} + \int_{\Omega} F \chi_{M_{-t}} dx \right) dt \\
&= \int_{\partial\Omega} \lambda \gamma u^+ dH_{n-1} - \int_{\partial\Omega} \mu \gamma u^- dH_{n-1} - \int_{\Omega} u \operatorname{div} \sigma_0 dx \\
&= \int_{\partial\Omega} \lambda \gamma u^+ dH_{n-1} - \int_{\partial\Omega} \mu \gamma u^- dH_{n-1} \\
&\quad - \int_{\partial\Omega} \sigma_0 \cdot \nu \gamma u H_{n-1} + \int_{\Omega} \sigma_0 \cdot \nabla u dx \\
&\geq \Phi(\nabla u, \gamma u).
\end{aligned}$$

Here we have used Green's formula in the last equality. This completes the proof.

By Lemma 2.5 and a version of Hahn-Banach theorem ([3; Corollary 2.2 in p.114]), there is a linear functional ξ on U satisfying $\Phi \leq \xi$ on V and $\xi \leq \rho$ on U . The next lemma is directly proved.

LEMMA 2.6. *If $\cup_{x \in \Omega} \Gamma(x)$ is bounded, then ξ is continuous on U with respect to the canonical norm topology.*

By Lemma 2.6, there is $\sigma \in L^\infty(\Omega; R^n)$ and $\eta \in L^\infty(\partial\Omega)$ such that

$$\xi(v, \alpha) = \int_{\Omega} \sigma \cdot v dx + \int_{\partial\Omega} \eta \alpha dH_{n-1}$$

for all $(v, \alpha) \in U$. However, from the inequality $\xi(v, \alpha) \leq \rho(v)$ for all $\alpha \in L^\infty(\partial\Omega)$, η must be 0.

LEMMA 2.7. Assume that $\cup_{x \in \Omega} \Gamma(x)$ is bounded. Then the vector field σ obtained above is a solution to (P).

PROOF: We set $\Omega_0 = \{x \in \Omega \mid 0 \notin \Gamma(x) - \sigma(x)\}$. Then Ω_0 is a measurable set. Assume that the measure of Ω_0 is positive. Since $\hat{K} = \{\phi \in L^\infty(\Omega; R^n) \mid \phi(x) \in \Gamma(x) - \sigma(x)\}$ is a weakly* closed convex set and does not contain 0, there is $\varphi \in L^1(\Omega; R^n)$ such that $\sup_{\phi \in \hat{K}} \int_{\Omega} \varphi \cdot \phi dx < 0$. Therefore

$$\rho(\varphi) = \sup_{\phi \in \hat{K}} \int_{\Omega} \varphi \cdot (\phi + \sigma) dx < \int_{\Omega} \varphi \cdot \sigma dx = \xi(\varphi, 0).$$

This is a contradiction since $\xi \leq \rho$ on U . Thus $\sigma(x) \in \Gamma(x)$ for almost all $x \in \Omega$.

Next we prove $\operatorname{div} \sigma = F$. If $u \in C_0^\infty(\Omega)$, then $\gamma u = 0$ so that

$$\Phi(\nabla u, \gamma u) = \int_{\Omega} \sigma_0 \cdot \nabla u dx \leq \xi(\nabla u, 0) = \int_{\Omega} \sigma \cdot \nabla u dx.$$

It follows that

$$\int_{\Omega} \sigma_0 \cdot \nabla u dx = \int_{\Omega} \sigma \cdot \nabla u dx$$

for all $u \in C_0^\infty(\Omega)$. This implies that $\operatorname{div} \sigma = \operatorname{div} \sigma_0 = F$ in a distribution sense.

Finally we prove that $\lambda \leq \sigma \cdot \nu \leq \mu$ H_{n-1} -a.e. on $\partial\Omega$. Since $\operatorname{div} \sigma = F \in L^n(\Omega)$, $\sigma \cdot \nu$ is defined as a function in $L^\infty(\partial\Omega)$ and the inequality $\Phi(\nabla u, \gamma u) \leq \int_{\Omega} \sigma \cdot \nabla u dx$ implies that

$$\int_{\partial\Omega} \lambda \gamma u^+ - \mu \gamma u^- dH_{n-1} \leq \int_{\partial\Omega} \gamma u \sigma \cdot \nu dH_{n-1}.$$

For any $\alpha \in L^1(\partial\Omega)$, there is $u \in W^{1,1}(\Omega)$ such that $\alpha = \gamma u$ by Gagliardo (1957). Thus for any nonnegative function $\alpha \in L^1(\partial\Omega)$, we have

$$\begin{aligned} \int_{\partial\Omega} \lambda \alpha dx &\leq \int_{\partial\Omega} \sigma \cdot \nu \alpha dH_{n-1}, \\ - \int_{\partial\Omega} \mu \alpha dx &\leq - \int_{\partial\Omega} \sigma \cdot \nu \alpha dH_{n-1}. \end{aligned}$$

Accordingly, $\lambda \leq \sigma \cdot \nu \leq \mu$ H_{n-1} -a.e. on $\partial\Omega$. This completes the proof.

PROOF OF THEOREM 2.1: The first statement follows from Lemma 2.3 and the second statement follows from Lemma 2.7.

3. Supply - Demand theorem

Let A, B be disjoint Borel subsets of $\partial\Omega$ and a, b be Borel measurable functions on A, B respectively. Then (SD) in §1 should be written in the following concrete form:

$$\begin{aligned}
 \text{(SD)} \quad & \text{Find } \sigma \in L(\Omega; R^n) \\
 & \text{such that } \sigma(x) \in \Gamma(x) \text{ for a.e. } x \in \Omega, \\
 & \operatorname{div} \sigma = 0 \text{ a.e. on } \Omega, \\
 & \sigma \cdot \nu = 0 \text{ } H_{n-1}\text{-a.e. on } \partial\Omega - (A \cap B), \\
 & -\sigma \cdot \nu \leq a \text{ } H_{n-1}\text{-a.e. on } A, \\
 & \sigma \cdot \nu \geq b \text{ } H_{n-1}\text{-a.e. on } B.
 \end{aligned}$$

By setting $\lambda = -a$ on A , $\lambda = b$ on B , $\lambda = 0$ elsewhere on $\partial\Omega$ and $\mu = \max(\lambda, 0)$, Theorem 2.1 implies

THEOREM 3.1. *Assume that (H1), (H2) hold and that $\cup_{x \in \Omega} \Gamma(x)$ is bounded. Then (SD) has a solution if and only if*

$$\text{(G)} \quad C(S) \geq \int_{B \cap \partial^* S} b dH_{n-1} - \int_{A \cap \partial^* S} a dH_{n-1} \text{ for all } S \in Q.$$

Finally we refer to a relation between (SD) and a max-flow problem of Strang's type (MFS) which has been used in the proof of Lemma 2.2 with the boundary condition $\sigma \cdot \nu = 0$. Now let f be an arbitrary function in $L^\infty(\partial\Omega)$ which satisfies the conservation law $\int_{\partial\Omega} f dH_{n-1} = 0$. Then for (Ω, Γ, f) , (MFS) with $F = 0$ is stated as follows:

$$\begin{aligned}
 \text{(MFS)} \quad & \text{Maximize } \lambda \\
 & \text{subject to } (\lambda, \sigma) \in R \times L^\infty(\Omega; R^n), \\
 & \sigma(x) \in \Gamma(x) \text{ a.e. } x \in \Omega, \\
 & \operatorname{div} \sigma = 0 \text{ a.e. on } \Omega, \sigma \cdot \nu = \lambda f \text{ a.e. on } \partial\Omega,
 \end{aligned}$$

and the corresponding min-cut problem (MCS) is

$$\begin{aligned}
 \text{(MCS)} \quad & \text{Minimize } C(S)/L(S) \\
 & \text{subject to } S \subset \Omega, \chi_S \in BV(\Omega), L(S) > 0,
 \end{aligned}$$

where $L(S) = \int_{\partial\Omega \cap \partial^* S} f dH_{n-1}$. Then we have

PROPOSITION 3.2. Assume that (H1) and (H2) hold.

(1) Assume that (G) implies the existence of solutions to (SD) for any disjoint Borel subsets A, B of $\partial\Omega$ and $a \in L^\infty(A), b \in L^\infty(B)$. Then $MFS = MCS$ and (MFS) has an optimal solution for any $f \in L^\infty(\partial\Omega)$ satisfying the conservation law.

(2) Conversely if $MFS = MCS$ and (MFS) has an optimal solution for any $f \in L^\infty(\partial\Omega)$ satisfying the conservation law, then (G) implies the existence of solutions to (SD) for any disjoint Borel subsets A, B of $\partial\Omega$ and $a \in L^\infty(A), b \in L^\infty(B)$ such that $\int_A adH_{n-1} = \int_B bdH_{n-1}$.

It is known that there is an example with $MFS < MCS$ if Γ is unbounded. (See [7].) Thus Proposition 3.2 (1) shows that there is an example of (SD) such that $\cup_{x \in \Omega} \Gamma(x)$ is bounded, condition (G) is satisfied and (SD) has no solution.

Acknowledgement

The author is grateful to Professor Yamasaki for his valuable advise, which is essential in proving Theorem 2.1.

References

- [1] Gale D. (1957) A theorem on flows in networks, Pacific J. Math. 7: 1073 - 1082
- [2] Kohn R. and Temam R. (1983) Dual spaces of stresses and strains, with applications to Hencky plasticity, Appl. Math. Optim. 10 : 1 - 35
- [3] König H. On some basic theorems in convex analysis, pp.107 - 144 in modern Applied Mathematics, ed. by B.Korte. North-Holland, Amsterdam, 1982
- [4] Maz'ja W. (1985) Sobolev spaces, Springer-Verlag, Berlin-New York
- [5] Neumann M.M. (1984) A Ford-Fulkerson type theorem concerning vector-valued flows in infinite networks, Czechoslovak Math. J. 34: 156-162
- [6] Nozawa R. (1990) Max-flow min-cut theorem in an anisotropic network, Osaka J. Math. 27 : 805 - 842
- [7] Nozawa R. (1994) Examples of max-flow and min-cut problems with duality gaps in continuous networks, Mathematical Programming 63 : 213 - 234
- [8] Oettli W. and Yamasaki M. (1994) On Gale's feasibility theorem for certain infinite networks, Arch. Math. (Basel) 62:378 - 384