

Title	COMPLETE INTEGRABILITY OF HAMILTONIAN SYSTEMS AND DIFFERENTIAL GALOIS GROUPS (Lie Groups, Geometric Structures and Differential Equations : One Hundred Years after Sophus Lie)
Author(s)	Ramis, J.P.
Citation	数理解析研究所講究録 (2000), 1150: 43-44
Issue Date	2000-04
URL	<a href="http://hdl.handle.net/2433/64063">http://hdl.handle.net/2433/64063</a>
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Kyoto Sophus Lie Conference SL 101 Lecture 12/1999

## COMPLETE INTEGRABILITY OF HAMILTONIAN SYSTEMS AND DIFFERENTIAL GALOIS GROUPS

by J.P. Ramis

The subject of the lecture is a recent new approach of the old problem of integrability of systems of classical mechanics based upon *differential Galois theory and algebraic groups*. (The lecture is mainly based upon a joint work with *Juan Morales* from Barcelona.)

Roughly speaking an hamiltonian system with  $n$  degrees of freedom is integrable if there exists sufficiently many first integrals (integrals of motion) such that its integration can be reduced to quadratures. Here we are mainly interested in *completely integrable systems* in the classical Liouville sense (i.e. there must exist  $n$  first integrals generically independent and in involution). The investigation of such systems was a central subject during the last century and stimulated the appearance of the theory of Lie groups. Single out integrable systems among all the Hamiltonian systems remain a big open question. H. Poincaré proved that the existence of global integrals of motion is exceptional, a fortiori completely integrable systems are even more exceptional and only a small number of such systems are known. Recently the situation was perfectly stated by Perelomov (1990): *To find a general criterion for complete integrability seems at present a hopeless task*. In this lecture we will present such a criterion. It will sound quite abstract, but it is (surprisingly even for me...) easy to apply to actual problems: it gives new elegant solutions of some classical problems (as Lagrange top) and allows us to solve very easily a lot of open problems.

1. Ordinary differential equations, first integrals, linearized (variational) equations.
2. Hamiltonian systems (the symplectic formulation). Completely integrable systems.
3. A first (weak) formulation of our theorem: if an hamiltonian system is completely integrable, then the Lie algebra of the Zariski closure in the complex linear symplectic group  $Sp(2n; C)$  of the monodromy group of a variational equation along a solution (in complexified time) is *abelian*.
4. Basics about differential Galois theory
5. Our main theorem: if an hamiltonian system is completely integrable,

then the Lie algebra of the differential Galois group of the variational equation along any solution is *abelian*.

#### 6. Applications.

Recipe for applications. Examples of solutions of classical open problems:

Collinear three body problems with homogeneous potentials, Bianchi IX Cosmological model, spring pendulum... (by J.Morales and J.P. Ramis), Collinear four bodies problems with potential  $1/r^2$  (solution of the first open problem after Jacobi and Moser-Calogero results, by Emmanuelle Juillard-Tosel 1999), non integrability of the planar Newton three bodies problem near the equilateral Lagrange solution by Delphine Boucher and Alexei Tsygvintsev 1999), Henon-Heiles systems...