

Title	Problem of Fenchel on the complex projective plane and representations of the 3rd braid group (Fundamental Groups and Algebraic Functions)
Author(s)	Matsuno, Takanori
Citation	数理解析研究所講究録 (2001), 1182: 80-82
Issue Date	2001-01
URL	http://hdl.handle.net/2433/64579
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Problem of Fenchel on the complex projective plane and representations of the 3rd braid group

Takanori MATSUNO

Department of Liberal Arts and Sciences
 Osaka Prefectural College of Technology

1 Abstract

We denote by \mathbf{P}^2 the complex projective plane. Let $C = \{(X_0 : X_1 : X_2) \in \mathbf{P}^2 \mid X_2 X_0^2 - X_1^3 = 0\}$ be a curve of \mathbf{P}^2 . Let $L_\infty = \{(X_0 : X_1 : X_2) \in \mathbf{P}^2 \mid X_2 = 0\}$ be the line of \mathbf{P}^2 , which we call line at infinity. C is a rational curve of degree 3 with a cusp at $(0 : 0 : 1)$. C and L_∞ are tangent at $(1 : 0 : 0)$. Let e_1, e_2 be positive integers greater than 1. Put $D = e_1 C + e_2 L_\infty$. We consider the following problem and give here a partial answer by constructing representations of the 3rd braid group.

Fenchel's Problem Give a condition on the pair (e_1, e_2) for the existence of a finite Galois covering $\pi : X \rightarrow \mathbf{P}^2$ which branches at D .

2 Elementary facts

We choose a point $p_0 \in \mathbf{P}^2 - \{C \cup L_\infty\}$ and fix it. The fundamental group $\pi_1(\mathbf{P}^2 - \{C \cup L_\infty\}, p_0)$ is isomorphic to $\langle \alpha, \beta, \delta \mid \alpha\beta\alpha = \beta\alpha\beta = \delta^{-1} \rangle$ the 3rd braid group. This group is isomorphic to $\langle \gamma, \delta \mid \gamma^3 = \delta^2 \rangle$. This isomorphism is given by $\gamma \mapsto (\alpha\beta)^{-1}, \delta \mapsto (\alpha\beta\alpha)^{-1}$. We identify α (resp. β , resp. δ) with a closed path in $\mathbf{P}^2 - \{C \cup L_\infty\}$ which rounds counterclockwise direction once around non-singular points P_α of C (resp. P_β of C , resp. P_δ of L_∞). Let J be the smallest normal subgroup of $\pi_1(\mathbf{P}^2 - \{C \cup L_\infty\}, p_0)$ which contains α^{e_1} and δ^{e_2} . There is a finite Galois covering which branches at D if and only if there is a normal subgroup K of $\pi_1(\mathbf{P}^2 - \{C \cup L_\infty\}, p_0)$ of finite index with $J \subset K$, which satisfies the following conditions: (1) If $\alpha^k \in K$ then $k \equiv 0 \pmod{e_1}$ and (2) If $\delta^l \in K$ then $l \equiv 0 \pmod{e_2}$. However it is difficult to look for such a K .

Let G be a finite group generated by two elements A, B , which satisfy the relation $ABA = BAB, A^{e_1} = B^{e_1} = 1, (ABA)^{e_2} = 1$. Obviously A and B are conjugate to each other. If there is a finite group G as above, we have a surjective homomorphism $\Phi : \pi_1(\mathbf{P}^2 - \{C \cup L_\infty\}, p_0) \rightarrow G$. Then the kernel of Φ corresponds to a finite Galois covering $\pi : X \rightarrow \mathbf{P}^2$ which branches at D .

Put $Q = ABA$. It is easy to see:

Lemma 2.1 *If G is abelian, then G is a cyclic group.*

Since Q^2 is an element of the center of G ,

Lemma 2.2 *If the order of Q is odd, then G is abelian (G is a cyclic group).*

Hence we have:

Theorem 2.1 *If e_2 is odd, then any covering $\pi : X \rightarrow \mathbf{P}^2$ which branches at D is cyclic.*

Trivially we have:

Proposition 2.1 For given odd number e_2 , if $e_2 \equiv 0 \pmod{3}$ put $e_1 = e_2/3$, otherwise put $e_1 = e_2$. Then there exists $\pi : X \rightarrow \mathbf{P}^2$ which branches at D .

It is well-known (see for example [1]):

Lemma 2.3 For given positive integer n there is a finite group G generated by two elements \hat{Q} of order 2 and \hat{R} of order 3 with $\hat{Q}\hat{R}$ of order n .

By putting $\hat{Q} = \hat{A}\hat{B}\hat{A}$ and $\hat{R} = \hat{A}\hat{B}$, we have:

Theorem 2.2 If e_2 is 2, then for any positive integer e_1 greater than 1 there is a covering $\pi : X \rightarrow \mathbf{P}^2$ which branches at D .

Let D be as before and let $D' = e_1'C + e_2'L_\infty$. Let e_j'' be the LCM $\langle e_j, e_j' \rangle$ ($j = 1, 2$) and put $D'' = e_1''C + e_2''L_\infty$.

By constructing the fiber product, we have:

Proposition 2.2 If there is a covering $\pi : X \rightarrow \mathbf{P}^2$ which branches at D and there is a covering $\pi' : X' \rightarrow \mathbf{P}^2$ which branches at D' , then there is a covering $\pi'' : X'' \rightarrow \mathbf{P}^2$ which branches at D''

3 Cyclic extension

We denote by S_n the symmetric group of n letters. Let $\hat{G} \subset S_r$ be a finite group generated by two permutations \hat{Q}, \hat{R} , which satisfy the relation $\hat{Q}^2 = \hat{R}^3 = 1$. Then \hat{Q} is a product of cycles of length 2 with no common letters and \hat{R} is a product of cycles of length 3 with no common letters.

We may assume \hat{G} has the following properties. (1)transitivity: For each letters x, y there is a permutation of \hat{G} which maps x to y . (2)simplicity: If a permutation of \hat{G} fixes a letter, then it is the unit element of \hat{G} .

Now by showing examples, we give a method to construct a cyclic extension $G \subset S_{r,q}$ of $\hat{G} \subset S_r$ by an element of its center.

The case $r = 3$. Put $\hat{Q} = (a\ b)$ and $\hat{R} = (a\ b\ c)$. In this case $\hat{G} = S_3$ and non-abelian. We need to assume q is odd. Put

$$Q = \begin{pmatrix} a_1 & a_2 & \dots & a_q & b_1 & b_2 & \dots & b_{q-1} & b_q \\ b_1 & b_2 & \dots & b_q & a_2 & a_3 & \dots & a_q & a_1 \end{pmatrix} \begin{pmatrix} c_1 & c_2 & \dots & c_p & c_{p+1} & c_{p+2} & \dots & c_q \\ c_{p+2} & c_{p+3} & \dots & c_q & c_1 & c_2 & \dots & c_{p+1} \end{pmatrix}$$

$$R = \begin{pmatrix} a_1 & a_2 & \dots & a_q & b_1 & b_2 & \dots & b_q & c_1 & c_2 & \dots & c_{q-1} & c_q \\ b_1 & b_2 & \dots & b_q & c_1 & c_2 & \dots & c_q & a_2 & a_3 & \dots & a_q & a_1 \end{pmatrix}$$

Then

$$F = Q^2 = R^3 = (a_1 \dots a_q)(b_1 \dots b_q)(c_1 \dots c_q)$$

and

$$A = R^{-1}Q = (a_1\ c_{p+1}\ a_{p+2}\ c_1 \dots)$$

where $q = 2p + 1$. The order of A is $2q$.

Let G be a finite group generated by two permutations Q, R . F is a center of G . In a natural way we have the following exact sequence:

$$1 \rightarrow \langle F \rangle^G \rightarrow G \rightarrow \hat{G} \rightarrow 1$$

where $\langle F \rangle^G$ is a subgroup of G generated by F . $\langle F \rangle^G$ is a cyclic group of order q . Then we can have a surjective homomorphism $\Phi : \pi_1(\mathbf{P}^2 - \{C \cup L_\infty\}, p_0) \rightarrow G$. Hence we have:

Theorem 3.1 *If q is odd, then there is a finite Galois covering $\pi : X \rightarrow \mathbf{P}^2$ which branches at $2qC + 2qL_\infty$*

The case $r = 4$. Put $\hat{Q} = (a b)$ and $\hat{R} = (b c d)$. In this case $\hat{G} \subset S_4$ and non-abelian. For the extension we need to assume the $\text{LCM}\langle 6, q \rangle = 1$. In a similar way, we have:

Theorem 3.2 *If q is as above, then there is a finite Galois covering $\pi : X \rightarrow \mathbf{P}^2$ which branches at $4qC + 2qL_\infty$*

The case $r = 12$. Put $\hat{Q} = (a j)(b d)(c h)(e l)(f i)(g k)$ and $\hat{R} = (a b c)(d e f)(g h i)(j k l)$. In this case $\hat{G} \subset S_{12}$ and non-abelian. In a similar way, we have:

Theorem 3.3 *There is a finite Galois covering $\pi : X \rightarrow \mathbf{P}^2$ which branches at $3q(q-1)C + 2qL_\infty$*

References

- [1] R.H.Fox, *On Fenchel's conjecture about F -groups*, Mat. Tidsskrift, vol B (1952) 61-65
- [2] M.Namba, *Branched coverings and algebraic functions*, Research Notes in Math. (1987) vol 161 Pitman-Longman