


Title	Studies on Thiourea Derivatives of Fatty Acids
Author(s)	Oda, Ryohei; Suzuki, Tokushige
Citation	京都大学化学研究所報告 (1952), 28: 72-72
Issue Date	1952-03-30
URL	http://hdl.handle.net/2433/74394
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

22. Studies on Thiourea Derivatives of Fatty Acids

Ryohei Oda and Tokushige Suzuki

(Oda Laboratory)

It is already known, that alkylthiuronium salt is produced when alkylhalide is condensed with thiourea in alcohol (Org. Synth. collected Vol. II, 1045) and also that pseudothiouthydantoin is obtained from monochloroacetic acid and thiourea (Org. Synth. 27, 71).



We cannot, however, find any publication in which the above mentioned reactions were performed with halogenated oleic acid ester (9,10-dibromstearic acid ester) and α -bromstearic acid. Therefore the authors intended to investigate this problem and obtained the following results:

- (1) The reaction product from methyl 9,10-dibromstearate and thiourea.

Yield	saponification value	N-content
15g from 20g 9,10-dibromstearate	298.5	9.89%
Calculated as $\text{CH}_3(\text{CH}_2)_7\text{CH}(\text{S})\text{CH}(\text{S})\text{COOH}_3$		
$\begin{array}{c} \text{S} \quad \text{S} \\ \quad \\ \text{C} \quad \text{C} \\ \diagdown \quad / \\ \text{HBr} \cdot \text{NH}_2 \quad \text{NHNHNH}_2 \cdot \text{HBr} \end{array}$	299.8	9.97%

- (2) The reaction product from α -bromstearic acid and thiourea.

Yield	S-content	N-content
17g from 20g α -bromstearic acid	10.25%	8.07%
Calculate as $\text{C}_{16}\text{H}_{33}\text{-CH-C=O}$		
$\begin{array}{c} \text{S} \quad \text{NH} \\ \quad \\ \text{C} \\ \\ \text{NH} \end{array}$	9.55%	8.18%

In both cases, 20g of the starting material was dissolved in 60g butanol and was stirred with 7g thiourea in a 500cc round bottom flask for 10 hours after cooling the reaction products were separated and purified.