

|             |   |
|-------------|---|
| Title       | Theoretical Consideration for the Mechanism of the Probe Gas for High Vacuum Leak-Detection |
| Author(s)   | Ikegami, Hidetsugu; Kokame, Jun; Uemura, Yosiaki  |
| Citation    | Bulletin of the Institute for Chemical Research, Kyoto University (1954), 32(5): 216-216    |
| Issue Date  | 1954-09-31  |
| URL         | <a href="http://hdl.handle.net/2433/75459">http://hdl.handle.net/2433/75459</a>             |
| Right       |   |
| Type        | Departmental Bulletin Paper   |
| Textversion | publisher   |

## Theoretical Consideration for the Mechanism of the Probe Gas for High Vacuum Leak-Detection

Hidetsugu IKEGAMI, Jun KOKAME and Yosiaki UEMURA

(K. Kimura Laboratory)

In order to analyze the mechanism of the leak-detection with the probe gas, the substitution sensitivity factor  $\phi$  (J. Blears, J. H. Leck, *Journ. of Sci. Instr.* Sup. No. 1, 22, (1952)) for the ionization gauge (apparent change of air pressure in the vacuum caused by covering leak with probe gas) (mean air pressure before probing) was calculated. It was assumed that, Poiseuille's law is applicable for the gas flow through the leak (J. Blears : *ibid* ; S. Dushman : "Scientific Foundation of Vacuum Technique", p. 84, (John Wiley & Sons, New York, 1949); A. Guthrie, R. K. Wakering : "Vacuum Equipment & Techniques," p. 22, (McGraw-Hill Book Co., New York, 1949)) and the ionization cross-section of the gas molecule is built up as the sum of the contributions from the component atoms (H. S. W. Massey and E. H. S. Burhop : "Electronic Ionic Impact Phenomena," p. 191, (Oxford, at the Clarendon Press, 1952)).

The values of  $\phi$  calculated for  $H_2$ ,  $CO_2$ ,  $C_4H_{10}$  and coal gas were in agreement with the observed values approximately. The theory was, however, invalid for the "vapour" with low vapor pressure at room temperature such as  $C_6H_6$ ,  $CCl_4$  etc. The reason of these disagreements is mainly ascribable to the "imperfection" of vapors; this effect reduces the  $\phi$ -value in a way for which Poiseuille's law is of no quantitative use. So, the more rigorous treatment for vapours would be desirable.

(Read at the semi-annual meeting of the Institute on June 11, 1954)

## On the Constitution of Zinc-Ferrite. (II)

Isao KUSHIMA and Tsuyoshi AMARUMA

(Sawamura Laboratory)

In the previous report, the constitution of  $Fe_2O_3$ - $ZnFe_2O_4$  system was reported. In this paper explanation is given on the constitution of the  $ZnFe_2O_4$ - $ZnO$  system and the formation of magnetite in the  $Fe_2O_3$ - $ZnO$  system.

$Fe_2O_3$  and  $ZnO$  are mixed in various molar ratio ( $ZnO:Fe_2O_3$ ), and the mixtures are pressed in the mould under 80 kg./cm<sup>2</sup>. pressure and heated at 1000°, 1100° and 1250°C for 3 hours.

For each sintered sample, Debye-Scherrer X-ray photograph was taken, and the magnetic properties were measured by the ballistic galvanometer method. After these experiments