

Title	Primary Processes in the Flash Photolysis of Dimethyl Carbonate Vapor II: Discrimination of the Hydrogen Producing Primary Processes (Commemoration Issue Dedicated to Professor Rempei Gotoh On the Occasion of his Retirement)
Author(s)	Ibuki, Toshio; Takezaki, Yoshimasa
Citation	Bulletin of the Institute for Chemical Research, Kyoto University (1969), 47(4): 239-245
Issue Date	1969-12-16
URL	<a href="http://hdl.handle.net/2433/76306">http://hdl.handle.net/2433/76306</a>
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

## Primary Processes in the Flash Photolysis of Dimethyl Carbonate Vapor II


### Discrimination of the Hydrogen Producing Primary Processes

Toshio IBUKI and Yoshimasa TAKEZAKI\*

Received May 12, 1969


In succession to the previous paper on the primary steps of direct photolysis of dimethyl carbonate vapor by flash method (1849 Å) at room temperature, in which the details of the hydrogen producing processes could not be elucidated, further investigation has been made with the intention to discriminate hydrogen formation processes.

From the isotopic compositions of hydrogen from the mixtures of  $(\text{CH}_3\text{O})_2\text{CO}$ — $(\text{CD}_3\text{O})_2\text{CO}$ , measured mass-spectrometrically, it has been shown that about 32% of the produced hydrogen is formed intramolecularly. The following hydrogen producing primary processes explain the isotopic distribution of hydrogen reasonably :



#### INTRODUCTION

In the previous paper,<sup>1)</sup> we proposed the following primary steps in the flash photolysis of dimethyl carbonate (DMC) vapor at 1849 Å on the basis of the product analyses :



However, it has not been clear whether the hydrogen is produced as an atom or through intramolecular elimination in the primary steps. The photolysis of  $(\text{CH}_3\text{O})_2\text{CO}$ — $(\text{CD}_3\text{O})_2\text{CO}$  mixtures must reveal this matter, *i.e.*, the existence of only  $\text{H}_2$  and  $\text{D}_2$  will be a proof that hydrogen is formed by intramolecular elimination, and on the other hand the presence of an appreciable amount of HD will reflect the atom formation in the system. Thus the isotopic composition measurement of hydrogen has been conducted on the flash photolysis of  $(\text{CH}_3\text{O})_2\text{CO}$ — $(\text{CD}_3\text{O})_2\text{CO}$  mixtures, the results of which is presented in this paper.

#### EXPERIMENTALS

The apparatus, reaction system and optical arrangement were essentially

\* 伊吹 紀男, 竹崎 嘉真: Laboratory of High Pressure Chemistry, Institute for Chemistry, Institute for Chemical Research, Kyoto University, Uji, Kyoto.

identical with those described previously.<sup>1)</sup> Three hydrogen isotopes, H<sub>2</sub>, HD and D<sub>2</sub> were analysed quantitatively by means of CEC Model 21-401 mass-spectrometer, where the average value of the sensitivities of H<sub>2</sub> and D<sub>2</sub> was adopted as the sensitivity of HD.

Dimethyl carbonate was purified by repeated bulb-to-bulb distillation. Gas chromatogram showed no impurity.

Dimethyl carbonate-*d*<sub>6</sub> was prepared<sup>2-4)</sup> by passing 0.3 mole of carbonyl chloride into 0.25 mole of boiling methanol-*d*<sub>4</sub> (E. Merk A. G. Darmstadt, Germany, 99 atomic %). After addition of 0.1 mole of methanol-*d*<sub>4</sub> and reflux for 10 hours in order to convert methyl chloroformate-*d*<sub>3</sub> produced as a by-product into dimethyl carbonate-*d*<sub>6</sub>, the products were fractionally distilled. Distillate in the boiling range of 89.0~90.0°C was further purified by repeated bulb-to-bulb distillation under vacuum. No impurity was detected by gas chromatography with a flame ionization detector.

Since the amounts of H<sub>2</sub> and/or HD formed from partially deuterated dimethyl carbonate in the prepared dimethyl carbonate can be obtained as the ratio to the amount of D<sub>2</sub> in the flash photolysis of that sample alone, mass-spectrometric purity test of the prepared dimethyl carbonate-*d*<sub>6</sub> was not performed.

## RESULTS AND DISCUSSION

The results are summarized in Table 1 and Fig. 1. As may be seen from the isotopic analyses, the hydrogen formed in the flash photolysis is rich in the unmixed ones, though some HD were detected. And as shown in the last column, there is a large deviation of the value of  $[\text{HD}]^2/[\text{H}_2][\text{D}_2]$  from the equilibrium constant for H<sub>2</sub>-D<sub>2</sub> system, *i.e.*, 3.25 at 25°C<sup>5)</sup>.

Table 1. Hydrogen Isotope Analyses in the Flash photolysis of (CH<sub>3</sub>O)<sub>2</sub>CO—(CD<sub>3</sub>O)<sub>2</sub>CO Mixtures.

$\frac{[\text{DMC}]}{[\text{DMC-}d_6]}$	Total Press. (mm Hg)	H <sub>2</sub>	HD (mole %)	D <sub>2</sub>	$\frac{[\text{HD}]^2}{[\text{H}_2][\text{D}_2]}$
0	22.7	0	8.25	91.75	
0.228	24.5	16.4	27.9*	55.7	0.92
0.580	24.0	33.3	34.7*	32.0	1.06
1.00	21.0	45.4	33.7*	20.9	1.09
2.08	23.0	64.6	25.5*	9.9	1.00
3.04	24.1	73.0	20.6*	6.4	0.95
3.94	24.8	76.6	18.0*	5.4	0.89

\* As the HD was produced 0.09 relative to D<sub>2</sub>=1.00 in the products of the photolysis of the prepared DMC-*d*<sub>6</sub> alone, the HD from the partially deuterated dimethyl carbonate was corrected by subtracting 9% of D<sub>2</sub> from the observed amounts of the HD in other runs.

This can be interpreted most simply in terms of the formation of both hydrogen atom and molecule in our system. Since HD should be formed intermolecularly in the succeeding steps and H<sub>2</sub> and D<sub>2</sub> can result from both inter- and intramolecular reactions, it may be natural to assume that hydrogen is

## Flash Photolysis of Dimethyl Carbonate II




Fig. 1. Variation of the yields of hydrogen isotopes with  $[\text{DMC}]/[\text{DMC-d}_6]$ .





formed partly by molecular detachment and partly as an atom in the primary process, *i.e.*,



where  $2a + b = 6$ , from a stoichiometric relation.

There can be two sources of HD: (1) D or H atom abstraction by H or D atom from dimethyl carbonate- $d_6$  or dimethyl carbonate, respectively, and (2) recombination reaction of H and D atoms.

The hydrogen isotope production may be represented by the following reaction mechanism:






where  $\varepsilon_H$  or  $\varepsilon_D$  is the absorption coefficient of ordinary dimethyl carbonate or deuterated dimethyl carbonate, respectively, and parentheses represent the concentration of the compound. M is DMC and/or DMC- $d_6$ , and  $k$ 's are rate constants.

Reaction (5)~(7) may be excluded from the following considerations.

The values of rate constants in the above scheme are not available, but  $k_1$  may be roughly estimated as follows.

The rate constants of next reactions (8)~(10) are known:



$$\log_{10} (k_8/1 \text{ mole}^{-1} \text{ sec}^{-1}) = 9.53 - 6800/2.303RT,^{61}$$

$$\log_{10} (k_9/1 \text{ mole}^{-1} \text{ sec}^{-1}) = 7.628 + 1/2 \log T - 11300/2.303RT,^{7)}$$

$$\log_{10} (k_{10}/1 \text{ mole}^{-1} \text{ sec}^{-1}) = 7.38 - 8140/2.303RT.^{8)}$$

If we assume that the ratio of the rates of abstraction of H from ethane and from methanol is the same for hydrogen atom and methyl- $d_3$  radical, we get  $k_{11} = k_8 k_{10}/k_9 = 2.16 \times 10^5 \text{ l mole}^{-1} \text{ sec}^{-1}$  at 300°K. Further if the rate of hydrogen abstraction from methoxy group of dimethyl carbonate is equal to that of methanol, we obtain  $k_1 = k_{11}$ .

Now, the rate constant of reaction (5) for M=DMC, though not available, must be of the order of  $3 \times 10^{10} \text{ l}^2 \text{ mole}^{-2} \text{ sec}^{-1}$  by analogy to the value obtained for water as the third body.<sup>9)</sup> Then the relative rate of reaction (5) to (1),  $R_5/R_1$ , is  $k_5(\text{H})/k_1$ .

The concentration of hydrogen atom should be much less than  $7 \times 10^{-7}$  atom /l/flash because the decomposition is lower than 5%, then we have  $R_5/R_1 \ll 0.1$ . This means that the extent of recombination reactions, (5)~(7), must be, if not at all, much less than 10% in the hydrogen producing steps.

Moreover, if reaction (I), (II) and (5)~(7) are predominant sources of hydrogen isotopes, the following relation can be deduced under steady-state condition :

$$\frac{R(\text{H}_2)R(\text{D}_2)}{R(\text{HD})^2} = \frac{a}{b} \cdot \frac{R(\text{H}_2) + R(\text{D}_2)}{R(\text{HD})} + \text{constant},$$

where constant is equal to  $\frac{1}{K_e} \cdot \left(2 - \frac{a}{b} + 1\right)^2 - \left(\frac{a}{b}\right)^2$  and  $K_e$  is a equilibrium constant for  $\text{H}_2$ - $\text{D}_2$  system.




Fig. 2. Plot of  $R(\text{H}_2)R(\text{D}_2)/R(\text{HD})^2$  against  $(R(\text{H}_2) + R(\text{D}_2))/R(\text{HD})$ .

When our experimental data are plotted in the form given by this equation, we obtain that  $a/b$  is equal to 0.13 from the gradient, and the "constant" is 0.646 (see Fig. 2), accordingly  $K_e$  becomes 2.42. This value deviates largely from the equilibrium constant, 3.25 at 25°C and 3.18 even at 0°C,<sup>5)</sup> suggesting that the atom recombination is not the major process. Therefore, reactions (5)~(7) may be omitted in the above reaction scheme, and the hydrogen isotopes must result from reaction (I), (II) and (1)~(4).

Now, assuming steady-state behavior we get the following rate relation :

Flash Photolysis of Dimethyl Carbonate II

$$\frac{R(\text{H}_2)}{R(\text{D}_2)} = \frac{\varepsilon_H}{\varepsilon_D} \cdot X \cdot \frac{a + \frac{b\alpha X}{1 + \alpha X}}{a + \frac{b}{1 + \alpha X}}, \quad (\text{A})$$

where  $X$  represents the ratio of  $[\text{DMC}]/[\text{DMC-}d_6]$ , and only primary isotope effect is assumed *i.e.*, that  $k_1/k_2 = k_4/k_3 = \alpha$ .<sup>8,10-12)</sup>

The values of  $\alpha$  and  $\varepsilon_H/\varepsilon_D$  are not available, but since  $\alpha$  is a constant, mathematically there must exist the unique value of  $X$  which satisfies  $\alpha X = 1$ . For the value of  $X = 1/\alpha$ , eq. (A) is rewritten in the following form:

$$\frac{\varepsilon_H}{\varepsilon_D} = \alpha \frac{R(\text{H}_2)}{R(\text{D}_2)}. \quad (\text{B})$$

Thus, assuming any certain value  $\alpha_0$ , we can fix the corresponding single value,  $X_0 = 1/\alpha_0$  from Fig. 1, and the ratio of absorption coefficients,  $\varepsilon_H/\varepsilon_D$ , is obtained by inserting the assumed  $\alpha_0$  and the observed value of  $(R(\text{H}_2)/R(\text{D}_2))_0$  into eq. (B).

On the other hand, we have

$$\frac{R(\text{H}_2) + R(\text{D}_2)}{R(\text{HD})} = \left(1 + \frac{a}{b}\right) \left( \frac{\alpha X}{1 + \varepsilon_D/\varepsilon_H} + \frac{1}{1 + \varepsilon_H/\varepsilon_D} \cdot \frac{1}{X} \right) + \frac{a}{b}. \quad (\text{C})$$

Using assumed  $\alpha_0$  and the value of  $\varepsilon_H/\varepsilon_D$  obtained from eq. (B), we can get the value of  $a/b$  from both gradient and intercept in eq. (C). Then  $\alpha_0$  can be estimated by trial and error method until the values derived by both procedures







Fig. 3. Plot of  $(R(\text{H}_2) + R(\text{D}_2))/R(\text{HD})$  against  $\left\{ \frac{\alpha X}{1 + \varepsilon_D/\varepsilon_H} + \frac{1}{1 + \varepsilon_H/\varepsilon_D} \cdot \frac{1}{X} \right\}$ .

coincide. Such value of  $\alpha_0$  is found to be 1.02, and  $\epsilon_H/\epsilon_D$  and  $a/b$  are 2.14 and 0.48, respectively. This means that the fraction of the total hydrogen formed molecularly is about  $0.48/(1+0.48)=0.32$ , because one hydrogen atom forms one hydrogen molecule in the above scheme.

From  $2a+b=6$  and  $a/b=0.48$  we have  $(\text{CH}_3\text{O})_2\text{CO}+h\nu \longrightarrow 3\text{CO}+3/2\text{H}_2+3\text{H}$ , as an apparent stoichiometric relation, but actually the following primary processes must be involved:



where  $p$ ,  $q$  and  $r$  are the respective probabilities with which reactions (12), (13) and (14) occur, and other hydrogen atom producing steps can be ruled out from the energetic considerations.


Since  $p+q+r=1$  and  $a/b=(3p+2q+r)/(2q+4r)\simeq 1/2$ , we get  $p=r-1/2$  and  $q=3/2-r$ . Since  $p$  and  $q$  can not be negative, we obtain  $3/4 \geq r \geq 1/2$ .

We have estimated that the hydrogen producing primary processes amount to 40% of all primary steps,<sup>11</sup> then the extent of the reaction (14) becomes to be from 20 to 30%.

Since dimethyl carbonate vapor absorbs the light shorter than 2,000 Å,<sup>13)</sup> the 1849 Å line lies near the threshold of absorption of dimethyl carbonate. Then it may not be unreasonable that  $\epsilon_H/\epsilon_D$  has the value 2.14 at 1849 Å, as is supposed from the instances that the absorption coefficients of the deuterated hydrocarbons are about one half of that of the ordinary ones near the threshold of its absorption.<sup>14-17)</sup>

## CONCLUSION

From the results in the previous paper<sup>1)</sup> and the hydrogen isotopic composition of the hydrogen from the mixtures of  $(\text{CH}_3\text{O})_2\text{CO}-(\text{CD}_3\text{O})_2\text{CO}$ , the fraction of the hydrogen formed molecularly is about 0.32 of the total hydrogen, and in the flash photolysis of dimethyl carbonate vapor the following primary processes can explain the observed reaction products and the isotopic distribution in the hydrogen:



## ACKNOWLEDGMENT

The authors wish to express their thanks to Professor Kumasaburo Kodera of Faculty of Science, Kyoto University, for giving facilities in the mass spectral analyses.

## Flash Photolysis of Dimethyl Carbonate II

### REFERENCES

- (1) T. Ibuki and Y. Takezaki, *this Bulletin*, **45**, 406 (1967).
- (2) E. Muller, "Methoden der Organischen Chemie", Vol. 8, Georg Thieme Verlag, Stuttgart, 1952, p. 106.
- (3) F. D. Chattaway and E. Saerens, *J. Chem. Soc.*, 708 (1920).
- (4) H. Lux, *Chem. Ber.*, **62**, 1824 (1929).
- (5) H. C. Urey, *J. Chem. Soc.* 562 (1947).
- (6) M. R. Berlie and D. J. LeRoy, *Disc. Faraday Soc.*, **14**, 50 (1953).
- (7) (a) J. R. McNesby and A. S. Gordon, *J. Am. Chem. Soc.*, **77**, 4719 (1955). (b) M. H. J. Wijnen, *J. Chem. Phys.*, **23**, 1357 (1955).
- (8) R. Shaw and J. C. J. Thynne, *Trans. Faraday Soc.*, **62**, 104 (1966).
- (9) W. Steiner, *ibid.*, **31**, 962 (1935).
- (10) W. M. Jackson, J. R. McNesby, and B. de B. Darwent, *J. Chem. Phys.*, **37**, 1610 (1962).
- (11) P. Gray and A. A. Herod, *Trans. Faraday Soc.*, **64**, 1568 (1968).
- (12) P. Gray, A. Jones, and J. C. J. Thynne, *ibid.*, **61**, 474 (1965).
- (13) A. S. Gordon and W. P. Norrish, *J. Phys. Chem.*, **69**, 3013 (1965).
- (14) H. Okabe and D. A. Becker, *J. Chem. Phys.*, **39**, 2549 (1963).
- (15) H. Okabe and J. R. McNesby, *ibid.*, **37**, 1340 (1962).
- (16) C. L. Currie, H. Okabe, and J. R. McNesby, *J. Phys. Chem.*, **67**, 1494 (1963).
- (17) P. G. Wilkinson and R. S. Mulliken, *J. Chem. Phys.*, **23**, 1895 (1955).