

Title	Activation Cross Sections of Zirconium and Molybdenum Isotopes for 14.6 MeV Neutrons (Commemoration Issue Dedicated to Professor Takuji Yanabu on the Occasion of his Retirement)
Author(s)	Fujino, Yoshitaka; Hyakutake, Mikio; Kumabe, Isao
Citation	Bulletin of the Institute for Chemical Research, Kyoto University (1982), 60(2): 205-207
Issue Date	1982-08-31
URL	http://hdl.handle.net/2433/76974
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

NOTE

Activation Cross Sections of Zirconium and Molybdenum Isotopes for 14.6 MeV Neutrons

Yoshitaka FUJINO[†], Mikio HYAKUTAKE and Isao KUMABE*

Received March 12, 1982

About 80% of the total energy in the $D+T$ fusion is carried off by the 14 MeV neutrons. The investigations on the interaction of 14 MeV neutrons with structural materials, therefore, are very important for the design of the fusion reactor. Neutron data are needed for the calculations of tritium breeding, radiation shielding, radiation damage effects, neutron multiplication, isotope production, etc.

Zirconium and molybdenum are the expected structural materials that will be used in fusion reactors. Many workers¹⁾ have measured activation cross sections for 14 MeV neutrons on zirconium and molybdenum. However many of their results often show gross disagreements with one another. The gross disagreement seems to result from that most previous measurements have been made with β -ray counting or with γ -ray counting by NaI (TI) detector. This drawback is obviated with the use of a Ge(Li) detector, because its high resolution available for γ -ray identification makes it possible to obtain reliable and precise cross sections. In the present paper, therefore, the (n, p) , (n, α) and $(n, 2n)$ reaction cross sections on Zr and Mo have been measured at 14.6 MeV by the activation technique using a Ge(Li) detector.

Experimental procedures were similar to those of a previous investigation,²⁾ so that they are described only briefly here. Since values for the half-lives of the product activities of interest range from 1 min to 10 d, several samples of Zr and Mo were prepared and each of them was irradiated for periods from 1 min to 2 h, depending on the half-lives of the product activities of interest. Metal samples of about 1 gr were used. Thin Al-foils of about 50 mg were placed in the front and back of each sample. The sample and the monitor Al-foils were covered with Cd-plates of 0.5 mm thick and was then irradiated at about 9 mm behind a Ti-T neutron source.

The neutron-yield variation was monitored by counting the associated α -particles with a Si-detector. After the end of irradiation, the γ -activities of the samples together with the Al-monitor foils were measured with a 60 cm³ coaxial Ge(Li) detector. The γ -spectra from the Ge(Li) detector were automatically stored on a magnetic tape of a USC-3 on-line computer through an ADC. The activities were identified by both their γ energies and their half-lives. Irradiations and measurements of short-lived activities such as ^{91m}Mo and ^{89m}Zr were repeated for several times in order to obtain good counting statistics.

* 藤野吉隆, 百武幹雄, 隈部 功: Department of Nuclear Engineering, Kyushu University, Fukuoka.

† Present Address: Ohnojo City Municipal Office, Ohnojo-shi, Fukuoka-ken.

The $^{27}\text{Al}(n, \alpha)^{24}\text{Na}$ reaction with a cross section of $114.5 \pm 4.8 \text{ mb}^{31)}$ was used as the monitor for long-lived reaction products. The $^{27}\text{Al}(n, p)^{27}\text{Mg}$ reaction with a cross section of $68.0 \pm 4.8 \text{ mb}$ was used for short-lived reaction products. This cross section for $^{27}\text{Al}(n, p)$ reaction has been obtained in the present work.

Typical gamma decay curve is shown in Fig. 1. Table I shows the results obtained from the present work, together with the half-lives of the products, the γ -ray energies and the number of γ quanta emitted per disintegration used in the calculations of the cross sections.




Fig. 1. Gamma decay curve of $^{89\text{m}}\text{Zr}$ from $^{92}\text{Mo}(n, \alpha)$ reaction

Table 1. Cross sections for (n, p) , (n, α) and $(n, 2n)$ reactions with 14.6 MeV neutrons from the present work

Reaction	$T_{1/2}$	E_r (keV)	η (%)	cross section (mb)	
				experimental	predicted
$^{90}\text{Zr}(n, p)^{90\text{m}}\text{Y}$	3.1 h	482	91	9.0 ± 0.8	
$^{91}\text{Zr}(n, p)^{91\text{m}}\text{Y}$	50.3 m	551	95	21.4 ± 2.2	
$^{92}\text{Zr}(n, p)^{92}\text{Y}$	3.53 h	934	14	20.1 ± 1.5	17.9
$^{94}\text{Zr}(n, p)^{94}\text{Y}$	20.3 m	918	73	10.7 ± 1.1	8.14
$^{92}\text{Mo}(n, p)^{92\text{m}}\text{Nb}$	10.16 d	934	99	68 ± 6	
$^{96}\text{Mo}(n, p)^{96}\text{Nb}$	23.4 h	569	59	18.1 ± 1.4	24.9
$^{97}\text{Mo}(n, p)^{97\text{g}}\text{Nb}$	72 m	665	98	19.2 ± 1.4	
$^{98}\text{Mo}(n, p)^{98}\text{Nb}$	51 m	720	75	3.6 ± 0.3	11.6
$^{90}\text{Zr}(n, \alpha)^{87\text{m}}\text{Sr}$	2.83 h	388	80	4.1 ± 0.3	
$^{94}\text{Zr}(n, \alpha)^{91}\text{Sr}$	9.67 h	1025	30	6.3 ± 0.5	3.91
$^{96}\text{Zr}(n, \alpha)^{93}\text{Sr}$	8.3 m	888	23.7	2.4 ± 0.3	2.20
$^{92}\text{Mo}(n, \alpha)^{89\text{m}}\text{Zr}$	4.18 m	588	87	5.6 ± 0.5	29.9
$^{92}\text{Mo}(n, \alpha)^{89\text{g}}\text{Zr}$	78.4 h	909	99	20.1 ± 1.6	
$^{90}\text{Zr}(n, 2n)^{89\text{m}}\text{Zr}$	4.18 m	588	87	86 ± 8	
$^{90}\text{Zr}(n, 2n)^{89\text{g}}\text{Zr}$	78.4 h	909	99	805 ± 58	
$^{92}\text{Mo}(n, 2n)^{91\text{m}}\text{Mo}$	66 s	658	54	11.8 ± 1.2	
$^{94}\text{Mo}(n, 2n)^{93\text{m}}\text{Mo}$	6.9 h	685	100	2.4 ± 0.2	
$^{100}\text{Mo}(n, 2n)^{99}\text{Mo}$	67 h	740	12	1420 ± 100	

η : Intensity of γ -rays per disintegration

Activation Cross Sections on Zr and Mo Isotopes for 14.6 MeV Neutrons

The empirical formulas proposed by Levkovskii⁴⁾ are perhaps the most successful, or the most widely used formulas for predicting (n, p) and (n, α) cross sections at 14~15 MeV. However rather large discrepancies between experimental cross sections and those calculated by the Levkovskii formulas⁴⁾ for both the (n, p) and (n, α) reactions are observed for nuclei with relatively small or large values of asymmetry parameter $(N-Z)/A$ for a given values of atomic number Z . Kumabe and Fukuda⁵⁾ have reparameterized the Levkovskii formulas over three limited mass regions by carrying out least-squares fits to experimental cross sections. The empirical formulas are as follows.

$$\sigma_{np} = \begin{cases} 21.8A \exp[-34(N-Z)/A] & (40 \leq A \leq 62) \\ 0.75A^2 \exp[-43.2(N-Z)/A] & (63 \leq A \leq 89) \\ 0.75A^2 \exp[-45(N-Z)/A] & (90 \leq A \leq 160) \end{cases}$$

$$\sigma_{n\alpha} = \begin{cases} 51.0\sqrt{A} \exp[-30(N-Z)/A] & (30 \leq A \leq 60) \\ 55.0\sqrt{A} \exp[-33(N-Z)/A] & (61 \leq A \leq 105) \\ 7.6 \times 10^{-4}A^3 \exp[-40(N-Z)/A] & (106 \leq A \leq 150) \end{cases}$$

The (n, p) and (n, α) cross section values predicted from above formulas are also shown in Table I. The predicted values except for $^{98}\text{Mo}(n, p)$ and $^{94}\text{Zr}(n, \alpha)$ reactions are in fairly good agreement with the present experimental cross sections.

REFERENCES

- (1) CINDA 81, An index to the literature on microscopic neutron data, (May, 1981), IAEA, Vienna.
- (2) T. Sato, Y. Kanda and I. Kumabe, *J. Nucl. Sci. Technol.* **12**, 681 (1975).
- (3) Y. Kanda and R. Nakasima, JAERI-1207 (1972).
- (4) V. N. Levkovskii, *Sov. J. Nucl. Phys.* **18**, 361 (1974).
- (5) I. Kumabe and K. Fukuda, NEANDC (J)-56/U, 45 (1978).