

Title	Chern-Simons variation and Hida-Mazur theory(Topology, Complex Analysis and Arithmetic of Hyperbolic Spaces)
Author(s)	Morishita, Masanori; Terashima, Yuji
Citation	数理解析研究所講究録 (2007), 1571: 23-30
Issue Date	2007-10
URL	http://hdl.handle.net/2433/81293
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Chern-Simons variation and Hida-Mazur theory

Masanori Morishita (Kyushu Univ.)
 Yuji Terashima (Tokyo Inst. of Tech.)

We would like to discuss the variation of $SL_2(\mathbb{C})$ Chern-Simons invariants over the deformation space of hyperbolic structures on a knot complement in analogy with Hida-Mazur theory on the deformation of Galois representations and modular p -adic L -functions. The motivation and idea are coming from the analogy between knot theory and number theory, and so let us start to recall the basic analogies between knots and primes.

1. Analogies

knot $K : S^1 = K(\mathbb{Z}, 1) \hookrightarrow S^3 = \mathbb{R}^3 \cup \{\infty\}$	\leftrightarrow	prime $\text{Spec}(\mathbb{F}_p) = K(\hat{\mathbb{Z}}, 1) \hookrightarrow \text{Spec}(\mathbb{Z}) \cup \{\infty\}$
tube neighborhood V_K ∂V_K $D_K = \pi_1(\partial V_K)$	\leftrightarrow	p -adic integers $\text{Spec}(\mathbb{Z}_p)$ p -adic numbers $\text{Spec}(\mathbb{Q}_p)$ $D_p = \pi_1^{\text{ét}}(\text{Spec}(\mathbb{Q}_p))$
$1 \rightarrow \langle m_K \rangle \rightarrow D_K \rightarrow \langle l_K \rangle \rightarrow 1$ l_K : longitude of K m_K : meridian of K $[m_K, l_K] = 1$	\leftrightarrow	$1 \rightarrow I_p \rightarrow D_p \rightarrow \langle \sigma_p \rangle \rightarrow 1$ σ_p : Frobenius over p I_p : inertia gr. over p , $I_p^t = \langle \tau_p \rangle$ τ_p : monodromy over p , $\tau_p^{p-1}[\tau_p, \sigma_p] = 1$

Here, I_p^t denotes the maximal tame quotient of I_p . In general, we call an element of a quotient of I_p a monodromy over p .


$X_K = S^3 \setminus K$ knot group $G_K = \pi_1(X_K)$	\leftrightarrow	$X_p = \text{Spec}(\mathbb{Z}[1/p])$ prime group $G_p = \pi_1^{\text{ét}}(X_p)$
infinite cyclic cover $X_K^\infty \rightarrow X_K$ $\text{Gal}(X_K^\infty/X_K) = \langle \alpha \rangle$	\leftrightarrow	\mathbb{Z}_p -cover $X_p^\infty = \text{Spec}(\mathbb{Z}[1/p, \sqrt[p^\infty]{1}]) \rightarrow X_p$ $\text{Gal}(X_p^\infty/X_p) = \langle \gamma \rangle$
Alexander module/polynomial $\Delta_K(t) = \det(t - \alpha H_1(X_K^\infty))$ analytic torsion = Reidemeister torsion	\leftrightarrow	Iwasawa module/polynomial $I_p(T) = \det(T - (\gamma - 1) H_1(X_p^\infty))$ Iwasawa main conjecture

For more analogies and details, we refer to [Mo]. In fact, there are close analogies between Alexander-Fox theory and Iwasawa theory. From the variational point of view, one sees that:

considering the infinite cyclic cover $X_K^\infty \rightarrow X_K$ (resp. \mathbb{Z}_p -cover $X_p^\infty \rightarrow X_p$)
is equivalent to

considering the deformation of GL_1 -representations of the knot group G_K (resp. prime group G_p).

In number theory, there is a non-abelian generalization of the classical Iwasawa theory along this variational viewpoint, due to mainly H. Hida and B. Mazur, namely the deformation theory of GL_n -representations of G_p and the theory of associated arithmetic invariants. Our motivation was to find an analogue of Hida-Mazur theory in the context of knot theory, which would be a natural non-abelian generalization of Alexander-Fox theory:



2. Deformation of hyperbolic structures on a knot complement and of modular Galois representations

We start to recall some general notions in group representations.

knot: For a knot $K \subset S^3$ with $G_K := \pi_1(S^3 \setminus K)$ and $n \geq 1$, we set

$$\begin{aligned} \mathfrak{X}_K^n &= \text{Hom}(G_K, GL_n(\mathbb{C})) // GL_n(\mathbb{C}) \\ &:= \text{Hom}_{\mathbb{C}\text{-alg}}((R_K^n)^{G_K}, \mathbb{C}), \end{aligned}$$

where R_K^n denotes the tautological n -dimensional representation ring on which G_K acts by the conjugation via the tautological representation $G_K \rightarrow GL_n(R_K^n)$, and $(R_K^n)^{G_K}$ stands for the invariant subring. The set \mathfrak{X}_K^n is a complex affine variety, called the character variety of n -dimensional representations of G_K .

prime: For a prime $\text{Spec}(\mathbb{F}_p)$, the prime group G_p is profinite and hence the naive analogue of the character variety does not provide a good moduli. Thus, following Mazur ([Ma1]), we consider “infinitesimal deformations” of a given residual representation

$$\bar{\rho} : G_p \longrightarrow GL_n(\mathbb{F}_p).$$

Namely, the pair (R, ρ) is called a deformation of $\bar{\rho}$ if

$$\left\{ \begin{array}{l} \bullet R \text{ is a complete noetherian local ring with residue field } R/m_R = \mathbb{F}_p \\ \bullet \rho : G_p \rightarrow GL_n(R) \text{ is a continuous representation with } \rho \bmod m_R = \bar{\rho}. \end{array} \right.$$

In the rest of this note, we assume for simplicity that $p > 2$ and $\bar{\rho}$ is absolutely irreducible. A fundamental theorem by Mazur is:

Theorem 1 ([Ma]). *There is a universal deformation (R_p^n, ρ_p^n) of $\bar{\rho}$ so that any deformation of (R, ρ) is obtained up to a certain conjugacy via a \mathbb{Z}_p -algebra homomorphism $R_p^n \rightarrow R$.*

We then define the universal deformation space $\mathfrak{X}_p^n(\bar{\rho})$ of $\bar{\rho}$ by

$$\mathfrak{X}_p^n(\bar{\rho}) := \text{Hom}_{\mathbb{Z}_p\text{-alg}}(R_p^n, \mathbb{C}_p)$$

where \mathbb{C}_p stands for the p -adic completion of an algebraic closure of \mathbb{Q}_p , and $\mathfrak{X}_p^n(\bar{\rho})$ is regarded as a rigid analytic space. We write $\rho_\varphi := \varphi \circ \rho_p^n : G_p \rightarrow GL_n(\mathbb{C}_p)$ for a $\varphi \in \mathfrak{X}_p^n(\bar{\rho})$.

We will discuss analogies between \mathfrak{X}_K^n and $\mathfrak{X}_p^n(\bar{\rho})$ and some invariants defined on them for the cases of $n = 1$ and 2.

$n = 1$: The GL_1 -theory is simply a restatement of the analogy between Alexander-Fox theory and Iwasawa theory:

$$\begin{array}{ccc} R_K^1 = \Lambda_{\mathbb{C}} = \mathbb{C}[t^{\pm 1}] & & R_p^1 = \hat{\Lambda} = \mathbb{Z}_p[[T]] \\ \mathfrak{X}_K^1 \simeq \mathbb{C}^\times & \leftrightarrow & \mathfrak{X}_p^1(\bar{\rho}) \simeq D_p^1 = \{z \in \mathbb{C}_p \mid |z|_p < 1\} \\ \chi \mapsto \chi(\alpha) & & \varphi \mapsto \rho_\varphi(\gamma) - 1 \\ (\text{Gal}(X_K^\infty/X_K) = \langle \alpha \rangle = \mathbb{Z}) & & (\text{Gal}(\mathbb{Q}^\infty/\mathbb{Q}) = \langle \gamma \rangle = \mathbb{Z}_p) \end{array}$$

$$\begin{array}{ccc} \text{invariants on } \mathfrak{X}_K^1 : & & \text{invariants on } \mathfrak{X}_p^1(\bar{\rho}) : \\ \text{twisted Alexander poly. (analytic torsion)} & \leftrightarrow & \text{twisted Iwasawa poly. (} p\text{-adic } L\text{-function)} \\ \text{for a repr. } \rho : G_K \rightarrow GL_n(\mathbb{C}) & & \text{for a repr. } \rho : G_p \rightarrow GL_n(\mathbb{Z}_p) \\ \Delta_{K,\rho}(t) \ (\tau_{K,\rho}) & & I_{p,\rho}(T) \ (L_p(\rho, s)) \\ \text{describes the variation of} & & \text{describes the variation of} \\ H^1(G_K, \rho \otimes \chi), \ \chi \in \mathfrak{X}_K^1 & & \text{Sel}(G_p, \rho \otimes \rho_\varphi), \ \varphi \in \mathfrak{X}_p^1(\bar{\rho}) \end{array}$$

Here, $\text{Sel}(G_p, M)$ denotes the Selmer group for a G_p -module M (a subgroup of $H^1(G_p, M)$ with a local condition).

$n = 2$: The GL_2 -theory is concerned with hyperbolic geometry and Chern-Simons gauge theory in the knot side and Hida-Mazur theory and p -adic gauge theory in the prime side.

knot: We assume that K is a hyperbolic knot. Since G_K has a trivial center and any representation $G_K \rightarrow PGL_2(\mathbb{C}) = PSL_2(\mathbb{C})$ can be lifted to $G_K \rightarrow SL_2(\mathbb{C})$, we may consider only $SL_2(\mathbb{C})$ -representations without losing generality. So, we set

$$\mathfrak{X}_K^2 := \text{Hom}_{\text{gr}}(G_K, SL_2(\mathbb{C})) // SL_2(\mathbb{C}).$$

Note that the restriction of $[\rho] \in \mathfrak{X}_K^2$ to D_K is conjugate to an upper triangular representation:

$$\rho|_{D_K} \simeq \begin{pmatrix} \chi_\rho & * \\ 0 & \chi_\rho^{-1} \end{pmatrix}.$$

Let ρ° be a lift of the holonomy representation associated to the hyperbolic structure on $S^3 \setminus K$ and let $\mathfrak{X}_K^{2,\circ}$ be the irreducible component of \mathfrak{X}_K^2 containing $[\rho^\circ]$. The following theorem was shown by W. Thurston:

Theorem 2 ([T]). *The map $\Phi_K : \mathfrak{X}_K^{2,\circ} \rightarrow \mathbb{C}$ defined by $\Phi_K([\rho]) := \text{tr}(\rho(m_K))$ is bianalytic in a neighborhood of $[\rho^\circ]$. In particular, $\mathfrak{X}_K^{2,\circ}$ is a complex algebraic curve.*

Let m and l be functions on $\mathfrak{X}_K^{2,\circ}$ defined by $m(\rho) = \chi_\rho(m_K)$ and $\chi_\rho(l_K)$ respectively.

Theorem 3 ([NZ]). *Let $x := \log m(\rho)$ ($\log m(\rho^\circ) = 0$) and suppose $\partial V_K = \mathbb{C}^\times / q^{\mathbb{Z}}$. Then we have*

$$\left. \frac{dl}{dx} \right|_{x=0} = \frac{1}{2} \frac{\log q}{2\pi\sqrt{-1}}.$$

prime: We assume that $\bar{\rho}$ is a mod p representation associated to an ordinary modular elliptic curve E over \mathbb{Q} which corresponds to an ordinary Hecke eigenform f of weight 2: $\bar{\rho} = \rho_E \bmod p = \rho_f \bmod p$, $\rho_E = \rho_f : G_p \rightarrow GL_2(\mathbb{Z}_p)$. Here a representation $\rho : G_p \rightarrow GL_2(A)$ is called ordinary if the restriction of ρ to D_p is conjugate to an upper triangular representation:

$$\rho|_{D_p} \simeq \begin{pmatrix} \chi_{\rho,1} & * \\ 0 & \chi_{\rho,2} \end{pmatrix}, \quad \chi_{\rho,1}|_{I_p} = 1.$$

Compared with the knot case, it is natural to impose the ordinary condition to deformations of $\bar{\rho}$.

We have the following fundamental:

Theorem 4 (1) ([H1,2]). *There is a universal ordinary modular deformation $(R_p^{2,o,m}, \rho_p^{2,o,m})$ ($R_p^{2,o,m}$ is called the p -adic Hecke-Hida ring) of $\bar{\rho}$ such that any ordinary modular deformation (R, ρ) of $\bar{\rho}$ is obtained via a \mathbb{Z}_p -algebra homomorphism $R_p^{2,o,m} \rightarrow R$.*

(2) ([Ma]). *There is a universal ordinary deformation $(R_p^{2,o}, \rho_p^{2,o})$ of $\bar{\rho}$ such that any ordinary deformation (R, ρ) of $\bar{\rho}$ is obtained via a \mathbb{Z}_p -algebra homomorphism $R_p^{2,o} \rightarrow R$.*

By the universality of $(R_p^{2,o}, \rho_p^{2,o})$, we have a \mathbb{Z}_p -algebra homomorphism

$$R_p^{2,o} \rightarrow R_p^{2,o,m}$$

which we assume in the following to be an isomorphism. In fact, this assumption is satisfied under a mild condition owing to the works of A. Wiles etc.

We then define the universal ordinary deformation space of $\bar{\rho}$ by

$$\mathfrak{X}_p^{2,o}(\bar{\rho}) := \text{Hom}_{\mathbb{Z}_p\text{-alg}}(R_p^{2,o}, \mathbb{C}_p)$$

which may be regarded as an (infinitesimal) analog of $\mathfrak{X}_K^{2,o}$. As an analogue of Theorem 2, we have:

Theorem 5. *Take an element $\gamma \in I_p$ which is mapped to a generator of $\text{Gal}(\mathbb{Q}^\infty/\mathbb{Q})$ where \mathbb{Q}^∞ is the unique \mathbb{Z}_p -extension of \mathbb{Q} . The map $\Phi_p : \mathfrak{X}_p^{2,o} \rightarrow \mathbb{C}_p$ defined by $\Phi_p(\varphi) := \text{tr}(\rho_\varphi(\gamma))$ is bianalytic in a neighborhood of φ_f where $\varphi_f \circ \rho_p^{2,o} = \rho_f$.*

Remark. The analogy between the structures of $\mathfrak{X}_K^{2,o}$ and $\mathfrak{X}_p^{2,o}(\rho)$ was first pointed out by Kazuhiro Fujiwara.

The following theorem by Greenberg and Stevens may be seen as an analog of Neumann-Zagier's theorem 3.

Theorem 6 ([GS]). *Suppose that E is split multiplicative at p . Write*

$$\rho_p^{2,o}|_{D_p} \simeq \begin{pmatrix} \chi_1 & * \\ 0 & \chi_2 \end{pmatrix}, \quad \chi_1|_{I_p} = 1$$

and set $a_p := \chi_1(\sigma_p) : \mathfrak{X}_p^{2,o} \rightarrow \mathbb{C}_p$, and let $E(\mathbb{C}_p) = \mathbb{C}_p^\times/q^{\mathbb{Z}}$. Then we have

$$\frac{da_p}{d\rho}|_{\rho=2} = -\frac{1}{2} \frac{\log_p(q)}{\text{ord}_p(q)} (= \text{Mazur-Tate-Teitelbaum's } \mathcal{L}\text{-invariant}).$$

Next, we discuss some analogies between invariants defined on $\mathfrak{X}_K^{2,o}$ and $\mathfrak{X}_p^{2,o}(\rho)$.

prime: A typical invariant on $\mathfrak{X}_p^{2,o}(\bar{\rho})$ is a p -adic modular L -function $L_p(\rho, s)$, $\rho \in \mathfrak{X}_p^{2,o}$, $s \in \overline{\mathbb{Z}_p}$ ([GS]). Geometrically, $L_p(\rho, s)$ is given as a section of a rigid analytic line bundle \mathfrak{L}_p of modular symbols:

$$\begin{array}{c} \mathfrak{L}_p \\ \downarrow \\ \mathfrak{X}_p^{2,o} \end{array} \quad L_p(\rho, s) \text{ is a section } (s \text{ fixed})$$

We note that the value $L_p(\rho, 0) = L_p(f_\rho, 0)$ at $s = 0$ (f_ρ being a modular form corresponding to ρ) is given by $r_p(\{u, v\})(\omega) \cdot c$, where $r_p : K_2(C_\rho) \rightarrow H_{DR}^1(C_\rho/\mathbb{Q}_p)$ (C_ρ being a modular curve) is the p -adic regulator.

knot: Take a small affine open $\mathfrak{X} \subset \mathfrak{X}_K^{2,o}$ containing ρ^o if necessary, and let

$$L_K(\rho) := -2\pi^2 \text{CS}(\rho) + \sqrt{-1} \text{Vol}(\rho)$$

be the $SL_2(\mathbb{C})$ Chern-Simons invariant. Our theorem is

Theorem 7. *There is a holomorphic line bundle \mathfrak{L}_K with holomorphic connection on \mathfrak{X} such that $L_K(\rho)$ is given by a flat section:*

$$\begin{array}{c} \mathfrak{L}_K \\ \downarrow \\ \mathfrak{X} \end{array} \quad L_K(\rho) \text{ is a flat section}$$

For the construction of \mathfrak{L}_K , we apply S. Bloch's geometric construction of a tame symbol ([Bl]). Let H be the 3×3 Heisenberg group:

$$H(R) := \left\{ \begin{pmatrix} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \mid a, b, c \in R \right\} \quad (R : \text{a commutative ring})$$

The complex manifold $P := H(\mathbb{Z}) \backslash H(\mathbb{C})$ is a principal \mathbb{C}^\times -bundle over $\mathbb{C}^\times \times \mathbb{C}^\times$ by the map

$$P \rightarrow \mathbb{C}^\times \times \mathbb{C}^\times; \quad \begin{pmatrix} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \mapsto (\exp(2\pi\sqrt{-1}a), \exp(2\pi\sqrt{-1}b))$$

and 1-form $\theta = dc - adb$ gives a connection on P . Let $T(l, m^2) : \mathfrak{X} \rightarrow \mathbb{C}^\times \times \mathbb{C}^\times$ be a holomorphic map defined by $T(l, m^2)(\rho) := (l(\rho), m^2(\rho))$ and define

$$\mathfrak{L}_K := T(l, m^2)^*(P, \theta).$$

Then the flat section is given by

$$S(\rho^\circ) + \log l(\rho^\circ) \int_{\rho^\circ}^{\rho} d \log m^2 + \int_{\rho^\circ}^{\rho} d \log l d \log m^2 = L_K(\rho).$$

To see why $L_K(\rho)$ is seen as an analog of $L_p(\rho, 0)$, we give a cohomological interpretation of the above construction. Let $r_\infty : K_2(\mathfrak{X}) \rightarrow H^1(\mathfrak{X}, \mathbb{R})$ be the Beilinson regulator. We consider the natural map $\iota : H_D^2(\mathfrak{X}, \mathbb{Z}(2)) \rightarrow H_D^2(\mathfrak{X}, \mathbb{R}(2)) \hookrightarrow H^1(\mathfrak{X}, \mathbb{R})$ where H_D^* stands for the Deligne cohomology ([Br]). It is known that $H^2(\mathfrak{X}, \mathbb{Z}(2))$ is interpreted as the group of isomorphism classes of holomorphic line bundle on \mathfrak{X} with holomorphic connection ([ibid]), and so \mathfrak{L}_K is regarded as an element of $H^2(\mathfrak{X}, \mathbb{Z}(2))$. Then we can show that

$$\iota(\mathfrak{L}_K) = r_\infty(\{l, m^2\})$$

which is reminiscent of the connection between $L_p(\rho, 0)$ and the p -adic regulator.

Remark. Kirk and Klassen ([KK]) also constructed a line bundle E_K over $\mathfrak{X}_K^{2,0}$ so that $\mathfrak{L}_K(\rho)$ is regarded as a section. Though we have not seen the connection between E_K and \mathfrak{L}_K yet, our construction using Deligne cohomology seems to be natural conceptually.

Now, compared with the prime side, we may expect that

there should be a 2-variable L -function $L_K(\rho, s)$, $\rho \in \mathfrak{X}$, $s \in \mathbb{C}$ such that $L_K(\rho)$ would be a dominant term (special value) of $L_K(\rho, s)$ at $s = 0$.

Here is a candidate for such a L -function. Let M_ρ be the hyperbolic deformation of $M = S^3 \setminus K$ with holonomy ρ . Then M_ρ is a spin manifold with $Spin(3) = SU(2)$ -principal bundle $Spin(M_\rho) \rightarrow M_\rho$. Let D_ρ be the corresponding Dirac operator acting on $C^\infty(Spin(M_\rho) \otimes (\mathbb{C}^2)_\rho)$ and we define the spectral zeta function by

$$L_K(\rho, s) := \sum_{\lambda} \pm (\pm \lambda)^s, \quad \pm = \text{sign}(\text{Re}(\lambda)), \quad \text{Re}(s) \gg 0$$

where λ 's run over eigenvalues of D_ρ . Note that D_ρ may not be self-adjoint (though its symbol is self-adjoint) and so λ may be imaginary. For a closed hyperbolic 3-manifold, Jones-Westbury ([JW],[Y]) showed that $L_K(\rho, s)$ is continued as a meromorphic function to \mathbb{C} and the equality

$$L_K(\rho) = 2\pi^2 L_K(\rho, 0).$$

It is desirable to extend this equality for a non-closed hyperbolic 3-manifold.

Finally, we note the following

Theorem 8 (cf. [MT]). $L_K(\rho)$ gives a variation of mixed Hodge structure (V, W_*, F^*) on \mathfrak{X} defined by:

$$V = \mathbb{Z}^3,$$

$$V = W_0 \supset W_{-1} = \mathbb{Z}v_2 \oplus \mathbb{Z}v_3 \supset W_{-2} = \mathbb{Z}v_3 \text{ where}$$

$$\begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} := \begin{pmatrix} 1 & \log l(\rho^\circ) & S(\rho^\circ) \\ 0 & 1 & \log m^2(\rho^\circ) \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \int_{\rho^\circ}^\rho d \log l & \int_{\rho^\circ}^\rho d \log l d \log m^2 \\ 0 & 1 & \int_{\rho^\circ}^\rho d \log m^2 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} e_1 \\ (2\pi\sqrt{-1})e_2 \\ (2\pi\sqrt{-1})^2 e_3 \end{pmatrix}.$$

Here $\{e_1, e_2, e_3\}$ is a standard basis of V , and $W_0/W_{-1} = \mathbb{Z}(0)$, $W_{-1}/W_{-2} = \mathbb{Z}(1)$, $W_{-2} = \mathbb{Z}(2)$.

$V = F^{-2} \supset F^{-1} = \mathbb{Z}e_1 \oplus \mathbb{Z}e_2 \supset F^0 = \mathbb{Z}e_1$ with $\nabla F^{i-1} \subset \Omega^1 \otimes F^i$ so that

$$\nabla v := dv - v \begin{pmatrix} 1 & d \log l & 0 \\ 0 & 1 & d \log m^2 \\ 0 & 0 & 1 \end{pmatrix}.$$

References

- [Bl] S. Bloch, The dilogarithm and extensions of Lie algebras, Algebraic K-theory, Evanston 1980 (Proc. Conf., Northwestern Univ., Evanston, Ill., 1980), pp. 1–23, Lecture Notes in Math., 854, Springer, Berlin-New York, 1981.
- [Br] J.-L. Brylinski, Loop spaces, characteristic classes and geometric quantization, Progress in Mathematics, 107, Birkhäuser Boston, Inc., Boston, MA, 1993.
- [GS] R. Greenberg, G. Stevens, p -adic L -functions and p -adic periods of modular forms, Invent. Math. 111 (1993), no. 2, 407–447.
- [H1] H. Hida, Iwasawa modules attached to congruences of cusp forms, Ann. Sci. École Norm. Sup. (4) 19, (1986), 231–273.
- [H2] H. Hida, Galois representations into $GL_2(\mathbb{Z}_p[[X]])$ attached to ordinary cusp forms, Invent. Math. 85, (1986), 545–613.
- [JW] J. Jones, B. Westbury, Algebraic K-theory, homology spheres, and the η -invariant, Topology 34 (1995), no. 4, 929–957.
- [KK] P. Kirk, E. Klassen, Chern-Simons invariants of 3-manifolds decomposed along tori and the circle bundle over the representation space of T^2 . Comm. Math. Phys. 153, (1993), no. 3, 521–557.
- [Ma] B. Mazur, Deforming Galois representations, In: Galois groups over \mathbb{Q} , Math. Sci. Res. Inst. Publ., 16, Springer, (1989), 385–437.
- [Mo] M. Morishita, Analogies between knots and primes, 3-manifolds and number rings, to appear in Sugaku Expositions, AMS. (Analogies between prime numbers and knots, (Japanese) Sugaku 58, (2006), no. 1, 40–63.)
- [MT] M. Morishita, Y. Terashima, Geometry of polysymbols, preprint, 2006.
- [NZ] W. Neumann, D. Zagier, Volumes of hyperbolic three-manifolds, Topology 24 (1985), no. 3, 307–332.
- [T] W. Thurston, The geometry and topology of 3-manifolds, Lect. Note, Princeton, 1977.
- [Y] T. Yoshida, The η -invariant of hyperbolic 3-manifolds, Invent. Math. 81, (1985), no. 3, 473–514.

M.M: Graduate School of Mathematics, Kyushu University, 6-10-1, Hakozaki, Higashi-ku, Fukuoka, 812-8581 Japan. e-mail: morisita@math.kyushu-u.ac.jp

Y.T: Department of Mathematics, Tokyo Institute of Technology, 2-12-1 Oh-okayama, Meguro-ku, Tokyo 152-8551, Japan. e-mail: tera@math.titech.ac.jp