

Title	Relative Intrinsic Distance and Hyperbolic Imbedding(HOLOMORPHIC MAPPINGS, DIOPHANTINE GEOMETRY and RELATED TOPICS : in Honor of Professor Shoshichi Kobayashi on his 60th Birthday)
Author(s)	Kobayashi, Shoshichi
Citation	数理解析研究所講究録 (1993), 819: 239-242
Issue Date	1993-01
URL	http://hdl.handle.net/2433/83144
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Relative Intrinsic Distance and Hyperbolic Imbedding

Shoshichi Kobayashi *

December 1, 1992

Let Y be a complex space and X a complex subspace with compact closure \bar{X} . Let d_X and d_Y denote the intrinsic pseudo-distances of X and Y , respectively, (see [3]). We say that X is *hyperbolically imbedded* in Y if, for every pair of distinct points p, q in the closure $\bar{X} \subset Y$, there exist neighborhoods U_p and U_q of p and q in Y such that $d_X(U_p \cap X, U_q \cap X) > 0$. (In applications, X is usually a relatively compact open domain in Y .) It is clear that a hyperbolically imbedded complex space X is hyperbolic. The condition of hyperbolic imbedding says that the distance $d_X(p_n, q_n)$ remains positive when two sequences $\{p_n\}$ and $\{q_n\}$ in X approach two distinct points p and q of the boundary $\partial X = \bar{X} - X$. The concept of hyperbolic imbedding was first introduced in Kobayashi [3] to obtain a generalization of the big Picard theorem. The term "hyperbolic imbedding" was first used by Kiernan [2].

We shall now introduce a pseudo-distance $d_{X,Y}$ on \bar{X} so that X is hyperbolically imbedded in Y if and only if $d_{X,Y}$ is a distance.

Let $\mathcal{F}_{X,Y}$ be the family of holomorphic maps $f: D \rightarrow Y$ such that $f^{-1}(X)$ is either empty or a singleton. Thus, $f \in \mathcal{F}_{X,Y}$ maps all of D , with the exception of possibly one point, into X . The exceptional point is of course mapped into \bar{X} .

We define a pseudo-distance $d_{X,Y}$ on \bar{X} in the same way as d_Y , but using only chains of holomorphic disks belonging to $\mathcal{F}_{X,Y}$:

$$(1) \quad d_{X,Y}(p, q) = \inf_{\alpha} l(\alpha), \quad p, q \in \bar{X},$$

*During the preparation of this paper the author was at Technische Universität Berlin, supported by the Alexander von Humboldt-Stiftung.

where the infimum is taken over all chains α of holomorphic disks from p to q which belong to $\mathcal{F}_{X,Y}$. If p or q is in the boundary of X , such a chain may not exist. In such a case, $d_{X,Y}(p,q)$ is defined to be ∞ . For example, if X is a convex bounded domain in \mathbb{C}^n , any holomorphic disk passing through a boundary point of X goes outside the closure \overline{X} , so that $d_{X,\mathbb{C}^n}(p,q) = \infty$ if p is a boundary point of X . On the other hand, if X is Zariski-open in Y , any pair of points p, q in $\overline{X} = Y$ can be joined by a chain of holomorphic disks belonging to $\mathcal{F}_{X,Y}$, so that $d_{X,Y}(p,q) < \infty$.

Since

$$\text{Hol}(D, X) \subset \mathcal{F}_{X,Y} \subset \text{Hol}(D, Y),$$

we have

$$(2) \quad d_Y \leq d_{X,Y} \leq d_X,$$

where the second inequality holds on X while the first is valid on \overline{X} .

For the punctured disk $D^* = D - \{0\}$, we have

$$(3) \quad d_{D^*,D} = d_D.$$

The inequality $d_{D^*,D} \geq d_D$ is a special case of (2). Using the identity map $\text{id}_D \in \mathcal{F}_{D^*,D}$ as a holomorphic disk joining two points of D yields the opposite inequality.

Let $X' \subset Y'$ be another pair of complex spaces with $\overline{X'}$ compact. If $f: Y \rightarrow Y'$ is a holomorphic map such that $f(X) \subset X'$, then

$$(4) \quad d_{X',Y'}(f(p), f(q)) \leq d_{X,Y}(p, q) \quad p, q \in \overline{X}.$$

We can also define the infinitesimal form $F_{X,Y}$ of $d_{X,Y}$ in the same way as the infinitesimal form F_Y of d_Y , again using $\mathcal{F}_{X,Y}$ instead of $\text{Hol}(D, Y)$.

Theorem. *A complex space X is hyperbolically imbedded in Y if and only if $d_{X,Y}(p, q) > 0$ for all pairs $p, q \in \overline{X}$, $p \neq q$.*

Proof. From $d_{X,Y} \leq d_X$ it follows that if $d_{X,Y}$ is a distance, then X is hyperbolically imbedded in Y .

Let E be any length function on Y . In order to prove the converse, it suffices to show that there is a positive constant c such that $cE \leq F_{X,Y}$ on \overline{X} . Suppose that there is no such constant. Then there exist a sequence of tangent vectors v_n of \overline{X} , a sequence of holomorphic maps $f_n \in \mathcal{F}_{X,Y}$ and a sequence of tangent vectors e_n of D with Poincaré length $\|e_n\| \searrow 0$ such that $f_n(e_n) = v_n$. Since D is homogeneous, we may assume that e_n is a vector at the origin of D .

In constructing $\{f_n\}$, instead of using the fixed disk D and varying vectors e_n , we can use varying disks D_{R_n} and a fixed tangent vector e at the origin with $R_n \nearrow \infty$. (We take e to be the vector d/dz at the origin of D , which has the Euclidean length 1. Let $|e_n|$ be the Euclidean length of e_n , and $R_n = 1/|e_n|$. Instead of $f_n(z)$ we use $f_n(|e_n|z)$.) Let $\mathcal{F}_{X,Y}^{R_n}$ be the family of holomorphic maps $f: D_{R_n} \rightarrow Y$ such that $f^{-1}(X)$ is either empty or a singleton. Having replaced D, e_n by D_{R_n}, e , we may assume that $f_n \in \mathcal{F}_{X,Y}^{R_n}$ and $f_n(e) = v_n$. We want to show that a suitable subsequence of $\{f_n\}$ converges to a nonconstant holomorphic map $f: \mathbb{C} \rightarrow \overline{X}$.

By applying Brody's lemma [1] to each f_n and a constant $0 < c < \frac{1}{4}$ we obtain holomorphic maps $g_n \in \text{Hol}(D_{R_n}, Y)$ such that

- (a) $g_n^* E^2 \leq c R_n^2 ds_{R_n}^2$ on D_{R_n} and the equality holds at the origin 0;
- (b) $\text{Image}(g_n) \subset \text{Image}(f_n)$.

Since g_n is of the form $g = f_n \circ \mu_{r_n} \circ h_n$, where h_n is an automorphism of D_{R_n} and μ_{r_n} , ($0 < \mu_{r_n} < 1$, is the multiplication by r_n , each g_n is also in $\mathcal{F}_{X,Y}$.

Now, as in the proof of Brody's theorem [1] we shall construct a nonconstant holomorphic map $h: \mathbb{C} \rightarrow Y$ to which a suitable subsequence of $\{g_n\}$ converges. In fact, since

$$g_n^* E^2 \leq c R_n^2 ds_{R_n}^2 \leq c R_m^2 ds_{R_m}^2 \quad \text{for } n \geq m,$$

the family $\mathcal{F}_m = \{g_n|_{D_{R_m}}, n \geq m\}$ is equicontinuous for each fixed m . Since the family $\mathcal{F}_1 = \{g_n|_{D_{R_1}}\}$ is equicontinuous, the Arzela-Ascoli theorem implies that we can extract a subsequence which converges to a map $h_1 \in \text{Hol}(D_{R_1}, Y)$. (We note that this is where we use the compactness of \overline{X} .) Applying the same theorem to the corresponding sequence in \mathcal{F}_2 , we extract a subsequence which converges to a map $h_2 \in \text{Hol}(D_{R_2}, Y)$. In this way we obtain maps $h_k \in \text{Hol}(D_{R_k}, Y)$, $k = 1, 2, \dots$ such that each h_k is an extension of h_{k-1} . Hence, we have a map $h \in \text{Hol}(\mathbb{C}, Y)$ which extends all h_k .

Since $g_n^* E^2$ at the origin 0 is equal to $(c R_n^2 ds_{R_n}^2)_{z=0} = 4cdz d\bar{z}$, it follows that

$$(h^* E^2)_{z=0} = \lim_{n \rightarrow \infty} (g_n^* E^2)_{z=0} = 4cdz d\bar{z} \neq 0,$$

which shows that h is nonconstant.

Since $g_n^* E^2 \leq c R_n^2 ds_{R_n}^2$, in the limit we have

$$h^* E^2 \leq 4cdz d\bar{z}.$$

By suitably normalizing h we obtain

$$h^* E^2 \leq dz d\bar{z} \quad \text{with the equality holding at } z = 0.$$

We may assume that $\{g_n\}$ itself converges to h . Since h is the limit of $\{g_n\}$, clearly $h(\mathbf{C}) \subset \overline{X}$. Let p, q be two points of $h(\mathbf{C})$, say $p = h(a)$ and $q = h(b)$. Taking a subsequence and suitable points a, b we may assume that $g_n(a), g_n(b) \in X$. Then $\lim g_n(0) = p$ and $\lim g_n(a) = q$ and

$$d_X(g_n(a), g_n(b)) \leq d_{D_{R_n}}(a, b) \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

contradicting the assumption that X is hyperbolically imbedded in Y . Q.E.D.

This relative distance $d_{X,Y}$ simplifies the proof of the big Picard theorem as formulated in [3].

Bibliography

1. R. Brody, Compact manifolds and hyperbolicity, *Trans. Amer. Math. Soc.* 235 (1978), 213-219.
2. P.J. Kiernan, Hyperbolically imbedded spaces and the big Picard theorem, *Math. Ann.* 204 (1973), 203-209.
3. S. Kobayashi, *Hyperbolic Manifolds and Holomorphic Mappings*, Marcel Dekker, New York, 1970.

Department of Mathematics
 University of California
 Berkeley, CA 94720, USA
 e-mail: kobayash@math.berkeley.edu