

Title	A Lower Bound of the Expected Maximum Number of Edge-disjoint s-t Paths on Probabilistic Graphs
Author(s)	CHENG, Peng; MASUYAMA, Shigeru
Citation	数理解析研究所講究録 (1993), 833: 80-90
Issue Date	1993-04
URL	<a href="http://hdl.handle.net/2433/83414">http://hdl.handle.net/2433/83414</a>
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

# A Lower Bound of the Expected Maximum Number of Edge-disjoint s-t Paths on Probabilistic Graphs

程 鵬                      増山 繁  
*Peng CHENG*              *Shigeru MASUYAMA*

豊橋技術科学大学知識情報工学系  
Department of Knowledge-Based Information Engineering,  
Toyohashi University of Technology,  
Toyohashi-shi 441, Japan

## Abstract

For a probabilistic graph  $(G = (V, E, s, t), p)$ , where  $G$  is an undirected graph with specified source vertex  $s$  and sink vertex  $t$  ( $s \neq t$ ) in which each edge has independent failure probability and each vertex is assumed to be failure-free, and  $p = (p(e_1), \dots, p(e_{|E|}))$  is a vector consisting of failure probabilities  $p(e_i)$ 's of all edges  $e_i$ 's in  $E$ , we consider the problem of computing the expected maximum number  $\Gamma_{(G,p)}$  of edge-disjoint s-t paths. It has been known that this computing problem is NP-hard even if  $G$  is restricted to several classes like planar graphs, s-t out-in bitrees and s-t complete multi-stage graphs. In this paper, for a probabilistic graph  $(G = (V, E, s, t), p)$ , we propose a lower bound of  $\Gamma_{(G,p)}$  and show the necessary and sufficient conditions by which the lower bound coincides with  $\Gamma_{(G,p)}$ . Furthermore, we also give a method of computing the lower bound of  $\Gamma_{(G,p)}$  for a probabilistic graph  $(G = (V, E, s, t), p)$ .

## 1 Introduction

We consider a probabilistic graph  $(G = (V, E, s, t), p)$ , where  $G$  is an undirected graph with specified source vertex  $s$  and sink vertex  $t$  ( $s \neq t$ ) in which each edge has independent failure probability and each vertex is assumed to be failure-free, and  $p = (p(e_1), \dots, p(e_{|E|}))$  is a vector consisting of failure probabilities  $p(e_i)$ 's of all edges  $e_i$ 's in  $E$ . The expected maximum number  $\Gamma_{(G,p)}$  of edge-disjoint s-t paths (namely, s-t paths having no edge in common) in a probabilistic graph  $(G, p)$  is useful for network reliability analysis. Note that the problem of computing s, t-connectedness [1,3], namely, probability that there exists at least one operative s-t path, is a special case of computing  $\Gamma_{(G,p)}$  in a probabilistic graph  $(G, p)$ .

However, it is known that the problem of computing  $\Gamma_{(G,p)}$  in a probabilistic graph  $(G, p)$  is NP-hard, even if  $G$  is restricted to several classes, e.g., planar graphs, s-t out-in bitrees and s-t complete multi-stage graphs [2]. Thus, for estimating  $\Gamma_{(G,p)}$ , it is interesting for us to find its lower bound in a probabilistic graph  $(G, p)$ .

In this paper, we define a lower bound of  $\Gamma_{(G,p)}$  using an s-t path number function of  $G$  for a probabilistic graph  $(G, p)$ , and give the necessary and sufficient conditions by which this lower bound coincides with  $\Gamma_{(G,p)}$  and a method of computing this lower bound. This paper is organized as follows:

Graph theoretic terminologies used throughout this paper are described in section 2. A lower bound of  $\Gamma_{(G,p)}$  in a probabilistic graph  $(G, p)$  is defined in section 3. Section 4 shows the necessary and sufficient conditions by which this lower bound coincides with  $\Gamma_{(G,p)}$ . Furthermore, we suggest a method of computing the lower bound in section 5.

## 2 Preliminaries

### 2.1 Graph Theoretic Terminologies

A two-terminal undirected graph  $G = (V, E, s, t)$  consists of a finite vertex set  $V$  and a set  $E$  of pairs of vertices, called edges, where  $s$  and  $t$ , called *source* and *sink*, respectively, are two specified distinct vertices of  $V$ . For an edge  $(u, v)$ , the two vertices  $u$  and  $v$  are said to be end vertices of  $(u, v)$ , and  $(u, v)$  is said to be incident to  $u$  and  $v$ .

In  $G = (V, E, s, t)$ , an  $x$ - $y$  path  $\pi$  of length  $k$  from vertex  $x$  to vertex  $y$  is an alternating sequence of vertices  $v_i \in V$  ( $0 \leq i \leq k$ ) and edges  $(v_{i-1}, v_i) \in E$  ( $1 \leq i \leq k$ ),

$$\pi : (x =)v_0, (v_0, v_1), v_1, \dots, v_{k-1}, (v_{k-1}, v_k), v_k(= y),$$

where vertices  $v_i$ 's ( $0 \leq i \leq k$ ) are distinct. i.e., a path denotes a simple path throughout this paper. For short, we also denote an  $x$ - $y$  path  $\pi$  by

$$\pi : (x =)v_0, v_1, \dots, v_{k-1}, v_k(= y).$$

The vertices  $v_1, \dots, v_{k-1}$  are called its internal vertices and the vertices  $v_0(= s), v_k(= t)$  are called its end vertices. Let  $V(\pi), E(\pi)$  denote the set of all vertices and the set of all edges on an  $x$ - $y$  path  $\pi$ , respectively. The set of all  $x$ - $y$  paths in  $G$  is denoted by  $P_{xy}(G)$ . Paths  $\pi_1, \dots, \pi_r$  are called *internal vertex-disjoint paths* if they have no vertex in common except their end vertices.  $s$ - $t$  paths  $\pi_1, \dots, \pi_r$  are called *edge-disjoint  $s$ - $t$  paths* if any two of them have no edge in common, and the maximum number of edge-disjoint  $s$ - $t$  paths in  $G$  is denoted by  $\lambda_{st}(G)$ .

A graph  $G_1 = (V_1, E_1)$  is a subgraph of  $G = (V, E, s, t)$ , if  $V_1 \subseteq V$  and  $E_1 \subseteq E$  hold. If  $G_1$  is a subgraph of  $G$ , other than  $G$  itself, then  $G_1$  is a proper subgraph of  $G$ . For a subset  $E' \subseteq E$ , the subgraph derived from  $G$  by deleting all edges of  $E'$  is denoted by  $G - E' (= (V, E - E', s, t))$ . A subset  $E' (\subseteq E)$  is called an  *$s$ - $t$  edge-cutset* if  $G - E'$  has no  $s$ - $t$  path. An  $s$ - $t$  path  $\pi$  is an  *$s$ - $t$  edge-cut-path* if  $E(\pi)$  is an  $s$ - $t$  edge-cutset. An  $s$ - $t$  edge-cutset with the minimum cardinality among  $s$ - $t$  edge-cutsets of  $G$  is said to be *minimum*. By well-known Menger's theorem [4],  $\lambda_{st}(G)$  is equal to the cardinality of a minimum  $s$ - $t$  edge-cutset of  $G$  for any  $G$ .

### 2.2 Probabilistic Graph

A probabilistic graph, denoted by  $(G = (V, E, s, t), p)$ , or  $(G, p)$ , for short, is defined as follows:

- (i)  $G = (V, E, s, t)$  is a two-terminal graph, where each edge  $e$  of  $E$  is in either of the following two states: failed or operative (not failed), having known independent failure probability  $p(e)$ ,  $0 \leq p(e) \leq 1$  (or operative probability  $q(e) = 1 - p(e)$ ), and each vertex is assumed to be failure-free.
- (ii)  $p$  is a vector consisting of all edge failure probabilities  $p(e)$ 's in  $E$ .

For a probabilistic graph  $(G = (V, E, s, t), p)$ , let a subgraph  $G - U (\subseteq E)$  correspond to an event  $\mathcal{E}_U$  that all edges of  $U$  are failed and all edges of  $E - U$  are operative. Clearly, the probability  $\rho(G - U)$  of arising a subgraph  $G - U (\subseteq E)$  is computed by the following formula.

$$\rho(G - U) = \prod_{e \in U} p(e) \prod_{e \in E - U} q(e) (= 1 - p(e)).$$

Furthermore,  $\sum_{U \subseteq E} \rho(G - U) = 1$  holds.

Now, we define the *expected maximum number*  $\Gamma_{(G,p)}$  of edge-disjoint  $s$ - $t$  paths in a probabilistic graph  $(G = (V, E, s, t), p)$  as follows:

$$\Gamma_{(G,p)} \equiv \sum_{U \subseteq E} \lambda_{st}(G - U) \rho(G - U). \quad (1)$$

It is known that the problem of computing  $\Gamma_{(G,p)}$  for a probabilistic graph  $(G,p)$  is NP-hard, even if  $G$  is restricted to several special classes like planar graphs, s-t out-in bitrees and s-t multi-stage complete graphs, etc. [2]. Thus, it is interesting for us to consider a lower bound of  $\Gamma_{(G,p)}$  for estimating it.

### 3 A Lower Bound of $\Gamma_{(G,p)}$

We define a lower bound of the expected maximum number of edge-disjoint s-t paths in a probabilistic graph.

An *s-t path number function*  $f$  of  $G = (V, E, s, t)$  is a one-to-one integral function  $f : P_{st}(G) \mapsto \{1, \dots, l\}$ . The s-t path  $\pi$  with  $f(\pi) = k$  is said to be the *s-t path of number k*, and denoted by  $\pi_k$ . The s-t path with the minimum number in  $G - E' (\subseteq E)$  with respect to  $f$  is denoted by  $\pi_{m(G-E',f)}$ .

First, we give the following procedure **FEDP** to find edge-disjoint s-t paths in  $G = (V, E, s, t)$ .

#### Procedure **FEDP**

**Input** A graph  $G = (V, E, s, t)$  and an s-t path number function  $f$  of  $G$ .

**Output** The set of edge-disjoint s-t paths  $FEDP(G, f)$ .

BEGIN

$G' := G; FEDP(G, f) := \phi;$

WHILE  $P_{st}(G') \neq \phi$  DO

BEGIN

Find  $\pi_{m(G',f)}$  from  $P_{st}(G')$ ;

$FEDP(G, f) := FEDP(G, f) \cup \{\pi_{m(G',f)}\};$

$G' := G' - E(\pi_{m(G',f)})$

END;

Output  $FEDP(G, f)$

END. □

It is clear that  $FEDP(G, f)$  obtained by **FEDP** is a set of edge-disjoint s-t paths in  $G$ . Namely, the following formula holds.

$$|FEDP(G, f)| \leq \kappa_{st}(G), \text{ for any } G, f. \quad (2)$$

For a probabilistic graph  $(G = (V, E, s, t), p)$  and an s-t path number function  $f$  of  $G$ , we now define the value  $\underline{\Gamma}_{(G,f,p)}$  as follows:

$$\underline{\Gamma}_{(G,f,p)} \equiv \sum_{U \subseteq E} |FEDP(G - U, f)| \rho(G - U). \quad (3)$$

By formulas (1),(2),(3),  $\underline{\Gamma}_{(G,f,p)}$  is a lower bound of  $\Gamma_{(G,p)}$ , namely, the following formula holds.

$$\underline{\Gamma}_{(G,f,p)} \leq \Gamma_{(G,p)}, \text{ for any } G, f, p.$$

### 4 Necessary and Sufficient Conditions

In this section, we give the necessary and sufficient conditions by which  $\underline{\Gamma}_{(G,f,p)}$  coincides with  $\Gamma_{(G,p)}$  in a probabilistic graph  $(G, p)$ .

#### 4.1 A Necessary and Sufficient Condition of an s-t Path Number Function

By formulas (1),(2),(3), the following Theorem 4.1 immediately holds.

**Theorem 4.1.** Given  $(G = (V, E, s, t), p)$ , then  $\Gamma_{(G,f,p)} = \Gamma_{(G,p)}$  holds iff  $G$  has an s-t path number function  $f$  satisfying the following formula.

$$|FEDP(G - U, f)| = \lambda_{s,t}(G - U), \text{ for any } U \subseteq E. \quad (4)$$

□

**Definition 4.1.** An s-t path number function  $f$  of  $G$  is called *exact* if  $f$  satisfies formula (4). □

A graph  $G = (V, E, s, t)$  is said to be *s-t k-edge-connected* if  $\lambda_{s,t}(G) = k$  holds. A graph  $G$  is said to be  *$\pi$ -edge-cut* if  $\pi$  is an s-t edge-cut-path in  $G$ . A graph  $G$  is said to be  *$\pi$ -edge-cut s-t 2-edge-connected* if  $\pi$  is an s-t edge-cut-path of  $G$  and  $G$  is s-t 2-edge-connected. A  $\pi$ -edge-cut s-t 2-edge-connected graph  $G = (V, E, s, t)$  is *minimal*, if  $G - \{e\}$  for any  $e \in E - E(\pi)$  is not  $\pi$ -edge-cut s-t 2-edge-connected. For example, the graph  $G$  shown in Fig.1 is a  $\pi$ -edge-cut s-t 2-edge-connected graph, where  $\pi : v_0(= s), v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8, v_9(= t)$ . But it is not minimal as  $G - \{e\}$  is  $\pi$ -edge-cut s-t 2-edge-connected. Furthermore, the set of all  $\pi$ -edge-cut s-t 2-edge-connected subgraphs of an s-t path  $\pi$  of  $G$  is denoted by  $\mathcal{W}(G, \pi)$ . For example, in the graph  $G$  given in Fig.1,  $\mathcal{W}(G, \pi) = \{G - \{e = (u_1, u_2)\}, G - \{(u_1, v_4), (u_2, v_5), (v_3, v_5)\}\}$ . Clearly, the following Lemma 4.1 holds.




Fig.1 A  $\pi$ -edge-cut s-t 2-edge-connected graph.

**Lemma 4.1.** If  $\lambda_{s,t}(G) \geq 2$  holds and an s-t path  $\pi$  of  $G$  is an s-t edge-cut-path, then  $\mathcal{W}(G, \pi) \neq \phi$  holds. □

**Lemma 4.2.** In a graph  $G = (V, E, s, t)$ , if there exists an s-t path  $\pi$  satisfying  $\mathcal{W}(G, \pi) = \phi$ , then the following formula holds.

$$\lambda_{s,t}(G - E(\pi)) = \lambda_{s,t}(G) - 1.$$

**Proof.** Clearly,  $\lambda_{s,t}(G - E(\pi)) \leq \lambda_{s,t}(G) - 1$  holds. Assume that  $\lambda_{s,t}(G - E(\pi)) < \lambda_{s,t}(G) - 1$  holds. By this assumption, there exists a minimum s-t edge-cutset  $E^*$  in  $G - E(\pi)$  that satisfies  $|E^*| \leq \lambda_{s,t}(G) - 2$  by Menger's Theorem [4]. Consider graph  $G - E^*$ , and it is clear that all s-t paths in  $G - E^*$  share at least one edge of  $E(\pi)$ , i.e.,  $\pi$  is an s-t edge-cut-path of  $G - E^*$ . Furthermore, let  $E'$  be a minimum s-t edge-cutset of  $G - E^*$ . As  $E' \cup E^*$  is an s-t edge-cutset of  $G$ ,  $|E' \cup E^*| = |E'| + |E^*| \geq \lambda_{s,t}(G)$  holds. By  $|E^*| \leq \lambda_{s,t}(G) - 2$ , we obtain  $|E'| = \lambda_{s,t}(G - E^*) \geq 2$ , contradicting the fact that  $\mathcal{W}(G, \pi) \neq \phi$  holds by Lemma 4.1. □

We now prove the following Theorem 4.2.

**Theorem 4.2.** In a graph  $G = (V, E, s, t)$ , an s-t path number function  $f$  of  $G$  is exact iff for any  $U \subseteq E$  with  $P_{st}(G - U) \neq \phi$ ,  $\mathcal{W}(G - U, \pi_{m(G-U, f)}) = \phi$  holds.

**Proof.** Necessity: Assume that an s-t path number function  $f$  of  $G$  is exact and that for some  $U \subseteq E$  with  $P_{st}(G - U) \neq \phi$ ,  $\mathcal{W}(G - U, \pi_{m(G-U, f)}) \neq \phi$  holds. By  $\mathcal{W}(G - U, \pi_{m(G-U, f)}) \neq \phi$ ,  $G - U$  has a subgraph  $G' \in \mathcal{W}(G - U, \pi_{m(G-U, f)})$ .  $\lambda_{st}(G') = 2$  holds by the definition of  $\mathcal{W}(G - U, \pi_{m(G-U, f)})$ . As  $\pi_{m(G-U, f)}$  is the s-t path with the minimum number of  $G'$  and an s-t edge-cut-path of  $G'$ , we have  $FEDP(G', f) = \{\pi_{m(G-U, f)}\}$  by **FEDP**. Hence,  $|FEDP(G', f)| (= 1) < \lambda_{st}(G') (= 2)$  holds, contradicting the fact that  $f$  is exact.

Sufficiency: Assume that for any  $U \subseteq E$  with  $P_{st}(G - U) \neq \phi$ ,  $\mathcal{W}(G - U, \pi_{m(G-U, f)}) = \phi$  holds. Then it is easy to prove that for any  $U \subseteq E$ ,  $|FEDP(G - U, f)| = \lambda_{st}(G - U)$  holds by iteratively applying Lemma 4.2.  $\square$

## 4.2 A Necessary and Sufficient Condition of s-t Paths

**Definition 4.2.** (*Prohibitive s-t Path Set*)

Let  $P (\subseteq P_{st}(G))$  be a subset of the set of all s-t paths of  $G$ . If, for each s-t path  $\pi$  of  $P$ , there is a  $\pi$ -edge-cut s-t 2-edge-connected subgraph  $G_\pi \in \mathcal{W}(G, \pi)$  in  $G$  that satisfies  $P_{st}(G_\pi) \subseteq P$ , then  $P$  is called a *prohibitive s-t path set*.  $\square$

**Procedure TEST**

Input: A graph  $G = (V, E, s, t)$ .

Output: Either an s-t path number function  $f$  of  $G$  or a subset  $P$  of  $P_{st}(G)$ .

BEGIN

$P := P_{st}(G); i := 1; Q := \{ \pi \in P_{st}(G) \mid \mathcal{W}(G, \pi) = \phi \};$

WHILE  $Q \neq \phi$  DO

BEGIN

$P := P - Q;$

REPEAT

Select an s-t path  $\pi$  from  $Q;$

$f(\pi) := i; i := i + 1; Q := Q - \{\pi\}$

UNTIL  $Q = \phi;$

$Q := \{ \pi \in P \mid P_{st}(G_\pi) \not\subseteq P, \text{ for all } G_\pi \in \mathcal{W}(G, \pi) \}$

END;

IF  $P = \phi$  THEN output  $f$  ELSE output  $P$

END.  $\square$

Clearly, the following Lemma 4.3 holds by Definitions 4.1 and 4.2.

**Lemma 4.3.** If **TEST** outputs an s-t path number function  $f$  of  $G$ , then  $f$  is exact, when a graph  $G = (V, E, s, t)$  is input. If **TEST** outputs a subset  $P$  of  $P_{st}(G)$ , then  $P$  is a prohibitive s-t path set, when a graph  $G = (V, E, s, t)$  is input.  $\square$

If there is a prohibitive s-t path set  $P (\subseteq P_{st}(G))$  where  $G = (V, E, s, t)$ , then there does not exist any exact s-t path number function  $f$ . Otherwise, if  $G$  has an exact s-t path number function  $f$ , and suppose  $\pi_m$  be the s-t path of the minimum number with respect to  $f$  among  $P$ . By Definition 4.2,

there is  $G_{\pi_m} \in \mathcal{W}(G, \pi_m)$  in  $G$  that satisfies  $P_{st}(G_{\pi_m}) \subseteq P$ . Thus,  $\pi_m$  is also the  $s$ - $t$  path of the minimum number with respect to  $f$  in  $G_{\pi_m}$ . Therefore, by FEDP,  $FEDP(G_{\pi_m}, f) = 1 < \lambda_{st}(G_{\pi_m}) = 2$  holds. This leads to a contradiction that  $f$  is an exact  $s$ - $t$  path number function of  $G$ . Hence, by Theorem 4.2 and Lemma 4.3, the following Theorem 4.3 holds.

**Theorem 4.3.** In a graph  $G = (V, E, s, t)$ ,  $G$  has an exact  $s$ - $t$  path number function iff it contains no prohibitive  $s$ - $t$  path set as its  $s$ - $t$  path subset.  $\square$

### 4.3 Characterization of Graph Having a Prohibitive $s$ - $t$ Path Set

A graph is connected if there is a path connecting each pair of vertices and otherwise disconnected. A connected component of  $G$  is a maximal connected subgraph, which is simply called a component. If there exist vertices  $x$  and  $y$ ,  $x \neq v$  and  $y \neq v$  such that all the paths connecting  $x$  and  $y$  have  $v$  as an internal vertex, then  $v$  is an *articulation vertex*. A two-terminal connected graph is said to be  *$s, t$  non-separable* if its subgraph obtained by removing  $s, t$  is connected. In the following discussion, we assume that  $G$  is an  $s, t$  non-separable two-terminal connected graph, unless otherwise specified.

**Definition 4.3.** ( *$s$ - $t$  2-edge-connected Articulation Vertex*)

A vertex  $v$  is said to be an  *$s$ - $t$  2-edge-connected articulation vertex* of  $G$ , if  $v$  is an  $s$ - $t$  articulation vertex of  $G$  and there exist both two edge-disjoint  $s$ - $v$  paths and two edge-disjoint  $v$ - $t$  paths in  $G$ .  $\square$

For example, in the graph illustrated in Fig.2(a), vertices  $u, v, w$  are  $s$ - $t$  2-edge-connected articulation vertices of  $G$ .




Fig.2 An illustration of separation of  $G$  at an  $s$ - $t$  2-edge-connected articulation vertex.

**Definition 4.4.** (*Separation of  $G$  at an  $s$ - $t$  2-edge-connected Articulation Vertex*)

Assume that  $G$  has an  $s$ - $t$  2-edge-connected articulation vertex  $v$ . The following sequence of operations is said to be *separation of  $G$  at an  $s$ - $t$  2-edge-connected articulation vertex  $v$* .

- (i) The two components  $C_1$  and  $C_2$  are obtained by removing  $v$  from  $G$ .
- (ii)  $v$  is connected to  $C_1$  (or  $C_2$ ) with all edges  $(u, v)$ 's of  $G$  having one end vertex  $u$  in  $C_1$  (or  $C_2$ ).
- (iii) Note that  $C_1$  contains either of  $s, t$ . If  $C_1$  contains  $s$  (or  $t$ ) then let  $s$  (or  $t$ ) be  $s_1$  (or  $t_1$ ) and let  $v$  be  $t_1$  (or  $s_1$ ).  $s_2$  and  $t_2$  are similarly defined for  $C_2$ .  $\square$

For example, the two graphs illustrated in Fig.2(b),(c) are obtained by separation of the graph given in Fig.2(a) at an  $s$ - $t$  2-edge-connected articulation vertex  $v$ .

**Definition 4.5. (Prohibitive Graph)**

A graph  $G$  is said to be a *prohibitive graph*, if  $G$ , or one of the graphs derived from  $G$  by separations of  $G$  at all  $s$ - $t$  2-edge-connected articulation vertices in  $G$  is homeomorphic to the graph shown in Fig.3.  $\square$

The two graphs illustrated in Fig.2(a),(b) are both prohibitive graphs. But the graph given in Fig.2(d), although it contains a subgraph homeomorphic to the graph shown in Fig.3, is not a prohibitive graph as the vertex  $u$  is not its  $s$ - $t$  2-edge-connected articulation vertex and it is not homeomorphic to the graph shown in Fig.3. It is easy to verify that for a prohibitive graph  $G$ ,  $P_{s,t}(G)$  is a prohibitive  $s$ - $t$  path set. Thus, we immediately obtain the following Lemma 4.4.




Fig.3 A prohibitive graph.

**Lemma 4.4.** If  $G$  contains a prohibitive graph as its subgraph, then it also has a prohibitive  $s$ - $t$  path set as its  $s$ - $t$  path subset.  $\square$

Now, we show that if  $G$  has a prohibitive  $s$ - $t$  path set as its  $s$ - $t$  path subset, then it contains a prohibitive graph as its subgraph. For our aim, we need more definitions.

**Definition 4.6. (Attachment Vertex [5],[6])**

An *attachment vertex* of a subgraph  $G_1$  in  $G$  is a vertex of  $G_1$  incident in  $G$  with some edge not belonging to  $G_1$ .  $\square$

**Definition 4.7. (Bridges [5],[6])**

Let  $J$  be a fixed subgraph of  $G$ . A subgraph  $G_1$  of  $G$  is said to be  *$J$ -detached* in  $G$  if all its attachment vertices are in  $J$ . We define a *bridge* of  $J$  in  $G$  as any subgraph  $B$  that satisfies the following three conditions:

- (i)  $B$  is not a subgraph of  $J$ .
- (ii)  $B$  is  $J$ -detached in  $G$ .
- (iii) No proper subgraph of  $B$  satisfies both (i) and (ii).  $\square$

**Definition 4.8. (Degenerate and Proper Bridges. Nucleus of a Bridge [5],[6])**

An edge  $e = (u, v)$  of  $G$  not belonging to  $J$  but having both end vertices in  $J$  is referred to as a *degenerate bridge*.

Let  $G^-$  be the graph derived from  $G$  by deleting the vertices of  $J$  and all edges incident to them.



Let  $C$  be any component of  $G^-$ . Let  $B$  be the subgraph of  $G$  obtained from  $C$  by adjoining to it each edge of  $G$  having one end vertex in  $C$  and the other end vertex in  $J$  and adjoining also the end vertices in  $J$  of all such edges. The subgraph  $B$  satisfies the conditions (i),(ii),(iii) in Definition 4.7 and is a bridge. Such a bridge is called to be *proper*. The component  $C$  of  $G^-$  is the *nucleus* of  $B$ .  $\square$

For the graph  $G$  shown in Fig.4, let  $J$  be an  $s$ - $t$  path  $\pi : v_0(=s), v_1, v_2, v_3, v_4, v_5, v_6(=t)$ , then all vertices on  $\pi$  other than  $v_4$  are all attachment vertices of  $\pi$  in  $G$ .  $B_1, B_2, B_3$  are proper bridges of  $\pi$  in  $G$  and  $B_4$  is a degenerate bridge of  $\pi$  in  $G$ . By Definitions 4.6,4.7, the following Lemma 4.5 obviously holds.




Fig.4 An illustration of attachment vertices, bridges and nuclei.

**Lemma 4.5.** Let  $\pi$  be an  $s$ - $t$  path of  $G$ . If there is a proper bridge  $B$  of  $\pi$  in  $G$ , then any two vertices  $u, v$  in  $B$  are connected by a path consisting of edges and vertices only in the nucleus of  $B$ .  $\square$

Let  $\gamma : v_0, v_1, \dots, v_{k-1}, v_k$  be a path from  $v_0$  to  $v_k$  of  $G$ . If  $0 \leq i < j \leq k$ , then the sequence  $v_i, v_{i+1}, \dots, v_{j-1}, v_j$  is a subpath of  $\gamma$ , and denoted by  $\gamma[v_i, v_j]$ .

**Definition 4.9.** (*Path Avoiding s-t Path  $\pi$* )

Let  $\pi$  be an  $s$ - $t$  path of  $G$ . For two vertices  $v_i, v_j$  in  $V(\pi)$ , a path between  $v_i$  and  $v_j$  consisting of edges not in  $E(\pi)$  and vertices not in  $V(\pi)$  except  $v_i, v_j$  is said to be *avoiding  $\pi$* .  $\square$

For example, the path  $v_1, u_1, u_2, v_5$  is avoiding the  $s$ - $t$  path  $\pi$  in the graph  $G$  illustrated in Fig.1.

**Definition 4.10.** (*Order Relation with Respect to an s-t Path  $\pi$* )

Let  $\pi : v_0(=s), v_1, \dots, v_{k-1}, v_k(=t)$  be an  $s$ - $t$  path of  $G$ . We define an *order relation*  $<_{\pi}$  on  $V(\pi)$  with respect to  $\pi$  as follows: For any  $v_i, v_j$  ( $0 \leq i, j \leq k$ ),  $v_i <_{\pi} v_j$  holds iff  $i < j$  holds. If  $v_i <_{\pi} v_j$ ,  $v_i$  ( $v_j$ ) is said to be to the left (right) of  $v_j$  ( $v_i$ ).  $\square$

**Definition 4.11.** (*Intersection Vertex of Two Paths  $\pi, \alpha$* )

Let  $\pi, \alpha$  be two paths of  $G$ . A vertex  $v$  is called an *intersection vertex* of  $\pi, \alpha$  if  $\pi$  and  $\alpha$  have at least three distinct edges incident to  $v$ . The set of all intersection vertices of  $\pi, \alpha$  is denoted by  $V_{\pi\alpha}$ .  $\square$

In the graph  $G$  given in Fig.1, for two  $s$ - $t$  paths  $\pi$  and  $\alpha : v_0(=s), v_1, u_1, u_2, v_6, v_7, v_9(=t)$ , we have  $V_{\pi\alpha} = \{v_1, v_6, v_7, v_9\}$ .

**Definition 4.12.** (*Interlacing Subpaths*)

Suppose that  $G$  has an s-t path  $\pi : v_0(= s), v_1, \dots, v_{k-1}, v_k(= t)$  satisfying  $\mathcal{W}(G, \pi) \neq \phi$ . Let  $G_\pi \in \mathcal{W}(G, \pi)$  be a minimal  $\pi$ -edge-cut s-t 2-edge-connected subgraph of  $G$ . Let  $\alpha, \beta$  be two edge-disjoint s-t paths of  $G_\pi$ . Let  $V_{\pi\alpha} = \{x_1, x_2, \dots, x_p\} (\subseteq V(\pi))$  be the set of all intersection vertices of  $\pi, \alpha$ , where  $x_1 <_\pi x_2 <_\pi \dots <_\pi x_p$ . Let  $V_{\pi\beta} = \{y_1, y_2, \dots, y_q\} (\subseteq V(\pi))$  be the set of all intersection vertices of  $\pi, \beta$ , where  $y_1 <_\pi y_2 <_\pi \dots <_\pi y_q$ . Let  $V_{\pi\alpha\beta} = \{z_1, \dots, z_r\} (\subseteq V(\pi))$  be the set of all vertices which  $\pi, \alpha, \beta$  have in common, where  $z_1 <_\pi z_2 <_\pi \dots <_\pi z_r$ . Subpaths  $\alpha[x_i, x_{i+1}]$  of  $\alpha$  avoiding  $\pi$  and  $\beta[y_j, y_{j+1}]$  of  $\beta$  avoiding  $\pi$ , where either  $x_i <_\pi y_j$  or  $y_j <_\pi x_i$ , are said to be *interlacing subpaths*, if the subpath  $\pi[x_i, y_{j+1}]$  ( $\pi[y_j, x_{i+1}]$ ) contains no vertex of  $V_{\pi\alpha\beta}$  when  $x_i <_\pi y_j$  ( $y_j <_\pi x_i$ ).  $\square$

In the graph  $G$  given in Fig.1, for two edge-disjoint s-t paths;  
 $\alpha : v_0(= s), v_1, u_1, v_4, v_5, u_2, v_6, v_7, v_9(= t)$ ,  $\beta : v_0(= s), w_1, v_2, v_3, v_5, v_6, v_8, v_9(= t)$ ,  
 we have  $V_{\pi\alpha} = \{v_1, v_4, v_5, v_6, v_7, v_9\}$ ,  $V_{\pi\beta} = \{v_0, v_2, v_3, v_5, v_6, v_8\}$ ,  $V_{\pi\alpha\beta} = \{v_0, v_5, v_6, v_9\}$ . And subpaths  $\alpha[v_1, v_4]$  and  $\beta[v_0, v_2]$  are interlacing subpaths, and  $\alpha[v_7, v_9]$  and  $\beta[v_6, v_8]$  are also interlacing paths. But  $\alpha[v_1, v_4]$  and  $\beta[v_6, v_8]$  are not interlacing subpaths as  $v_5, v_6 \in V_{\pi\alpha\beta}$  are on  $\pi[v_0, v_8]$ .

In order to show that if graph  $G$  has a prohibitive s-t path set  $P(\subseteq P_s(G))$ , then  $G$  must contain a prohibitive graph as its subgraph, we can prove the following Lemma 4.6 and Lemma 4.7.

**Lemma 4.6.** Suppose that  $G$  has a prohibitive s-t path set  $P$ . Then there is an s-t path  $\pi$  of  $P$  whose proper bridge  $B$  in  $G$  contains two interlacing subpaths  $\alpha[x_i, x_{i+1}]$  of  $\alpha$  and  $\beta[y_j, y_{j+1}]$  of  $\beta$  with respect to  $\pi$  in  $G_\pi$ , where  $G_\pi$  is a minimal  $\pi$ -edge-cut s-t 2-edge-connected subgraph of  $G$ , and  $\alpha, \beta$  are two edge-disjoint s-t paths in  $G_\pi$ .

**Sketch of Proof.** Let  $P$  be a prohibitive s-t path set of  $G$ . We can find the s-t path  $\pi$  of  $P$  satisfying the following *condition I* by using the following *procedure I*.

*Condition I:* There is a proper bridge  $B$  of  $\pi$  in  $G$  such that  $B$  contains interlacing subpaths  $\alpha[x_i, x_{i+1}]$  of  $\alpha$  and  $\beta[y_j, y_{j+1}]$  of  $\beta$  with respect to  $\pi$  in  $G_\pi$ , where  $G_\pi$  is a minimal  $\pi$ -edge-cut s-t 2-edge-connected subgraph of  $G$ , and  $\alpha, \beta$  are two edge-disjoint s-t paths in  $G_\pi$ .

*Procedure I:* Let  $\pi$  be an s-t path of  $P$ . Let  $B$  be a proper bridge of  $\pi$  in  $G$ . We do the following *Loop* iteratively.

*Loop:* If  $\pi$  satisfies *Condition I* then end. Otherwise, we can find an s-t path  $\pi'$  of  $P$  such that there is a bridge  $B'$  of  $\pi'$  in  $G$  whose nucleus contains the nucleus of  $B$  and there are more vertices in the nucleus of  $B'$  than in the nucleus of  $B$ . Let  $B, \pi$  be  $B', \pi'$ , respectively.

Note that, in each loop, the nucleus of  $B$  increases at least by one vertex. Thus the loop will end in at most  $|V|$  times, where  $V$  is the set of vertices in  $G$ .  $\square$




Fig.5 An illustration of the proof of Lemma 4.7.

**Lemma 4.7.** Suppose that  $G$  has an s-t path  $\pi$  satisfying  $\mathcal{W}(G, \pi) \neq \phi$ . Let  $\alpha, \beta$  be two edge-disjoint s-t paths of  $G_\pi \in \mathcal{W}(G, \pi)$ . Let  $V_{\pi\alpha} = \{x_1, x_2, \dots, x_p\}$ ,  $V_{\pi\beta} = \{y_1, y_2, \dots, y_q\}$  and  $V_{\pi\alpha\beta} = \{z_1, \dots, z_r\}$

be defined as in Definition 4.12. If a bridge  $B$  of  $\pi$  in  $G$  contains interlacing subpaths  $\alpha[x_i, x_{i+1}]$  of  $\alpha$  and  $\beta[y_j, y_{j+1}]$  of  $\beta$  in  $G_\pi$  with respect to  $\pi$ , then  $G$  contains a prohibitive graph as its subgraph.

**Sketch of Proof.** By the known conditions given in this lemma, we construct a prohibitive graph as its subgraph.

By Lemma 4.5, there is a path  $\pi_{uv}$  between an internal vertex  $u$  on  $\alpha[x_i, x_{i+1}]$  and an internal vertex  $v$  on  $\beta[y_j, y_{j+1}]$  consisting of edges and vertices only in the nucleus of bridge  $B$ , i.e.,  $\pi_{uv}$  is vertex-disjoint path with  $\pi$  except  $u, v$ . See Fig.5. Thus, we can also find a prohibitive graph as subgraph of  $G$  independently of the way how the path  $\pi_{uv}$  is traced.  $\square$

By Theorem 4.3 and Lemmas 4.5, 4.6, 4.7, the following Theorem 4.4 holds.

**Theorem 4.4.** In a probabilistic graph  $(G, p)$ ,  $\underline{\Gamma}_{(G, f, p)} = \Gamma_{(G, p)}$  holds iff  $G$  contains no prohibitive graph as its subgraph.  $\square$

## 5 A Method of Computing the Lower Bound

Given a probabilistic graph  $(G, p)$  and an s-t path number  $f$  of  $G$ , we show a method of computing the lower bound  $\underline{\Gamma}_{(G, f, p)}$ . We first wish to recall the procedure **FEDP** and the definition of  $\underline{\Gamma}_{(G, f, p)}$  in section 3.

For a probabilistic graph  $(G = (V, E, s, t), p)$  and an s-t path number function  $f$  of  $G$ , let  $\mathcal{U}_{f, \pi_i}$  denote the set of all  $U \subseteq E$  for which s-t path  $\pi_i$  is selected as a member of edge-disjoint s-t paths  $FEDP(G - U, f)$ . Let  $p(\mathcal{E}_U)$  be the probability of the event  $\mathcal{E}_U$  that all edges of  $U$  are failed and all edges of  $E - U$  are operative, and  $p(\mathcal{E}_{f, \pi_i})$  is the probability of the event that at least one event  $\mathcal{E}_U$ , for all  $U \in \mathcal{U}_{f, \pi_i}$ , arises in  $(G, p)$ . Thus, we have

$$\begin{aligned}
 \underline{\Gamma}_{(G, f, p)} &= \sum_{U \subseteq E} |FEDP(G - U, f)| \rho(G - U) \\
 &= \sum_{i=1}^{|P_{s,t}(G)|} \sum_{U \in \mathcal{U}_{f, \pi_i}} \rho(G - U) \\
 &= \sum_{i=1}^{|P_{s,t}(G)|} \sum_{U \in \mathcal{U}_{f, \pi_i}} p(\mathcal{E}_U) \\
 &= \sum_{i=1}^{|P_{s,t}(G)|} p(\mathcal{E}_{f, \pi_i}). \tag{5}
 \end{aligned}$$

We can compute the lower bound  $\underline{\Gamma}_{(G, f, p)}$  by formula (5) instead of formula (3).

## 6 Concluding Remarks

For a probabilistic graph, we proposed a lower bound for estimating the expected maximum number of edge-disjoint s-t paths. The necessary and sufficient conditions with respect to both s-t path number function and graph construction, where this lower bound coincides with the expected maximum number of edge-disjoint s-t paths, are clarified. A method of computing this lower bound is also given, although by this computing method the lower bound does not seem to be efficiently computed for a general probabilistic graph.

However, for a probabilistic one-layered  $s$ - $t$  graph, (a two-terminal graph where the subgraph obtained by deleting its  $s, t$  is exactly a simple path. Fig.6 illustrates an example of one-layered  $s$ - $t$  graph.) as it satisfies the necessary and sufficient conditions and the number of all its  $s$ - $t$  paths is a polynomial function in the number of its vertices, the lower bound based on its exact  $s$ - $t$  path number function can efficiently be computed by the computing method shown in section 5, i.e., the expected maximum number of edge-disjoint  $s$ - $t$  paths in a probabilistic one-layered  $s$ - $t$  graph can efficiently be computed. Detailed description of these proofs is lengthy and to be reported elsewhere.




Fig.6 A one-layered  $s$ - $t$  graph.

## References

- [1] M. O. Ball: "Computational complexity of network reliability analysis: An overview", *IEEE Trans. Reliability*, Vol.R-35, pp230-239 (1986).
- [2] P. Cheng and S. Masuyama: "Problem of computing the expected maximum number of vertex-disjoint  $s$ - $t$  paths on probabilistic graphs", *Research Report COMP92-1, Inst. Electron. Infor. Comm. Eng. Japan*, pp1-8, (1992) (in Japanese).
- [3] C. J. Colbourn: *The Combinatorics of Network Reliability*, Oxford University Press (1987).
- [4] K. Menger: "Zer allgemeinen kurventheorie", *Fund. Math.* Vol 10, pp.96-115(1927).
- [5] B. Mishra and R. E. Tarjan: "A linear-time algorithm for finding a ambitus", *Algorithmica*, 7, pp.521-554(1992).
- [6] W. T. Tutte: "Bridges and hamiltonian circuits in planar graph", *Aequationes mathematicae*, 15, pp.1-13(1977).