

ICESluvan, a 94-Kilobase Mosaic Integrative Conjugative Element Conferring Interspecies Transfer of VanB-Type Glycopeptide Resistance, a Novel Bacitracin Resistance Locus, and a Toxin-Antitoxin Stabilization System

Eva K. Bjørkeng,^{a*} Erik Hjerde,^b Torunn Pedersen,^c Arnfinn Sundsfjord,^{a,c} Kristin Hegstad^{a,c}

Research Group for Host-Microbe Interactions, Department of Medical Biology, Faculty of Health Sciences, University of Tromsø, Tromsø, Norway^a; Molecular Biosystems Research Group, Department of Chemistry, Faculty of Science and Technology, University of Tromsø, Tromsø, Norway^b; Reference Centre for Detection of Antimicrobial Resistance, Department of Microbiology and Infection Control, University Hospital of North Norway, Tromsø, Norway^c

A 94-kb integrative conjugative element (ICESluvan) transferable to *Enterococcus faecium* and *Enterococcus faecalis* from an animal isolate of *Streptococcus lutetiensis* consists of a mosaic of genetic fragments from different Gram-positive bacteria. A variant of ICESluvan was confirmed in *S. lutetiensis* from a patient. A complete Tn5382/Tn1549 with a *vanB2* operon is integrated into a streptococcal ICESde3396-like region harboring a putative bacteriophage exclusion system, a putative agglutinin receptor precursor, and key components of a type IV secretion system. Moreover, ICESluvan encodes a putative MobC family mobilization protein and a relaxase and, thus, in total has all genetic components essential for conjugative transfer. A 9-kb element within Tn5382/Tn1549 encodes, among others, putative proteins similar to the TnpX site-specific recombinase in *Faecali-bacterium* and VanZ in *Paenibacillus*, which may contribute to the detected low-level teicoplanin resistance. Furthermore, ICESluvan encodes a novel bacitracin resistance locus that is associated with reduced susceptibility to bacitracin when transferred to *E. faecium*. The expression of a streptococcal *pezAT* toxin-antitoxin-encoding operon of ICESluvan in *S. lutetiensis*, *E. faecium*, and *E. faecalis* was confirmed by reverse transcription (RT)-PCR, indicating an active toxin-antitoxin system which may contribute to stabilizing ICESluvan within new hosts. Junction PCR and DNA sequencing confirmed that ICESluvan excised to form a circular intermediate in *S. lutetiensis*, *E. faecalis*, and *E. faecium*. Transfer between *E. faecalis* cells was observed in the presence of helper plasmid pIP964. Sequence analysis of the original *S. lutetiensis* donor and enterococcal transconjugants showed that ICESluvan integrates in a site-specific manner into the C-terminal end of the chromosomal tRNA methyltransferase gene *rumA*.

Horizontal gene transfer is a key factor in bacterial evolution, and mobile genetic elements (MGEs) play an important role in the dissemination and persistence of antimicrobial resistance in enterococci. Genome sequence analysis and comparative genome hybridizations of seven *Enterococcus faecium* isolates from various sources have revealed large differences in genome size, mostly due to the variable presence of mobile genetic elements. The *E. faecium* pan-genome is considered to be unrestricted in size (1), implying that excess genes, such as those involved in environmental persistence, colonization, and virulence, can easily be incorporated into the *E. faecium* gene pool. Up to 38% of the *E. faecium* genome may be noncore, and differences in gene content indicate that gain and loss of genes as important in the evolution of *E. faecium* (1, 2, 3). These findings are consistent with the presence of more than 25% mobile or foreign DNA in *Enterococcus faecalis* V583 (3, 4), suggesting extensive genome plasticity.

The first high-level vancomycin-resistant enterococci (VRE) were described in 1988 (5, 6). VRE have since become an increasing nosocomial problem both in Europe and the United States (7, 8). The origin of vancomycin resistance determinants is not known, although soil bacteria seem to represent a rich and assorted reservoir of genes closely related to *vanA* (9). Moreover, analyses of genomes from invasive *vanB*-positive *E. faecium* isolates and *vanB*-positive anaerobic gut commensals demonstrate that *vanB* resistance in enterococci commonly arises through gene transfer from members of other bacterial genera in the human

gastrointestinal tract (10). The *vanB* cluster is mainly found in *E. faecium* and *E. faecalis*, though it has also been described in isolates of *vanB*-resistant *Enterococcus gallinarum* (11, 12, 13, 14), *Enterococcus hirae* (15), *Enterococcus durans* (16), *Enterococcus casseliflavus* (17), *Staphylococcus* (18), *Streptococcus* (19, 20, 21, 22), *Eggerthella*, *Clostridium*, *Ruminococcus* (23, 24, 25, 26, 27), and *Atopobium* (27).

The *vanB* gene cluster has a conserved gene order and can be divided into three genetic subtypes, *vanB1*, *vanB2*, and *vanB3* (28, 29, 30, 31, 32). As an integral part of the conjugative transposon Tn5382/Tn1549, *vanB2* is the most-widespread subtype in clinically important enterococci (30, 33, 34, 35, 36, 37, 38). Tn5382/Tn1549 is able to support transfer of the *vanB2* operon from *Clos-*

Received 27 November 2012 Accepted 24 September 2013

Published ahead of print 27 September 2013

Address correspondence to Kristin Hegstad, Kristin.Hegstad@uit.no.

* Present address: Eva K. Bjørkeng, Department of Chemistry, Faculty of Science and Technology, University of Tromsø, Tromsø, Norway.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/JB.02165-12>.

Copyright © 2013, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.02165-12

The authors have paid a fee to allow immediate free access to this article.

TABLE 1 Bacterial strains used in this study and their relevant characteristics

Strain	Species	Properties ^a	MIC mg liter ⁻¹			
			Vancomycin	Teicoplanin	Bacitracin	Reference(s)
5-F9	<i>S. lutetiensis</i>	Van ^r veal calf strain with <i>vanB2</i> Tn5382 chromosomally located	≥256	3	≥256	19, 20
NEM760	<i>S. lutetiensis</i>	Van ^r human patient strain with <i>vanB2</i> chromosomally located	≥256	4	≥256	22, 42
4-C11	<i>S. gallolyticus</i>	Van ^r veal calf strain with nontransferable <i>vanB2</i> Tn5382 and <i>vanA</i>	≥256	32	24	19, 20
4-G10	<i>S. gallolyticus</i>	Van ^r veal calf strain with nontransferable <i>vanB2</i> Tn5382 and <i>vanA</i>	≥256	64	≥256	19, 20
C68	<i>E. faecium</i>	<i>vanB2</i> Tn5382 type strain				40
BM4105-RF	<i>E. faecium</i>	Rif ^r Fus ^r plasmid-free recipient	1,5	0,5	48	46
MM5-F9a	<i>E. faecium</i>	Transconjugant resulting from mating of 5-F9 and BM4105-RF	32	3	≥256	20
OG1-RF	<i>E. faecalis</i>	Rif ^r Fus ^r recipient derived from OG1	4	0,25	24	73
OG5-F9a	<i>E. faecalis</i>	Transconjugant resulting from mating of 5-F9 and OG1-RF	≥256	3	32	This study
JH2-2	<i>E. faecalis</i>	Rif ^r Fus ^r plasmid-free recipient derived from JH2	3	1	16	74
JH5-F9a	<i>E. faecalis</i>	Transconjugant resulting from mating of 5-F9 and JH2-2	≥256	3	8	20
UV202	<i>E. faecalis</i>	Rif ^r Fus ^r plasmid-free recipient, recombination-deficient derivative of JH2-2	1,5	0,5	128	75
UV5-F9d	<i>E. faecalis</i>	Transconjugant resulting from mating of 5-F9 and UV202	≥256	3	192	20
BM4110	<i>E. faecalis</i>	Str ^r plasmid-free recipient strain derived from JH2	1,5	0,25	96	76
BM4110 pIP964	<i>E. faecalis</i>	Str ^r recipient strain derived from JH2 containing plasmid pIP964 (pCF10 <i>rep agg</i> ⁺ Tra ⁺ , 65 kb)	1,5	0,38	≥256	76
ICESluvan BM4110 pIP964	<i>E. faecalis</i>	1st-generation transconjugant resulting from mating of donor 5-F9 and BM4110 pIP964	≥256	3	≥256	This study
ICESluvan pIP964 JH2-2	<i>E. faecalis</i>	2nd-generation transconjugant resulting from mating of ICESluvan BM4110 pIP964 and JH2-2	≥256	3	≥256	This study
ICESluvan pIP964 UV202	<i>E. faecalis</i>	2nd-generation transconjugant resulting from mating of ICESluvan BM4110 pIP964 and UV202	≥256	3	192	This study

^a Van^r, vancomycin resistant; Rif^r, rifampin resistant; Fus^r, fusidic acid resistant; Str^r, streptomycin resistant.

tridium to enterococci in the intestinal environment (39), but the transposon is more often transferred as a part of larger chromosomal elements or plasmids (30, 35, 37, 40, 41).

An approximately 100-kb transferable chromosomal element containing *vanB2* has been described in a patient isolate of *Streptococcus lutetiensis* previously designated *Streptococcus bovis* bio-type II with *vanB3* (22, 42) and in the plasmid-free *S. lutetiensis* strain 5-F9 from animal feces (20). The *vanB2* cluster of the 5-F9 strain has previously been shown to be an integral part of a Tn5382/Tn1549 element (20) and was transferred to *E. faecium* and *E. faecalis* at relatively high frequencies (10^{-7} to 10^{-5}). In contrast, the element was unable to transfer between *E. faecium* and *E. faecalis* strains (20, 42), which suggests a coresident transfer system in *Streptococcus* (42). Retransfer of the ~100-kb element from the patient isolate between *E. faecalis* strains, including to a recombination-deficient recipient, was obtained in the presence of a conjugative helper plasmid (42), and the ~100-kb element appeared to integrate in a site-specific manner (20, 42). Transfer of this ~100-kb chromosomal element from the *S. lutetiensis* strain that contains no visible plasmids into enterococci suggests self-encoded transfer functions consistent with an integrative and conjugative element (ICE) (20). The transfer mechanism has not been resolved.

Here, we describe the sequencing, annotation, and functional analysis of the 94-kb element designated ICESluvan, originally detected in *S. lutetiensis* from animal feces in the Netherlands. A variant of this element was confirmed in the *S. lutetiensis* patient isolate from France. The element encodes VanB-type vancomycin resistance on conjugative transposon Tn5382/Tn1549, a VanZ homologue which may contribute to the detected low-level resistance to teicoplanin, a novel bacitracin locus which seems to contribute to bacitracin resistance in *S. lutetiensis* and *E. faecium*, and a streptococcal toxin-antitoxin (TA) *pezAT* locus, as well as all essential components for conjugative transfer. ICESluvan excises to form a circular intermediate in *S. lutetiensis*, *E. faecalis*, and *E. faecium* and integrates in a site-specific manner. However, transfer between enterococci required a helper plasmid. The expression of the *pezAT* operon indicates a functional toxin-antitoxin system.

MATERIALS AND METHODS

Bacterial strains and construction of BAC clones. The bacterial strains used in this study and their relevant characteristics are given in Table 1. Briefly, genomic DNA from the *E. faecium* transconjugant MM5-F9a was used to generate a bacterial artificial chromosome (BAC) library with an average insert size of 90 kb (MWG Biotech AG). MM5-F9a was obtained by transfer of the ICESluvan element from *S. lutetiensis* 5-F9, isolated from

the feces of a veal calf in the Netherlands, to recipient *E. faecium* BM4105-RF by filter mating (20). The *vanB2*-containing strains *S. lutetiensis* NEM760, *Streptococcus gallolyticus* 4-C11, and *S. gallolyticus* 4-G10 were included to search for ICESlvan in streptococci. *E. faecalis* transconjugants OG5-F9a, JH5-F9a, and UV5-F9d and laboratory strains OG1-RF, JH2-2, BM4110, and UV202 were used for various studies of ICESlvan as indicated below. Total DNA from *E. faecium* C68 (40) was used as the template for probe synthesis.

Selection and sequencing of BAC clones. Sequential dot blot and Southern hybridizations were used to identify BAC clones positive for both *vanB2* and Tn5382/Tn1549. PCR-based probes were made (PCR digoxigenin [DIG] probe synthesis kit; Boehringer Mannheim) for *vanB* using the consensus primers 5'-CAAAGCTCCGAGCTTGATG-3' and 5'-TGCATCCAAGCACCCGATAATAC-3' (29) and for Tn5382 using primers 5'-GTTCTTATCCGAGGTGGTATT-3' and 5'-ACGCCATGCTATTACTCCGGC-3' (40). Hybridizations were detected using the DIG luminescent detection kit (Boehringer Mannheim). Plasmid DNA was isolated from BAC clones that scored positive for one or both probes ($n = 24$). Clones were grown overnight in yeast extract-tryptone (YT) medium containing 12.5 $\mu\text{g ml}^{-1}$ chloramphenicol, plasmid replication was induced (Epicentre induction solution; Epicentre), and BAC DNA was purified using the E.Z.N.A BAC/PAC kit (Omega Bio-Tek). NotI (New England BioLabs)-digested plasmid DNA was separated on a 1.2% agarose gel, 5 to 15 s, 6 V/cm, 16 h, at 15°C on a CHEF-DR III (Bio-Rad) and analyzed by Southern hybridization. Five clones positive for both probes and with an average insert size of 90 kb were selected for sequencing.

Libraries of the BAC inserts of each selected clone were constructed with the use of multiplex identifiers (MIDs) and sequenced to an average depth of 91.5-times coverage using 454-GS-FLX technology. The BAC inserts were *de novo* assembled individually with Newbler (Roche).

End sequencing of the BAC inserts provided additional geographical information for contig ordering. End sequencing and extension of contigs were set up using extracted BAC DNA and BigDye 3.1 for cycle sequencing and an ABI prism 377 genetic analyzer (Applied Biosystems). The cycle sequencing program was as follows: initial denaturation at 96°C for 5 min followed by 99 cycles of 96°C for 30 s, 52°C for 10 s, and 60°C for 4 min (primers available on request).

DNA extraction and sequencing of transconjugants. Genomic DNA from transconjugants was obtained using a bacterial DNA kit (E.Z.N.A) with the following modifications: lysis was performed at 30°C for 40 min using 1.5 mg lysozyme and 100 U mutanolysin in a total volume of 220 μl . Direct genome sequencing of the transconjugants was performed as for BAC end sequencing.

Gap closure and annotation. Abacas (<http://abacas.sourceforge.net/index.html>) was used to map contigs from the *de novo* assembly against *Streptococcus suis* BM407 (FM252032) and Tn1549 (AF192329). The sequence was annotated using Artemis software (43). Homology comparisons to nonredundant protein databases were performed with BLAST (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) and FASTA (<http://www.ebi.ac.uk/Tools/fasta33/index.html>) software. Protein motifs were identified using Pfam (<http://pfam.sanger.ac.uk/search>) and Prosite (<http://au.expasy.org/prosite/>). Transmembrane domains were identified with TMHMM (<http://www.cbs.dtu.dk/services/TMHMM/>), and signal sequences were identified with SignalP version 3.0 (44). Comparison of the large mobile elements was facilitated using the Artemis Comparison Tool (ACT) (45), which enabled the visualization of BLASTN and TBLASTX comparisons between these sequences.

Verification of sequence assembly by PCRs. To verify the sequence assembly, over 40 PCRs with an average product size of 2.5 kb were performed in ICESlvan (see Table S1 in the supplemental material), using genomic DNA of transconjugants *E. faecium* MM5-F9a and OG5-F9a and *E. faecalis* JH5-F9a as templates. Some of these primers were used to confirm the presence of ICESlvan regions (see Table S1) in streptococci (*S. lutetiensis* 5-F9 and NEM760 and *S. gallolyticus* 4-C11 and 4-G10) and to

sequence the bacitracin locus in the *E. faecalis* transconjugants OG5-F9a and JH5-F9a.

RNA extraction for expression analyses. The expression of the *pezAT* operon was analyzed by semiquantitative reverse transcription (RT)-PCR. Total RNA was extracted from stationary-phase cultures using an RNeasy minikit (Qiagen) with an extended lysis step of 1 h with 10 mg lysozyme and 10 U mutanolysin in a total volume of 200 μl . DNA contamination was removed by DNase I treatment, and RNA quality and quantity examined by gel analyses and spectrophotometric measurements prior to cDNA synthesis. Reverse transcription of 100 ng total RNA was performed using SuperScript II (Invitrogen). For each sample, a control reaction (without reverse transcriptase) with no enzyme was included. PCR was performed using primer sets specific for each of the genes (*pezAF* [5'-TGACCTGGAATCGTCTTTCC-3'] and *pezAR* [5'-TGGCCGAATTGATTAACACA-3'] and *pezTF* [5'-TTTTCTGCAAATCCCTTG-3'] and *pezTR* [5'-GACAAAGCGGAGCAGGTAAG-3']). In addition, PCR using primers *pezTF* and *pezAR* was performed. Amplicons were visualized on an agarose gel and confirmed by sequencing using BigDye 3.1 technology (Applied Biosystems).

Verification of ICESlvan circularization. Circularization of the ICESlvan element was examined by PCR using primers located on the very end of the element (see Fig. 2), as follows: 5'-AAGGGAAAATTAGC GATACC-3' (nucleotides [nt] 93779 to 93798), 5'-CTGCTGCAAATGT AAATGGC-3' (nt 93383 to 93402), and 5'-TCCAACAGACATTCACA TCAA-3' (nt 130 to 151). PCR products were confirmed by sequencing.

Transfer of ICESlvan. Transfer of the ICESlvan element to *E. faecalis* strains BM4110 and JH2-2 with pIP964 and retransfer from these first-generation transconjugants to recipients (*E. faecalis* strains JH2-2, UV202, and BM4110) without this helper plasmid were performed by filter mating as previously described (33), with some modifications. Briefly, donor and recipient strains were grown to an A_{600} of 0.6 and mixed in a ratio of 1:1 in 2 ml of culture. The suspension was pelleted, resuspended in 150 μl brain heart infusion (BHI), and used for filter mating before application on selective agar plates using 8 mg liter⁻¹ vancomycin, 1,000 mg liter⁻¹ streptomycin, 20 mg liter⁻¹ rifampin, and 10 mg liter⁻¹ fusidic acid. A control experiment (*E. faecalis* JH5-F9a \times BM4110) with recipients containing no helper plasmid was also carried out. The presence of the *vanB* gene of ICESlvan was confirmed by PCR (29), and ICESlvan insertion in the transconjugant genome was confirmed by visual inspection of the SmaI pulsed-field gel electrophoresis (PFGE) pattern. The presence of the pheromone-responsive helper plasmid pIP964 (46) was determined by PCRs that detect the pCF10 replication initiation gene *rep*_{pCF10} as described by Jensen et al. (47), and the presence of the pheromone plasmid's conserved aggregation substance (*agg*) was detected using primers 5'-GC TCGTGGTGATGTTCTTTC-3' and 5'-CTTTTCTACCACTAATGGCT CTAC-3'.

PFGE analyses. Total DNA was digested with SmaI (New England BioLabs) and analyzed by PFGE as previously described (48), with some modifications. Briefly, 5 μl lysozyme (100 mg ml⁻¹) and 2 μl mutanolysin (10 U μl^{-1}) were added to the agarose plugs before molding, as well as to the lysis buffer.

Antimicrobial susceptibility testing. Susceptibility testing was performed using MIC test strips for vancomycin, teicoplanin (bioMérieux), and bacitracin (Liofilchem) on *S. lutetiensis* 5-F9, *E. faecalis* JH5-F9a, JH2-2, OG5-F9a, and OG1-RF, and *E. faecium* MM5-F9a and BM4105-RF according to the manufacturers' recommendations.

Nucleotide sequence accession number. The whole sequence of ICESlvan is available in GenBank under the accession number HE963029.

RESULTS AND DISCUSSION

Gap closure and ICESlvan mosaic structure. The five selected BAC clones were sequenced to an average depth of 91.5-times coverage and assembled individually into an average of 28 contigs (>500 bp). Mapping of contigs against reference sequences and




FIG 1 Genetic organization of the 94,189-bp ICES_{Luvan} element and linear DNA comparison against Tn1549, the 64-kb ICES_{Sde3396} element from *S. dysgalactiae* subsp. *equisimilis*, and regions from *S. suis* 89/1591, *S. mitis* B6, and *Faecalibacterium prausnitzii* A2-165. In ICES_{Luvan}, coding sequences (CDSs) are colored according to predicted functions. In the other genetic elements, black arrows indicate CDSs with no assigned functions and white arrows indicate CDSs with putative assigned functions. Gene identifiers for Tn1549, *S. dysgalactiae* subsp. *equisimilis*, *S. suis* 89/1591, *S. mitis* B6, and *F. prausnitzii* A2-165 are captured from records with GenBank accession numbers AF192329, EU142041, NZ_AAFA03000001, FN568063, and NZ_ACOP00000000, respectively. For simplicity, the CDS prefixes are omitted in the figure. Gene names are indicated if available from annotation. Disrupted genes are linked via dashed lines, and both disrupted and pseudogenes are indicated as crosshatched arrows. For the comparison, red shading between the genetic elements represents regions with similarity: light red indicates regions with >65% amino acid sequence identity; red indicates regions with >90% amino acid sequence identity; and dark red indicates regions with >99% amino acid sequence identity.

end sequencing of BAC inserts provided scaffold information for the gap closure that resulted in one large, 94-kb contig (GenBank accession number HE963029) containing 86 open reading frames (ORFs) (Fig. 1; see also Table S2 in the supplemental material). Nine smaller contigs of >1 kb with BLAST similarities to unculturable organisms and transposases were discarded from further analysis. Extensive control PCRs (see Table S1) verified the assembly for *E. faecium* transconjugant MM5-F9a and confirmed the presence of the ICES_{Luvan} element in *E. faecalis* transconjugants OG5-F9a and JH5-F9a (Table 1). Some primer combinations used for MM5-F9a did not give PCR products with OG5-F9a and JH5-F9a. However, the presence of these regions was confirmed by altering the combination of the primers (denoted by plus signs in Table S1). Some of the PCRs worked at one time point and not at another, and when we sequenced the bacitracin locus of the *E. faecalis* transconjugants, we found that some of the primer combinations did not give a PCR product even though both primers matched the sequence 100%. Thus, we think there might be other reasons than nucleotide differences, possibly related to the primary or secondary DNA sequence, why PCR products were not achievable for all transconjugants in some specific regions.

Interestingly, homology searches of the 86 predicted ORFs showed that the ICES_{Luvan} element contains genes similar to 24 ORFs (ORF4 to -13, ORF15 to -17, ORF60 to -66, and ORF71 to -74) of ICES_{Sde3396} found in the beta-hemolytic *Streptococcus dysgalactiae* subsp. *equisimilis* NS3396 (49). Furthermore, ICES_{Luvan} ORF1 to -3, ORF14, ORF69, and ORF75 to -86 showed similarities to ORFs from three non-beta-hemolytic streptococci, *Streptococcus pneumoniae*, *S. suis*, and *Streptococcus mitis*.

A complete *vanB2* Tn5382/Tn1549 (ORF18 to -30 and ORF39 to -59) putative conjugative transposon was integrated into the partial ICES_{Sde3396}-like sequence in ORF17, similar to ICES_{Sde3396}_54. The *vanB2* Tn5382/Tn1549 sequence of ICES_{Luvan} was 99% iden-

tical at the nucleotide level to the fully sequenced Tn1549 (Fig. 1; see also Table S2 in the supplemental material) (30). Five more ORFs were identified in the Tn5382/Tn1549 element inserted in ICES_{Luvan} compared to those from previous reports (see Table S2). Identical sequences were also present in Tn1549 (30) but have not been annotated previously. In addition, a 9-kb contig located within ORF30 in Tn5382/Tn1549 was flanked by directly repetitive sequences with 291 of 292 nucleotides identical. Annotation of the 9-kb contig revealed a TnpX site-specific recombinase (ORF31), a pseudotransposase of the IS30 family (ORF33), a putative VanZ family protein (ORF34), a putative mobilization protein (ORF36), and a DNA primase (ORF37), as well as hypothetical proteins (ORF32, -35, and -38) with 40 to 96% amino acid identity to putative proteins in *Faecalibacterium*, *Lactobacillus*, and *Paenibacillus* species, respectively (see Table S2).

The mosaic pattern of the ICES_{Luvan} element, which consists of genes from several different streptococcal species, *Faecalibacterium*, enterococci, and possibly other intestinal bacteria (see Table S2 in the supplemental material), indicate that mobilization and recombination events have been key factors in the assembly of this element. A mosaic pattern was also observed for ICES_{Sde3396}, which is a montage of genes derived from group A, B, and G *Streptococcus* organisms, in addition to genes acquired from non-streptococcal Gram-positive bacteria, such as *Listeria innocua* and *E. faecalis* (49).

ICES_{Luvan} variant confirmed in an *S. lutetiensis* isolate from a patient in France. During examination of mixed fecal samples from 556 veal calf herds in the Netherlands, four vancomycin-resistant *Streptococcus* isolates were found. In addition to *S. lutetiensis* 5-F9, *S. gallolyticus* isolates 4-C11 and 4-G10 were positive for the *vanB* gene and found to have a *vanB2* cluster integrated in Tn5382/Tn1549 (20). Furthermore, *S. lutetiensis* NEM760, isolated from a stool swab of a patient in France, showed transfer of

an approximately 100-kb element containing *vanB2* (22, 42) that inserted into the same *Sma*I PFGE fragment as ICESlvan in both *E. faecium* BM4105-RF- and *E. faecalis* JH2-derived recipients (20, 22, 42). Since 4-C11, 4-G10, and NEM760 are likely candidates to contain ICESlvan or variants of this element, we have examined these isolates by using primers selected to confirm the ICESlvan sequence assembly (see Table S1 in the supplemental material). The circa-100-kb element found in NEM760 was indeed very similar to ICESlvan, with some differences in the right end of the element, while 4-C11 and 4-G10 showed the presence of Tn5382/Tn1549 and adjacent regions but not the left or right end of ICESlvan, which is not surprising since transfer of *vanB* was not achieved for these isolates (19, 20). The presence of highly similar variants of ICESlvan capable of interspecies transfer in two *S. lutetiensis* isolates from completely different environments implies that ICESlvan is a successful element that may contribute to resistance in enterococci.

ICESlvan type IV secretion system components. A putative transfer system (ORF11 to -16 and ORF79 to -80) (see Table S2 in the supplemental material) was identified in the ICESlvan element, in addition to those encoded by Tn5382/Tn1549 (ORF21 to -23, ORF26, and ORF43 and -44) (see Table S2) (30). ORF11 to -15 were 94 to 99% identical to ICESde3396_59 to ICESde3396_55 (49). The ORFs display the same genetic order in the ICESlvan element and belong to a type IV secretion system. Three factors have been recognized as important for DNA transfer by type IV secretion system in Gram-positive bacteria. These are murein hydrolase (50), which is involved in controlled local degradation of the peptidoglycan, making space for the formation of a mating channel, the VirB4 ATPase, which provides energy for the translocation, and the VirD4 coupling protein, which links the DNA transfer intermediate to the mating channel (50, 51). In the ICESlvan element, we identified two putative *virB4* genes (ORF15 and ORF26) encoding ATPases, a putative *virB6* gene (ORF13) encoding a putative membrane protein that has previously been shown to interact with several other Vir proteins mediating DNA substrate transfer through the cytoplasmic membrane channel (50), and two putative *virD4* genes (ORF11 and ORF21) encoding coupling proteins of the TraG/TraD family. ORF11 is truncated by a frameshift after codon 465, but ORF21 encodes a putative coupling protein which belongs to the same coupling-protein family. Moreover, ICESlvan ORF16 encodes a putative *N*-acetylmuramoyl L-alanine amidase that may aid in local degradation of the peptidoglycan by hydrolyzing the amide bond between the *N*-acetylmuramic acid side chain and L-alanine of the short peptide (52). Genes encoding putative mobilization proteins of the MobC family (ORF79 and ORF44) and relaxases (ORF80 and ORF43) were also identified in ICESlvan. Taken together, this indicates that the ICESlvan element contains most genes necessary for mobilization and conjugative transfer. Furthermore, ICESlvan is transferable by conjugation from *S. lutetiensis* organisms with no visible plasmids, strongly suggesting that its transfer system is intact (20).

ICESlvan circularization, transfer, and integration. A putative site-specific serine recombinase (ORF86) (see Table S2 in the supplemental material) that can be involved in the genomic exit and integration of mobile genetic elements was found at the right flank of the ICESlvan. Although it lacks 1 or 20 amino acids in the C-terminal end compared to the sequences of its closest homologues, SsuiDRAFT_2393, identified in *S. suis* (98% amino acid

identity), and SP70585_1107, identified in *S. pneumoniae* (95% amino acid identity), we hypothesize that the putative site-specific recombinase functions as normal in a transfer situation. This hypothesis was supported by circularization of the ICESlvan element both in the *S. lutetiensis* donor 5-F9 and the *Enterococcus* transconjugants, as shown by PCR and confirmed by sequence analyses (Fig. 2). Furthermore, we have transferred ICESlvan from *S. lutetiensis* 5-F9 to both *E. faecium* and *E. faecalis* (20). Retransfer was only possible when ICESlvan was transferred into recipients that contain the helper plasmid pIP964 (Tra⁺), which is restricted to *E. faecalis*. We did not test broad-host-range plasmids to achieve retransfer between *E. faecium* strains. Retransfer from *E. faecalis* BM4110 ICESlvan pIP964 to *E. faecalis* JH2-2 or UV202 (Fig. 3, lanes 4 to 6) and from JH2-2 ICESlvan pIP964 to BM4110 (data not shown) was confirmed by growth on selective plates, ICESlvan- and pIP964-specific PCRs, *Sma*I PFGE analysis (Fig. 3), and hybridization with *vanB* probe (data not shown) showing transconjugant patterns with an approximately 100-kb enlargement of one of the two 240-kb fragments compared to the recipient patterns (Fig. 3, lanes 4 to 6). However, transconjugant BM4110 ICESlvan pIP964 had gained an extra copy of ICESlvan, as shown by the presence of a fragment around 220 kb in Fig. 3, lane 4, which hybridized with *vanB* (data not shown). The presence of more than one copy of ICESlvan has been shown before in other JH-derived transconjugants (20). Both *Sma*I PFGE patterns and DNA sequencing show that the ICESlvan element is integrated in a site-specific manner into the recipient chromosome, although the *E. faecium* and *E. faecalis* integration sites within the C-terminal end of tRNA methyltransferase gene *rumA* showed some sequence differences (Fig. 4). The same integration site was identified within each species for all transconjugants tested (data not shown). The ICESlvan integration site in *S. lutetiensis* is also within the *rumA* gene. However, the right flank of ICESlvan in *S. lutetiensis* showed an extended 5.7-kb region identical to part of the *E. faecalis* G1-01247 *vanG* operon which was not transferred together with ICESlvan into the enterococci (see Fig. S1 in the supplemental material). Interestingly, ICESlvan ORF1 is a truncated homolog identical to the 13-amino-acid C-terminal part of *S. pneumoniae rumA* (see Table S2). This C-terminal part is in the same translational frame as the N-terminal part of the enterococcal *rumA*, and together, they form a recombined full-length *rumA* gene that is likely to be functional.

Since ICESlvan was shown to transfer from *Streptococcus* to *Enterococcus* but retransfer between enterococci required a helper plasmid, we hypothesize that a host factor is necessary for transfer. This has been shown for SXT elements in *Vibrio cholerae*, which require the host factors IHF and Fis for excision, recombination, and conjugation (53). Host factors may be chromosomally encoded and not on the MGE itself, leading to the inability of the MGE to conjugate in their absence. Transfer of the *E. faecalis* V583 pathogenicity island (PAI), containing the features of an ICE, by an ICE-independent mechanism has been reported. Characterization of V583 PAI transconjugants showed cotransfer of selectable markers representing virtually all regions of the chromosome, including a vancomycin resistance transposon, capsule genes, and alleles which are used for multilocus sequence typing. PAI transfer was dependent upon helper plasmid function in the donors (54).

Putative bacteriophage exclusion system. The ICESlvan ORF17 located upstream and ORF60 to -64 located downstream from Tn5382/Tn1549 are 81 to 97% identical to ICESde3396_54




FIG 2 Integrated (A) and circular (B) forms of ICESluvan in transconjugants *E. faecium* MM5-F9a, *E. faecalis* OG5-F9a, and *E. faecalis* JH5-F9a, as well as donor *S. lutetiensis* 5-F9. The left and right ends of ICESluvan are colored gray and black, respectively, while the native chromosomal DNA is shown by hatching. The sequences of the 20-bp ends of ICESluvan are also shown. Black and gray arrows indicate the positions and directions of primers for the left and right end of ICESluvan, respectively. The positions of primers designed from native *E. faecium* MM5-F9a regions are indicated with black-and-white arrows. PCR and sequencing confirmed both the integrated and the circular form of ICESluvan in *E. faecium* MM5-F9a, *E. faecalis* OG5-F9a, *E. faecalis* JH5-F9a, and *S. lutetiensis* 5-F9.

to ICESde3396_48. ORF60 and ORF17 are homologous to ICESde3396_54 and ICESde3396_53, encoding the abortive infection proteins AbiGI and AbiGII, respectively. The *Lactococcus lactis* *abiG* resistance genes confer complete resistance to ϕ 712 phages (936 phage species) and partial resistance to ϕ c2 phages (2 species) (55). Most Abi systems are plasmid encoded and are widespread in bacteria (56). The *abi* genes are also known to be involved in increased stress tolerance of the host. These genes are commonly found in *Streptococcus* (57). Transposon Tn5382/Tn1549 was inserted in the C-terminal end after codon 251 of *abiGII* in ICESluvan (Fig. 1), thereby disrupting this gene. Since this leads to an AbiGII protein lacking 30 amino acids compared to its closest homologue, it is not known whether these *abi* genes encode a functional system.

Putative virulence determinants. The ICESluvan ORF61 was 95% identical at the nucleotide level to ICESde3396_52, which encodes a putative agglutinin receptor precursor protein that harbors an LPXTG motif known to be common among several virulence factors in enterococci (58). Human salivary agglutinin has

previously been shown to interact with streptococci in a calcium-dependent reaction for oral bacterial aggregation (59, 60). The ICESluvan putative agglutinin receptor precursor is located next to a putative Ca^{2+} binding protein (ORF62). Studies have shown that transformation of nonaggregating *E. faecalis* with the streptococcal surface antigen SSP-5 confers an aggregation-positive phenotype in the presence of saliva agglutinin (61).

Resistance to glycopeptides and bacitracin. The ICESluvan element encodes vancomycin resistance mediated by a *vanB2* cluster (ORF48 to -54) located in Tn5382/Tn1549. *S. lutetiensis* 5-F9 and the transconjugants, as well as *S. lutetiensis* NEM760 and *S. gallolyticus* 4-C11 and 4-G10, accordingly expressed resistance to vancomycin, whereas the recipient strains *E. faecium* BM4105-RF and *E. faecalis* OG1-RF, JH2-2, UV202, BM4110, and BM4110 pIP964 were susceptible to vancomycin according to EUCAST clinical breakpoints for enterococci (MIC, ≤ 4 mg liter⁻¹) (Table 1). In addition, ORF34 on the 9-kb element inserted into Tn5382/Tn1549 encoded a protein with 40% amino acid similarity to a putative VanZ family protein from *Paenibacillus* (see Table S2 in the supplemental material).




FIG 3 PFGE of SmaI-digested total DNA of recipient, donor, and transconjugants from filter matings. Lanes 1 and 9, low-range PFGE marker; lane 2, ICESlvan donor *S. lutetiensis* 5-F9; lane 3, recipient BM4110 containing pIP964; lane 4, 1st-generation transconjugant/2nd-generation donor BM4110 ICESlvan pIP964; lane 5, 2nd-generation transconjugant JH2-2 ICESlvan pIP964; lane 6, 2nd-generation transconjugant UV202 ICESlvan pIP964; lane 7, recipient JH2-2; lane 8, recipient UV202. Vancomycin, streptomycin, and rifampin and fusidic acid resistance are indicated by plus signs opposite Van^r, Str^r, and Rif^r Fus^r, respectively. The presence of ICESlvan was verified by *vanB* PCR and visual inspection of the SmaI PFGE pattern, where ICESlvan insertion in a 240-kb band results in replacement of one of the 240-kb double bands with a 340-kb band, as shown previously for JH2-derived recipients (20). The presence of the pheromone-responsive plasmid pIP964 was verified by *rep_{PCF10}* and *agg* PCRs.

The *vanZ* gene, contributing to teicoplanin resistance, is normally located within the *vanA* gene cluster, whereas *vanB* usually only confers resistance to vancomycin. The functionality of the VanZ homologue was supported by Etest results showing MICs of 3 or 4 mg liter⁻¹ to teicoplanin (low-level resistance according to EU-

CAST clinical breakpoints for enterococci) for *S. lutetiensis* NEM760 and 5-F9 and the 5-F9 transconjugants, which was at least 3-fold higher than the MICs of the corresponding susceptible recipient strains (BM4105-RF and UV202 MIC, 0.5 mg liter⁻¹; JH2-2 MIC, 1 mg liter⁻¹; BM4110 pIP964 MIC, 0.38 mg liter⁻¹; and OG1-RF MIC, 0.25 mg liter⁻¹) (Table 1). The finding of this *vanZ* gene illustrates the dynamics and the potential of the *Streptococcus* and *Enterococcus* genomes to acquire and collect genes which may increase their ability for environmental adaptation.

The streptococcal ICES_{Sde3396} encodes resistance toward cadmium and arsenic (49). These resistance genes were not identified in ICESlvan. Rather, ORF81 to -84 encode an ABC transporter (ATPase and permease) and a putative two-component system (histidine kinase and response regulator) resembling proteins BceA, -B, -R, and -S (previously designated MbrABCD), which are involved in bacitracin resistance in *S. mutans* through active efflux of bacitracin (GenBank accession number AB078507) (62, 63). ORF81 encodes 249 amino acids with 48% identity to the original *S. mutans* BceA (250 amino acids) and 82% identity to a putative BceA of *S. mitis* (GenBank accession number EFN94736) (245 amino acids), ORF82 encodes 672 amino acids with 29% identity to the original *S. mutans* BceB (667 amino acids) and 67% identity to a putative ABC transporter permease of *S. mitis* (GenBank accession number EFN94737) (671 amino acids), ORF83 encodes 524 amino acids with >20% identity to BceS (249 amino acids) and 65% identity to a putative membrane protein with a histidine kinase domain of *S. mitis* (GenBank accession number EFN94738) (524 amino acids), and ORF84 encodes 198 amino acids with 24% identity to BceR (223 amino acids) and 75% identity to a putative response regulator of *S. mitis* (GenBank accession number EFN94739) (198 amino acids). Compared to the *bceABRS* locus of *S. mutans*, ICESlvan holds a different gene order of the putative response regulator and histidine kinase. The putative bacitracin locus of ICESlvan shows about 20% amino acid identity to components encoded by a *bcrRABD* locus conferring high-level bacitracin resistance in *E. faecalis*, which show yet another synteny with the regulator gene upstream from the ATPase, permease, and kinase genes (64). Phenotypic testing indeed shows at




FIG 4 (A) Operon into which ICESlvan has been integrated in *E. faecium* MM5-F9a compared to the corresponding region in the fully sequenced *E. faecalis* OG1-RF (accession number CP002621). Dark gray shading indicates regions of similarity between compared CDSs. The *tRNA* (uracil-5)-methyltransferase *rumA* disrupted by the insertion is colored light gray, and the insertion is indicated with a dotted line. (B) Sequence comparison of the insertion regions of ICESlvan in *E. faecium* MM5-F9a and *E. faecalis* OG5-F9a and the corresponding regions in their recipient strains, *E. faecium* BM4105-RF and *E. faecalis* OG1-RF, respectively. Vertical lines indicate identical nucleotides. Left- and right-end sequences of ICESlvan are shown in uppercase. Amino acid residues of CDSs that have been disrupted upon insertion of ICESlvan are marked (capital and boldface), together with their respective codons. #, stop codon.

least a 5-fold-increased MIC for bacitracin associated with transfer of ICES_l from *S. lutetiensis* 5-F9 to *E. faecium*. The *E. faecium* transconjugant MM5-F9a expressed a bacitracin MIC of ≥ 256 mg liter⁻¹, while the MIC of the corresponding *E. faecium* recipient BM4105-RF was 48 mg liter⁻¹, implying functionality of the putative bacitracin locus of ICES_l as a novel bacitracin resistance locus. On the other hand, the *E. faecalis* transconjugants and their corresponding recipients showed similar low or high MICs to bacitracin, except for the second-generation transconjugant JH2-2 ICES_l pIP964, which after transfer from the high-bacitracin-MIC background in *E. faecalis* BM4110 pIP964 showed at least a 16-fold-increased MIC compared to that of JH2-2 (Table 1). Undecaprenyl pyrophosphate phosphatase has recently been shown to account for the low-level resistance to bacitracin (MICs of 32 to 48 mg liter⁻¹) in both laboratory (JH2-2) and clinical (V583) strains of *E. faecalis* (65), while variants of the transferable *bcrRABD* locus may be responsible for the high-level bacitracin resistance (MIC ≥ 128 mg liter⁻¹) in enterococci (64, 66). Alternative primer combinations had to be used to confirm the bacitracin locus region of ICES_l (see Q38 in Table S1 in the supplemental material) in *E. faecalis* JH5-F9a and OG5-F9a compared to that in *E. faecium* MM5-F9a, indicating some sequence differences. However, sequencing *E. faecalis* JH5-F9a and OG5-F9a revealed identical nucleotides of the bacitracin locus (nt 85669 to 91794) compared to the sequence of MM5-F9a (GenBank accession number HE963029). Thus, host-related factors of *E. faecalis* and *E. faecium* may be a possible explanation for the lack of increased bacitracin MICs when ICES_l was introduced directly into *E. faecalis* strains with low-level resistance to bacitracin.

ICES_l expresses a *pezAT* TA system. ICES_l ORF75 and ORF76 are 93 and 89% identical at the nucleotide level to the *pezAT* plasmid maintenance system previously described in *S. pneumoniae* (67), *S. agalacticae* (57, 67), and *S. suis* (57). The *pezAT* gene cassette consists of two genes, encoding an epsilon antitoxin and a zeta toxin, respectively. Toxin-antitoxin systems are traditionally known as plasmid addiction systems that ensure stable maintenance of plasmids in a bacterial population (68). However, during recent years, these gene loci have been found on MGEs and shown to promote the maintenance of ICE elements in *V. cholerae* (69, 70). Experimental evidence also indicated that TA systems are involved in the stress response, enabling the cell to survive hostile growth conditions (71, 72). Analyses of the DNA sequence upstream from the *pezAT* locus showed the presence of a putative -10 promoter sequence (TATAAT) 34 bp upstream from an indicated start codon and a -35 promoter sequence (GT GCGTT) 19 bp from the putative Pribnow box. In addition, a putative ribosome-binding site (AGGAG) was shown 12 bp upstream from the putative start codon. All except the -35 sequence were identical to the *S. pneumoniae* sequences (67). RT-PCR analyses revealed the expression of both *pezAT* genes (data not shown) in all of the donor and transconjugant strains tested. Figure 5 shows the PCR products of both genes. Moreover, both genes were detected in a single transcript (Fig. 5), confirming that *pezA* and *pezT* constitute an operon, as in *S. pneumoniae* (67). PCR products were not detected in the absence of reverse transcriptase or in the recipient strain (BM4105-RF). All sequences were identical to the sequence found in our *E. faecium* transconjugant. These observations suggest that this is a functional toxin-antitoxin system




FIG 5 Expression of *pezAT* toxin-antitoxin loci in original donor *S. lutetiensis* 5-F9, transconjugants *E. faecalis* JH5-F9a, *E. faecium* MM5-F9a, and *E. faecalis* OG5-F9a, and recipient strain *E. faecium* BM4105-RF tested with RT-PCR (+RT). As DNA contamination controls, reaction mixtures with no reverse transcriptase added were included (-RT). L, kb ladder.

that may contribute to the stabilization of ICES_l after integration into new hosts.

In conclusion, our report describes the complete genetic structure of a 94-kb element, named ICES_l, originally detected in *S. lutetiensis* from veal calf feces in the Netherlands. A variant of ICES_l is confirmed for an *S. lutetiensis* isolate from a patient in France. ICES_l encodes glycopeptide and bacitracin resistance loci, a putative bacteriophage exclusion system, a putative virulence gene, and components necessary for conjugative transfer, as well as a toxin-antitoxin stabilization system. This element is a novel ICE, since it forms a circular intermediate, is self-transferable from streptococci to enterococci, and integrates into the chromosome in a site-specific manner.

ACKNOWLEDGMENT

This work was supported by research grants from the Norwegian Research Council (projects no. 165997 and 183653/S19), Northern Norway Regional Health Authority Medical Research Program, and the European Commission (LSHE-CT-2007-03410 ACE).

We thank Tracy Munthali Lunde, Christina Borch-Pedersen, and Trine Tessem for their excellent technical assistance. We also thank Eirik W. Lundblad and Johanna E. Sollid for critical reading of the manuscript.

REFERENCES

1. Van Schaik W, Top Riley JD, Boekhorst J, Vrijenhoek J, Schapendonk C, Hendrickx A, Nijman I, Bonten M, Tettelin H, Willems R. 2010. Pyrosequencing-based comparative genome analysis of the nosocomial pathogen *Enterococcus faecium* and identification of a large transferable pathogenicity island. *BMC Genomics* 11:239. doi:10.1186/1471-2164-11-239.
2. Lam MM, Seemann T, Bulach DM, Gladman SL, Chen H, Haring V, Moore RJ, Ballard S, Grayson ML, Johnson PD, Howden BP, Stinear TP. 2012. Comparative analysis of the first complete *Enterococcus faecium* genome. *J. Bacteriol.* 194:2334–2341.
3. van Schaik W, Willems RJ. 2010. Genome-based insights into the evolution of enterococci. *Clin. Microbiol. Infect.* 16:527–532.
4. Paulsen IT, Banerjee L, Myers GSA, Nelson KE, Seshadri R, Read TD, Fouts DE, Eisen JA, Gill SR, Heidelberg JF, Tettelin H, Dodson RJ, Umayam L, Brinkac L, Beanan M, Daugherty S, DeBoy RT, Durkin S, Kolonay J, Madupu R, Nelson W, Vamathevan J, Tran B, Upton J, Hansen T, Shetty J, Khouri H, Utterback T, Radune D, Ketchum KA, Dougherty BA, Fraser CM. 2003. Role of mobile DNA in the evolution of vancomycin-resistant *Enterococcus faecalis*. *Science* 299:2071–2074.
5. Uttley AH, Collins CH, Naidoo J, George RC. 1988. Vancomycin-resistant enterococci. *Lancet* i:57–58.
6. Leclercq R, Derlot E, Duval J, Courvalin P. 1988. Plasmid-mediated resistance to vancomycin and teicoplanin in *Enterococcus faecium*. *N. Engl. J. Med.* 319:157–161.
7. Werner G, Coque TM, Hammerum AM, Hope R, Hryniewicz W, Johnson A, Klare I, Kristinsson KG, Leclercq R, Lester CH, Lillie M, Novais C, Olsson-Liljequist B, Peixe LV, Sadowy E, Simonsen GS, Top Vuopio-Varkila JJ, Willems RJ, Witte W, Woodford N. 2008. Emer-

- gence and spread of vancomycin resistance among enterococci in Europe. *Euro Surveill.* 13:1–11.
8. Galloway-Pena JR, Nallapareddy SR, Arias CA, Eliopoulos GM, Murray BE. 2009. Analysis of clonality and antibiotic resistance among early clinical isolates of *Enterococcus faecium* in the United States. *J. Infect. Dis.* 200:1566–1573.
 9. Guardabassi L, Agersø Y. 2006. Genes homologous to glycopeptide resistance *vanA* are widespread in soil microbial communities. *FEMS Microbiol. Lett.* 259:221–225.
 10. Howden BP, Holt KE, Lam MM, Seemann T, Ballard S, Coombs GW, Tong SY, Grayson ML, Johnson PD, Stinear TP. 2013. Genomic insights to control the emergence of vancomycin-resistant enterococci. *mBio* 4(4): e00412–13. doi:10.1128/mBio.00412-13.
 11. Ishii Y, Ohno A, Kashitani S, Iwata M, Yamaguchi K. 1996. Identification of VanB-type vancomycin resistance in *Enterococcus gallinarum* from Japan. *J. Infect. Chemother.* 2:102–105.
 12. Simonsen GS, Andersen BM, Digranes A, Harthug S, Jacobsen T, Lingaas E, Natas OB, Olsvik Ø, Ringertz SH, Skulberg A, Syversen G, Sundsfjord A. 1998. Low faecal carrier rate of vancomycin resistant enterococci in Norwegian hospital patients. *Scand. J. Infect. Dis.* 30:465–468.
 13. Liassine N, Frei R, Jan I, Auckenthaler R. 1998. Characterization of glycopeptide-resistant enterococci from a Swiss hospital. *J. Clin. Microbiol.* 36:1853–1858.
 14. Schooneveldt JM, Marriott RK, Nimmo GR. 2000. Detection of a *vanB* determinant in *Enterococcus gallinarum* in Australia. *J. Clin. Microbiol.* 38:3902.
 15. Torres C, Tenorio C, Portillo A, Garcia M, Martinez C, Del CR, Ruiz-Larrea F, Zarazaga M. 2003. Intestinal colonization by *vanA*- or *vanB2*-containing enterococcal isolates of healthy animals in Spain. *Microb. Drug Resist.* 9(Suppl 1):S47–S52.
 16. Jenney A, Franklin C, Liolios L, Spelman D. 2000. *Enterococcus durans vanB*. *J. Antimicrob. Chemother.* 46:515.
 17. Ameri S, Talebi M, Rahimi F, Pourshafie MR, Ebrahimipour G. 2009. The homogeneity of *vanB* gene cluster among enterococcal isolates in Iran. *Lett. Appl. Microbiol.* 48:157–161.
 18. Roberts MC, Soge OO, Giardino MA, Mazengia E, Ma G, Meschke JS. 2009. Vancomycin-resistant *Enterococcus* spp. in marine environments from the West Coast of the U. S. A. *J. Appl. Microbiol.* 107:300–307.
 19. Mevius D, Devriese L, Butaye P, Vandamme P, Verschure M, Veldman K. 1998. Isolation of glycopeptide resistant *Streptococcus gallolyticus* strains with *vanA*, *vanB*, and both *vanA* and *vanB* genotypes from faecal samples of veal calves in The Netherlands. *J. Antimicrob. Chemother.* 42:275–276.
 20. Dahl KH, Sundsfjord A. 2003. Transferable *vanB2* Tn5382-containing elements in fecal streptococcal strains from veal calves. *Antimicrob. Agents Chemother.* 47:2579–2583.
 21. Poyart C, Quesne G, Coulon S, Berche P, Trieu-Cuot P. 1998. Identification of streptococci to species level by sequencing the gene encoding the manganese-dependent superoxide dismutase. *J. Clin. Microbiol.* 36: 41–47.
 22. Poyart C, Quesne G, Trieu-Cuot P. 2002. Taxonomic dissection of the *Streptococcus bovis* group by analysis of manganese-dependent superoxide dismutase gene (*sodA*) sequences: reclassification of '*Streptococcus infantarius* subsp. *coli*' as *Streptococcus lutetiensis* sp. nov. and of *Streptococcus bovis* biotype 11.2 as *Streptococcus pasteurianus* sp. nov. *Int. J. Syst. Evol. Microbiol.* 52:1247–1255.
 23. Stinear TP, Olden DC, Johnson PD, Davies JK, Grayson ML. 2001. Enterococcal *vanB* resistance locus in anaerobic bacteria in human faeces. *Lancet* 357:855–856.
 24. Ballard SA, Pertile KK, Lim M, Johnson PDR, Grayson ML. 2005. Molecular characterization of *vanB* elements in naturally occurring gut anaerobes. *Antimicrob. Agents Chemother.* 49:1688–1694.
 25. Ballard SA, Grabsch EA, Johnson PDR, Grayson ML. 2005. Comparison of three PCR primer sets for identification of *vanB* gene carriage in feces and correlation with carriage of vancomycin-resistant enterococci: interference by *vanB*-containing anaerobic bacilli. *Antimicrob. Agents Chemother.* 49:77–81.
 26. Domingo MC, Huletsky A, Giroux R, Boissinot K, Picard FJ, Lebel P, Ferraro MJ, Bergeron MG. 2005. High prevalence of glycopeptide resistance genes *vanB*, *vanD*, and *vanG* not associated with enterococci in human fecal flora. *Antimicrob. Agents Chemother.* 49:4784–4786.
 27. Marvaud JC, Mory F, Lambert T. 2011. *Clostridium clostridioforme* and *Atopobium minutum* clinical isolates with *vanB*-type resistance in France. *J. Clin. Microbiol.* 49:3436–3438.
 28. Evers S, Courvalin P. 1996. Regulation of VanB-type vancomycin resistance gene expression by the VanS(B)-VanR(B) two-component regulatory system in *Enterococcus faecalis* V583. *J. Bacteriol.* 178:1302–1309.
 29. Dahl KH, Simonsen GS, Olsvik Ø, Sundsfjord A. 1999. Heterogeneity in the *vanB* gene cluster of genomically diverse clinical strains of vancomycin-resistant enterococci. *Antimicrob. Agents Chemother.* 43:1105–1110.
 30. Garnier F, Taourit S, Glaser P, Courvalin P, Galimand M. 2000. Characterization of transposon Tn1549, conferring VanB-type resistance in *Enterococcus* spp. *Microbiology* 146:1481–1489.
 31. Gold HS, Unal S, Cercenado E, Thauvin-Eliopoulos C, Eliopoulos GM, Wennersten CB, Moellering RC, Jr. 1993. A gene conferring resistance to vancomycin but not teicoplanin in isolates of *Enterococcus faecalis* and *Enterococcus faecium* demonstrates homology with *vanB*, *vanA*, and *vanC* genes of enterococci. *Antimicrob. Agents Chemother.* 37:1604–1609.
 32. Patel R, Uhl JR, Kohner P, Hopkins MK, Steckelberg JM, Kline B, Cockerill FR, III. 1998. DNA sequence variation within *vanA*, *vanB*, *vanC-1*, and *vanC-2/3* genes of clinical *Enterococcus* isolates. *Antimicrob. Agents Chemother.* 42:202–205.
 33. Dahl KH, Lundblad EW, Røkenes TP, Olsvik Ø, Sundsfjord A. 2000. Genetic linkage of the *vanB2* gene cluster to Tn5382 in vancomycin-resistant enterococci and characterization of two novel insertion sequences. *Microbiology* 146:1469–1479.
 34. McGregor KF, Nolan C, Young HK, Palepou MF, Tysall L, Woodford N. 2001. Prevalence of the *vanB2* gene cluster in *vanB* glycopeptide-resistant enterococci in the United Kingdom and the Republic of Ireland and its association with a Tn5382-like element. *Antimicrob. Agents Chemother.* 45:367–368.
 35. Zheng B, Tomita H, Inoue T, Ike Y. 2009. Isolation of VanB-type *Enterococcus faecalis* strains from nosocomial infections: first report of the isolation and identification of the pheromone-responsive plasmids pMG2200, encoding VanB-type vancomycin resistance and a Bac41-type bacteriocin, and pMG2201, encoding erythromycin resistance and cytolysin (Hly/Bac). *Antimicrob. Agents Chemother.* 53:735–747.
 36. Demertzi E, Palepou MF, Kaufmann ME, Avlami A, Woodford N. 2001. Characterisation of *vanA* and *vanB* elements from glycopeptide-resistant *Enterococcus faecium* from Greece. *J. Med. Microbiol.* 50:682–687.
 37. Hanrahan J, Hoyer C, Rice LB. 2000. Geographic distribution of a large mobile element that transfers ampicillin and vancomycin resistance between *Enterococcus faecium* strains. *Antimicrob. Agents Chemother.* 44: 1349–1351.
 38. Valdezate S, Labayru C, Navarro A, Mantecon MA, Ortega M, Coque TM, Garcia M, Saez-Nieto JA. 2009. Large clonal outbreak of multidrug-resistant CC17 ST17 *Enterococcus faecium* containing Tn5382 in a Spanish hospital. *J. Antimicrob. Chemother.* 63:17–20.
 39. Launay A, Ballard SA, Johnson P, Grayson M, Lambert T. 2006. Transfer of vancomycin resistance transposon Tn1549 from *Clostridium symbiosum* to *Enterococcus* spp. in the gut of gnotobiotic mice. *Antimicrob. Agents Chemother.* 50:1054–1062.
 40. Carias LL, Rudin SD, Donskey CJ, Rice LB. 1998. Genetic linkage and cotransfer of a novel, *vanB*-containing transposon (Tn5382) and a low-affinity penicillin-binding protein 5 gene in a clinical vancomycin-resistant *Enterococcus faecium* isolate. *J. Bacteriol.* 180:4426–4434.
 41. Dahl KH, Røkenes TP, Lundblad EW, Sundsfjord A. 2003. Nonconjugative transposition of the *vanB*-containing Tn5382-like element in *Enterococcus faecium*. *Antimicrob. Agents Chemother.* 47:786–789.
 42. Poyart C, Pierre C, Quesne G, Pron B, Berche P, Trieu-Cuot P. 1997. Emergence of vancomycin resistance in the genus *Streptococcus*: characterization of a *vanB* transferable determinant in *Streptococcus bovis*. *Antimicrob. Agents Chemother.* 41:24–29.
 43. Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, Barrell B. 2000. Artemis: sequence visualization and annotation. *Bioinformatics* 16:944–945.
 44. Bendtsen JD, Nielsen H, von Heijne G, Brunak S. 2004. Improved prediction of signal peptides: SignalP 3.0. *J. Mol. Biol.* 340:783–795.
 45. Carver TJ, Rutherford KM, Berriman M, Rajandream MA, Barrell BG, Parkhill J. 2005. ACT: the Artemis Comparison Tool. *Bioinformatics* 21:3422–3423.
 46. Poyart C, Trieu-Cuot P. 1994. Heterogeneric conjugal transfer of the pheromone-responsive plasmid pIP964 (IncHly1) of *Enterococcus faecalis*

- in the apparent absence of pheromone induction. *FEMS Microbiol. Lett.* 122:173–179.
47. Jensen LB, Garcia-Migura L, Valenzuela AJ, Lohr M, Hasman H, Aarestrup FM. 2010. A classification system for plasmids from enterococci and other Gram-positive bacteria. *J. Microbiol. Methods* 80:25–43.
 48. Saeedi B, Hallgren A, Jonasson J, Nilsson LE, Hanberger H, Isaksson B. 2002. Modified pulsed-field gel electrophoresis protocol for typing of enterococci. *APMIS* 110:869–874.
 49. Davies MR, Shera J, Van Domselaar GH, Sriprakash KS, McMillan DJ. 2009. A novel integrative conjugative element mediates genetic transfer from group G *Streptococcus* to other beta-hemolytic streptococci. *J. Bacteriol.* 191:2257–2265.
 50. Alvarez-Martinez CE, Christie PJ. 2009. Biological diversity of prokaryotic type IV secretion systems. *Microbiol. Mol. Biol. Rev.* 73:775–808.
 51. Grohmann E, Muth G, Espinosa M. 2003. Conjugative plasmid transfer in Gram-positive bacteria. *Microbiol. Mol. Biol. Rev.* 67:277–301.
 52. Layec S, Decaris B, Leblond-Bourget N. 2008. Diversity of Firmicutes peptidoglycan hydrolases and specificities of those involved in daughter cell separation. *Res. Microbiol.* 159:507–515.
 53. McLeod SM, Burrus V, Waldor MK. 2006. Requirement for *Vibrio cholerae* integration host factor in conjugative DNA transfer. *J. Bacteriol.* 188:5704–5711.
 54. Manson JM, Hancock LE, Gilmore MS. 2010. Mechanism of chromosomal transfer of *Enterococcus faecalis* pathogenicity island, capsule, antimicrobial resistance, and other traits. *Proc. Natl. Acad. Sci. U. S. A.* 107:12269–12274.
 55. O'Connor L, Coffey A, Daly C, Fitzgerald GF. 1996. AbiG, a genotypically novel abortive infection mechanism encoded by plasmid pCI750 of *Lactococcus lactis* subsp. *cremoris* UC653. *Appl. Environ. Microbiol.* 62:3075–3082.
 56. Chopin MC, Chopin A, Bidnenko E. 2005. Phage abortive infection in lactococci: variations on a theme. *Curr. Opin. Microbiol.* 8:473–479.
 57. Holden MT, Hauser H, Sanders M, Ngo TH, Cherevach I, Cronin A, Goodhead I, Mungall K, Quail MA, Price C, Rabbinowitsch E, Sharp S, Croucher NJ, Chieu TB, Mai NT, Diep TS, Chinh NT, Kehoe M, Leigh JA, Ward PN, Dowson CG, Whatmore AM, Chanter N, Iversen P, Gottschalk M, Slater JD, Smith HE, Spratt BG, Xu J, Ye C, Bentley S, Barrell BG, Schultsz C, Maskell DJ, Parkhill J. 2009. Rapid evolution of virulence and drug resistance in the emerging zoonotic pathogen *Streptococcus suis*. *PLoS One* 4:e6072. doi:10.1371/journal.pone.0006072.
 58. Hendrickx APA, Willems RJJ, Bonten MJM, van Schaik W. 2009. LPxTG surface proteins of enterococci. *Trends Microbiol.* 17:423–430.
 59. Malamud D, Appelbaum B, Kline R, Golub EE. 1981. Bacterial aggregating activity in human saliva: comparisons of bacterial species and strains. *Infect. Immun.* 31:1003–1006.
 60. Rundegren J. 1986. Calcium-dependent salivary agglutinin with reactivity to various oral bacterial species. *Infect. Immun.* 53:173–178.
 61. Demuth DR, Berthold P, Leboy PS, Golub EE, Davis CA, Malamud D. 1989. Saliva-mediated aggregation of *Enterococcus faecalis* transformed with a *Streptococcus sanguis* gene encoding the SSP-5 surface antigen. *Infect. Immun.* 57:1470–1475.
 62. Tsuda H, Yamashita Y, Shibata Y, Nakano Y, Koga T. 2002. Genes involved in bacitracin resistance in *Streptococcus mutans*. *Antimicrob. Agents Chemother.* 46:3756–3764.
 63. Ouyang J, Tian XL, Versey J, Wishart A, Li YH. 2010. The BceABRS four-component system regulates the bacitracin-induced cell envelope stress response in *Streptococcus mutans*. *Antimicrob. Agents Chemother.* 54:3895–3906.
 64. Manson JM, Keis S, Smith JM, Cook GM. 2004. Acquired bacitracin resistance in *Enterococcus faecalis* is mediated by an ABC transporter and a novel regulatory protein, BcrR. *Antimicrob. Agents Chemother.* 48:3743–3748.
 65. Shaaly A, Kalamorz F, Gebhard S, Cook GM. 2013. Undecaprenyl pyrophosphate phosphatase confers low-level resistance to bacitracin in *Enterococcus faecalis*. *J. Antimicrob. Chemother.* 68:1583–1593.
 66. Matos R, Pinto VV, Ruivo M, Lopes MF. 2009. Study on the dissemination of the *bcrABDR* cluster in *Enterococcus* spp. reveals that the BcrAB transporter is sufficient to confer high-level bacitracin resistance. *Int. J. Antimicrob. Agents* 34:142–147.
 67. Khoo SK, Loll B, Chan WT, Shoeman RL, Ngoo L, Yeo CC, Meinhart A. 2007. Molecular and structural characterization of the PezAT chromosomal toxin-antitoxin system of the human pathogen *Streptococcus pneumoniae*. *J. Biol. Chem.* 282:19606–19618.
 68. Gerdes K, Christensen SK, Lobner-Olesen A. 2005. Prokaryotic toxin-antitoxin stress response loci. *Nat. Rev. Microbiol.* 3:371–382.
 69. Wozniak RA, Waldor MK. 2009. A toxin-antitoxin system promotes the maintenance of an integrative conjugative element. *PLoS Genet.* 5:e1000439. doi:10.1371/journal.pgen.1000439.
 70. Szekeres S, Dauti M, Wilde C, Mazel D, Rowe-Magnus DA. 2007. Chromosomal toxin-antitoxin loci can diminish large-scale genome reductions in the absence of selection. *Mol. Microbiol.* 63:1588–1605.
 71. Christensen SK, Mikkelsen M, Pedersen K, Gerdes K. 2001. RelE, a global inhibitor of translation, is activated during nutritional stress. *Proc. Natl. Acad. Sci. U. S. A.* 98:14328–14333.
 72. Christensen SK, Pedersen K, Hansen FG, Gerdes K. 2003. Toxin-antitoxin loci as stress-response-elements: ChpAK/MazF and ChpBK cleave translated RNAs and are counteracted by tmRNA. *J. Mol. Biol.* 332:809–819.
 73. Dunny GM, Brown BL, Clewell DB. 1978. Induced cell aggregation and mating in *Streptococcus faecalis*: evidence for a bacterial sex pheromone. *Proc. Natl. Acad. Sci. U. S. A.* 75:3479–3483.
 74. Jacob AE, Hobbs SJ. 1974. Conjugal transfer of plasmid-borne multiple antibiotic resistance in *Streptococcus faecalis* var. *zymogenes*. *J. Bacteriol.* 117:360–372.
 75. Yagi Y, Clewell DB. 1980. Recombination-deficient mutant of *Streptococcus faecalis*. *J. Bacteriol.* 143:966–970.
 76. Courvalin P, Carlier C. 1986. Transposable multiple antibiotic resistance in *Streptococcus pneumoniae*. *Mol. Gen. Genet.* 205:291–297.