

1 **-Congenital anomalies of the kidney and the urinary tract: a Murmansk County Birth**

2 **Registry study**

3 **Running title: Congenital urinary anomalies in Murmansk County**

4 Vitaly A. Postoev<sup>1,2</sup>, Andrej M. Grjibovski<sup>2-4</sup>, Anton A. Kovalenko<sup>1,2</sup>, Erik Eik Anda<sup>1</sup>, Evert

5 Nieboer<sup>5</sup>, Jon Øyvind Odland<sup>1,6</sup>

6 **Full affiliations**

7 <sup>1</sup>- Department of Community Medicine, UiT- The Arctic University of Norway, Tromsø, Norway

8 <sup>2</sup>- International School of Public Health, Northern State Medical University, Arkhangelsk,

9 Russia

10 <sup>3</sup>- Department of International Public Health, Norwegian Institute of Public Health, Oslo,

11 Norway

12 <sup>4</sup>- Department of Preventive Medicine, International Kazakh-Turkish University, Turkestan,

13 Kazakhstan

14 <sup>5</sup>- Department of Biochemistry and Biomedical Sciences, McMaster University, Hamilton, ON,

15 Canada

16 <sup>6</sup>- School of Health Systems and Public Health, Faculty of Health Sciences, University of

17 Pretoria, South Africa

18  
19 **Corresponding author**

20 Vitaly Postoev:

21 Mail address: 163061 Troickij av, 51 NSMU, ISPHA, office 1252

22 e-mail: [vpo000@post.uit.no](mailto:vpo000@post.uit.no)

23 tel. +79214704363

24  
25  
26  
27  
28

29 **Abstract**

30 **Background**

31 Congenital anomalies of the kidney and the urinary tract (CAKUTs) are relatively common birth  
32 defects. The combined prevalence in Europe was 3.3 per 1000 in 2012. The risk factors for these  
33 anomalies are not clearly identified. The aims of our study were to calculate the birth  
34 prevalences of urinary malformations in Murmansk County during 2006-2011 and to investigate  
35 related prenatal risk factors.

36 **Methods**

37 The Murmansk County Birth Registry was the primary source of information and our study  
38 included 50 936 singletons in the examination of structure, prevalence and proportional  
39 distribution of CAKUTs. The multivariate analyses of risk factors involved 39 322 newborns.

40 **Results**

41 The prevalence of CAKUTs was 4.0 per 1000 newborns [95%CI: 3.4-4.5] and did not change  
42 during the study period. The most prevalent malformation was congenital hydronephrosis  
43 (14.2% of all cases). Diabetes mellitus or gestational diabetes [OR = 4.77, 95%CI: 1.16-19.65],  
44 acute infections while pregnant [OR = 1.83, 95%CI: 1.14-2.94], the use of medication during  
45 pregnancy [OR = 2.03, 95%CI: 1.44-2.82], and conception during the summer [OR = 1.75,  
46 95%CI 1.15-2.66] were significantly associated with higher risk of CAKUTs.

47 **Conclusions**

48 The overall four-fold enhancement of the occurrence of urinary malformations in Murmansk  
49 County for the 2006-2011 period showed little annual dependence. During pregnancy, use of  
50 medications, infections, pre-existing diabetes mellitus or gestational diabetes were associated  
51 with increased risk of these anomalies, as was conception during summer. Our findings have  
52 direct applications in improving prenatal care in Murmansk County and establishing targets for  
53 prenatal screening and women's consultations.

54 **Key words:** Congenital anomalies of the kidney and the urinary tract, risk factors, Russia, the  
55 Murmansk County Birth Registry

56

## 57 **Introduction**

58 Congenital anomalies of the kidney and the urinary tract (CAKUTs) constitute one of the most  
59 common groups of birth defects. In 2012, the prevalence in Europe was 3.3 per 1000 births  
60 (EUROCAT, 2013). They represent a highly heterogeneous group, with a few being  
61 incompatible with life while minor abnormalities can be asymptomatic for long periods of time  
62 (Barakat and Drougas, 1991). Anomalies of the kidney are more common than those of the lower  
63 urinary tract and often lead to chronic renal failure (Barakat and Drougas, 1991). Collectively,  
64 CAKUTs account for 30–50% of cases of renal failure among children worldwide (dos Santos  
65 Junior et al., 2014).

66 The development of human kidneys is completed by weeks 34-36 of gestation (Hei and Yi,  
67 2014), and thus are amenable to teratogens during the whole pregnancy. The role of maternal and  
68 environmental factors in development of CAKUTs is not well established. Previous studies have  
69 reported associations of CAKUTs with a number of maternal factors, including: pre-existing  
70 diabetes mellitus (DM) or gestational diabetes (GD) (Kalter, 2003; Sheffield et al., 2002;  
71 Shnorhavorian et al., 2011); febrile illness during pregnancy (Abe et al., 2003), advance age  
72 (Shnorhavorian et al., 2011); renal disease (Shnorhavorian et al., 2011); as well as medication  
73 use (especially non-steroid anti-inflammatory drugs and angiotensin converting enzyme  
74 inhibitors; Boubred et al., 2006). Studies of the effect on CAKUTs from tobacco use and alcohol  
75 consumption have yielded controversial results in that some show negative associations, while  
76 others do not (Kallen, 1997; Li et al., 1996; Shnorhavorian et al., 2011). The evidence for  
77 seasonal variation is also contradictory in that increased risk of CAKUTs has been reported for  
78 both newborns conceived (Luteijn et al., 2014) or delivered during the summer months (Feldman  
79 et al., 2009).

80 The prevalence of CAKUTs in Russia is estimated at the population and country level through  
81 mandatory reporting of birth defects of its three most severe forms, namely renal agenesis or  
82 dysgenesis, epispadias and bladder extrophy. The total prevalence of these three forms in Russia  
83 was 0.1 per 1000 newborns in 2011 (RIPCS, 2011). Studies of urinary birth defects in different  
84 regions of Russia report prevalences of all CAKUTs that vary from 1.1 per 1000 in Saratov to  
85 5.7 per 1000 in Izhevsk (Andreeva, 2008). Moreover, CAKUTs represented 10-15% of all birth  
86 defects (Andreeva, 2008; Trefilov, 2006) and were responsible for 65% of cases of chronic renal  
87 failure (Molchanova, 2004). By contrast, we could not identify any Russian studies that  
88 explored risk factors for this group of congenital defects.

89 The population-based birth registry in Murmansk County recorded all anomalies diagnosed from  
90 22 weeks of gestation until hospital discharge and thus provides unique opportunities for  
91 research. Previously we reported a high prevalence of musculoskeletal and urinary tract  
92 malformations in the town of Monchegorsk, which is located in Murmansk County (Postoev et  
93 al., 2015a). The overall perinatal prevalence of CAKUTs in Monchegorsk increased from 4.4 per  
94 1000 in 1993-2002 (that of all BD was 38.7 per 1000) to 19.1 in 2003-2011 (46.3 per 1000 for  
95 all BD). Thus, the CAKUTs accounted for 41% of all registered birth defects. An increase in the  
96 prevalence of CAKUTs has also been reported by the European Surveillance of Congenital  
97 Anomalies (EUROCAT), but to a lesser extent (from 1.5/1000 in 1980 to 3.6/1000 in 2011)  
98 (EUROCAT, 2013). The cystic kidney prevalence reported by the International Clearinghouse  
99 for Birth Defects Surveillance and Research (ICBDSR) for Norway increased from 0.1/1000 in  
100 1974 to 0.6/1000 in 2010 (ICBDSR, 2013). This trend could partly be explained by  
101 implementation of prenatal diagnostics in 1990s in Western countries and in the early 2000s in  
102 Russia. Pertinent visualisation allowed 81.8% of CAKUTs to be detected prior to birth in Europe  
103 (Wiesel et al., 2005), while in the Murmansk region it was 42% (Postoev et al., 2015b).  
104 Nevertheless, the exact reasons that explain the observed increases remain unknown. Factors  
105 such as smoking and alcohol consumption during pregnancy and genital/urinary infection may  
106 have contributed, as their prevalences increased considerably in Russia during a period of  
107 transition.

108 The aims of this study were to calculate the birth prevalence of CAKUTs in Murmansk County  
109 and to investigate the perinatal risk factors for their occurrence.

## 110 **Materials and Methods**

### 111 SUBJECTS AND DATA SOURCES

112 The study population of this registry-based study included all newborns delivered in Murmansk  
113 County and registered in the MCBR in the period 2006-2011 (total N=52 086). The MCBR was  
114 also the primary source of information about associated risk factors.

115 Murmansk County is situated in the Kola Peninsula of Northwest Russia, of which a significant  
116 part is located above the Arctic Circle. The region has 766 300 inhabitants, of whom 92.6% are  
117 urbanites (Murmanskstat, 2015a). Its population has decreased by 30 000 during the last five  
118 years. There were 9017 newborns in the County in 2014 and corresponds to a birth rate of  
119 11.7/1000, which is similar to that reported for the whole of Russia (Murmanskstat, 2015b).

120 The MCBR was established in 2005 and includes all births in Murmansk County as of January 1,  
121 2006. The registered data were systematically collected in the county's 15 delivery departments  
122 and included the following specific sources: the medical histories of the mothers; delivery  
123 journals and records; birth records; and interviews with the mothers conducted by a physician or  
124 midwife. The MCBR database contains data about paternal age, ethnicity, residence, and  
125 occupation; maternal civil status, education, height and weight; data about previous pregnancies  
126 and diseases before pregnancy; course of the current pregnancy (including complications, intake  
127 of supplements, smoking habits, any abuse, diseases during pregnancy, and results of prenatal  
128 screening); delivery type and post-delivery maternal complications; and the status of the  
129 newborn, his/her gestational age and any medical conditions during the early neonatal period.  
130 These recorded sources have previously been described in detail (Anda et al., 2008; Anda et al.,  
131 2011), and their quality was determined to be satisfactory for epidemiological research.

#### 132 STUDY VARIABLES

133 The diagnoses of congenital malformations were based on the International Statistical  
134 Classification of Diseases and Related Health Problems 10<sup>th</sup> revision (ICD-10), which includes  
135 birth defects in Chapter XVII. The CAKUTs fall under the codes Q60-Q64. In spite of their  
136 different clinical manifestations, CAKUTs have a common mesoderm origin, and manifest  
137 during similar critical periods of embryogenesis (Moore et al., 2003; dos Santos Junior et al,  
138 2014). Consequently, investigation of common risk factors for this group of malformations  
139 seems pertinent and any malformation coded as a CAKUT was thus included in our analysis.  
140 Diagnoses were recorded up to the departure of the newborns from the birth clinics.

141 Prenatal ultrasound screening in Russia was established by national law in 2000, and includes  
142 ultrasonography at gestational ages of 10-14, 20-24 and 30-32 weeks and biochemical tests of  
143 blood alpha-fetoprotein and chorionic gonadotropin at gestational age 11-14 weeks (Ministry of  
144 Health of Russian Federation, 2000). It is offered by obstetricians to every pregnant woman in  
145 compliance with a national law, and is free of charge. Postnatal ultrasonography supplemented  
146 routine examinations and any prenatal suspicion of an anomaly of the urinary system was  
147 confirmed by examination after birth. Fetal deaths were followed up by autopsy examinations.

148 Potential risk factors were selected based on a detailed literature review and pertinent  
149 information was taken directly from the MCBR database. All continuous variables were recoded  
150 as categorical. Maternal age was divided into three groups, specifically less than 18, 18-35, and  
151 older than 35 years, while the father's age was dichotomized with a cut-off point at 35 years. The

152 international classification for maternal body mass index (BMI) was adopted, with underweight  
153 defined as  $<18.5 \text{ kg/m}^2$ ,  $18.5\text{-}24.9 \text{ kg/m}^2$  for normal weight, and  $\geq 25 \text{ kg/m}^2$  as being overweight  
154 or obese. BMIs were calculated according to maternal anthropometry at the first antenatal visit.  
155 Yes/no dichotomization with the absence of a factor as the reference category was adopted for  
156 the following: occurrence or use during pregnancy of DM or GD, consumption of multivitamins  
157 or folic acid intake, cigarette smoking and infections; and evidence of alcohol abuse, chronic  
158 genital or urinary infections before pregnancy. The season of conception was calculated using  
159 the first day of the last menses.

## 160 DATA ANALYSES

161 We used the registered two- and three-digit level ICD-10 codes to analyse the classification of  
162 CAKUTs, their prevalences and proportional distributions. Newborns with multiple  
163 malformations were included in the numerator of these estimates when a CAKUT was present  
164 among the diagnoses. All prevalence estimates are presented with their 95% confidence intervals  
165 (CI). There were 52 806 pregnancy outcomes registered in the MCBR, of which 1 313 were  
166 excluded due to missing or incorrect information about gestational age and 99 due to missing  
167 information about birth defects. All cases of multiple pregnancy ( $n = 458$ ) were also precluded  
168 from the analyses. Thus, 50 936 singletons were included in the investigation of structure,  
169 prevalence and proportional distribution of CAKUTs.

170 The prevalence of CAKUTs was estimated for newborns with and without a consideration of  
171 possible risk factors (or for maternal characteristics with multiple values), and compared  
172 statistically using the chi-square test. Adjusted malformation risks associated with the selected  
173 predictors were analysed by multiple logistic regression using CAKUTs as a binary outcome.  
174 Risk ratios were approximated by odds ratios extracted from the regression model. There were  
175 39 322 newborns (185 cases) without any missing information for the variables of interest, and  
176 these were included in the logistic regression analysis. Detailed information about missing  
177 variables is presented in Figure 1. The final regression model was established by employing the  
178 backward stepwise regression model, using the likelihood ratio method for inclusion of all  
179 studied factors and probability criteria for removal of 0.1. All analyses were performed using  
180 IBM SPSS 21.0 software package.

## 181 ETHICAL CONSIDERATIONS

182 The Committee for Research Ethics at the Northern State Medical University (Arkhangelsk,  
183 Russia) and the Regional Committee for medical and health research ethics (Tromsø, Norway)  
184 approved the current study.

## 185 **Results**

186 There were 203 registered newborns with CAKUTs in Murmansk County in 2006-2011. The  
187 prevalence at birth was 4.0 per 1000 newborns (95%CI 3.4-4.5) and it did not change over time  
188 (p for linear trend = 0.26). Nevertheless, there was some variation in the birth prevalence: from  
189 2.4 per 1000 newborns (95% CI 1.3-3.4) in 2006 to 5.6 per 1000 newborns (95% CI 4.0-7.1) in  
190 2008.

191 More than half of the malformations were diagnosed as “other congenital anomalies of kidney”.  
192 Congenital hydronephrosis was the most prevalent malformation and represented 14.2% of all  
193 registered CAKUTs. Multiple anomalies of the kidney or urinary system affected every tenth  
194 newborn with CAKUTs (see Table 1).

195 Paternal age, maternal BMI, proportion of supplement intake (multivitamin, folic acid), as well  
196 as the fraction of mothers who smoked or consumed alcohol was not significantly different  
197 between newborns with and without malformations (Table 2).

198 The prevalence of CAKUTs at birth was significantly higher among newborns whose mothers  
199 had genitourinary infections before the pregnancy (p = 0.02), or any infection during the current  
200 pregnancy (p < 0.001) and mothers who took medications when being pregnant (p = 0.001).  
201 Similarly, the occurrence of CAKUTs was different between newborns conceived in different  
202 seasons (p = 0.001), with the highest prevalence for newborns conceived during the summer  
203 (Table 2).

204 The prevalence of CAKUTs among newborns whose mothers suffered from DM or GD was  
205 higher but not significantly so (p = 0.09). The latter likely reflects the low proportion (0.2%) of  
206 mothers with these conditions.

207 Based on the multivariate analysis results summarized in Table 3, diabetes mellitus or gestational  
208 diabetes (adjusted OR = 4.77), infections during pregnancy (adjusted OR = 2.03), the use of any  
209 medication during pregnancy (adjusted OR = 1.83) and conception during summer (adjusted OR  
210 = 1.75) were the only variables significantly associated with CAKUTs.

## 211 **Discussion**

### 212 PREVALENCE OF CAKUTS

213 The estimated all-year birth prevalence of CAKUT at birth in Murmansk County was higher than  
214 that for the combined data for all EUROCAT member countries for the same period (Dolk et al.,  
215 2010). This difference reached statistical significance for 2008 ( $p=0.0008$ ) and 2009 ( $p=0.003$ ):  
216 *5.6 versus 3.4* and *5.4 versus 3.5*, respectively (EUROCAT, 2013).

217 The most prevalent group was unspecified malformations. Since some of the latter were not  
218 confirmed after discharge from the maternity wards, our results could have been subject to  
219 overestimation. In addition, some minor anomalies might be revealed later in life even when the  
220 level of prenatal diagnosis of CAKUTs is high. For example, 73% of all congenital  
221 hydronephrosis cases were diagnosed prior to birth in Europe in 1995-2004 (Garne et al., 2009).  
222 In our previous study (Postoev et al., 2015b), the prenatal detection rate for CAKUTs in  
223 Monchegorsk was 42.1% for the 2000-2007 period. The possibility for underestimating the  
224 prevalence when using data at birth is illustrated by Caiulo et al., 2012, who showed that the  
225 prevalence of CAKUTs among children at the age of two months was close to 1 % based on  
226 mass ultrasound screening.

227 Our observed prevalences of renal agenesis and congenital hydronephrosis were lower and that  
228 of cystic kidney disease higher relative to the EUROCAT registry data. The latter indicate that  
229 the proportion of pregnancy terminations due to kidney malformation varied from 4.9% for  
230 congenital hydronephrosis to 68.2% for renal agenesis (EUROCAT, 2013). The lower  
231 prevalence of severe malformation in our data set might be explained by the exclusion of  
232 pregnancies under 22 weeks of gestation.

### 233 DIAGNOSTIC ISSUES

234 The estimation of birth defect prevalences depend on the diagnostic quality and experience of the  
235 clinicians performing the ultrasound examination. There were no national guidelines for the  
236 influence of gestational age in assessments of urodynamic and parenchymal changes (Adamenko  
237 et al., 2008), nor strict ultrasound criteria for pyelectasis and hydronephrosis. Consequently, it is  
238 not surprising that diagnoses of CAKUTs may be done using different clinical norms. Our data  
239 supports this perspective, as the prevalence of CAKUTs in Murmansk County varied between  
240 districts and hospitals. Such inconsistency can lead to over diagnosis of some CAKUTs during  
241 perinatal screening, especially congenital hydronephrosis and non-specified malformations.



243 The risk factors identified in the current study (see Table 3) have been reported previously,  
244 although they appear to differ in the duration and extent of their influence. The discussion that  
245 follows explores this.

246 Amri et al. (1999) demonstrated a potential harmful effect of hyperglycemia on kidney  
247 development in rats involving *in vivo* and *in vitro* experiments. Prospective cohort and case-  
248 control studies have also assessed this impact (Sheffield et al., 2002; Banhidly et al., 2010;  
249 Shnorhavorian et al., 2011; Dart et al., 2015). Relative to GD, the risk of CAKUTs was higher  
250 for DM (Shnorhavorian et al., 2011). Due to the small number of cases, it was not possible in our  
251 study to consider DM and GD separately. Nevertheless, the combined variable showed a strong  
252 association with CAKUTs. The magnitude of the observed prevalence of newborns with  
253 CAKUTs among mothers with diabetes was 1.6%, which is four-fold lower than that reported in  
254 a Texas (USA) study conducted in 1991-2000 (Shnorhavorian et al., 2011).

255 The significant association of CAKUTs with maternal infections during pregnancy is supported  
256 by other studies (Abe et al., 2003; Lukomska et al., 2012). These studies differ from the current  
257 assessment by limiting the exposure to the first trimester of pregnancy rather than considering  
258 the whole pregnancy duration. We opted for this approach because nephrogenesis is not limited  
259 to the first 12 weeks of pregnancy. Explanations of the teratogenic effect of maternal infectious  
260 may include the associated hyperthermia. In a summary of their animal experiments, Edwards et  
261 al., (1995) document the occurrence of structural malformations of the kidney (e.g., hypoplasia  
262 and agenesis) in offspring of mothers with extensive exposure to hyperthermia during pregnancy.  
263 Based on the Atlanta Birth Defects Case-control Study, Erickson (1991) reports robust  
264 associations between all defects and maternal febrile illnesses (specifically, “any fever” or “flu”  
265 experienced by the study respondents during the 4 months prior to conception and the first  
266 trimester). Unfortunately, we could not separate infectious diseases by presence or absence of  
267 fever, because no such information was available in the registry database. Moreover, prevalence  
268 of non-febrile infections could have been underestimated due to underreporting by mothers.  
269 Thus, the effect of infections observed in the present work should be considered as a combined  
270 effect of infectious agents and hyperthermia. Even though the emphasis in birth defect risk-factor  
271 studies has been on febrile infections in the context of hyperthermia as the teratogen, there is  
272 good evidence that maternal non-febrile infections and chronic diseases do indeed pose potential  
273 risks of birth defects and susceptibility to disease after birth (e.g., Erickson, 1991; Dong et al.,  
274 2015; Lee et al., 2015).

275 Infections during pregnancy may be expected to be closely related with the intake of certain  
276 medications, and this is perhaps this is reflected in the nearly two-fold risk of CAKUTs among  
277 newborns of mothers reporting infections and use of medications during pregnancy. This risk is  
278 similar in magnitude to those reported for antibiotics and antipyretics medications (Abe et al.,  
279 2003; Lukomska et al., 2012). We refrained from dividing the drugs into pharmacological  
280 groups, because most women (12 452 or 31.8%) consumed more than one medication due to  
281 pregnancy complications or maternal illness that existed prior to pregnancy; the most frequently  
282 used were dipyridamolum, drotaverinum and dydrogesteronum. Higher risk of CAKUTs has also  
283 been identified for aspirin-containing drugs (Abe et al., 2003) and angiotensin-converting  
284 enzyme inhibitors (Ratnapalan and Koren, 2002), but these were not frequently administrated in  
285 our study and this obviated any further analysis.

286 We found that the highest prevalences of CAKUTs occurred for newborns conceived during the  
287 months March-August, and this is supported by our logistic regression analysis (see Table 3).  
288 Luteijn et al., (2014) reported that urinary defects among European newborns peaked in July and  
289 indicated that this seasonality was driven by congenital hydronephrosis; it made up roughly 40%  
290 of the urinary defects. While Carton (2012) found the same seasonal variation with a peak in July  
291 for renal agenesis or hypoplasia based on the data for 1 967 654 livebirths in New York State,  
292 USA. Since Murmansk County is within or near the Arctic Circle, the summer season is brief  
293 and relatively cold; moreover, this is an unique arctic daylight regimens during the year (long in  
294 summer and short in winter). A recent study from Norway (Hwang et al., 2013), which includes  
295 data for regions with similar climate features, did not show seasonal variation in the prevalence  
296 of CAKUTs. However, it did identify peaks in March and February, respectively for respiratory  
297 defects and Down syndrome. Generally speaking, there is some evidence (see Hwang et al., 2013  
298 and references therein) that observed seasonal variation may reflect environmental factors (e.g.,  
299 different concentrations of teratogenic pollutants in air or water), meteorological conditions  
300 (polar night and day), and variation in the frequency of acute maternal infections. In our data set,  
301 adjustment for infections during pregnancy indirectly identified potential influences of the first  
302 two factors, but this could not be quantified due to the absence of such information in the  
303 MCBR.

304 The impact of maternal chronic genital and urinary tract infections has not been routinely  
305 assessed in CAKUTs studies. However, Shnorhavorian et al. (2011) report a five-fold increase in  
306 CAKUTs in association with pre-existing maternal renal disease. Our own findings indicate a  
307 near significant association (see Table 3). We presume that this reflects insufficient statistical

308 power because of a rather high proportion of missing data for this variable. Related issues are the  
309 misclassification of this risk factor due to different diagnostic practices and an inherent  
310 reluctance to report it (i.e., non-differential bias). Inclusion of missing values as non-exposure  
311 lead to increased risk ratios. Moreover, we also expect misclassification to have occurred  
312 between this pre-pregnancy variable and “acute infections during pregnancy”.

#### 313 STRENGTHS AND WEAKNESSES

314 The current study is the first population-based assessment of the epidemiology of CAKUTs and  
315 their risk factors in North-West of Russia that employs a register database shown to be of  
316 acceptable validity (Anda et al., 2008). The MCBR provides information not only on outcome,  
317 but also about exposure risk factors during pregnancy. It was established in accordance with  
318 international standards, and minimizes the possibility of selection bias. These attributes lead to  
319 good generalizability. Nevertheless, our study has some limitations.

320 The ascertainment of the CAKUT cases is possibly incomplete, because only those diagnosed  
321 during the stay in maternity houses were included. As indicated earlier, there were possibilities  
322 for misclassification due to the high proportion of non-specified malformations.

323 Missing values constituted an important issue in our multivariable analysis, as for some of the  
324 variables this exceeded than 10% (see Figure 1). These necessary exclusions from the logistic  
325 regression analysis might have biased our risk ratio estimates, which could have resulted in an  
326 underestimation of the risk ratios. As indicted above, a high proportion of missing data could  
327 have resulted in an underestimation of the risk ratios.

328 We sought to control confounding in estimating the influence of CAKUT risk factors by  
329 considering only well-established variables in the logistic regression analysis. We also avoided  
330 the inclusion in the model of all independent variables as categorical, as this could potentially  
331 lead to imperfect adjustment and bias due to residual confounding (Rothman et al., 2008). The  
332 breakdown into more than two categories of maternal age, body mass index and season are other  
333 cases in point.

334 The likelihood of underreporting of socially sensitive information such as maternal smoking also  
335 needs to be mentioned. Furthermore, data about the consumption of alcohol were not provided  
336 by the mothers, but was based on documented evidence of abuse provided by physicians.  
337 Although these factors were not significant predictors, they could have led to non-differential  
338 misclassification of exposure by attenuating the CAKUT risk estimates among smoking mothers.

## 339 **Conclusions**

340 The prevalence of CAKUTs in Murmansk County during 2006-2011 was 4.0 per 1000 newborns  
341 without significant changes over the observation period. The usage of medications during  
342 pregnancy, maternal diabetes mellitus or gestational diabetes, maternal infections during the  
343 pregnancy, and conception during June-August was associated with increased risk of CAKUTs.  
344 These findings have direct applications in improving prenatal care in Murmansk County and  
345 establishing targets for prenatal screening and consultations with affected women; they constitute  
346 a framework for future research on the teratogenic effects of medications and infections.

## 347 **Acknowledgements**

348 The authors thank the office staff of the Murmansk County Birth Registry and Rod  
349 Wolstenholme, executive officer, Section for Dissemination Services at Faculty of Health  
350 Sciences, The Arctic University of Norway.

## 351 **References**

- 352 Abe K, Honein MA, Moore CA. 2003. Maternal febrile illnesses, medication use, and the risk of  
353 congenital renal anomalies. *Birth Defects Res A Clin Mol Teratol* 67(11):911-918.
- 354 Adamenko O, Khalepa Z, Kotova L. 2008. Ultrasound monitoring in prenatal diagnostic of  
355 urinary system malformations and in identification of management tactic for children (in  
356 Russian). *Pediatrics* 3:13-17.
- 357 Amri K, Freund N, Vilar J, Merlet-Benichou C, Lelievre-Pegorier M. 1999. Adverse effects of  
358 hyperglycemia on kidney development in rats: in vivo and in vitro studies. *Diabetes*  
359 48(11):2240-2245.
- 360 Anda EE, Nieboer E, Voitov AV, Kovalenko AA, Lapina YM, Voitova EA, Kovalenko LF,  
361 Odland JO. 2008. Implementation, quality control and selected pregnancy outcomes of  
362 the Murmansk County Birth Registry in Russia. *Int J Circumpolar Health* 67(4):318-334.
- 363 Anda EE, Nieboer E, Wilsgaard T, Kovalenko AA, Odland JO. 2011. Perinatal mortality in  
364 relation to birthweight and gestational age: a registry-based comparison of Northern  
365 Norway and Murmansk County, Russia. *Paediatr Perinat Epidemiol* 25(3):218-227.
- 366 Andreeva LP. 2008. Prophylactic of congenital and hereditary diseases (in Russian). *Russian*  
367 *Medical Journal* 1:3-7.

- 368 Banhidy F, Acs N, Puho EH, Czeizel AE. 2010. Congenital abnormalities in the offspring of  
369 pregnant women with type 1, type 2 and gestational diabetes mellitus: a population-based  
370 case-control study. *Congenit Anom* 50(2):115-121.
- 371 Barakat AJ, Drougas JG. 1991. Occurrence of congenital abnormalities of kidney and urinary  
372 tract in 13,775 autopsies. *Urology* 38(4):347-350.
- 373 Boubred F, Vendemmia M, Garcia-Meric P, Buffat C, Millet V, Simeoni U. 2006. Effects of  
374 maternally administered drugs on the fetal and neonatal kidney. *Drug Saf* 29(5):397-419.
- 375 Caiulo VA, Caiulo S, Gargasole C, Chiriaco G, Latini G, Cataldi L, Mele G. 2012. Ultrasound  
376 mass screening for congenital anomalies of the kidney and urinary tract. *Pediatr Nephrol*  
377 27(6):949-953.
- 378 Caton AR. 2012. Exploring the seasonality of birth defects in the New York State Congenital  
379 Malformations Registry. *Birth Defects Res Part A Clin Mol Teratol* 94(6):424-437.
- 380 Dart AB, Ruth CA, Sellers EA, Au W, Dean HJ. 2015. Maternal Diabetes Mellitus and  
381 Congenital Anomalies of the Kidney and Urinary Tract (CAKUT) in the Child. *Am J*  
382 *Kidney Dis* 65(5):684-691.
- 383 Dolk H, Loane M, Garne E. 2010. The prevalence of congenital anomalies in Europe. *Adv Exp*  
384 *Med Biol* 686:349-364.
- 385 Dong DY, Binongo JN, Kancherla V. 2015. Maternal Chlamydia infection during pregnancy and  
386 risk of cyanotic congenital heart defects in the offspring. *Matern Child Health*. [Epub  
387 ahead of print]
- 388 dos Santos Junior AC, de Miranda DM, Simoes e Silva AC. 2014. Congenital anomalies of the  
389 kidney and urinary tract: an embryogenetic review. *Birth Defects Res C Embryo Today*  
390 102(4):374-381.
- 391 Edwards MJ, Shiota K, Smith MS, Walsh DA. 1995. Hyperthermia and birth defects. *Reprod*  
392 *Toxicol* 9(5):411-425.
- 393 Erickson JD. 1991. Risk factors for birth defects: data from the Atlanta Birth Defects Case-  
394 Control Study. *Teratology* 43(1):41-51.
- 395
- 396 European Surveillance of Congenital Anomalies. 2013. Prevalence tables 2008–2012.
- 397 Feldman I, Friger M, Wiznitzer A, Mazor M, Holcberg G, Sheiner E. 2009. Is oligohydramnios  
398 more common during the summer season? *Arch Gynecol Obstet* 280:3-6.
- 399 Garne E, Loane M, Wellesley D, Barisic I, Eurocat Working G. 2009. Congenital  
400 hydronephrosis: prenatal diagnosis and epidemiology in Europe. *J Pediatr Urol* 5(1):47-  
401 52.
- 402 Hei MY, Yi ZW. 2014. Environmental factors for the development of fetal urinary  
403 malformations. *World J Pediatr* 10(1):17-23.

404 Hwang B-F, Magnus P, Jaakkola JJK. 2013. Seasonal variation of birth defects in Norway.  
405 BioMedicine 3(2):95-101.

406 International clearinghouse for birth defects surveillance and research (ICBDSR). 2013. Annual  
407 Report 2012.

408 Kallen K. 1997. Maternal smoking and urinary organ malformations. *Int J Epidemiol* 26(3):571-  
409 574.

410 Kalter H. 2003. Maternal diabetes mellitus and infant malformations. *Obstet Gynecol*  
411 101(4):815; author reply 815-816.

412 Lee BK, Magnusson C, Gardner RM, Blomström Å, Newschaffer CJ, Burstyn I, Karlsson H,  
413 Dalman C. 2015. Maternal hospitalization with infection during pregnancy and risk of  
414 autism spectrum disorders. *Brain Behav Immun.* 44:100-5.

415 Li DK, Mueller BA, Hickok DE, Daling JR, Fantel AG, Checkoway H, Weiss NS. 1996.  
416 Maternal smoking during pregnancy and the risk of congenital urinary tract anomalies.  
417 *Am J Public Health* 86(2):249-253.

418 Lukomska A, Baumert M, Krzych LJ, Sypniewska K. 2012. Estimation of the risk factors of the  
419 congenital urinary system defects. *Cent Eur J Med* 7(1):85-90.

420 Luteijn JM, Dolk H, Addor MC, Arriola L, Barisic I, Bianchi F, Calzolari E, Draper E, Garne E,  
421 Gatt M, Haeusler M, Khoshnood B, McDonnell B, Nelen V, O'Mahony M, Mullaney C,  
422 Queisser-Luft A, Rankin J, Tucker D, Verellen-Dumoulin C, de Walle H, Yevtushok L.  
423 2014. Seasonality of congenital anomalies in Europe. *Birth Defects Res A Clin Mol*  
424 *Teratol* 100(4):260-269.

425 Ministry of Health of Russian Federation (2000). Order № 457 28.12.2000. "About  
426 improvement of perinatal diagnostic and prophylactic congenital diseases among  
427 children" (in Russian).

428 Molchanova EA, Volov, A.L., Kabak, M.M. 2004. The first results from implemented Russian  
429 register of children with chronic renal failure (in Russian). *Nephrology and dialysis* 5:4.

430 Moore KL, Persaud TVN. 2003. *The developing human: clinically oriented embryology*. 7th ed.  
431 Philadelphia, Pa.: Saunders. xv, 560 p. p.

432 Murmansk State Statistical Office. 2015a Murmansk County in Figures. Population.  
433 [http://murmanskstat.gks.ru/wps/wcm/connect/rosstat\\_ts/murmanskstat/resources/07f65b0](http://murmanskstat.gks.ru/wps/wcm/connect/rosstat_ts/murmanskstat/resources/07f65b00490af5f6b4eeb4e321b4d743.pdf)  
434 [0490af5f6b4eeb4e321b4d743.pdf](http://murmanskstat.gks.ru/wps/wcm/connect/rosstat_ts/murmanskstat/resources/07f65b00490af5f6b4eeb4e321b4d743.pdf)

435 Murmansk State Statistical Office. 2015b Murmansk County in Figures. Indicators of vital  
436 statistics  
437 [http://murmanskstat.gks.ru/wps/wcm/connect/rosstat\\_ts/murmanskstat/resources/61e8fc0](http://murmanskstat.gks.ru/wps/wcm/connect/rosstat_ts/murmanskstat/resources/61e8fc00490af40fb492b4e321b4d743.pdf)  
438 [0490af40fb492b4e321b4d743.pdf](http://murmanskstat.gks.ru/wps/wcm/connect/rosstat_ts/murmanskstat/resources/61e8fc00490af40fb492b4e321b4d743.pdf)

439 Postoev VA, Nieboer E, Grijbovski AM, Odland JO. 2015a. Prevalence of birth defects in an  
440 Arctic Russian setting from 1973 to 2011: a register-based study. *Reprod Health* 12:3.

441 Postoev VA, Nieboer E, Grijbovski AM, Odland JO. 2015b. Changes in detection of birth  
442 defects and perinatal mortality after introduction of prenatal screening in the Kola  
443 Peninsula (North-West Russia): combination of two birth registries -*submitted*

444 Ratnapalan S, Koren G. 2002. Taking ACE inhibitors during pregnancy. Is it safe? *Can Fam*  
445 *Physician* 48:1047-1049.

446 Russian institute of pediatrics and children surgery. 2011. Annual report about birth defect  
447 monitoring in Russia (in Russian). Moscow.

448 Rothman KJ, Greenland S, Lash TL. 2008. *Modern epidemiology*. 3rd ed. Philadelphia: Wolters  
449 Kluwer Health/Lippincott Williams & Wilkins. x, 758 p. p.

450 Sheffield JS, Butler-Koster EL, Casey BM, McIntire DD, Leveno KJ. 2002. Maternal diabetes  
451 mellitus and infant malformations. *Obstet Gynecol* 100(5 Pt 1):925-930.

452 Shnorhavorian M, Bittner R, Wright JL, Schwartz SM. 2011. Maternal risk factors for congenital  
453 urinary anomalies: results of a population-based case-control study. *Urology* 78(5):1156-  
454 1161.

455 Trefilov AA. 2006. Prevalence of congenital anomalies of kidney by the results of mass  
456 ultrasound screening (in Russian). *Pediatric pharmacology* 2:44-47.

457 Wiesel A, Queisser-Luft A, Clementi M, Bianca S, Stoll C, the EUROSCAN Study Group.  
458 2005. Prenatal detection of congenital renal malformations by fetal ultrasonographic  
459 examination: an analysis of 709,030 births in 12 European countries. *Eur J Med Genet*  
460 48:131–144.

461

462

463

464

465

466

467

468 **Table legend**

469 **TABLE 1.** Diagnosis-specific Frequencies, Prevalences (per 1000 newborns) and Distribution  
470 (%) of CAKUTs Among Newborns in Murmansk County (2006-2011)

471 **TABLE 2.** Univariate Comparisons of the Prevalence of CAKUTs by Maternal Socio-  
472 demographic, Anthropometric and Lifestyle Characteristics in Murmansk County (2006-2011)

473 **TABLE 3.** Risk Factors Associated with CAKUTs (results of the multivariate analysis)

474 **Figure legend**

475 **FIGURE 1.** Number of Births Included in the Analyses and Information about Missing  
476 Variables

477



**TABLE 1.** *Diagnosis-specific Frequencies, Prevalences (per 1000 newborns) and Distribution (%) of CAKUTs Among Newborns in Murmansk County (2006-2011)*

ICD-10 code	Birth defect	Distribution		Prevalence (95%CI) per 1000 newborns
		n	%	
Q 60	Renal agenesis and other reduction defects of kidney	4	2.0	0.08 (0.03-0.20)
Q 61	Cystic kidney disease	21	10.3	0.41 (0.27-0.63)
Q 62	Congenital obstructive defects of renal pelvis and congenital malformations of ureter	39	19.2	0.77 (0.56-1.05)
Q62.0	Congenital hydronephrosis	29	14.3	0.57 (0.40-0.82)
Q 63	Other congenital malformations of kidney	108	53.2	2.13 (1.76-2.56)
Q 64	Other congenital malformations urinary system	7	3.5	0.14 (0.07-0.28)
	Multiple anomalies of kidney or urinary system	24	11.8	0.47 (0.32-0.70)
Total		203	100	4.00 (3.43-4.54)

**TABLE 2.** *Univariate Comparisons of the Prevalence of CAKUTs by Maternal Socio-demographic, Anthropometric and Lifestyle Characteristics in Murmansk County (2006-2011)*

Characteristics	N	%	Prevalence of CAKUTs		p-Value <sup>1</sup>
			n	Per 1000 newborns	
<b>Age of mother:</b>					0.39
less 18 years	721	1.4	5	6.9	
18-35 years	45706	89.7	178	3.9	
older 35 years	4508	8.9	20	4.4	
<b>Age of father:</b>					0.10
18-35 years	37092	79.6	143	3.9	
older 35 years	9477	20.4	48	5.1	
<b>Maternal BMI:</b>					0.42
less 18.5 kg/m <sup>2</sup>	3133	6.3	10	3.2	
18.5-24.9 kg/m <sup>2</sup>	32801	65.7	140	4.3	
more than 25kg/m <sup>2</sup>	13976	28.0	50	3.6	
<b>Use of medications during pregnancy:</b>					0.001
yes	39125	76.8	176	4.5	
no	11811	23.2	27	2.3	
<b>Diabetes mellitus or gestational diabetes:</b>					0.09
yes	124	0.2	2	16.1	
no	50812	99.8	201	4.0	
<b>Multivitamin intake during the pregnancy:</b>					0.34
yes	46900	92.5	191	4.1	
no	3824	7.5	11	2.9	
<b>Folic acid intake during the pregnancy:</b>					0.16
yes	12965	25.6	43	3.3	
no	37619	74.4	159	4.2	
<b>Cigarette smoking during the pregnancy:</b>					0.52
yes	9169	18.3	39	4.3	
no	40836	81.7	155	3.8	
<b>Evidence of alcohol abuse:</b>					0.17
yes	187	0.4	2	10.7	
no	50545	99.6	200	4.0	
<b>Chronic sex tract or urinal infections before pregnancy:</b>					0.02
yes	10992	25.2	61	5.5	
no	32564	74.8	124	3.8	
<b>Infections during the pregnancy:</b>					<0.001
yes	7345	14.4	56	7.6	
no	43591	85.6	147	3.4	
<b>Season of conception<sup>2</sup>:</b>					0.001
Winter	12671	24.9	37	2.9	
Spring	12383	24.3	56	4.5	
Summer	12734	25.1	71	5.6	
Autumn	13077	25.7	39	3.0	

<sup>1</sup>- Significant p-values indicate that differences in prevalence exist between the categories of the indicated

characteristics.

<sup>2</sup>The months of the year are grouped by seasons as follows: December-February (Winter); March-May (Spring); June-August (Summer); and September-November (Autumn)

**TABLE 3.** *Risk Factors Associated with CAKUTs (results of the multivariate analysis)*

<b>Characteristics<sup>1</sup></b>	<b>Adjusted OR</b>	<b>95%CI</b>
Use of medications during pregnancy	1.83	1.14-2.94
Diabetes mellitus or gestational diabetes	4.77	1.16-19.65
Chronic sex tract or urinal infections before pregnancy	1.34	0.97-1.84
Infections during the pregnancy	2.03	1.44-2.82
Season of conception		
Winter	1.00	-
Spring	1.45	0.94-2.25
Summer	1.75	1.15-2.66
Autumn	0.86	0.51-1.35

<sup>1</sup>- Only those variables that remained in the final regression model (backward stepwise regression) are presented.

