

Molecular Cloning of Bone Morphogenetic Protein (BMP) Gene from the Planarian *Dugesia japonica*

Hidefumi Orii*, Kentaro Kato, Kiyokazu Agata
and Kenji Watanabe

*Laboratory of Regeneration Biology, Department of Life Science, Faculty of Science,
Himeji Institute of Technology, Harima Science Garden City, Akou, Hyogo 678-1297, Japan*

ABSTRACT—BMP (Bone Morphogenetic Protein) acts as a morphogen for dorso-ventral patterning and organogenesis in both vertebrate and invertebrate development. A cDNA encoding BMP (named *Djbmp*) has been cloned and sequenced from the planarian *Dugesia japonica*. The mature form of DjBMP which was deduced from the cDNA sequence was composed of 114 amino acid residues. The position of seven cysteine residues of the mature DjBMP was highly conserved among the TGF- β superfamily. DjBMP had high similarity to human BMP-2A (50% amino acid identity), BMP-4 (49%) and *Drosophila decapentaplegic* protein (48%), indicating that DjBMP belongs to DVR (decapentaplegic-Vg1-related) group. The expression pattern in intact and regenerating planarians revealed by whole mount *in situ* hybridization suggested that the DjBMP plays a role not only in dorso-ventral but also in mid-lateral body patterning.

INTRODUCTION

Transforming growth factor- β (TGF- β) superfamily is a large group including biological active proteins such as activin, inhibin, growth/differentiation factor (GDF) and bone morphogenetic proteins (BMPs) (Horgan *et al.*, 1994; Kingsley, 1994). One of the members, BMP-4, was found to act as a morphogen for ventralization in vertebrate development (Dale *et al.*, 1992; Jones *et al.*, 1992; Dosch *et al.*, 1997). In *Drosophila decapentaplegic* (*dpp*) gene which encodes a BMP-4 homologue has been known to play a key role in determination of dorsalization during the early development (Irish and Gelbart, 1987). The idea that the dorso-ventral (D-V) axis in deuterostomes including vertebrate corresponds to the ventro-dorsal axis in protostomes including arthropod is supported by molecular evidence that DPP and BMP-4 are functionally similar in inducing ventral structure in frog (deuterostome) and dorsal structure in fly (protostome), respectively (Holley *et al.*, 1995; Horgan, 1995; De Robertis and Sasai, 1996).

The freshwater planarian *Dugesia japonica*, belonging to the Phylum Plathelminthes has powerful activity for regeneration (for review; Baganà *et al.*, 1994). We are interested in how body plan is established during the planarian regeneration. As for the antero-posterior axis in planarians, the expression of Hox/HOM-C genes may be important (Orii *et al.*, in preparation). On the other hand, we have no information on their D-V patterning. In this study, we focused on a planarian BMP-4/DPP homologue as a candidate molecule in D-V pat-

terning. The molecular cloning of the BMP gene and its expression in intact and regenerating planarians are reported.

MATERIALS AND METHODS

Organisms

All planarians in this study were derived from one worm of *Dugesia japonica* collected in Irima river in Gifu, Japan and maintained clonally in our laboratory (clonal population GI) (Orii *et al.*, 1993). Amputated worms were regenerated in autoclaved tap water at about 22°C.

cDNA cloning and sequencing

A cDNA library (4×10^6 in size) was constructed from poly A⁺ RNA of whole worms using λ ZAPII vector (Umesono *et al.*, 1997). The mixture (10 μ l) for PCR (polymerase chain reaction) contained 1 \times Taq buffer (10 mM Tris-HCl pH 8.3, 50 mM KCl, 1.5 mM MgCl₂), Takara Taq polymerase (Takara, 0.025 U/ μ l), dNTPs (0.25 mM each), *D. japonica* total DNA (about 40 ng/ μ l) and a set of degenerate primers (1 pmole/ μ l each): a forward primer 5'-GGITGG(A/G/C)AIGA(C/T)TGG(G/A)TIGCICC-3' and a reverse primer 5'-ACIA(G/A)IGT(T/C)TGIA(C/G)IA(T/C)IGC(G/A)TG(G/A)TT-3' corresponding to amino acids GW(N/D/Q)DW(I/V)(I/V)AP and NHA(I/V)VQTLV, respectively. The reaction conditions were as follows: an initial denaturation at 94°C for 1 min followed by 40 cycles of 94°C for 1 min, 45°C for 1 min, 72°C for 1 min and a final elongation step at 72°C for 5 min. The PCR product (about 120 bp) was extracted from 6% acrylamide gel, amplified again, cloned into pT7blue T vector (Novagen) and sequenced. From the sequence of the PCR product corresponding to BMP, we designated a specific forward primer 5'-CGTATCCGTTATC-AGATAATTTTAA-3' (arrow in Fig. 1) and screened the cDNA library by PCR with the primer and M13 universal primer 5'-CGCCAGGGTTT-TCCCAGTCACGAC-3' (Watanabe *et al.*, 1997). Out of 11 positives, one clone carrying the longest insert (pDjBMP17) has been sequenced as to both strands by using Thermo Sequenase cycle sequencing kit (Amersham) and automatic DNA sequencer DSQ1000L (Shimadzu).

* Corresponding author: Tel. +81-7915-8-0187;
FAX. +81-7915-8-0187.

Southern blot hybridization

D. japonica total DNA (10 µg) was digested with *EcoRI* or *SpeI*, subjected to 0.8% agarose gel electrophoresis, and blotted on Hybond N⁺ membrane (Amersham). The membrane were prehybridized in 6 × SSC, 5 × Denhardt's solution, 1%SDS, 100 µg/ml salmon testes DNA at 65°C for 1 hr and hybridized in the same solution with probe. pDjBMP17 DNA was labeled by using random primed labeling kit (Takara) with α-[³²P]-dCTP (Amersham, ~3000 Ci/mmol) and used as the hybridization probe. The membrane was washed twice in 2 ×

SSC, 0.1% SDS at 65°C for 30 min and exposed to film with an intensifying screen.

Whole mount *in situ* hybridization

The digoxigenin (DIG) labeled anti-sense and sense RNA probes were synthesized with T7 RNA polymerase and T3 RNA polymerase, respectively, and used for hybridization without alkaline hydrolysis (Umesono *et al.*, 1997; Agata *et al.*, 1998).

```

10      20      30      40      50      60      70      80      90
AAATTATTACTCATTAAACATACATTTTTTTGTTATTTTCTATTCAAATTCACAAAATCAAATGGCACAGCGATTTCGTTTTATGAAAAG
K L L L I N I H F F C Y F L F K F T K S N G T S D S F Y E K

100     110     120     130     140     150     160     170     180
AAACTCGGTAACAAAATCACGAATCCATCAGAAAAAGTACGCAGTAATTTGCAAAAAGATTTTATTGTTAAACATGGGACTTAAACCGGAC
K L G N K I T N P S E K V R S N L Q K I L L L N M G L K P D

190     200     210     220     230     240     250     260     270
GAAGTAGTGGACAGTGAATAATTTGCCGAGTTATCGGAAAAATACCACATTTTCGTTCTGATTTTATGAAACTACTCTATGTTAAATAC
E V V D S E N F A E L S E K L P H F V P D F M K L L Y V K Y

280     290     300     310     320     330     340     350     360
CGATCTAATTTATACAATTTTCTGTGAAATGAAGGAAAACCGAAAACATTTAGGACCAGTTGAAATGATTCATTGCCAAAGATTAAGA
R S N L Y N F S V E I E G K P K H L G P V E M I H C Q R L R

370     380     390     400     410     420     430     440     450
AATATCCAAACATGGTCTTCGGTTGGTATTTCATAAATTAATATCAAGTCATTTACGAAATcaAAAACACTAGTTATCCATTAGCAagAGAA
N I Q T W S S V G I H K L I F K S F T K S K T S Y P L A R E

460     470     480     490     500     510     520     530     540
ttGCGAATAAATAAAAGATTGTTTCCGAGAATTATTCATGAATCTCCAACAATGAATTCAGTGAATTATATATGATCATAATGGTTTTT
L R I N K R L F P R I I H E S P N N E F S E L Y M I I M V F

550     560     570     580     590     600     610     620     630
GATGATCGTTTTAAATTTGTTAAAGAAAAATTAATTAATTCAAAGTCATTTAATAAGCAAGGATGGATTTCAATAGATATTTCAAATAT
D D R F K L L K E K L I N S K S F N K Q G W I S I D I S K Y

640     650     660     670     680     690     700     710     720
TTACAAAATTCAGTGTTTTTTATCAAAAACGTTTGGTTTTATGGAAAAACACAAATCAATTTTCAAACGAGTCAATGTAAAAATATCAATA
L Q N S V F F I K T F G L W K K H K S I F K R V N V K Y Q I

730     740     750     760     770     780     790     800     810
CAACAATCTCATCAATTGTATTGTATTATGCAAAATATTGTGACATTTTACAGTAATAAAGTTTCGACCATCATCAAATAAACAATATGAA
Q Q S H Q L Y L Y Y A N I V T F Y S N K V R P S S N K Q Y E

820     830     840     850     860     870     880     890     900
TCGACAATTTCCGAAACTGAAAATCTACATTTATCTAGAGTTAAGCGTGATACGTCAGGATACATGCCAGGCCATGAAGAAGACTGTCAA
S T I S E T E N L H L S R V K R D T S G Y M P G H E E D C Q

910     920     930     940     950     960     970     980     990
AGATATCCTTTAATGTACTTTTAAAGAAGTTGGATGGAGCAAAATGGATAATTCACCCCAAACATAATGCTTATTATTTGTAAGGT
R Y P L I V T F K E V G W S K W I I A P Q N Y N A Y Y C K G

1000    1010    1020    1030    1040    1050    1060    1070    1080
AATTGTCGGTATCCGTTATCAGATAATTTAATGCAACAAATCATGCAATTTTCAGTTATTAGTGCATGGTTTAAAGACCTTTCTATT
N C P Y P L S D N F (N) (A) (T) N H A I I Q L L V H G L K D L S I

1090    1100    1110    1120    1130    1140    1150    1160    1170
CCAAAACCGTGTGTGTGCCTTATCACTTACACCCAGAACTTTATTATATTAAACAATGAAGGCGATGCACTTTTGCCTGAATTCAAA
P K P C C V P Y H L H P E T L L Y L N N E G D A L L R E F K

1180    1190    1200    1210    1220    1230    1240    1250    1260
GATATGTCGGTCAAGTGTCTTCTGTGCTGTTGACAAGTAAAATTCACAATTTTTCGATTTTGTATACCTCAACTGTAAC TAAGATTCT
D M S V S S C S C R *

1270    1280    1290    1300    1310    1320    1330    1340    1350
AAAACTTGTTGATCAATTTTTCTTCTGCTTTTTTTATTTGACTTTTCATTTGGTGAAAACAAAAGAAAATTTTTTAAACGGAAAAA
AAAAA
1360
AAAAA

```

Fig. 1. Nucleotide sequence of *Djbmp* cDNA (upper) and the deduced amino acid sequence (lower). The first methionine is double underlined. Seven cysteine residues conserved among the members of TGF-β superfamily are in bold. The box shows the consensus sequence of proteolytic cleavage. The predicted mature form is underlined. The potential asparagine-linked glycosylation sites are circled. The arrow shows the specific primer used for PCR screening of the cDNA library. The amino acids are shown with single letter abbreviations.

RESULTS AND DISCUSSION

We have aligned many members of TGF- β superfamily to designate a set of degenerate primers for PCR (see Materials and Methods). Using the primers, PCR was performed with *D. japonica* total DNA. Cloning and sequencing analysis revealed that only one PCR fragment had similarity with BMP genes. A cDNA library was screened by the stepwise dilution method with a specific primer designed for the sequence of PCR product (Watanabe *et al.*, 1997). The positive cDNA clone with the longest insert (pDjBMP17) was sequenced (Fig. 1). The longest open reading frame (ORF) from the 5' end of the insert to the stop codon (nucleotide 1201-1203) was found. Although the first methionine in the ORF was found at nucleotide 163-165, it did not seem to be the initiation codon, because the upstream region of the methionine of the ORF did not contain any stop codons in frame and the N-terminal region of the deduced protein translated from the methionine did not contain many hydrophobic amino acid residues, which may serve as signal sequence of secreted peptide such as BMP precursor (Nishimatsu *et al.*, 1992). As the deduced protein was similar to the members of BMP group on the whole, we designated it DjBMP (*Dugesia japonica* BMP). In comparison with other proteins belonging to TGF- β superfamily, the pro-protein of DjBMP might be processed into mature form by cleavage at the carboxyl end of R (amino acid 286) of RVKR corresponding to the cleavage consensus sequence RXXR (Panganiban *et al.*, 1990). The position of seven cysteine residues conserved among the members of TGF- β superfamily was also confirmed in the protein. Figure 2 shows the comparison of mature protein of DjBMP with other TGF- β related proteins from the first cysteine to the carboxyl terminal. In this region, DjBMP protein was similar to human BMP-2A (50% amino acid identity), BMP-4 (49%), sea urchin univin (50%), *Drosophila* DPP (48%), *Drosophila* 60A (47%), human BMP-5 (44%), TGF- β 1 (33%) and inhibin- β a (33%). A putative N-linked glycosylation site (NXT/S) was found at amino acid 341-343 (NAT), whose position was also conserved among 60A and DPP subclasses. These results indicate that DjBMP belongs to DVR (decapentaplegic-Vg1-related) group of TGF- β superfamily (Kingsley, 1994). It was very difficult to classify DjBMP more precisely even by the phylogenetic sequence analysis with Genetics Computer Group (GCG) program (Madison, Wisconsin), because the sequences of members of TGF- β superfamily are various for their length.

Genomic Southern hybridization probed with the *Djbmp* cDNA revealed that the *Djbmp* was a single copy (Fig. 3). PCR using genomic DNA as a template indicated that there was no intron in the region encoding mature protein as well as vertebrate BMP-4 gene (Kurihara *et al.*, 1993) (data not shown).

To investigate the expression pattern of the gene, whole mount *in situ* hybridization was performed in intact worms (Fig. 4A, B and C). The *Djbmp* was expressed, though very weakly, in dorsal cells (Fig. 4A and B). No signal was detected with sense probe (data not shown). We never observed any sig-

nals in the ventral side (Fig. 4C). The expression was stronger in the medial region. There has been no report of morphologically special cells whose distribution is the same as that of *Djbmp* expressing cells. Unfortunately, we could not identify what kind of cells expressed *Djbmp*, because of the sensitivity of our *in situ* hybridization method on paraffin embedded sections. In addition to sequence comparison (Fig. 2), the expression pattern suggests that DjBMP may be a homologue of DPP/BMP-4 in *Drosophila* and vertebrates. DjBMP may play a key role as a dorsal forming or anti-neurogenic factor in the planarian as well as DPP in *Drosophila* (Sasai *et al.*, 1995; Wilson and Hemmati-Brivanlou, 1995). To elucidate the role of DjBMP in D-V patterning, one should investigate the expression pattern of *Djbmp* during regeneration of dorsal and ventral parts. However, it is impossible to divide a flat worm into dorsal and ventral pieces.

The expression along the dorso-medial region suggests that DjBMP also plays some role in mid-lateral patterning of the dorsal side. The planarian *Dugesia japonica* has high regeneration ability and a piece of the body can regenerate completely. The left or right marginal fragment without medial region can also regenerate to a whole animal. We investigated the expression pattern of *Djbmp* during regeneration from the lateral piece without the cells expressing *Djbmp*. One day after amputation, the cells expressing *Djbmp* appeared dispersedly and very weakly only on the dorsal side (Fig. 4D). On the second or third day, the positive cells were distributed mainly in the dorso-medial region (Fig. 4E). It was not until five days after amputation that a pair of small eyes and a pharynx appeared, indicating the formation of the medial structure. *Djbmp* expressing cells were distributed mainly on the curved dorso-medial region at this stage. Seven days after amputation, distinct eyes and pharynx were regenerated and the expressing cells were more clearly distributed on the dorso-medial region (Fig. 4F). The expression pattern is represented schematically in Fig. 4G. The expression before differentiation of bilateral eyes suggests that DjBMP is involved in mid-lateral patterning during regeneration. As we could not monitor the behavior of single cells during regeneration, we do not know if the expressing cells move into the midline during regeneration or not. The change of expression pattern during regeneration suggests that dynamic morphogenesis occurs throughout the remaining tissue after amputation rather than in the restricted regenerating region called 'blastema'. In other words, the planarian regeneration could be referred to as 'morphallaxis' rather than 'epimorphosis'. Furthermore, the quick response of the expression after amputation (after only 1 day) also suggests that it may be involved in initial D-V patterning which occurs before dorsal cell differentiation.

It is interesting that the *Djbmp* was expressed in both intact and regenerating worms, because *dpp/Bmp-4* gene is expressed and functions in early embryogenesis of fly and vertebrates. It is probable that *Djbmp* is expressed in early embryogenesis as well as regeneration. If so, the mechanism of body planning in early embryogenesis may be maintained and utilized for regeneration in planarians. Furthermore, the




Fig. 2. A comparison of the amino acid sequence of the DjBMP with other members of TGF-β superfamily. Primers and regions used for PCR are shown. All sequence data were obtained from DNA data bank of Japan. Gaps in the alignment are represented by a dots. Amino acid residues identical to DjBMP are represented by a dash. Accession numbers are as follows: human TGF-β1 (hTGFb1), Swiss-Prot/P01137; human TGF-β3 (hTGFb3), Swiss-Prot/P10600; human inhibin-βa (hINH-Ba), PIR/B24248; human inhibin-βb (hINH-Bb), Swiss-Prot/P09529; human BMP-2A (hBMP2A), PIR/B37278; human BMP-3 (hBMP3), Swiss-Prot/P12645; human BMP-4 (hBMP4), PIR/C37278; human BMP-5 (hBMP5), PIR/A39263; human BMP-6 (hBMP6), PIR/B39263; human BMP-7 (hBMP7), PIR/C39263; human GDF-1 (hGDF1), PIR/C39364; human GDF-5 (hGDF5), PIR/JC2347; human Mullerian inhibiting substance (hMIS), PIR/A01397; *Drosophila* dpp (dDPP), PIR/A26158; *Drosophila* 60A (d60A), PIR/A43918; *Xenopus* Vg-1 (xVG1), PIR/A29619; sea urchin DVR-1 (suDVR1), PIR/S52408; sea urchin uninivin (suUNIVIN), GenBank/U10533; chicken dorsalin (cDORSALIN), GenBank/L12032; mouse nodal (mNODAL), PIR/S29718; ascidian BMPa (HrBMPa), Miya et al., 1996.




Fig. 3. Southern blot hybridization. Molecular size (*Hind*III digested λ DNA) is shown on the left.

expression in intact worm suggests that the molecule is important in maintenance of body plan and that the cellular and molecular events during regeneration themselves occur constantly in an intact body to maintain the body plan. This was also supported by the expression pattern of Hox/HOM-C genes along the antero-posterior axis of regenerating and intact body (Orii *et al.*, in preparation).

Recently, De Robertis and Sasai (1996) proposed a hypothetical ancestral and primitive bilateral animal, *Urbilateria*, from which the arthropod and the chordate lineages diverged 600 million years ago, with Hox gene complexes, D-V patterning system by *sog* (*short gastrulation*)/*chordin* and *dpp*

Bmp-4 and so on. In accordance with recent molecular evolutionary studies, the *Urbilateria* was divided into two groups, the *Deuterostomia* and the *Protostomia*. The *Protostomia* was further subdivided into the *Lophotrochozoa* and the *Ecdysozoa* during the evolution (for review; Balavoine and Adoutte, 1998). The Plathelminthes including *Dugesia japonica* has simple body plan and is grouped into the *Lophotrochozoa*. In this paper, we showed that the Plathelminthes also has BMP gene as well as the *Deuterostomia* (vertebrates *etc.*) and the *Ecdysozoa* (arthropoda, nematoda *etc.*). In addition to Hox gene complexes and *Djbmp*, the presence of *DjotxA*, *DjotxB* and *Djotp* which are planarian homologues to *orthodenticle* and *orthopedia* in *Drosophila* (Umesono *et al.*, 1997; Umesono *et al.*, 1998), strongly suggests that the basic body plan of the *Bilateria* including *Deuterostomia*, *Ecdysozoa* and *Lophotrochozoa*, are common. To date, the presence of BMP gene has not been reported in radiata hydrozoan such as hydra. BMP gene may be related to the establishment of D-V axis in animal evolution.

We have no information on other BMP genes in the planarian. However, it has been suggested that BMP-4 forms a heterodimer with BMP-7 to function in mesoderm induction in *Xenopus* (Suzuki *et al.*, 1997) and that DPP acts to establish D-V pattern by forming a heterodimer with SCREW, a member of 60A subfamily, in *Drosophila* (Arora *et al.*, 1994). DjBMP may also act with another unidentified BMP-like member in the planarian. In higher organisms, *tolloid/Bmp-1* gene product also regulates D-V patterning in relation to DPP/BMP and SOG/Chordin proteins (Marqués *et al.*, 1997; Piccolo *et al.*, 1997). To search for and analyze such molecules in planarians may help us to understand the evolution of the common mechanism of body patterning in the *Bilateria*.

ACKNOWLEDGMENTS

We thank Dr. Y. Umesono for whole mount *in situ* hybridization, Drs. T. Miya (Tokyo Inst. of Tech.) and N. Ueno (Natl. Inst. for Basic Biol.) for ascidian BMP sequence data, Dr. T. Katayama (Ushimado Marine Laboratory, Okayama Univ.) and members of our laboratory for encouragement and discussion.

This work was supported by Grant-in-Aid for Encouragement of Young Scientists to H.O. (no. 09780688), Special Coordination Funds for Promoting Science and Technology to K.A. and Grant-in-Aid for Scientific Research on Priority Areas to K.W. (no. 09275224)

The nucleotide sequence data reported in this paper will appear in the DDBJ/EMBL/GenBank nucleotide sequence databases with the accession number AB010966.

REFERENCES

- Agata K, Soejima Y, Kato K, Kobayashi C, Umesono Y, Watanabe K (1998) Structure of the planarian central nervous system (CNS) revealed by neuronal cell markers. *Zool Sci* 15: 433–440
- Arora K, Levine MS, O'Connor MB (1994) The screw gene encodes a ubiquitously expressed member of the TGF- β family required for specification of dorsal cell fates in the *Drosophila* embryo. *Genes Dev* 8: 2588–2601
- Baguña J, Saló E, Romero R, Garcia-Fernández J, Bueno D, Muñoz-Mármol AM, Bayascas-Ramírez JR, Casali A (1994) Regeneration and pattern formation in planarians: Cells, molecules and




Fig. 4. Whole mount *in situ* hybridization. Anterior dorsal (A), posterior dorsal (B) and anterior ventral (C) view of intact worm. Dorsal view of regenerating marginal piece 1 day (D), 3 days (E) and 7 days (F) after amputation. The region expressing *Djbmp* is shown by the arrowheads. Schematic representation of expression of *Djbmp* during regeneration of right marginal piece (G). Dots show the expression of *Djbmp*.

- genes. *Zool Sci* 11: 781–795
- Balavoine G, Adoutte A (1998) One or three Cambrian radiations? *Science* 280: 397–398
- Dale L, Howes G, Price BMJ, Smith JC (1992) Bone morphogenetic protein 4: a ventralizing factor in early *Xenopus* development. *Development* 115: 573–585
- De Robertis EM, Sasai Y (1996) A common plan for dorsoventral patterning in bilateria. *Nature* 380: 37–40
- Dosch R, Gawantka V, Delius H, Blumenstock C, Niehrs C (1997) Bmp-4 acts as a morphogen in dorsoventral mesoderm patterning in *Xenopus*. *Development* 124: 2325–2334
- Holley SA, Jackson PD, Sasai Y, Lu B, De Robertis EM, Hoffmann FM, Ferguson EL (1995) A conserved system for dorsal-ventral patterning in insects and vertebrates involving *sog* and *chordin*. *Nature* 376: 249–253
- Horgan BLM, Blessing M, Winnier GE, Suzuki N, Jones CM (1994) Growth factors in development: the role of TGF- β related polypeptide signalling molecules in embryogenesis. *Development* 1994 suppl: 53–60
- Horgan BLM (1995) Upside-down ideas vindicated. *Nature* 376: 210–211
- Irish VF, Gelbart WN (1987) The decapentaplegic gene is required for dorsal-ventral patterning of the *Drosophila* embryo. *Genes Dev* 1: 868–879
- Jones CM, Lyons KM, Lapan PM, Wright CVE, Horgan BLM (1992) DVR-4 (Bone Morphogenetic Protein-4) as a posterior-ventralizing factor in *Xenopus* mesoderm induction. *Development* 115: 639–647
- Kingsley DM (1994) The TGF- β superfamily: New members, new receptors, and new genetic tests of function in different organisms. *Genes Dev* 8: 133–146
- Kurihara T, Kitamura K, Takaoka K, Nakazato H (1993) Murine bone morphogenetic protein-4 gene: Existence of multiple promoters and exons for the 5'-untranslated region. *Biochem Biophys Res Commun* 192: 1049–1056
- Marqués G, Musacchio M, Shimell MJ, Wunnenberg-Stapleton K, Cho KWY, O'Connor M (1997) Production of a DPP activity gradient in the early *Drosophila* embryo through the opposing actions of the SOG and TLD proteins. *Cell* 91: 417–426
- Miya T, Morita K, Ueno N, Satoh N (1996) An ascidian homologue of vertebrate *BMPs-5-8* is expressed in the midline of the anterior neuroectoderm and in the midline of the ventral epidermis of the embryo. *Mech Dev* 57: 181–190
- Nishimatsu S-i, Suzuki A, Shoda A, Murakami K, Ueno N (1992) Genes for bone morphogenetic proteins are differentially transcribed in early amphibian embryos. *Biochem Biophys Res Commun* 186: 1487–1495
- Orii H, Agata K, Watanabe K (1993) POU-domain genes in planarian *Dugesia japonica*: The structure and expression. *Biochem Biophys Res Commun* 192: 1395–1402
- Panganiban GEF, Rashka KE, Neitzel MD, Hoffmann FM (1990) Biochemical characterization of the *Drosophila dpp* protein, a member of the transforming growth factor β family of growth factors. *Mol Cell Biol* 10: 2669–2677
- Piccolo S, Agius E, Lu B, Goodman S, Dale L, De Robertis EM (1997) Cleavage of Chordin by Xolloid metalloprotease suggests a role for proteolytic processing in the regulation of Spemann organizer activity. *Cell* 91: 407–416
- Sasai Y, Lu B, Steinbeisser H, De Robertis EM (1995) Regulation of neural induction by the Chd and Bmp-4 antagonistic patterning signals in *Xenopus*. *Nature* 376: 333–336
- Suzuki A, Kaneko E, Maeda J, Ueno N (1997) Mesoderm induction by BMP-4 and -7 heterodimers. *Biochem Biophys Res Commun* 232: 153–156
- Umesono Y, Watanabe K, Agata K (1997) A planarian *orthopedia* homolog is specifically expressed in the branch region of both the mature and regenerating brain. *Dev Growth Differ* 39: 723–727
- Umesono Y, Watanabe K, Agata K (1998) Distinct structural domains in the planarian brain defined by the expression of evolutionarily conserved homeobox genes. *Dev Genes Evol*, in press
- Watanabe K, Sakai F, Orii H (1997) Stepwise dilution screening of a cDNA library by polymerase chain reaction. *Anal Biochem* 252: 213–214
- Wilson PA, Hemmati-Brivanlou A (1995) Induction of epidermis and inhibition of neural fate by Bmp-4. *Nature* 376: 331–333

(Received July 23, 1998 / Accepted August 24, 1998)